

# Recent advances of in vitro culture systems for spermatogonial stem cells in mammals

Mahesh G. Sahare<sup>1</sup> | Suyatno<sup>2,3</sup> | Hiroshi Imai<sup>3</sup> 

<sup>1</sup>National Facility for Gene Function in Health and Disease, Indian Institute of Science, Education and Research, Pune, India

<sup>2</sup>Indonesian Agency for Agricultural Research and Development, Jakarta, Indonesia

<sup>3</sup>Laboratory of Reproductive Biology, Graduate School of Agriculture, Kyoto University, Kyoto, Japan

## Correspondence

Mahesh G. Sahare, National Facility for Gene Function in Health and Disease, Indian Institute of Science, Education and Research, Pune, India.

Email: mahesh@iiserpune.ac.in

## Funding information

This work was supported in part by the Grant in -Aid for Scientific Research (B) from Japan Society for the promotion of Science to H.I.

## Abstract

**Background:** Spermatogonial stem cells (SSCs) in the mammalian testis are unipotent stem cells for spermatozoa. They show unique cell characteristics as stem cells and germ cells after being isolated from the testis and cultured in vitro. This review introduces recent progress in the development of culture systems for the establishment of SSC lines in mammalian species, including humans.

**Methods:** Based on the published reports, the isolation and purification of SSCs, identification and characteristics of SSCs, and culture system for mice, humans, and domestic animals have been summarized.

**Results:** In mice, cell lines from SSCs are established and can be reprogrammed to show pluripotent stem cell potency that is similar to embryonic stem cells. However, it is difficult to establish cell lines for animals other than mice because of the dearth of understanding about species-specific requirements for growth factors and mechanisms supporting the self-renewal of cultured SSCs. Among the factors that are associated with the development of culture systems, the enrichment of SSCs that are isolated from the testis and the combination of growth factors are essential.

**Conclusion:** Providing an example of SSC culture in cattle, a rational consideration was made about how it can be possible to establish cell lines from neonatal and immature testes.

## KEYWORDS

cell lines, germ cells, gonocytes, pluripotent stem cells, spermatogonial stem cell


## 1 | INTRODUCTION

In mammals, spermatogenesis is a sequential, organized process of self-renewal and differentiation of spermatogonial stem cells (SSCs) that are found in the testis and that result in the continuous production of spermatozoa throughout the life of a man.<sup>1-4</sup> Spermatogenesis protects genomic integrity and plays an essential role in the preservation of the species and genetic diversity.<sup>5</sup> The processes in spermatogenesis are conserved among mammalian species. However, the transformation of spermatogenesis from self-renewing stem

cells to mature spermatozoa is completely different and unique among species. The process lasts 35 days in mice,<sup>6</sup> 74 days in humans,<sup>3</sup> and 63 days in cattle.<sup>7</sup> For the duration of this transformation, the SSCs undergo mitotic multiplication, meiotic recombination of genetic material and morphological changes into spermatozoa.<sup>8</sup> This is a highly productive process that begins at puberty in male animals and ultimately produces 100 million spermatozoa in adult men<sup>9</sup> and 6000 million spermatozoa in mature bulls.<sup>10</sup> Male fertility completely relies on the steady state of spermatogenesis in pubertal animals.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2018 The Authors. Reproductive Medicine and Biology published by John Wiley & Sons Australia, Ltd on behalf of Japan Society for Reproductive Medicine.



**FIGURE 1** Schematic diagram of the developmental origin of spermatogonia. During embryonic development, primordial germ cells differentiate into gonocytes and both cell types are called “embryonic primitive germ cells.” Gonocytes will migrate to the basal compartment of the seminiferous tubule and initiate spermatogenesis by producing spermatogonial stem cells (SSCs) ( $A_{\text{single}}$  or  $A_{\text{dark}}$  spermatogonia). These processes occur soon after birth in rodents but take several weeks in domesticated species and humans. The SSCs will self-renew and differentiate into their progenitors. Both the SSCs and their progenitors ( $A_{\text{pair}}$  and  $A_{\text{align}}$  or  $A_{\text{pale}}$  spermatogonia) also are called “male undifferentiated germ cells.” Finally, differentiating spermatogonia enter meiosis and produce mature sperm via spermatogenesis

The development of a culture system and successful establishment of SSC lines in rodents has attracted much attention from researchers. Although SSCs from many mammalian species have been shown to proliferate for more than 6 months in the seminiferous tubules of immunodeficient mice, no germ cell (GC) line has been established in most mammalian species, other than mice. It is still unknown whether this lack of cell line is related to the lack of knowledge regarding culture conditions and the factors regulating and maintaining SSCs in culture.

This review summarizes the recent progress in the development of the culture system and possible challenges in establishing a SSC line in human and livestock species.

## 2 | SPERMATOGONIAL STEM CELLS

Spermatogonial stem cells originate from gonocytes, they are a derivative of primordial germ cells (PGCs), which are cells from a germ line lineage that arises from the extraembryonic mesoderm at the posterior end of the primitive streak. They migrate to the urogenital ridge, which forms gonads.<sup>11</sup> The PGCs that cease their proliferation in the male genital ridge are called gonocytes. After birth, the gonocytes resume their proliferation, migrate to the basement membrane of the seminiferous tubules, and transform into SSCs. The transition of gonocytes to SSCs after birth occurs within 3 days in mice<sup>12</sup> and 20 weeks in bulls.<sup>13</sup>

The SSCs have a unique ability for both self-renewal and cell differentiation toward spermatogenesis (Figure 1). The existing self-renewal model of SSCs was originally proposed by Huckins<sup>1</sup> in rats and Oakberg<sup>6</sup> in mice. This model proposes that only  $A_{\text{single}}$  ( $A_{\text{s}}$ ) spermatogonia act as stem cells and give rise to committed cells that divide into  $A_{\text{pair}}$  ( $A_{\text{pr}}$ ) and  $A_{\text{align}}$  ( $A_{\text{al}}$ ) cells during spermatogenesis. The extended studies of the self-renewal model of  $A_{\text{s}}$  spermatogonia ( $A_{\text{s}}$  model) using genetic labeling, lineage tracing analysis, and live imaging have provided a striking observation that  $A_{\text{s}}$  spermatogonial cells represent heterogeneity<sup>14</sup> and showed that populations of  $A_{\text{pr}}$  and  $A_{\text{al}}$  SSCs change their behavior during regeneration and acquire stem cell potential. The actual cell number of SSCs having stem cell potential is very low, with ~2000 cells per testis, as calculated by using a pulse-labeling strategy<sup>14</sup> and ~3000 cells per testis by using a serial transplantation assay.<sup>15</sup> This number is very low, compared to the  $A_{\text{s}}$  model based on morphological characteristics,<sup>16</sup> which was estimated at ~35 000 cells per testis. These findings support the heterogeneity of  $A_{\text{s}}$  SSCs in states of morphological similarity. In humans, the spermatogonial renewal model was first proposed by Clemont in 1966.<sup>17</sup> The model postulates that the  $A_{\text{dark}}$  and  $A_{\text{pale}}$  spermatogonia, similar to  $A_{\text{pr}}$  and  $A_{\text{al}}$  in rodents, occur in the human testis and that the  $A_{\text{dark}}$  spermatogonia are mostly undifferentiated and reserved as stem cells, whereas the  $A_{\text{pale}}$  spermatogonia were renewing and were spermatogonia in the early stages of differentiation.

**TABLE 1** Germ cell transplant and transgenesis in domestic animals

Species	Donor-derived spermatogenesis	Reference
Pig (homologous)	Complete	Honaramooz, Megee, Dobrinski <sup>27</sup>
Goat (homologous)	Complete	Honaramooz, Behboodi, Megee, et al <sup>28</sup>
Cattle (autologous)	Complete	Izadyar <sup>29</sup>
Cattle (homologous)	Not demonstrated	
Cattle (homologous)	Not demonstrated	Hill, Dobrinski <sup>30</sup>
Goat	Complete with the integration of a transgene (adenovirus)	Honaramooz, Megee, Zeng, et al <sup>31</sup>

### 3 | SPERMATOGONIAL STEM CELL NICHE IN THE TESTIS

Adult stem cells can self-renew only in a specialized microenvironment called a niche, which provides architectural support, growth factors, and extrinsic stimuli for SSCs.<sup>18,19</sup> The SSCs reside in the basement of the seminiferous tubules and constitute a niche that is surrounded by Sertoli cells, Leydig cells, and peritubular myoid cells.<sup>20</sup> The Sertoli cells seem to play a particularly important role in the SSC niche because numerous factors, such as glial cell-derived neurotrophic factor (GDNF), fibroblast growth factor 2, kit ligand, activin A, and bone morphogenic protein 4 (BMP4), are produced by Sertoli cells and affect the self-renewal, proliferation, and differentiation of the

SSCs.<sup>21</sup> Recent evidence suggests that As, Apr, and Aal spermatogonia can be found along the peritubular blood vessels and are preferentially located in a specific compartment that serves as the niche.<sup>22,23</sup>

### 4 | IDENTIFICATION OF SPERMATOGONIAL STEM CELLS

#### 4.1 | Transplant assay of the isolated spermatogonial stem cells

The first transplant assay for the identification of SSCs in mice was performed by Brinster and Zimmermann.<sup>24</sup> The recipient mice are

**TABLE 2** Overview of spermatogonial markers in rodents, humans, and domestic animals

Molecular marker	Species		
	Human	Mouse	Cattle
VASA/DDX4	ND	+	+
		(Sakai, Noce, Yamashina <sup>39</sup> )	(Fujihara, Kim, Minami, Yamada, Imai <sup>40</sup> )
UCHL1	+	+	+
	(He, Kokkinaki, Jiang, Dobrinski, Dym <sup>43</sup> )	(Kwon, Kikuchi, Setsue, Ishii, Kyuwa, Yoshikawa <sup>44</sup> )	(Herrid, Davey, Hill <sup>45</sup> )
DBA	ND	ND	+
			(Izaydar <sup>49</sup> )
PLZF	ND	+	+
		(Buaas, Kirsh, Sharma, et al <sup>51</sup> )	(Reding, Stepnoski, Cloninger, Oatley <sup>52</sup> )
THY1	+	+	+
	(He, Kokkinaki, Jiang, Dobrinski, Dym <sup>43</sup> )	(Kubota, Avarbock, Brinster <sup>53</sup> )	(Reding, Stepnoski, Cloninger, Oatley <sup>52</sup> )
POUF1	ND	+	+
		(Pesce, Wang, Wolgemuth, Schöler <sup>56</sup> )	(Fujihara, Kim, Minami, Yamada, Imai <sup>40</sup> )
NANOG	ND	ND	ND
GFR $\alpha$ 1	+	+	+
	(He, Kokkinaki, Jiang, Dobrinski, Dym <sup>43</sup> )	(Naughton, Jain, Strickland, Gupta, Milbrandt <sup>57</sup> )	(Sahare, Kim, Otomo, et al <sup>58</sup> )
GFR125	ND	+	ND
		(Seandel, James, Shmelkov, et al <sup>60</sup> )	
RET	ND	+	ND
		(Naughton, Jain, Strickland, Gupta, Milbrandt <sup>57</sup> )	
ID4	ND	+	ND
		(Oatley, Brinster <sup>61</sup> )	
ITGA6	ND	+	+
		(Shinohara, Avarbock, Brinster <sup>62</sup> )	(de Barros, Worst, Saurin, Mendes, Assumpção, Visintin <sup>63</sup> )
ITGB1	ND	+	ND
		(Shinohara, Avarbock, Brinster <sup>62</sup> )	

+, expression of the protein in undifferentiated SSCs; ND, not determined.



considered to be a feasible way to restore infertility after cancer treatment.<sup>38</sup>

## 4.2 | Biochemical characterization of the spermatogonial stem cells

Defining the SSC populations by using biochemical markers that distinguish them from spermatogonia in other stages of differentiation was a great tool for the isolation of potential SSCs and the development of culture systems in rodents. In recent years, several molecular markers have been identified for SSCs in rodents (Table 2).<sup>39-63</sup> Most of these markers are expressed in progenitor SSCs, including As spermatogonia and undifferentiated spermatogonia (Apr and Aal spermatogonia). Traditionally, As spermatogonia have been included in the SSC population that self-renews in order to maintain a foundational stem cell pool and the transition to Apr spermatogonia represents the initial step of spermatogenesis.<sup>1,6</sup> Recent findings show that the SSC population is not limited to the As spermatogonia population.<sup>64</sup> Some progenitor SSCs also exhibit stem cell behavior.

Some of these markers are identified as SSC markers in domestic animals (Table 2) and are conserved among mammalian species. The markers, GPR125, GFR1, THY1, ZBTB16, SSEA-4, and PLZF, that have been identified for SSCs in rodents have also been characterized in human spermatogonia and more differentiated GCs.<sup>43,65,66</sup>

## 5 | IN VITRO CULTURE OF THE SPERMATOGONIAL STEM CELLS

### 5.1 | Isolation and enrichment of the spermatogonial stem cells

The isolation and enrichment of SSCs is the first step towards establishing GS cell lines. The isolation of SSCs is challenging because of their limited number in the testis. A two-step enzymatic digestion was first proposed by Davis and Schuetz,<sup>67</sup> which is the most widely used technique for the isolation of SSCs in rodents. For further enrichment of SSCs, different approaches, such as differential plating,<sup>68</sup> percoll gradient,<sup>23</sup> magnetic-activated cell sorting (MACS) or fluorescence-activated cell sorting (FACS) have been used independently or in combination. In livestock species, SSC isolation and enrichment methods have progressed during the last few years. Differential plating is one feasible method for the enrichment of SSCs, along with MACS and FACS for bovine SSCs.<sup>69</sup>

### 5.2 | Establishment of a culture system for germ cell lines

The limited number of SSCs in the testis<sup>4</sup> hampers studies that elucidate biological characteristics and for applying SSCs. One approach to solve this problem is to develop a culture system that supports the self-renewal of SSCs and maintains their GC and stem-cell potentials. Glial cell-derived neurotrophic factor was shown to be the first molecule that regulates the self-renewal and differentiation of mouse

SSCs.<sup>70</sup> Glial cell-derived neurotrophic factor signals act through the multicomponent receptor complex that is composed of GFR $\alpha$ -1 and RET tyrosine kinases in various cell types.<sup>71</sup> The GFR $\alpha$ -1 and RET also have been recognized as spermatogonial markers that are expressed in gonocytes, SSCs, and differentiated spermatogonia.<sup>72</sup> These coreceptors of GDNF-mediated signaling have been shown to be necessary for the self-renewal of GCs in rodents.<sup>57</sup> Subsequently, Nagano, Ryu, Brinster, Avarbock, and Brinster developed a short-term culture system that is supplemented with GDNF that improves the survival of GCs.<sup>15</sup> These cells complete spermatogenesis after transplant into the testis of immunodeficient mice. The long-term culture of SSCs is achieved by adding other growth factors and hormones in addition to GDNF.<sup>73</sup> These cells proliferate over a 2 year period (>10<sup>85</sup>-fold) in the presence of GDNF, while maintaining stable genetic and epigenetic properties and restoring spermatogenesis following transplant into the seminiferous tubules of infertile recipient mice. However, the growth factor requirements for the proliferation of GCs is strain-specific: in mice, the C57BL/6 and 129/Sv strains require fibroblast growth factor (FGF) and GDNF,<sup>74</sup> while strain DBA requires FGF, GDNF, and epidermal growth factor.<sup>75</sup> By using species-specific culture components, culture systems and GC lines have been established in rats,<sup>76,77</sup> hamsters,<sup>78</sup> and rabbits.<sup>79</sup>

Spermatogonial stem cells under appropriate culture conditions acquire embryonic stem (ES) cell-like characteristics called "multipotent GCs," which were first generated from GCs in the neonatal mouse testis without the introduction of any exogenous reprogramming factor.<sup>80</sup> These cell populations failed to form colonies following testicular transplants, which shows that they are devoid of GC potential and have the ability to differentiate into three germ layers. Later, successful evidence of the generation of a multipotent GS cell line was shown for adult mice.<sup>60,81</sup>

The successful translation of an in vitro culture of SSCs in rodents led to the establishment of a culture system for human SSCs from prepubertal and adult testes.<sup>82,83</sup> In humans, multipotent stem cell lines have been developed from SSCs by exposing the cells to ES cell culture conditions.<sup>84,85</sup> These cell lines can form a teratoma after they are injected into immunodeficient mice. These findings provide an important foundation for developing methods for the generation of autologous stem cell lines from human SSCs that have been collected from patients with cancer before the initiation of cancer treatment and the subsequent autologous transplant after cancer treatment could be a means for preserving the fertility of male patients with cancer.<sup>86</sup>

### 5.3 | Spermatogonial stem cell culture in livestock species

In livestock species, long-term culture systems for GCs and the establishment of multipotent GC lines could reduce the time and costs for producing transgenic animals and to preserve endangered species. These systems also could be an alternative for pronuclear microinjection and somatic cell cloning.<sup>87</sup> Although several attempts have been made to develop a culture system for livestock species, as shown in Table 3,<sup>40,58,88-94</sup> most of these studies achieved only short-term SSC

**TABLE 3** Overview of the culture conditions for spermatogonial stem cells in domestic species

Reference	Culture conditions	Age of donor	Culture term
<b>Cat</b>			
Izadyar, Den Ouden, Stout, et al <sup>88</sup>	Compare MEM and KSOM medium 0%-10% FCS	5 mo	MEM+2.5% FCS is effective for germ cell survival than KSOM, no expansion, showing differentiation during 150 days culture
Oatley, Reeves, McLean <sup>89</sup>	DMEMF + 10% FBS + GDNF	1-2 mo	2 wk
Aponte, Soda, van de Kant <sup>90</sup>	MEM +2.5% FCS + GDNF	4-6 mo	25 d, no passage, differentiation
Aponte, Soda, Teerds, Mizrak, van de Kant <sup>91</sup>	StemPro-SFM + GDNF, EGF, and FF	4-6 mo	25 d, no appearance of colonies after passage
Fujihara, Kim, Minami, Yamada, Imai <sup>40</sup>	DMEMF12 + 10% FCS	1-10 d	1.5 mo
Sahare, Kim, Otomo, et al <sup>58</sup>	DMEMF12 + 15% KSR on poly-L-lysine-coated dishes	1-10 d	>2 mo
<b>Pig</b>			
Dirami, Ravindranath, Pursel, Dym <sup>92</sup>	DMEMF12 + 10% FCS	2 mo	1 wk
Goel, Fujihara, Tsuchiya, et al <sup>93</sup>	DMEMF12 + 10% FCS	1-10 d	3 wk, reduction of germ cells every passage
Goel, Fujihara, Tsuchiya, et al <sup>94</sup>	StemPro SFM + GDNF, EGF, and FF	3-4 d	9 passages (30 d), reduction of germ cells every passage

DMEMF, Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12; EGF, epidermal growth factor; FCS, fetal calf serum; FF, feeder-free; GDNF, glial cell-derived neurotrophic factor; KSOM, potassium simplex optimized medium; KSR, knockout serum replacement; MEM, minimum essential medium; SFM, serum- and feeder-free medium.

cultures. The culture system for bovine SSCs has been demonstrated in the pre-pubertal testis<sup>29,49,89-91</sup> and the neonatal testis.<sup>40</sup> In pigs, cultured SSCs cannot survive more than 1 week<sup>50,92</sup>. In these studies, serum was used as an important component in the culture medium for the survival and self-renewal of SSCs. Some undefined factors in the serum induce cell differentiation, whereas others have detrimental effects on ES cells and GC survival in the culture.<sup>53,95</sup> In order to overcome this problem, serum-free culture systems have been developed for long-term cultures of SSCs in mice<sup>74,96</sup> and rats.<sup>77</sup> However, no long-term culture system for livestock species has been developed. In the authors' laboratory, growth factors, matrix substrates for culture dishes, and serum-free supplements have been examined in order to develop a defined system for culturing primitive GCs (gonocytes) from the neonatal bovine testis.<sup>58,97,98</sup> Poly-L-lysine is a suitable substrate for the selective inhibition of the growth of somatic cells and makes it possible to maintain gonocytes. Among the serum-free supplements that were tested, knockout serum replacement (KSR) in the culture medium supports the proliferation and survival of the gonocytes after sequential passages of the colonies. Under these optimized culture conditions that consist of 15% KSR on poly-L-lysine-coated dishes, the stem cell and GC potentials of cultured gonocytes can be maintained for more than 2 months. Subsequently, also developed was a culture system to maintain the SSCs from immature and adult testis in cattle.<sup>99</sup> H The SSCs from the immature testis are cultured under serum-free conditions in the presence of GDNF and bovine leukemia inhibitory factor-conditioning media. Established cell lines resemble ES-like cell properties and express both pluripotent and GC markers. However, the SSCs from the adult bovine testis are cultured in a low-serum concentration media that is supplemented

with 6-bromindirubin-3'-oxime, which is a small-molecule inhibitor of glycogen synthase kinase-3 $\alpha$  that leads to the activation of the wingless-type (Wnt)/ $\beta$ -catenin signaling pathway.<sup>100</sup> The established cell lines can be maintained under in vitro culture conditions for more than 3 months. This cell line has a normal karyotype and botryoidal morphology that is similar to the male GC lines from mouse SSCs. Taken together, this new finding provides a promising strategy to conserve GCs from livestock species at different stages of animal development.

## 6 | CONCLUSION

Recently, GCs with a GC lineage have been derived from ES cells<sup>101,102</sup> and induced pluripotent stem cells in mice.<sup>103-105</sup> The molecules that are involved in GC commitment, such as BMP4 and Wnt3, have been identified<sup>102,106</sup> and PGCs are induced from pluripotent stem cells under the control of these molecules and other cell differentiation-inducing factors.<sup>102,106</sup> In humans, PGCs also are induced in similar culture conditions.<sup>107-109</sup> The induced mouse PGCs can be maintained in a normal manner and differentiated into spermatozoa and oocytes with the ability to develop to term.<sup>110,111</sup> At this time, GC formation for the spermatozoa and oocytes was achieved under ex vivo conditions, in which somatic cells that were associated with spermatogenesis or oogenesis were cocultured and aggregated with the indicated PGC population.<sup>111</sup> Therefore, although additional studies are necessary in order to maintain and induce GCs in vitro, GS cell lines that have been established in some mammalian species might be candidates to produce spermatozoa and oocytes in vitro. These technologies in the near future will be helpful



for the retention of the fertility of patients before cancer therapy, the production of transgenic animals for human disease models, domestic animal improvement, and the conservation of endangered species.

## DISCLOSURES

**Conflict of interest:** The authors declare no conflict of interest. **Human rights statement and informed consent:** This article does not contain any studies with human subjects performed by any of the authors. **Animal studies:** The protocol for the research project, including the animal participants, was approved by a suitably constituted ethics committee.

## ORCID

Hiroshi Imai  <http://orcid.org/0000-0003-3702-2708>

## REFERENCES

- Huckins C. The spermatogonial stem cell population in adult rats: Evidence for a long-cycling population. *Cell Tissue Kinet.* 1971;4:335-349.
- Clermont Y. Kinetics of spermatogenesis in mammals: seminiferous epithelium cycle and spermatogonial renewal. *Physiol Rev.* 1972;52:198-236.
- Russell LD, Ettl RA, Hikim AP, Clegg ED, eds. Mammalian spermatogenesis. In: Russell LD, Ettl RA, Hikim AP, Clegg ED, eds. *Histological and Histopathological Evaluation of the Testis*. Florida: Cache River; 1990:1-40.
- Meistrich M, van Beek M. Spermatogonial stem cells. In: Desjardins C, Ewing LL, eds. *Cell and Molecular Biology of Testis*. New York: Oxford University Press; 1993:266-295.
- Wistuba J, Stukenbeorg BJ, Luetjens CM. Mammalian spermatogenesis. *Funct Dev Embryol.* 2007;1:99-117.
- Oakberg EF. A new concept of spermatogonial stem-cell renewal in the mouse and its relationship to genetic effects. *Mutat Res.* 1971;11:1-7.
- Hochereau MT, Courot M, Ortavanat R. Marquage des cellules germinales dubelger et du taureau par injection de thymidine tritium? Edans l'artkre spermatique. *Ann Biol Animale Biochim Biophys.* 1964;2:157-161.
- Ehmcke J, Hübner K, Schöler HR, Schlatt S. Spermatogonia: origin, physiology and prospects for conservation and manipulation of the male germ line. *Reprod Fertil Dev.* 2006;18:7-12.
- Sharpe R. Regulation of spermatogenesis. In: Neill EJD, ed. *The physiology of Reproduction*. New York: Raven Press; 1994:1363-1434.
- Amann RP, Kavanaugh JF, Griel LC Jr, Voglmayr JK. Sperm production of holstein bulls determined from testicular spermatid reserves, after cannulation of rete testis or vas deferens, and by daily ejaculation. *J Dairy Sci.* 1974;457:93-99.
- Lawson KA, Hage WJ. Clonal analysis of the origin of primordial germ cells in the mouse. *Ciba Found Symp.* 1994;182:68-84.
- McLean DJ, Friel PJ, Johnston DS, Griswold MD. Characterization of spermatogonial stem cell maturation and differentiation in neonatal mice. *Biol Reprod.* 2003;69:2085-2091.
- Curtis R, Amann P. Testicular development and establishment of spermatogenesis in Holstein bull. *J Anim Sci.* 1981;53:1645-1657.
- Nakagawa T, Nabeshima Y, Yoshida S. Article functional identification of the actual and potential stem cell compartments in mouse spermatogenesis. *Dev Cell.* 2007;12:195-206.
- Nagano M, Ryu B-Y, Brinster CJ, Avarbock MR, Brinster RL. Maintenance of mouse male germ line stem cells in vitro. *Biol Reprod.* 2003;68:2207-2214.
- Tegelenbosch RA, de Rooij DG. A quantitative study of spermatogonial multiplication and stem cell renewal in the C3H/101 F1 hybrid mouse. *Mutat Res.* 1993;290:193-200.
- Clermont Y. Spermatogenesis in man. A study of the spermatogonial population. *Fertil Steril.* 1966;17:705-721.
- Schofield R. The relationship between the spleen colony-forming cell and the haemopoietic stem cell. *Blood Cells.* 1978;4:7-25.
- Spradling A, Drummond-Barbosa D, Kai T. Stem cells find their niche. *Nature.* 2001;414:98-104.
- de Rooij DG. The spermatogonial stem cell niche. *Microsci Res Tech.* 2009;72:580-585.
- Boyer A, Yeh JR, Zhang X, et al. CTNNB1 signaling in sertoli cells downregulates spermatogonial stem cell activity via WNT4. *PLoS ONE.* 2012;7:e29764.
- Chiarini-Garcia H, Hornick JR, Griswold MD, Russell LD. Distribution of Type A spermatogonia in the mouse is not random. *Biol Reprod.* 2001;65:1179-1185.
- Yoshida S, Sukeno M, Nabeshima Y. A vasculature-associated niche for undifferentiated spermatogonia in the mouse testis. *Science.* 2007;317:1722-1726.
- Brinster RL, Zimmermann JW. Spermatogenesis following male germ-cell transplantation. *Proc Natl Acad Sci USA.* 1994;91:11298-11302.
- Clouthier DE, Avarbock MR, Maika SD, Hammer RE, Brinster RL. Rat spermatogenesis in mouse testis. *Nature.* 1996;381:418-421.
- Ogawa T, Dobrinski I, Avarbock MR, Brinster RL. Xenogeneic spermatogenesis following transplantation of hamster germ cells to mouse testes. *Biol Reprod.* 1999;60:515-521.
- Honaramooz A, Megee SO, Dobrinski I. Germ cell transplantation in pigs. *Biol Reprod.* 2002;66:21-28.
- Honaramooz A, Behboodi E, Megee SO, et al. Fertility and germline transmission of donor haplotype following germ cell transplantation in immunocompetent goats. *Mol Reprod Dev.* 2003;64:422-428.
- Izadyar F. Proliferation and differentiation of bovine type A spermatogonia during long-term culture. *Biol Reprod.* 2003;68:272-281.
- Hill JR, Dobrinski I. Male germ cell transplantation in livestock. *Reprod Fertil Dev.* 2006;18:13-18.
- Honaramooz A, Megee S, Zeng W, et al. Adeno-associated virus (AAV)-mediated transduction of male germ line stem cells results in transgene transmission after germ cell transplantation. *FASEB J.* 2008;22:374-382.
- Dobrinski I, Avarbock MR, Brinster RL. Transplantation of germ cells from rabbits and dogs into mouse testes. *Biol Reprod.* 1999;61:1331-1339.
- Dobrinski I, Avarbock MR, Brinster RL. Germ cell transplantation from large domestic animals into mouse testes. *Mol Reprod Dev.* 2000;57:270-279.
- Dobrinski I. Germ cell transplantation and testis tissue xenografting in domestic animals. *Reprod Sci.* 2005;89:137-145.
- Brinster RL. Germline stem cell transplantation and transgenesis. *Science.* 2002;296:2174-2176.
- Kanatsu-Shinohara M, Ikawa M, Takehashi M, et al. Production of knockout mice by random or targeted mutagenesis in spermatogonial stem cells. *Proc Natl Acad Sci USA.* 2006;103:8018-8023.
- Nagano M, Patrizio P, Brinster RL. Long-term survival of human spermatogonial stem cells in mouse testes. *Fertil Steril.* 2002;78:1225-1233.
- Kubota H, Brinster RL. Technology insight: in vitro culture of spermatogonial stem cells and their potential therapeutic uses. *Nat Clin Pract Endocr Metab.* 2006;2:99-108.
- Sakai Y, Noce T, Yamashina S. Cleavage-like cell division and explosive increase in cell number of neonatal gonocytes. *Dev Growth Differ.* 2004;46:15-21.

40. Fujihara M, Kim SM, Minami N, Yamada M, Imai H. Characterization and in vitro culture of male germ cells from developing bovine testis. *J Reprod Dev.* 2011;57:355-364.
41. Borjigin U, Davey R, Hutton K, Herrid M. Expression of promyelocytic leukaemia zinc-finger in ovine testis and its application in evaluating the enrichment efficiency of differential plating. *Reprod Fertil Dev.* 2010;22:733-742.
42. Goel S, Reddy N, Mandal S, Fujihara M, Kim S-M, Imai H. Spermatogonia-specific proteins expressed in prepubertal buffalo (*Bubalus bubalis*) testis and their utilization for isolation and in vitro cultivation of spermatogonia. *Theriogenology.* 2010;74:1221-1332.
43. He ZI, Kokkinaki M, Jiang J, Dobrinski I, Dym M. Isolation, characterization, and culture of human spermatogonia. *Biol Reprod.* 2010;82:363-372.
44. Kwon J, Kikuchi T, Setsuie R, Ishii Y, Kyuwa S, Yoshikawa Y. Characterization of the testis in congenitally ubiquitin carboxy-terminal hydrolase-1 (Uch-L1) defective (gad) mice. *Exp Anim.* 2003;52:1-9.
45. Herrid M, Davey RJ, Hill JR. Characterization of germ cells from prepubertal bull calves in preparation for germ cell transplantation. *Cell Tissue Res.* 2007;330:321-329.
46. Luo J, Megee S, Rath R, Dobrinski I. Protein gene product 9.5 is a spermatogonia-specific marker in the pig testis: application to enrichment and culture of porcine spermatogonia. *Mol Reprod Dev.* 2006;73:1531-1540.
47. Rodriguez-Sosa JR, Dobson H, Hahnel A. Isolation and transplantation of spermatogonia in sheep. *Theriogenology.* 2006;66:2091-2103.
48. Heidari B, Rahmati-Ahmadabadi M. Isolation, identification, and culture of goat spermatogonial stem cells using c-kit and PGP9.5 markers. *J Assist Reprod Genet.* 2012;29:1029-1038.
49. Izadyar F. Proliferation and differentiation of bovine Type A spermatogonia during long-term culture. *Biol Reprod.* 2002;68:272-281.
50. Goel S, Sugimoto M, Minami N, Yamada M, Kume S, Imai H. Identification, isolation, and in vitro culture of porcine gonocytes. *Biol Reprod.* 2007;137:127-137.
51. Buaas FW, Kirsh AL, Sharma M, et al. Plzf is required in adult male germ cells for stem cell self-renewal. *Nat Genet.* 2004;36:647-652.
52. Reding SC, Stepnoski AL, Cloninger EW, Oatley JM. THY1 is a conserved marker of undifferentiated spermatogonia in the pre-pubertal bull testis. *Reproduction.* 2010;39:893-903.
53. Kubota H, Avarbock MR, Brinster RL. Growth factors essential for self-renewal and expansion of mouse spermatogonial stem cells. *Proc Natl Acad Sci USA.* 2004;101:16489-16494.
54. Abbasi H, Tahmoospur M, Morteza S, Nasiri Z. THY1 as a reliable marker for enrichment of undifferentiated spermatogonia in the goat. *Theriogenology.* 2013;80:923-932.
55. Rafeeqi T, Kaul G. Isolation and enrichment of type A spermatogonia from pre-pubertal buffalo (*Bubalus bubalis*) testis. *Andrologia.* 2013;45:195-203.
56. Pesce M, Wang X, Wolgemuth DJ, Schöler H. Differential expression of the Oct-4 transcription factor during mouse germ cell differentiation. *Mech Dev.* 1998;71:89-98.
57. Naughton CK, Jain S, Strickland AM, Gupta A, Milbrandt J. Glial cell-line derived neurotrophic factor-mediated RET signaling regulates spermatogonial stem cell fate. *Biol Reprod.* 2006;74:314-321.
58. Sahare M, Kim SM, Otomo A, et al. Factors supporting long-term culture of bovine male germ cells. *Reprod Fertil Dev.* 2016;28:2039-2050.
59. Lee WY, Park HJ, Lee R, et al. Establishment and in vitro culture of porcine spermatogonial germ cells in low temperature culture conditions. *Stem Cell Res.* 2013;11:1234-1249.
60. Seandel M, James D, Shmelkov SV, et al. Generation of functional multipotent adult stem cells from GPR125+ germline progenitors. *Nature.* 2007;449:346-350.
61. Oatley JM, Brinster RL. The germline stem cell niche unit in mammalian testes. *Physiol Rev.* 2012;92:577-595.
62. Shinohara T, Avarbock MR, Brinster RL. Beta1- and alpha6-integrin are surface markers on mouse spermatogonial stem cells. *Proc Natl Acad Sci USA.* 1999;96:5504-5509.
63. de Barros FR, Worst RA, Saurin GC, Mendes CM, Assumpção ME, Visintin JA. alpha 6 integrin expression in bovine spermatogonial cells purified by discontinuous Percoll density gradient. *Reprod Domest Anim.* 2012;47:887-890.
64. Hara K, Nakagawa T, Enomoto H, et al. Mouse spermatogenic stem cells continually interconvert between equipotent singly isolated and syncytial states. *Cell Stem Cell.* 2014;14:658-672.
65. Wu X, Schmidt JA, Avarbock MR, et al. Prepubertal human spermatogonia and mouse gonocytes share conserved gene expression of germline stem cell regulatory molecules. *Proc Natl Acad Sci USA.* 2009;106:21672-21677.
66. Dym M, Kokkinaki M, He Z. Spermatogonial stem cells: mouse and human comparisons. *Birth Defects Res C Embryo Today.* 2009;87:27-34.
67. Davis JC, Schuetz AW. Separation of germinal cells from immature rat testes by sedimentation at unit gravity. *Exp Cell Res.* 1975;91:79-86.
68. Dym M, Jia MC, Dirami G, et al. Expression of c-kit receptor and its autophosphorylation in immature rat type A spermatogonia. *Biol Reprod.* 1995;52:8-19.
69. Herrid M, Davey RJ, Hutton K, Colditz IG, Hill JR. A comparison of methods for preparing enriched populations of bovine spermatogonia. *Reprod Fertil Dev.* 2009;21:393-399.
70. Meng X, Lindahl M, Hyvönen, ME, et al. Regulation of cell fate decision of undifferentiated spermatogonia by GDNF. *Science.* 2000;287:1489-1493.
71. Jing S, Wen D, Yu Y, et al. GDNF-induced activation of the ret protein tyrosine kinase is mediated by GDNFR-alpha, a novel receptor for GDNF. *Cell.* 1996;85:1113-11124.
72. Widenfalk J, Parvinen M, Lindqvist E, Olson L. Neurturin, RET, GFRalpha-1 and GFRalpha-2, but not GFRalpha-3, mRNA are expressed in mice gonads. *Cell Tissue Res.* 2000;299:409-415.
73. Kanatsu-Shinohara M, Ogonuki N, Iwano T, et al. Genetic and epigenetic properties of mouse male germline stem cells during long-term culture. *Development.* 2005;132:4155-4163.
74. Kubota H, Avarbock MR, Brinster RL. Culture conditions and single growth factors affect fate determination of mouse spermatogonial stem cells. *Biol Reprod.* 2004;71:722-731.
75. Kanatsu-Shinohara M, Miki H, Inoue K, et al. Long-term culture of mouse male germline stem cells under serum-or feeder-free conditions. *Biol Reprod.* 2005;72:985-991.
76. Hamra FK, Chapman KM, Nguyen DM, Williams-Stephens A, Hammer RE. Self renewal, expansion, and transfection of rat spermatogonial stem cells in culture. *Proc Natl Acad Sci USA.* 2005;102:17430-17435.
77. Ryu B, Kubota H, Avarbock MR, Brinster RL. Conservation of spermatogonial stem cell self-renewal signaling between mouse and rat. *Proc Natl Acad Sci USA.* 2005;102:14302-14307.
78. Kanatsu-Shinohara M, Muneto T, Lee J, Takenaka M, Chuma S. Long-term culture of male germline stem cells from hamster testes. *Biol Reprod.* 2008;78:611-617.
79. Kubota H, Wu X, Goodyear SM, Avarbock MR, Brinster RL. Glial cell line-derived neurotrophic factor and endothelial cells promote self-renewal of rabbit germ cells with spermatogonial stem cell properties. *FASEB J.* 2011;25:2604-2614.
80. Kanatsu-Shinohara M, Inoue K, Lee J, et al. Generation of pluripotent stem cells from neonatal mouse testis. *Cell.* 2004;119:1001-1012.
81. Guan K, Nayernia K, Maier LS, et al. Pluripotency of spermatogonial stem cells from adult mouse testis. *Nature.* 2006;440:1199-1203.



82. Sadri-Ardekani H, Mizrak SC, van Daalen SK, et al. Propagation of human spermatogonial stem cells in vitro. *JAMA*. 2009;302:2127-2134.
83. Sadri-Ardekani H, Akhondi MA, van der Veen F, Repping S, van Pelt AMM. In vitro propagation of human prepubertal spermatogonial stem cells. *JAMA*. 2011;305:2416-2418.
84. Golestaneh N, Kokkinaki M, Pant D, et al. Pluripotent stem cells derived from adult human testes. *Stem Cells Dev*. 2009;18:1115-1126.
85. Kossack N, Meneses J, Shefi S, et al. Isolation and characterization of pluripotent human spermatogonial stem cell-derived cells. *Stem Cells*. 2009;27:138-149.
86. Struijk RB, Mulder CL, Van der VF, Van Pelt AMM, Repping S. Restoring fertility in sterile childhood cancer survivors by autotransplanting spermatogonial stem cells: are we there yet? *Biomed Res Int*. 2013;903142. <https://doi.org/10.1155/2013/903142>
87. Dobrinski I. Transplantation of germ cells and testis tissue for the study of mammalian spermatogenesis. *Anim Reprod*. 2006;3:135-145.
88. Izadyar F, Den Ouden K, Stout TA, et al. Autologous and homologous transplantation of bovine spermatogonial stem cells. *Reproduction*. 2003;126:765-774.
89. Oatley JM, Reeves JJ, McLean DJ. Biological activity of cryopreserved bovine spermatogonial stem cells during in vitro culture. *Biol Reprod*. 2004;71:942-947.
90. Aponte PM, Soda T, van de Kant HJ. Basic features of bovine spermatogonial culture and effects of glial cell line-derived neurotrophic factor. *Theriogenology*. 2006;65:1828-1847.
91. Aponte PM, Soda T, Teerds KJ, Mizrak SC, van de Kant HJ. Propagation of bovine spermatogonial stem cells in vitro. *Reproduction*. 2008;136:543-557.
92. Dirami G, Ravindranath N, Pursel V, Dym M. Effects of stem cell factor and granulocyte macrophage-colony stimulating factor on survival of porcine type A spermatogonia cultured in KSOM. *Biol Reprod*. 1999;61:225-230.
93. Goel S, Fujihara M, Tsuchiya K, et al. Multipotential ability of primitive germ cells from neonatal pig testis cultured in vitro. *Reprod Fertil Dev*. 2009;21:696-708.
94. Goel S, Fujihara M, Tsuchiya K, et al. The expansion of mouse spermatogonial stem cells. *Proc Natl Acad Sci USA*. 2006;101:16489-16494.
95. Li P, Tong C, Mehrian-Shai R, et al. Germline competent embryonic stem cells derived from rat blastocysts. *Cell*. 2008;135:1299-1310.
96. Kanatsu-Shinohara M, Inoue K, Ogonuki N, Morimoto H, Ogura A, Shinohara T. Serum and feeder-free culture of mouse germline stem cells. *Biol Reprod*. 2011;84:97-105.
97. Kim SM, Fujihara M, Sahare M, Minami N, Yamada M, Imai H. Effects of extracellular matrices and lectin *Dolichos biflorus* agglutinin on cell adhesion and self-renewal of bovine gonocytes cultured in vitro. *Reprod Fertil Dev*. 2013;26:268-281.
98. Sahare M, Otomo A, Komatsu K, Minami N, Yamada M, Imai H. The role of signaling pathways on proliferation and self-renewal of cultured bovine primitive germ cells. *Reprod Med Biol*. 2015;14:17-25.
99. Suyatno, Kitamura Y, Ikeda S, Minami N, Yamada M, Imai H. Long-term culture of undifferentiated spermatogonia isolated from immature and adult bovine testis. *Mol Reprod Dev*. 2018; In press.
100. Sato N, Meijer L, Skaltsounis L, Greengard P, Brivanlou AH. Maintenance of pluripotency in human and mouse embryonic stem cells through activation of Wnt signaling by a pharmacological GSK-3-specific inhibitor. *Nat Med*. 2004;10:55-63.
101. Kerkis AI, Fonseca SA, Serafim RC, et al. In vitro differentiation of male mouse embryonic stem cells into both presumptive sperm cells and oocytes. *Cloning Stem Cells*. 2007;9:535-548.
102. Geijsen N, Horoschak M, Kim K, Gribnau J, Eggan K, Daley GQ. Derivation of embryonic germ cells and male gametes from embryonic stem cells. *Nature*. 2004;427:148-154.
103. Hayashi K, Ohta H, Kurimoto K, Aramaki S, Saitou M. Reconstitution of the mouse germ cell specification pathway in culture by pluripotent stem cells. *Cell*. 2011;146:519-532.
104. Yang S, Bo J, Hu H, et al. Derivation of male germ cells from induced pluripotent stem cells in vitro and in reconstituted seminiferous tubules. *Cell Prolif*. 2012;45:91-100.
105. Li YI, Ray D, Ye P. Identification of germ cell-specific genes in mammalian meiotic prophase. *BMC Bioinformatics*. 2013;14:72.
106. Sakurai M, Hayashi R, Kageyama T, Yamato M, Nishida K. Induction of putative stratified epithelial progenitor cells in vitro from mouse-induced pluripotent stem cells. *J Artif Organs*. 2011;14:58-66.
107. Park T, Galic Z, Conway A, et al. Derivation of primordial germ cells from human embryonic and induced pluripotent stem cells is significantly improved by coculture with human fetal gonadal cells. *Stem Cells*. 2009;27:783-795.
108. Bucay N, Yebra M, Cirulli V, et al. A novel approach for the derivation of putative primordial germ cells and Sertoli cells from human embryonic stem cells. *Stem Cells*. 2009;27:68-77.
109. Ishii T. Human iPS cell-derived germ cells: current status and clinical potential. *J Clin Med*. 2014;3:1064-1083.
110. Sato T, Katagiri K, Yokonishi T, et al. In vitro production of fertile sperm from murine spermatogonial stem cell lines. *Nat Commun*. 2011;2:472.
111. Hayashi K, Ogushi S, Kurimoto K, Shimamoto S, Ohta H, Saitou M. Offspring from oocytes derived from in vitro primordial germ cell-like cells in mice. *Science*. 2012;338:971-975.

**How to cite this article:** Sahare MG, Suyatno, Imai H. Recent advances of in vitro culture systems for spermatogonial stem cells in mammals. *Reprod Med Biol*. 2018;17:134-142. <https://doi.org/10.1002/rmb2.12087>