TITLE:

Long-term culture of undifferentiated spermatogonia isolated from immature and adult bovine testes

AUTHOR(S):

Suyatno; Kitamura, Yuka; Ikeda, Shuntaro; Minami, Naojiro; Yamada, Masayasu; Imai, Hiroshi

CITATION:

Suyatno ...[et al]. Long-term culture of undifferentiated spermatogonia isolated from immature and adult bovine testes. *Molecular Reproduction and Development* 2018, 85(3): 236-249

ISSUE DATE:

2018-03

URL:

<http://hdl.handle.net/2433/230134>

RIGHT:

This is the accepted version of the following article: [Suyatno, Kitamura Y, Ikeda S, Minami N, Yamada M, Imai H. Long-term culture of undifferentiated spermatogonia isolated from immature and adult bovine testes. *Mol Reprod Dev.* 2018;85:236–249], which has been published in final form at <https://doi.org/10.1002/mrd.22958>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.; The full-text file will be made open to the public on 13 MAR 2019 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。

determining region Y)-box 2; SSCs, spermatogonial stem cells; UCHL-1, ubiquitin carboxyl-terminal hydrolase isozyme L1.

Abstract

Undifferentiated spermatogonia eventually differentiate in the testis to produce haploid sperm. Within this cell population, there is a small number of spermatogonial stem cells (SSCs). SSCs are rare cells in the testis, and their cellular characteristics are poorly understood. Establishment undifferentiated cell line would provide an indispensable tool for studying their biological nature and spermiogenesis/spermatogenesis in vitro. However, there have been few reports on the long-term culture of undifferentiated spermatogonia in species other than rodents. Here, we report the derivation and long-term in vitro culture of undifferentiated spermatogonia cell lines from immature and adult bovine testes. Cell lines from immature testes were maintained in serum-free culture conditions in the presence of glial-cell-line-derived neurotrophic factor (GDNF) and bovine leukemia inhibitory factor (bLIF). These cell lines have embryonic stem (ES)-like cell morphology, express pluripotent-stem-cell-specific and germ-cell-specific markers at the protein and mRNA levels, and contributed to the inner cell mass (ICM) of embryos in the blastocyst stage. Meanwhile, cell lines established from adult testes were maintained in low-serum media in the presence of 6-bromoindirubin-3'-oxime (BIO). These cell lines have characteristics resembling those of previously reported male mouse germ cell lines as confirmed by their botryoidally aggregated morphology, as well as the expression of germ-cell-specific markers and pluripotent stem cell markers. These findings could be useful for the development of long-term culture of undifferentiated spermatogonia, which could

aid in conservation of species and improvement of livestock production through genome editing technology.

Keywords: adult male germ cells, cattle, spermatogonial stem cells, pluripotent stem cells.

1. Introduction

Spermatogenesis is achieved by a complex process, in which diploid SSCs proceed over time through a series of differentiation steps to produce haploid spermatozoa (de Rooij, 2001). Since SSCs are rare cells in immature and adult testes (Tegelenbosch & de Rooij, 1993), it is hard to isolate and study them and to distinguish them from their undifferentiated progeny. Currently, several markers are used to identify SSCs in mice: GDNF family receptor α -1 (GFR α -1) (Naughton et al., 2006), Thy-1 membrane glycoprotein (Thy-1) (Kubota, Avarbock, & Brinster, 2003) and inhibitor of DNA binding 4 (Id4) (Sun, Xu, Zhao, & Chen, 2015).

In domestic species, gonocytes, SSCs, and SSC progenitors are identified by expressions of *Dolichos biflorus* agglutinin (DBA) (Fujihara, Kim, Minami, Yamada, & Imai, 2011; Goel et al., 2007; Goel et al., 2010; Izadyar, Spierenberg, Creemers, den Ouden, & de Rooij, 2002), ubiquitin carboxyl-terminal hydrolase isozyme L1 (UCHL-1) (Fujihara, Kim, Minami, Yamada, & Imai, 2011; Goel et al., 2010; Heidari et al., 2012; Herrid, et al., 2007; Reding et al., 2010), and GFR α -1 (Costa et al., 2012; Lee et al., 2013; Oatley, de Avila, Reeve, & McLean, 2004;). However, the expression of biomolecular markers in germ cells from domestic species is heterogeneous. They expressed in most of undifferentiated germ cells in the testis (Zheng et al., 2014). This makes difficult to characterize SSCs from domestic animals. Another method used to recognize SSCs is to identify their localization in the testis; SSCs are located close to the basal membrane in the seminiferous tubules.

It has been shown that in vitro culture of neonatal mouse SSCs allows to male germline stem (GS) cells (Kanatsu-Shinohara et al., 2003) and ES-like cells (Kanatsu-Shinohara et al., 2004) to be produced. Moreover, in vitro culture of adult mouse SSCs generates GS cells and multipotent male germline stem (MGS) cells (Guan et al., 2006; Seandel et al., 2007). GS cells form botryoidally aggregated colonies, retain their germ cell characteristics, and contribute to the germ cell lineage (Kanatsu-Shinohara et al., 2003). By contrast, ES-like cells form tightly packed 3-dimensional colonies, express pluripotent stem cell markers, and contribute to chimeras (Kanatsu-Shinohara et al., 2004; Ko et al., 2009). MGS cells have tightly packed 3-dimensional colonies, express pluripotent stem cell markers, and develop into cardiac tissue in vitro and functional blood vessels in vivo (Seandel et al., 2007). Moreover, genetically modified cultured SSCs produced transgenic animals (Kanatsu-Shinohara et al., 2008). This suggests that cultured SSCs can be a useful tool for making genetic modifications through gene targeting and genome editing technologies.

Undifferentiated spermatogonia including SSCs from neonatal and pre-pubertal stage of domesticated animal such as pig (Zheng et al., 2013; Zhang et al., 2017) and goat (Pramod and Mitra, 2014) can also be cultured in vitro. In cattle, although gonocytes and SSCs isolated from neonatal and immature bovine testes can be maintained in culture (Aponte, Soda, van de Kant, & de Rooij, 2006; Fujihara, Kim, Minami, Yamada, & Imai, 2011; Izadyar et al., 2002; Oatley, Kaucher, Yang, Waqas, & Oatley, 2016; Sahare et al., 2016), long-term culture systems for SSCs from adult testes have been established only in mice.

Several growth factors including GDNF are employed to derive germ cell lines from immature bovine testes (Oatley, Kaucher, Yang, Waqas, & Oatley, 2016; Sahare et al., 2015). GDNF is produced by Sertoli cells and supports mouse SSC proliferation in vivo (Meng et al., 2000) and in vitro (Hasegawa, Namekawa, & Saga, 2013). In cattle, the addition of GDNF to

culture medium activates the mitogen-activated protein kinase (MAPK) signaling pathway and supports long-term proliferation of gonocyte isolated from neonatal and immature testes (Sahare et al., 2015). However, when the neonatal culture system was applied to culture adult bovine spermatogonia, they could not be maintained for a long time (Fujihara, Kim, Minami, Yamada & Imai, 2011). Hence, experimental evidence suggests that the establishment of male germ cell lines seems to be dependent on the age of the animal from which the cells are derived. The application of male germ cell lines in domestic species will be more comprehensive when it have been established from both immature and adult animals.

In this study, we developed different culture systems for bovine undifferentiated spermatogonia isolated from both immature and adult testes and characterized the cells. This finding demonstrates the feasibility of male germ cell culture in domestic species, which could facilitate progress in research related to transgenic animal production, genome editing technology for improvement of livestock production or conservation of endangered species.

2. Results

2.1 Differential gene expression in undifferentiated spermatogonia of immature and adult testes

The expression patterns of undifferentiated spermatogonia markers from immature and adult testes was compared. Differential expression patterns of both $GFR\alpha$ -1-positive and UCHL-1-positive cells were observed (Figure 1a-1e). The number of UCHL-1-positive cells was higher in immature testes ($57.11 \pm 7.93\%$) than in adult testes ($34.25 \pm 3.80\%$) ($n=6$, Figure 1c). A similar result was observed for another undifferentiated spermatogonia marker, $GFR\alpha$ -1, in immature testes ($37.61 \pm 4.94\%$) and in adult testes ($7.14 \pm 3.1\%$) (Figure 1d). These results were also confirmed by RT-PCR (Figure 1e).

119

120 **2.2 Enrichment for undifferentiated spermatogonia isolated from adult testes**

121 In adult testicular cell suspensions, two fractions that appeared at approximately
122 20% and 40% Percoll gradients were collected after gradient centrifugation. Cell
123 suspensions were seeded in pregelatinized dishes for 2 h to allow somatic cells to attach
124 to the dishes, and the floating cell suspensions were collected. DBA-positive cells were
125 identified in both the 20% and 40% Percoll fractions. In the preliminary experiments, the
126 isolated fractions were evaluated for colony formation by short-term culture. Cells that
127 were collected from the 20% fraction formed colonies and could be passaged for further
128 expanded culture. In contrast, cells from the 40% fraction did not form colonies and could
129 not proliferate after primary culture (Figure S1a-S1b). Therefore, only the cell suspension
130 from the 20% fractions was used for further studies. The number of SSCs in culture was
131 estimated by UCHL-1 and DBA staining (Figure 2a-2b). The percentage of UCHL-1-
132 positive cells in the freshly isolated cell suspensions was $10.03 \pm 3.22\%$ and significantly
133 increased to $27.71 \pm 10.46\%$ after Percoll gradient centrifugation, then further increased
134 to $36.81 \pm 11.52\%$ after negative selection on gelatin-coated dishes (Figure 2c-2d).
135 Similar results were obtained by using DBA staining (before Percoll: $15.31 \pm 2.87\%$; after
136 Percoll: $52.70 \pm 7.50\%$; after plating on gelatin-coated dishes: $63.62 \pm 4.60\%$) (Figure 2e).

137

138 **2.3 Derivation of cell lines from undifferentiated spermatogonia of immature bovine**
139 **testes**

140 Enriched populations of undifferentiated spermatogonia from immature bovine testes
141 were isolated from 3-month-old calves. Isolated cells were cultured in medium containing both
142 15% KSR and 1% FBS and supplemented with GDNF for the initial culture. After 5 to 7 days

in primary culture, the cultured cells formed botryoidally aggregated multicellular colonies. After the first passage, the culture medium was changed to DMEM/F12 medium supplemented with 20% KSR, and the cells were passaged every 5 days. In the later passages (passages 7 and 8), the cultures were passaged every 2 to 3 days owing to robust proliferation of undifferentiated spermatogonia. The cell lines formed botryoidally aggregated colonies in primary culture (Figure 3a, white arrow) and gradually changed their morphology to become dome-shaped ES cell-like colonies after several passages (Figure 3a, black arrow).

The effects of different growth factors on the proliferation of undifferentiated spermatogonia were also tested. We compared the effect of KSR media supplemented with GDNF only, bLIF only, and a combination of GDNF and bLIF on the number of colonies formed (Figure 3b) and the total number of cells (Figure 3c). The results suggested that there were no significant differences among different culture conditions (Figure 3b-3c). However, RT-PCR results revealed that the colonies cultured in the presence of bLIF or GDNF and bLIF strongly expressed *UCHL-1* after long-term culture (Figure 3d).

2.4 Characterization of cell lines from undifferentiated spermatogonia of immature bovine testes

RT-PCR analysis was performed on freshly isolated undifferentiated spermatogonia and 4 cultured cell lines at passage 8 and passage 19 (Figure 4a). Freshly isolated undifferentiated spermatogonia express pluripotency-associated genes (*NANOG*, *OCT4*, *SOX2*, and *KLF4*), as well as the germ cell marker *UCHL-1*. Cell lines also expressed *OCT4*, *SOX2* and *KLF4* and *UCHL-1* at passage 8. At the passage 19, however, only *KLF4* and *UCHL-1* were detected (Figure 4a). Immunofluorescent analysis was performed on the cell lines and detected a pluripotent marker (SOX2) and germ cell markers (UCHL-1 and DBA) (Figure 4b).

167

168 **2.5 Derivation of cell lines from undifferentiated spermatogonia of adult bovine testes**

169 Isolated undifferentiated spermatogonia were cultured in one of four different
170 media: 1) the basic medium (DMEM/F12 containing 15% KSR and 1% FBS), 2) the
171 basic medium supplemented with GDNF only, 3) the basic medium supplemented with
172 BIO only, or 4) the basic medium supplemented with both GDNF and BIO. After 7 days
173 of primary culture, colonies were observed (Figure 5a). There were no differences in the
174 total number of cells in the primary cultures (P0) among the different culture conditions
175 (Figure 5b). RT-PCR analysis showed that P0 colonies expressed the germ cell marker
176 gene *UCHL-1* (Figure 5c). In the basic medium or the basic medium with GDNF only,
177 however, enhanced proliferation of somatic cells was observed, and those cells
178 overwhelmed the growth of germ cell colonies at the first passages (Figure 5a). On the
179 other hand, the basic medium with BIO or the basic medium with BIO and GDNF
180 similarly allowed the grow of germ cell colonies (Figure 5a); therefore, only the basic
181 medium with BIO was used as a culture medium for further experiments.

182

183 **2.6 Characterization of cell lines from undifferentiated spermatogonia of adult bovine** 184 **testes**

185 Typical germ cell colonies were maintained in the presence of BIO (Figure 6a).
186 RT-PCR analysis performed on the colonies at passages 3, 8 and 10 showed that the
187 germ-cell-specific marker *UCHL-1* was stably expressed, and the stem-cell specific
188 markers *OCT4* and *KLF4* were also detected with a slightly different expression pattern
189 than *SOX2* (Figure 6b). Moreover, immunofluorescence analysis confirmed the
190 expression of DBA and GFR α -1 (Figure 6c). The same culture conditions were used to

culture enriched cell suspensions from a 17-year-old bull. Colonies appeared in the presence of BIO and were maintained for at least 3 passages (Figure S3a). The expression of *UCHL-1* was shown by RT-PCR at passage 2 (Figure S3b).

2.7 Karyotype analysis of cell lines from undifferentiated spermatogonia of immature and adult bovine testes

Karyotype analysis also showed normal metaphase spreads and chromosome number (2n=60) in SSC cell lines from immature testes (93.3%, 28 out of 30 spreads) and adult testes (96.7%, 29 out of 30 spreads), while 6.67% had 59 chromosomes in cell lines from immature testes and 3.33% had 59 chromosomes in cell lines from adult testes (Figure 7a-7b).

2.8 Stem cell potential of established cell lines

To examine the stem cell potential of cell lines from both immature and adult bovine testes, we tagged the cells with a long-term fluorescent tracer and allowed them to aggregate into embryos in the 8- to 16-cell stages (Figure 7c). Cell lines from immature testes were incorporated into the ICM of blastocysts (26.67%, Table 1), whereas cell lines from adult testes only scattered around embryos or formed cell clumps outside of embryos (Figure 7c), and a few of them incorporated specifically into the TE region of embryos (15.38, Table 1). In contrast, neither freshly isolated cells nor bovine embryonic fibroblasts (BEFs) were incorporated into the embryos. Bovine naïve-type iPS cells, which were previously established (Kawaguchi et al., 2015), were used as positive control. These cells were highly incorporated into the embryos (14.29% ICM only and 85.71% ICM and TE, Table 1).

3. Discussion

Cell suspensions from immature and adult bovine testes were purified by Percoll gradient centrifugation. The proportions of UCHL-1-positive and GFRa-1 positive cells in the cell suspensions from immature testes were higher and sufficient to enrich undifferentiated spermatogonia by Percoll gradient centrifugation (Figure 1c-1d). However, the population of undifferentiated spermatogonia in adult testicular cell suspension was significantly lower than that from immature testes (Figure 1c-1e). Differential plating on gelatin-coated dishes after Percoll gradient centrifugation was necessary to enrich undifferentiated spermatogonia from adult testes.

Undifferentiated spermatogonia from immature testes formed colonies in culture in the presence of GDNF and FBS; however, testicular somatic cells also proliferated and overgrew in the cultures after a few passages (Figure S4). By contrast, when undifferentiated spermatogonia were cultured under serum-free conditions in the presence of KSR medium containing both GDNF and bLIF, undifferentiated spermatogonia colonies were formed and maintained for a long period.

The colonies derived from immature testes undifferentiated spermatogonia were identified by their botryoidally aggregated cell morphology (Figure 3a) and then transformed to tightly packed 3-dimensional colonies, similar to mouse ES cells, after a month of culture (Figure 3a). These ES cell-like colonies were maintained for only 2 months (11 passages); however, the botryoidally aggregated colonies were more stable and could be maintained for more than 3 months under serum-free culture conditions. SSCs isolated from neonatal mouse testes start to form ES-like colonies within 4-7 weeks after initial culture (Kanatsu-Shinohara et al., 2004) and can subsequently be cultured under ES cell culture conditions while maintaining their ES-like properties (Kanatsu-Shinohara et al., 2004). ES-like bovine undifferentiated spermatogonia colonies were cultured on STO feeder cells, which is the same culture condition

used for previously established bovine naïve-type iPS cells (Kawaguchi et al., 2015); however, they could not be maintained under this condition.

The established cell lines from immature testes had mouse ES cell-like morphology in serum-free medium supplemented with GDNF and bLIF, and they expressed both germ-cell-specific (*UCHL-1* and *DBA*) and pluripotent stem cell markers (*OCT4*, *SOX2*, *KLF4*). Consistent with our results, other studies have reported that in vitro culture of gonocytes from neonatal bovine testes yields ES cell-like colonies (Sahare, Otomo, Komatsu, Minami, Yamada, & Imai, 2015) that express cell surface markers of ES cells (Li et al., 2016). In contrast, isolated undifferentiated spermatogonia from adult testes could be maintained in the medium only in the presence of BIO and formed colonies with botryoidally aggregated cell morphology, similar to typical mouse GS cells. All 4 established cell lines expressed the germ-cell-specific markers *UCHL-1*, *DBA*, and *GFR α -1* and the pluripotent stem cell markers *OCT4*, *SOX2*, and *KLF4*. BIO is an inhibitor of glycogen synthase kinase-3 α (GSK3) (Sato, Meijer, Skaltsounis, Greengard, & Brivanlou, 2004). Inhibition of GSK3 leads to the activation of the Wnt/ β -catenin signaling pathway. As recently reported in tree shrew (Li et al., 2017), Wnt/ β -catenin signaling pathway is involved in the maintenance of undifferentiated spermatogonia from adult testes during the early stage of in vitro culture. Moreover, Wnt signaling also supports regulation of the character as stem cells of mouse spermatogonial stem/progenitor cells (Golestaneh, Beauchamp, Fallen, Kokkinaki, Üren, & Dym, 2009) and promotes the proliferation of SSC progenitor cells (Takase & Nüsse, 2016).

In this study, undifferentiated spermatogonia isolated from immature testes could be maintained for long term in the basic medium with GDNF. Therefore, we hypothesized that once adult undifferentiated spermatogonia gained stem cell-like properties in the presence of BIO, BIO could be omitted from the culture and replaced by GDNF. After removal of BIO at 10 passages, colonies can be maintained in the basic medium with only GDNF (Figure S5).

Moreover, colonies could not be maintained in culture by the removal of BIO before 10 passages. This result suggests that during the first 10 passages in the presence of BIO, a fraction of undifferentiated spermatogonia in heterogeneous differentiated states in the adult testis may transform to more undifferentiated states such as SSCs. In contrast, undifferentiated spermatogonia from immature testes could be maintained in the presence or absence of BIO with no significant differences (data not shown).

To examine the stem cell potential of the established cell lines, we co-cultured three different cell lines with in vitro fertilized bovine embryos at 8 to 16 cell stage. Cell lines from immature testes were incorporated into the ICM region in 26.67% of blastocysts (Table 1 and Figure 7C); however, none of the freshly isolated undifferentiated spermatogonia could be incorporated into blastocysts. In contrast, the contribution of cell lines from adult testes was limited, consistent with a study showing that the contribution of mouse SSC-derived cell lines to chimeras is dependent on the age of the testes (Azizi et al., 2016). In our previous reports (Kawaguchi et al., 2015), established naïve-type bovine iPS cells were abundantly incorporated into both the ICM and TE regions of blastocysts after aggregation with 8-cell and 16-cell embryos. Taken together, the evidence shows that cells of cell lines derived from immature testes have partial stem cell potential, probably because of a heterogeneous stem cell population in the established cell lines or a cell population with both stem cell and germ cell characteristics, but those from adult testis have narrower limits on their potential as stem cells. However, further studies are necessary to examine the ability of the cell lines to contribute to chimeric fetuses and contribute to spermatogenesis through their germ cell potential in bovine testes.

To extend this culture system for aged animals or wild animals, we isolated undifferentiated spermatogonia from a 17-year-old bull that was a sire but was no longer used for semen collection. Isolated undifferentiated spermatogonia from that bull could be successfully cultured for 3 passages at most (Figure S3), and the population of undifferentiated

spermatogonia in the testis was very limited (Figure S1). Procedures for enrichment of the undifferentiated spermatogonia population and stimulation of undifferentiated spermatogonia proliferation are necessary for aged and seasonally breeding animals.

4. Materials and Methods

4.1 Ethical statement

All animal experiments were performed in accordance with the “Welfare Management of Livestock” standards of Gifu Prefectural Livestock Research Institute and the guidelines of the Kyoto University Livestock Farm and were approved by the Kyoto University Animal Care and Use Committee.

4.2 Isolation of undifferentiated spermatogonia from bovine testes

Undifferentiated spermatogonia were isolated by sequential enzymatic digestion methods as previously reported (Fujihara, Kim, Minami, Yamada, & Imai, 2011) with minor modifications. Briefly, the testes were collected from Japanese Black cattle of the following ages: 3 months old, 1 to 2 years old, and 17 years old. The collected testes were placed immediately into a tube and immersed in Dulbecco’s modified Eagle’s medium (DMEM)/F12 medium (GIBCO BRL Invitrogen, Carlsbad, CA, USA) supplemented with 100 IU mL⁻¹ penicillin (Sigma-Aldrich, St. Louis, MO, USA), 50 mg mL⁻¹ streptomycin (Sigma-Aldrich), 40 mg mL⁻¹ gentamicin sulfate (Sigma-Aldrich) and 15 mM HEPES (Wako, Osaka, Japan). The testes were transported to the laboratory at 4°C to 10°C within 24 h. To prepare cell suspensions, we minced the decapsulated testicular tissues and treated with a three-step enzymatic digestion method. The cells were incubated with the first collagenase solution for 50 min at 37°C, then washed three times with DMEM/F12. This was followed by incubation with a second collagenase solution for 50 min at 37°C, after which the cells were

washed again three times with DMEM/F12. The obtained suspensions were centrifuged, and the resulting cell pellets were incubated with a third collagenase solution for only 10 min at 37°C, washed three times and filtered through a 50- μ m nylon mesh.

4.3 Enrichment of isolated undifferentiated spermatogonia from immature bovine testes

Percoll gradient centrifugation was used for enrichment of undifferentiated spermatogonia isolated from immature testes. We used a Percoll gradient of 60%, 40%, and 20% Percoll (GE Healthcare Life Sciences, Sweden) as previously reported (Fujihara, Kim, Minami, Yamada, & Imai, 2011) with minor modifications. Cell suspensions were loaded on top of the Percoll gradients and centrifuged at 3,000 rpm for 30 min. After centrifugation, different cell fractions were observed. The germ-cell fraction was collected from the 20 ~ 40% Percoll fraction. The purity of the fraction was assessed by immunofluorescence staining of cell suspensions for GFR α -1 and UCHL-1.

4.4 Enrichment of isolated undifferentiated spermatogonia from adult bovine testes

Since undifferentiated spermatogonia are a rare cell population in the adult testis, the undifferentiated spermatogonia fraction of the filtered suspension of adult testicular cells was enriched by a two-step purification protocols. Percoll gradient centrifugation was performed as the first germ cell selection step, and the germ cells were collected from the 20 ~ 40% Percoll fraction. Subsequently, differential plating of the cell suspension on a gelatin-coated culture dish were used to remove any additional remaining somatic cells from the enriched Percoll fractions. The undifferentiated spermatogonia-rich fractions were plated on 0.1% pregelatinized dishes in DMEM/F12 media supplemented with 5% fetal bovine serum (FBS, Invitrogen) for 2 h. Most of the somatic cells attached to the culture dish, while the undifferentiated spermatogonia floated. Floating cells were collected and used for further

studies. The purity of the cell suspensions was assessed by immunofluorescence staining for DBA and UCHL-1.

4.5 Histological analysis of the testes

Isolated testes were fixed with Bouin's solution, embedded in paraffin using standard procedures, and sliced to a thickness of 6 μm . The paraffin sections were deparaffinized, rehydrated, and stained with hematoxylin and eosin.

4.6 Culture of undifferentiated spermatogonia isolated from immature bovine testes

Collected undifferentiated spermatogonia were cultured at 5×10^4 cells/cm² on poly-L-lysine (0.01%)-precoated wells in 24-well plates with basic media containing DMEM/F12, 100 $\mu\text{g/mL}$ penicillin (Sigma-Aldrich), 50 $\mu\text{g/mL}$ streptomycin (Sigma-Aldrich), 40 $\mu\text{g/mL}$ gentamycin sulfate (Sigma-Aldrich), 10 $\mu\text{g/mL}$ apotransferrin (Sigma-Aldrich), 10 $\mu\text{g/mL}$ insulin (Sigma-Aldrich), 110 $\mu\text{g/mL}$ sodium pyruvate (Sigma-Aldrich), 0.015% (v/v) sodium lactate (Sigma-Aldrich), 0.1% (v/v) non-essential amino acid solution (GIBCO BRL Invitrogen), and 0.01 mM β -mercaptoethanol (Wako). In the initial culture, the basic media were supplemented with 15% knockout serum replacement (KSR), 1% FBS and 20 ng/ml GDNF. After the first passage, the media were replaced with DMEM/F12 supplemented with 20% KSR and growth factors. Different growth factors—GDNF, bLIF, or both in conjunction—were added to find the optimal culture conditions for undifferentiated spermatogonia from immature testes. To establish cell lines, we employed a combination of GDNF and bLIF. The culture media were changed every 3 days.

4.7 Culture of undifferentiated spermatogonia isolated from adult bovine testes

Isolated and enriched undifferentiated spermatogonia from adult testes were cultured at a cell density of 5×10^4 cells/cm² on poly-L-lysine (0.01%)-precoated 24-well plates (Nunc) using DMEM/F12 supplemented with 15% KSR (Invitrogen) and 1% FBS. Culture conditions for undifferentiated spermatogonia were studied using four different culture media: the basic medium only, the basic medium supplemented with 20 ng/ml GDNF (Peprotech), the basic medium supplemented with 2 μ M of BIO, and the basic medium supplemented with both GDNF and BIO. After 6 to 7 days of primary culture, multicellular botryoidally aggregated colonies were observed in all cultures. The cells were passaged with 0.25% trypsin and 0.5 mM EDTA (Nacalai Tesque, Kyoto, Japan) and further cultured at 37°C in a humidified atmosphere with 5% CO₂. The culture media were changed every 3 days.

4.8 Immunocytochemical analysis of cultured undifferentiated spermatogonia

Immunocytochemical analysis on freshly isolated or cultured cells was performed as described previously (Kim, Fujihara, Sahare, Minami, Yamada, & Imai, 2014). Briefly, the cells were fixed with 4% paraformaldehyde, permeabilized, blocked, and incubated with the appropriate primary and secondary antibodies indicated below.

The following antibodies and dilutions were used: anti-DBA conjugated with fluorescein isothiocyanate (FITC) (1: 200; Vector Laboratories, Burlingame, CA, USA), anti-GFR α -1 (1:200; Santa Cruz Biotechnology, Santa Cruz, CA, USA), anti-UCHL-1 (PGP 9.5; 1:200; Santa Cruz Biotechnology), anti-SOX2 (1:200; Santa Cruz Biotechnology), and anti-GATA4 (1:200, Santa Cruz Biotechnology). All primary antibodies were incubated with the specimens overnight at 4°C. Then, the cells were washed three times with TBST (Sigma), incubated with the appropriate secondary antibodies—either Alexa Fluor 488 goat anti-rabbit IgG (Invitrogen) or Alexa Fluor 594 rabbit anti-goat (Invitrogen)—for 1 h at room temperature, and washed

three more times with TBST. The nuclei were counterstained with Hoechst 33342 (Sigma), and the specimens were mounted in 50% glycerol. Samples were observed under an immunofluorescence microscope (BX 50; Olympus, Tokyo, Japan).

4.9 RT-PCR

RNA was isolated from either freshly isolated testicular cells or cultured cells using TRIzol reagent (Ambion, Austin, TX, USA) according to the manufacturer's instructions. cDNA synthesis was performed using 1 µg of total RNA per 20 µL of PCR reaction mixture. Briefly, oligo(dT) primers were added to the isolated RNA and incubated for 5 min at 65°C. ReverTra Ace (Toyobo, Tokyo, Japan) was added to the RNA mixture for the reverse transcriptase reaction. The samples were incubated for 60 min at 42°C followed by 5 min at 99°C. The collected cDNA was stored at -20°C or immediately used for PCR. PCR amplification was performed using 1 µL of cDNA per 20 µL of PCR reaction mixture containing 2 mM MgCl₂, 0.25 mM dNTPs, 1 x PCR buffer, 5 pmol of each primer and 1 U of Taq DNA polymerase (ExTaq; TaKaRa, Tokyo, Japan). The primer sequences used for the amplification of specific genes are listed in Table 2. The PCR products were separated and visualized on 2% (w/v) agarose gels containing ethidium bromide.

4.10 Karyotype analysis

Karyotype analysis was performed using a previously described protocol (Campos, Sartore, Abdalla, & Rehen, 2009). Cultured cells were incubated in 0.1 µg/ml KaryoMAX Colcemid Solution (Invitrogen) for 3 hours. The cells were dissociated to a single-cell suspension using 0.25% trypsin-EDTA solution. Metaphase spreads were prepared by incubating single-cell suspensions in a prewarmed (37°C) hypotonic solution (KCl 75 mM) for 15 minutes and fixed in 3:1 (methanol (Wako):glacial acetic acid (Sigma)) fixative solutions overnight at 4°C.

Chromosomes were stained with VECTASHIELD mounting medium with DAPI (Vector) and observed under an immunofluorescence microscope (BX 50; Olympus, Tokyo, Japan). Thirty metaphase spreads from three independent cell lines were counted.

4.11 Aggregation of cell lines with bovine embryos

Bovine 1-cell embryos at 20 h post-insemination (hpi) and produced in vitro as described elsewhere (Ikeda, Kawahara-Miki, Iwata, Sugimoto, & Kume, 2017) were treated with pronase solution (0.5% [w/v] in PBS) to remove the zona pellucida. The zona-free 1-cell embryos were individually allocated to a well-of-the-well system (Vajta et al., 2008) using the LinKID micro25 culture dish (Dai Nippon Printing) with 50 μ L of modified synthetic oviduct fluid (mSOF) and cultured until 70 hpi. At 70 hpi, only the embryos that had developed to the 8-cell to 16-cell stages were left, and the culture medium was replaced with mSOF containing 5% (v/v) KSR, bLIF-conditioned medium (1:1000 dilution), and 20 ng/ml GDNF.

Cell lines were tagged with 0.1 μ L of Vybrant CFDA SE cell tracer dye (Invitrogen) according to the manufacturer's instructions. Subsequently, 10 to 20 cells from each of three independent cell lines were transferred to zona-free embryos at the 8-cell to 16-cell stages. After four to five days in culture, the embryos were observed under a fluorescence microscope. For subsequent examination, embryos were fixed with 4% PFA, prepared on slides, and counterstained with Hoechst 33342 (Sigma).

4.12 Statistical analysis

Pairwise comparisons of group means were conducted using Student's t-test. Multiple comparison analysis was carried out using ANOVA followed by the least significant difference test.

437 Acknowledgements

438 We thank Dr. K. Mukojima of the Gifu Prefectural Livestock Research Institute for providing
439 adult testes, and we thank Dr. Y. Hoshino, Mr. H. Yoshioka, and Ms. E. Itoyama of the Kyoto
440 University Livestock Farm for providing immature testes. We also thank Drs. Gabriela
441 Durcova-Hills and Dr. Sandeep Goel for discussion and critical reading of the manuscript.

442

443 References

444 Aponte, P.M., Soda, T., van de Kant, H.J.G., & de Rooij, D.G. (2006). Basic features of bovine
445 spermatogonial culture and effects of glial cell line-derived neurotrophic factor.
446 *Theriogenology* 65, 1828–1847.

447 Azizi, H., Conrad, S., Hinz, U., Asgari, B., Nanus, D., Peterziel, H., Hajizadeh Moghaddam,
448 A., Baharvand, H., & Skutella, T. (2016). Derivation of Pluripotent Cells from Mouse SSCs
449 Seems to Be Age Dependent. *Stem Cells International*, 8216312,
450 <https://doi.org/10.1155/2016/8216312>.

451 Campos, P.B., Sartore, R.C., Abdalla, S.N., & and Rehen, S.K. (2009). Chromosomal spread
452 preparation of human embryonic stem cells for karyotyping. *Journal of Visualized Experiments*,
453 no. 31, <https://doi.org/10.3791/1512>.

454 Costa, G.M.J., Avelar, G.F., Rezende-Neto, J.V., Campos-Junior, P.H.A., Lacerda, S.M.S.N.,
455 Andrade, B.S.C., Thomé, R.G., Hofmann, M.-C., and Franca, L.R. (2012). Spermatogonial
456 Stem Cell Markers and Niche in Equids. *PLOS ONE* 7 (8): e44091.
457 <https://doi.org/10.1371/journal.pone.0044091>.

458 de Rooij, DG. 2001. Proliferation and Differentiation of Spermatogonial Stem Cells.
459 *Reproduction* 121 (3):347–54.

- 460 Fujihara, M., Kim, S.-M., Minami, N., Yamada, M., and Imai, H. (2011). Characterization and
461 in vitro culture of male germ cells from developing bovine testis. *Journal of Reproduction and*
462 *Development*, 57, 355–364.
- 463 Goel, S., Sugimoto, M., Minami, N., Yamada, M., Kume, S., and Imai, H. (2007). Identification,
464 Isolation, and In Vitro Culture of Porcine Gonocytes. *Biology of Reproduction*, 77, 127–137.
- 465 Goel, S., Reddy, N., Mandal, S., Fujihara, M., Kim, S.-M., and Imai, H. (2010).
466 Spermatogonia-specific proteins expressed in prepubertal buffalo (*Bubalus bubalis*) testis and
467 their utilization for isolation and in vitro cultivation of spermatogonia. *Theriogenology*, 74,
468 1221–1232.
- 469 Golestaneh, N., Beauchamp, E., Fallen, S., Kokkinaki, M., Üren, A., and Dym, M. (2009). Wnt
470 signaling promotes proliferation and stemness regulation of spermatogonial stem/progenitor
471 cells. *Reproduction*, 138, 151–162.
- 472 Guan, K., Nayernia, K., Maier, L.S., Wagner, S., Dressel, R., Lee, J.H., Nolte, J., Wolf, F., Li,
473 M., Engel, W., Hasenfuss, G.. (2006). Pluripotency of spermatogonial stem cells from adult
474 mouse testis. *Nature*, 440, 1199–1203.
- 475 Hasegawa, K., Namekawa, S.H., and Saga, Y. (2013). MEK/ERK signaling directly and
476 indirectly contributes to the cyclical self-renewal of spermatogonial stem cells. *Stem Cells*, 31,
477 2517–2527.
- 478 Heidari, B., Rahmati-Ahmadabadi, M., Akhondi, M.M., Zarnani, A.H., Jeddi-Tehrani, M.,
479 Shirazi, A., Naderi, M.M., and Behzadi, B. (2012). Isolation, identification, and culture of goat
480 spermatogonial stem cells using c-kit and PGP9.5 markers. *Journal of Assisted Reproduction*
481 *and Genetics*, 29, 1029–1038.

- 482 Herrid, M., Davey, R.J., and Hill, J.R. (2007). Characterization of germ cells from pre-pubertal
483 bull calves in preparation for germ cell transplantation. *Cell and Tissue Research*, 330, 321–
484 329.
- 485 Ikeda, S., Kawahara-Miki, R., Iwata, H., Sugimoto, M., and Kume, S. (2017). Role of
486 methionine adenosyltransferase 2A in bovine preimplantation development and its associated
487 genomic regions. *Scientific Reports*, 7, 3800.
- 488 Izadyar, F., Spierenberg, G.T., Creemers, L.B., den Ouden, K., and de Rooij, D.G. (2002).
489 Isolation and purification of type A spermatogonia from the bovine testis. *Reproduction*, 124,
490 85–94.
- 491 Kanatsu-Shinohara, M., Ogonuki, N., Inoue, K., Miki, H., Ogura, A., Toyokuni, S., and
492 Shinohara, T. (2003). Long-term proliferation in culture and germline transmission of mouse
493 male germline stem cells. *Biology of Reproduction*, 69, 612–616.
- 494 Kanatsu-Shinohara, M., Inoue, K., Lee, J., Yoshimoto, M., Ogonuki, N., Miki, H., Baba, S.,
495 Kato, T., Kazuki, Y., Toyokuni, S., et al. (2004). Generation of Pluripotent Stem Cells from
496 Neonatal Mouse Testis. *Cell*, 119, 1001–1012.
- 497 Kanatsu-Shinohara, M., Kato, M., Takehashi, M., Morimoto, H., Takashima, S., Chuma, S.,
498 Nakatsuji, N., Hirabayashi, M., and Shinohara, T. (2008). Production of transgenic rats via
499 lentiviral transduction and xenogeneic transplantation of spermatogonial stem cells. *Biology of*
500 *Reproduction*, 79, 1121–1128.
- 501 Kawaguchi, T., Tsukiyama, T., Kimura, K., Matsuyama, S., Minami, N., Yamada, M., and
502 Imai, H. (2015). Generation of naïve bovine induced pluripotent stem cells using piggyBac
503 transposition of doxycycline-inducible transcription factors. *PLOS ONE*, 10, e0135403.

- 504 Kim, S.-M., Fujihara, M., Sahare, M., Minami, N., Yamada, M., and Imai, H. (2014). Effects
505 of extracellular matrices and lectin *Dolichos biflorus* agglutinin on cell adhesion and self-
506 renewal of bovine gonocytes cultured in vitro. *Reproduction, Fertility and Development*, 26,
507 268–281.
- 508 Ko, K., Tapia, N., Wu, G., Kim, J.B., Bravo, M.J.A., Sasse, P., Glaser, T., Ruau, D., Han, D.W.,
509 Greber, B., et al. (2009). Induction of pluripotency in adult unipotent germline stem cells. *Cell*
510 *Stem Cell*, 5, 87–96.
- 511 Kubota, H., Avarbock, M.R., and Brinster, R.L. (2003). Spermatogonial stem cells share some,
512 but not all, phenotypic and functional characteristics with other stem cells. *Proceedings of the*
513 *National Academy of Sciences, USA*, Proc. Natl. Acad. Sci. 100, 6487–6492.
- 514 Lee, K.H., Lee, W.Y., Kim, J.H., Yoon, M.J., Kim, N.H., Kim, J.H., Uhm, S.J., Kim, D.H.,
515 Chung, H.J., and Song, H. (2013). Characterization of GFR α -1-positive and GFR α -1-negative
516 spermatogonia in neonatal pig testis. *Reproduction in Domestic Animals*, 48, 954–960.
- 517 Li, B., Zhuang, M., Wu, C., Niu, B., Zhang, Z., Li, X., Wei, Z., Li, G., and Hua, J. (2016).
518 Bovine male germline stem-like cells cultured in serum- and feeder-free medium.
519 *Cytotechnology*, 68, 2145–2157.
- 520 Li, C.-H., Yan, L.-Z., Ban, W.-Z., Tu, Q., Wu, Y., Wang, L., Bi, R., Ji, S., Ma, Y.-H., Nie, W.-
521 H., et al. (2017). Long-term propagation of tree shrew spermatogonial stem cells in culture and
522 successful generation of transgenic offspring. *Cell Research*, 27, 241–252.
- 523 Meng, X., Lindahl, M., Hyvönen, M.E., Parvinen, M., Rooij, D.G. de, Hess, M.W.,
524 Raatikainen-Ahokas, A., Sainio, K., Rauvala, H., Lakso, M., et al. (2000). Regulation of cell
525 fate decision of undifferentiated spermatogonia by GDNF. *Science*, 287, 1489–1493.

- 526 Naughton, C.K., Jain, S., Strickland, A.M., Gupta, A., and Milbrandt, J. (2006). Glial cell-line
527 derived neurotrophic factor-mediated RET signaling regulates spermatogonial stem cell fate.
528 *Biology of Reproduction*, 74, 314–321.
- 529 Oatley, J.M., de Avila, D.M., Reeves, J.J., and McLean, D.J. (2004). Testis tissue explant
530 culture supports survival and proliferation of bovine spermatogonial stem cells. *Biology of*
531 *Reproduction*, 70, 625–631.
- 532 Oatley, M.J., Kaucher, A.V., Yang, Q.-E., Waqas, M.S., and Oatley, J.M. (2016). Conditions
533 for long-term culture of cattle undifferentiated spermatogonia. *Biology of Reproduction*, 95,
534 14. doi: 10.1095/bioreprod.116.139832.
- 535 Pramod, R.K., and Mitra, A. (2014). In vitro culture and characterization of spermatogonial
536 stem cells on Sertoli cell feeder layer in goat (*Capra hircus*). *Journal of Assisted Reproduction*
537 *and Genetics*, 31, 993–1001.
- 538 Reding, S.C., Stepnoski, A.L., Cloninger, E.W., and Oatley, J.M. (2010). THY1 is a conserved
539 marker of undifferentiated spermatogonia in the pre-pubertal bull testis. *Reproduction* 139,
540 893-903.
- 541 Sahare, M., Otomo, A., Komatsu, K., Minami, N., Yamada, M., and Imai, H. (2015). The role
542 of signaling pathways on proliferation and self-renewal of cultured bovine primitive germ cells.
543 *Reproductive Medicine and Biology*, 14, 17–25.
- 544 Sahare, M., Kim, S.-M., Otomo, A., Komatsu, K., Minami, N., Yamada, M., and Imai, H.
545 (2016). Factors supporting long-term culture of bovine male germ cells. *Reproduction Fertility*
546 *and Development*, 28, 2039–2050.

- 547 Sato, N., Meijer, L., Skaltsounis, L., Greengard, P., and Brivanlou, A.H. (2004). Maintenance
548 of pluripotency in human and mouse embryonic stem cells through activation of Wnt signaling
549 by a pharmacological GSK-3-specific inhibitor. *Nature Medicine*, 10, 55–63.
- 550 Seandel, M., James, D., Shmelkov, S.V., Falciani, I., Kim, J., Chavala, S., Scherr, D.S., Zhang,
551 F., Torres, R., Gale, N.W., et al. (2007). Generation of functional multipotent adult stem cells
552 from GPR125+ germline progenitors. *Nature*, 449, 346–350.
- 553 Sun, F., Xu, Q., Zhao, D., and Chen, C.D. (2015). Id4 Marks Spermatogonial Stem Cells in the
554 Mouse Testis. *Scientific Reports*, 5, 17594.
- 555 Takase, H.M., and Nusse, R. (2016). Paracrine Wnt/ β -catenin signaling mediates proliferation
556 of undifferentiated spermatogonia in the adult mouse testis. *Proceedings of the National*
557 *Academy of Sciences, USA.*, 113, E1489-1497.
- 558 Tegelenbosch, R.A., and de Rooij, D.G. (1993). A quantitative study of spermatogonial
559 multiplication and stem cell renewal in the C3H/101 F1 hybrid mouse. *Mutation Research*, 290,
560 193–200.
- 561 Vajta, G., Korösi, T., Du, Y., Nakata, K., Ieda, S., Kuwayama, M., and Nagy, Z.P. (2008). The
562 Well-of-the-Well system: an efficient approach to improve embryo development. *Reproductive*
563 *Biomedicine Online*, 17, 73–81.
- 564 Zhang, P., Chen, X., Zheng, Y., Zhu, J., Qin, Y., Lv, Y., and Zeng, W. (2017). Long-term
565 propagation of porcine undifferentiated spermatogonia. *Stem Cells Development*, 26, 1121–
566 1131.

567 Zheng, Y., Tian, X., Zhang, Y., Qin, J., An, J., and Zeng, W. (2013). In vitro propagation of
568 male germline stem cells from piglets. *Journal of Assisted Reproduction and Genetics*, 30, 945-
569 952.

570 Zheng, Y., Zhang, Y., Qu, R., He, Y., Tian, X., and Zeng, W. (2014). Spermatogonial stem
571 cells from domestic animals: progress and prospects. *Reproduction*, 147, 65-74.

572

573

574

575

576

577

578

579

580

581

582

583

584

Figure 1. Differential expression of GFR α -1 and UCHL-1 in cell suspensions isolated from immature and adult testes after Percoll gradient centrifugation. GFR α -1-positive cells isolated from (a) 3-month-old bovine testes and from (b) adult bovine testes. The percentages of (c) UCHL-1-positive and (d) GFR α -1-positive cells from 3-month-old and adult bovine testes. RT-PCR analysis of *UCHL-1* expression in germ cells isolated from 3-month-old or adult bovine testes (e). Higher magnifications are shown in the inset. Scale bars represent 50 μ m. n=6.

596

597 **Figure 2.** Enrichment for undifferentiated spermatogonia isolated from adult
 598 testes. (a) – (e) show UCHL-1-positive cells in freshly isolated cell suspensions. (a)
 599 UCHL-1-positive cells in the 20% Percoll fraction, (b) UCHL-1-positive cells in the 40%
 600 Percoll fraction, and (c) UCHL-1-positive cells after gelatin selection. Enrichment (%) for
 601 undifferentiated spermatogonia was estimated as either (d) the number of UCHL-1
 602 positive cells or (e) the number of DBA-positive cells. Higher magnifications are shown
 603 in the inset. Scale bars represent 50 μ m. n=4.

Figure 3. Colony formation of immature testis undifferentiated spermatogonia cultured under different conditions. Undifferentiated spermatogonia isolated from immature testes were cultured in medium containing GDNF, bLIF, or both. Morphology of the colonies cultured in medium containing GDNF and bLIF. White arrows indicate botryoidally aggregated colonies resembling GS cells, and black arrows indicate tightly packed 3-dimensional colonies resembling ES-like cells (a). Number of colonies/cm² after the first passage (P1) (b); total number of cells after P1 (c); RT-PCR analysis of *UCHL-1* at P19 in colonies cultured under different culture conditions (d). Scale bars represent 200 μm .

614

615 **Figure 4.** Characterization of cell lines from immature testis undifferentiated
616 spermatogonia cultured in medium containing GDNF and bLIF. RT-PCR analysis of
617 *NANOG*, *OCT3/4*, *KLF4*, *SOX2* (stem cell markers) and *UCLH-1* (a germ cell marker)
618 (a); immunofluorescence analysis of the expression of SOX-2, UCLH-1, and DBA (b);
619 STO feeder cells as negative control of immunofluorescence analysis (c).

Figure 5. Colony formation of undifferentiated spermatogonia from adult testis under different culture conditions. Undifferentiated spermatogonia isolated from adult testes were cultured in basic medium in the presence of GDNF, BIO, or both (a). Colonies appeared in primary culture (P0) and at the first passage (P1); (b) the total number of cells was counted at P0. (c) RT-PCR analysis of *UCHL-1* expression in P0 colonies cultured under different culture conditions. Scale bars represent 100 μ m.

Figure 6. Characterization of undifferentiated spermatogonia colonies in the presence of BIO. (a) A typical GS cell colonies at P10. (b) RT-PCR analysis of *UCHL-1*, *OCT3/4*, *KLF4*, and *SOX2* at P3, P8, and P10. (c) Immunofluorescence analysis of DBA and GFR α -1 expression.

640

641 **Figure 7.** Characterization of undifferentiated spermatogonia-derived cell
 642 lines from both immature and adult testes. (a) Karyotyping image of cells from
 643 immature and adult testis cell lines at passages 12 and 14. (b) Percentage of cells with
 644 normal chromosome number ($n=30$). (c) Cell lines from both immature and adult testes
 645 tagged with CFDA SE cell tracker were aggregated with host 8- to 16-cell in vitro
 646 fertilized embryos; naïve-type bovine iPS cells and embryos only were used as positive
 647 and negative controls, respectively. Black dashed circles indicated non-integrated cells;
 648 black arrows indicate the inner cell mass (ICM).

649 Table 1. Aggregation of undifferentiated spermatogonia derived cell lines into *in vitro* fertilized
650 bovine embryos

Donor cells	No. of aggregated embryos	No. of blastocysts developed	Chimeric blastocysts		
			No. integrated to ICM (%)	No. integrated to both ICM and TE (%)	No. integrated to TE (%)
3 month testes	37	15 (40.54)	4 (26.67)	0 (0.00)	0 (0.00)
Adult testes	35	13 (37.14)	0 (0.00)	0 (0.00)	2 (15.38)
Freshly isolated cells	13	5 (38.46)	0 (0.00)	0 (0.00)	0 (0.00)
BEF	19	8 (42.11)	0 (0.00)	0 (0.00)	0 (0.00)
Naïve-type bovine iPSCs	14	7 (50.00)	1 (14.29)	6 (85.71)	0 (0.00)
Embryo only	19	10 (52.63)	0 (0.00)	0 (0.00)	0 (0.00)

663 **Table 2.** Primer sequences used in this experiment

Gene name	Primer sequence (5' - 3')	
	Forward	Reverse
<i>UCHL-1</i>	ACCCCGAGATGCTGAACAAAG	CCCAATGGTCTGCTTCATGAA
<i>OCT4</i>	AGAGAAAGCGGACGAGTAT	AGTACAGAGTAGTGAAGTGAGG
<i>SOX2</i>	TTACCTCTTCTTCCCACTCC	TTCTTGCTGTCCTCCATTTC
<i>KLF4</i>	CCCACACAGGTGAGAAACCT	ATGTGTAAGGCGAGGTGGTC
<i>C-KIT</i>	GACCTGGAGGACTTGCTGAG	AGGGGCTGCTTCCTAAAGAG
<i>STRA8</i>	GATGGGAATGCAAACAGCTT	GTCCAGGAAACTTGCCACAT
<i>SYCP3</i>	TGACTTTGTTCCAGCAGTGG	ACTTTCGGACACTTGCCATC
<i>β-ACTIN</i>	TCCCTGGAGAAGAGCTACGA	ACATCTGCTGGAAGGTGGAC

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678 Supporting information

679
 680 **Figure S1.** Histology of the testes. Hematoxylin and eosin staining shows germ cells in the
 681 seminiferous tubules of testes from immature (3-month-old) and adult (1.8-year-old and 17-
 682 year-old) cattle.

Figure S2. Germ cells isolated from the testis of a 1.4-year-old bovine. DBA-positive germ cells isolated from the 1.4-year-old bovine testis in the 40% Percoll fraction after gelatin selection (a); morphology of germ cell colonies collected from the 40% Percoll fraction. Isolated cells were cultured in the basic medium (control) and in the presence of GDNF, BIO, and a combination of BIO and GDNF (b). Colony formation was examined in primary culture (P0) and after the first passage (P1).

Figure S3. Undifferentiated spermatogonia isolated from the testis of a 17-year-old bovine. Undifferentiated spermatogonia colonies after passages 1 and 3 in the presence of GDNF or BIO (a); RT-PCR gene expression analysis of the germ cell marker *UCHL-1* was examined in colonies at passage 2 in the presence of BIO (b).

Figure S4. Undifferentiated spermatogonia colonies collected from immature bovine testes cultured in the presence and absence of FBS at passage 1 (P1) and P2.

723

724 **Figure S5.** Morphology and immunofluorescence analysis of undifferentiated
 725 spermatogonia colonies maintained in GDNF only. Botryoidally aggregated cell morphology
 726 of colonies in the presence of GDNF (a), immunofluorescences analysis of UCHL-1, Sertoli
 727 cell marker GATA4, and DBA (b), RT-PCR analysis of *UCHL-1*, *C-KIT*, *STRA8*, and *SYCP3*
 728 at passage 18 (P18) (c).