

1 **Non-free ionic transport of sodium, magnesium, and calcium in streams of two**
2 **adjacent headwater catchments with different vegetation types in Japan**

3
4 Tomomi TERAJIMA ^{*a}, Mihoko MORIIZUMI ^{*b}, and Tomohiro NAKAMURA ^{*c}

5
6 ^{*a}: Disaster Prevention Research Institute, Kyoto University. Gokasho, Uji, Kyoto
7 611-0011, Japan; terajima@scs.dpri.kyoto-u.ac.jp

8 ^{*b}: Faculty of Agriculture, Ryukoku University. 1-5, Yokoya, Seta-Ohecho, Otsu,
9 Shiga 520-2194, Japan; moriizu@agr.ryukoku.ac.jp

10 ^{*c}: Suuri-Keikaku Co. Ltd. 2-5-4 Sarugaku-cho, Chiyoda, Tokyo 101-0064, Japan;
11 nakamura_tomohiro@sur.co.jp

12
13 Corresponding author: T. Terajima

14 Tel: +81-774-38-4628

15 Fax: +81-774-38-4118

16 E-mail address: terajima@scs.dpri.kyoto-u.ac.jp

17
18 **SUMMARY**

19 Sodium (Na), magnesium (Mg), calcium (Ca) are usually believed to occur mostly as
20 free ions in the fresh water and consequently little is known about their chemical
21 species. To understand the importance of non-free ionic fractions (NIF) of major
22 metals in freshwater streams, Na, Mg, Ca, silicon (Si), and fulvic acid-like materials
23 (FAM) were measured in streams of mountainous adjacent headwater catchments

1 dominated by different vegetation types (planted evergreen coniferous forest and
2 natural deciduous broadleaf forest). During both no rainfall periods and rainstorms, the
3 proportion of NIF relative to total elements was lower in the coniferous catchment than
4 in the deciduous catchment, although it sometimes accounted for half or more of the
5 total concentrations of Na, Mg, and Ca in both catchments. The solubility of metal
6 compounds was higher than the measured maximum concentrations of Na⁺, Mg²⁺, and
7 Ca²⁺ to the extent that inorganic bonding was hardly possible. During no rainfall
8 periods when FAM was slightly produced into the streams, the fluxes of NIF and Si
9 were highly correlated ($r > 0.92$, $p < 0.0001$, $n = 30$) in both catchments. During a
10 small rainstorm, the flux of NIF correlated weakly with that of Si but did not correlate
11 with that of FAM in both catchments. In contrast, during a heavy rainstorm, the flux of
12 NIF correlated strongly ($r \geq 0.83$, $p < 0.0001$, $n = 26$) with that of FAM in the
13 deciduous catchment where relatively deep soil water compared to near-surface water
14 was the predominant component of stream water. However, during the heavy rainstorm
15 in the coniferous catchment, only the flux of NIF originated in the quick-flow
16 component (i.e., surface or near-surface water) in stream water (Δ NIF) correlated
17 strongly ($r \geq 0.81$, $p < 0.0001$, $n = 22$) with that of FAM. These findings imply that
18 heavy rainstorms may enhance the bonding of the major metals with humic substances
19 mainly in the deciduous catchment; and also exhibit that, in the headwater catchments,
20 both water flow pathways resulted from the different vegetation types play a very
21 important role to promote the bonding of major metals with humic substances in
22 stream water.

23

1 **Key words:** fulvic acid, major metal, bonding, humic substance, organic–inorganic
2 interaction

3

4 **1. Introduction**

5 Major metals such as sodium (Na), magnesium (Mg), and calcium (Ca), which are
6 the vital elements involving the metabolism and physiological function of life, are
7 thought to occur mainly (> 98%) as free ions in freshwater environments (e.g.,
8 Mantoura *et al.*, 1978). On the contrary, the biogeochemical implication (i.e., bonding)
9 of the major metals, humic substances, and nano-scale phyllosilicates represented by
10 clay minerals (Wilson *et al.*, 2008) has been mentioned in particular in the laboratory
11 and modeling studies (e.g., Choppin and Shanbhag, 1981; Livens, 1991; Romkens *et*
12 *al.*, 1996; Majzik, A. and Tombacz, E., 2007; Takahashi *et al.*, 1999 and 2002). This
13 discrepancy regarding major metal species implies that little is known about their
14 transport forms (i.e., chemical species) in freshwater environments.

15 In contrast, as described by Tipping (1993), the information regarding the effect of
16 environmental factors (topography, geology, climate, hydrology, vegetation, etc.) on
17 metal bonding must be very useful to allow the calculation and modeling of chemical
18 speciation under natural condition in freshwater environments. Furthermore, the
19 insights related to the interaction between aquatic and terrestrial biogeochemistry will
20 also facilitate a better understanding of controls on humic substance production,
21 consumption, and flux across whole landscapes and biomes (McDowell, 2003).
22 However, the effects of environmental factors on bonding possibility of major metals
23 to humic substances and nano-scale phyllosilicates and consequently the transport in

1 bonding in stream water are as yet unclear.

2 Accordingly, characterizing the bonding of major metals to humic substances and
3 nano-scale phyllosilicates in field sites must provide useful information about the
4 chemical regimes in stream catchments, and besides, give a novel knowledge in
5 catchment hydrology. If the free ions of major metals combined with humic substances
6 and nano-scale phyllosilicates, the relationships between the fluxes of combined
7 materials (i.e., non-free ionic fraction of the major metals), humic substances, and
8 nano-scale phyllosilicates can be the good indices to explore broadly both the
9 implication as represented by the bonding of Na, Mg, and Ca and consequently the
10 characteristics of bonding resulted from the environmental factors.

11 Thus we measured Na, Mg, Ca, and silicon (Si) in streams of two mountainous
12 adjacent headwater catchments composed of differing vegetation. Fulvic acid-like
13 materials (FAM), representative of dissolved organic matter (DOM) in the streams that
14 we observed (Terajima and Moriizumi, 2013) and usually accounts for 60% to 99% of
15 DOM in stream or ground water (Thurman, 1985; Malcolm, 1990; Artinger *et al.*,
16 2000), were also measured to understand the interaction among the major metals, clay
17 minerals, and humic substances. Additionally, FAM and dissolved organic carbon
18 (DOC) was measured for soil extracts from both catchments to determine where FAM
19 and DOC are stored in the soil profile and examine water flow pathways along the
20 slopes. Subsequently we broadly estimated how stream water chemistry (i.e., bonding
21 possibility) was provided and how slope hydrology depending on differing vegetation
22 (i.e., environmental factor) affected the transport of major metals.

23

1 **2. Site description**

2 *2.1. Topography, geology, and climate*

3 The study area consisted of two mountainous headwater catchments with differing
4 vegetation in the Nariki catchment (35°50'N and 139°10'E at the rain gauge in Fig. 1a),
5 which is about 40 km west of central Tokyo. Slope gradients in the catchment range
6 between 35° and 45°, and Mt. Kuro-yama, at 842 m, is the highest point. The
7 catchment is underlain by sandstone and mudstone bedrock of Jurassic age that also
8 contains some chert. The rock layers dip 60 - 90° northeast and strike NW-SE (Inosato
9 *et al.*, 1980; Tokyo Prefectural Public Work Institute, 2002).

10 According to data from the Automated Meteorological Data Acquisition System
11 (AMeDAS) of the Japan Meteorological Agency, the average annual precipitation
12 between 1976 and 2009 at Ohme city (35°47'N, 139°18'E; 10 km southeast of the
13 Nariki catchment and 100 m in elevation) was 1487 mm, and the average minimum,
14 maximum, and annual air temperatures during the same period were -6.7 °C in late
15 January, 36.9 °C in mid-August, and 14.2 °C, respectively. Rainstorms occur mainly
16 from mid-June to late July (the rainy season), and in the typhoon season which lasts
17 from mid-August to mid-October. About 80% of the annual precipitation falls during
18 these periods. Dry conditions prevail in winter, from early December to early March,
19 although snow usually accumulates in the Nariki catchment to a maximum depth of 10
20 to 20 cm in February. The stream water temperature of both headwater catchments
21 ranged from 10 °C in winter to 23 °C in summer.

22

23 *2.2. Experimental catchments*

1 A first headwater stream (first-order stream; 70 m in length) on the north side of
2 the Nariki River (Figs. 1a and 1b) is densely surrounded by an unmanaged evergreen
3 coniferous forest composed of Japanese cedar (*Cryptomeria japonica*) and Hinoki
4 cypress (*Chamaecyparis obtusa*) which were planted in 1961 (Coniferous catchment:
5 1.29 ha, 100 m in relief; Fig. 2a). In 2007, the catchment contained 2000 to 2500 trees
6 ha⁻¹, most with trunk diameters of ≤ 20 cm. The forest canopy is mostly closed, with
7 little understory vegetation owing to the weak penetration of sunlight to the forest floor.
8 The litter layer on the forest floor is very thin, and mineral soil is exposed in some
9 places.

10 A second headwater stream (first-order stream; 70 m in length) on the south side of
11 the Nariki River (Fig. 1a and 1c) is surrounded by a natural deciduous broadleaf forest
12 (Deciduous catchment: 1.28 ha, 100 m in relief; Fig. 2b). The dominant tree species in
13 this catchment are oak (*Quercus serrata* and *Quercus mongolica*), beech (*Fagus*
14 *japonica*), chestnut (*Castanea crenatus*), and Japanese maple (*Acer palmatum*). The
15 understory vegetation is dense in comparison to that in the coniferous catchment, and
16 there is a thick litter layer on the forest floor.

17 The riparian zone (the flat bottomland along the streams) is very small and narrow
18 in both catchments (Figs. 1b and 1c). Thus the contribution of subsurface and
19 groundwater flow for the stream water generation, which had been involved in the
20 riparian zone before rainstorms, seems to be relatively small compared to those from
21 the slopes.

22 The boundary between soil and bedrock was determined by cone penetration test to
23 occur at $N_{10} = 50$, and the soil depth ($N_{10} \leq 50$) in both catchments was ≤ 3 m. The

1 structure of the soil profile (the soil horizons present and their thicknesses) is similar
2 along the slopes of both catchments. In both catchments, the soils are classified as
3 Cambisols (according to the classification by the Food and Agriculture Organization of
4 the United Nations), and soil pH ranges between 3.8 and 5.2 throughout the vertical
5 soil profile in both catchments. The strongly organic-rich soil horizons, the A₀ to AB
6 horizons, are up to 25 cm thick in both catchments, and a gravel rich mineral soil (B
7 horizon) is below 15 to 25 cm depth. The soil parent material is presumably derived
8 from upslope via soil creep or rock slides because many discrete angular stones and
9 sediment particles in various size are included and mixed intricately throughout the
10 soil profiles. The saturation hydraulic conductivity (K_{sat}) of the soil above 80 cm depth
11 in both catchments ranges mostly between 10^{-2} and 10^{-3} cm s⁻¹. Soil porosity in the
12 coniferous catchment gradually decreases with soil depth, ranging between 70% at 10
13 cm depth and 45% at 70 cm depth, whereas soil porosity in the deciduous catchment is
14 relatively constant, decreasing from 70% near the soil surface to 53% at 70 cm depth.
15 The gravitational water drainage capacity of the soil in the coniferous catchment is
16 below 5%, except between 0 and 5 cm depth, where it is 28%. In the deciduous
17 catchment, it is constant at 9% throughout the soil profile up to 50 cm depth (Hirano *et*
18 *al.*, 2008).

19 A unique high permeable and organic rich layer, the so-called biomat ($K_s > 10^{-2}$ cm
20 s⁻¹; porosity above 70%), consisting of a dense network of fine roots of Japanese cedar
21 and Hinoki cypress within the loose litter and root-permeated zone (Sidle *et al.*, 2007),
22 was common on the slope surface (above 0 cm depth) in the coniferous catchment. Its
23 thickness varied from a few centimeters to 20 cm in the coniferous catchment, but it

1 was rare in the deciduous catchment (Terajima and Moriizumi, 2013). Thus water flow
2 through the biomat (i.e., biomatflow) during rainstorms appeared predominantly in the
3 coniferous catchment (Hirano *et al.*, 2008).

4 On the basis of the above facts, the characteristics of the environmental factors in
5 the two headwater catchments are briefly summarized as follows: The same or similar
6 characteristics are geography, topography (i.e., catchment area, elevation, relief, slope,
7 stream order, and stream length), geology, soil (i.e., type and depth), and climate. In
8 contrast, the different characteristics are vegetation, soil profile (i.e., biomat and little
9 litter layer predominant in the coniferous catchment), and resultantly hillslope
10 hydrology (i.e., biomatflow predominant in the coniferous catchment).

11

12 **3. Instrumentation and data collection**

13 *3.1. Soil collection and soil water extraction*

14 We collected soil samples at slope surfaces and at 20 to 30 cm intervals up to 140
15 cm depth (within the limits of the possible to excavate manually) at the ridgetop, mid-,
16 and lower slopes in both catchments.

17 The collected soil samples (soil pH ranged between 3.8 and 5.2) were air-dried and
18 subsequently 3 g of the samples were shaken for 24 h in 60 mL of ultrapure water (pH
19 5.6, 18.2 MΩ cm, and dissolved organic carbon (DOC) < 0.01 μg L⁻¹) obtained from
20 an Advantec Japan ® water purification system (model number: RFU585DA). Then,
21 the soil extracts were filtered for the measurements of excitation-emission matrix
22 fluorescence spectrum (EEM spectrum) and DOC to obtain the information on the
23 storage of water extractable FAM and DOC in soil. The EEM analysis for soil extracts

1 is described in Section 3.4.4. The DOC concentration was measured with a Shimadzu
2 ® organic carbon analyzer (model number: TOC-V; Detection limit is 0.004 mg L⁻¹ and
3 measurement error was $\leq \pm 0.075$ mg L⁻¹).

4 The absolute values of the EEM spectrum in soil extracts are variable on account of
5 the measurement method based on the proportion of air-dried soil weight relative to
6 ultrapure water volume. Thus, the values of EEM spectrum in soil extracts are
7 impossible to compare directly with those in stream water.

8

9 *3.2. Measurement of rainfall and stream discharge*

10 A rain gauge (0.2 mm count⁻¹), which recorded precipitation at 5-min intervals,
11 was installed in an open space near the two catchments (710 m elevation; Fig. 1a)
12 where rainfall was unlikely to be intercepted by the tree canopies because of no
13 canopies beyond 45° in elevation. According to our past study (Nogi, 2007;
14 unpublished), the average annual rainfall interception by the tree canopy was about
15 18% and 16% of total rainfall in the coniferous and deciduous catchments, respectively.
16 Moreover, the concentration of Na⁺, Mg²⁺, and Ca²⁺ supplied by rainfall (through fall +
17 stem flow) into the forests in both catchments differed by < 1.0 mg L⁻¹. These things
18 imply that the flux of Na⁺, Mg²⁺, and Ca²⁺ into the forests was < 4.66×10⁻² mg s⁻¹ and
19 < 4.98×10⁻² mg s⁻¹ in the coniferous and deciduous catchments, respectively; mostly
20 similar in both catchments. In contrast, the flux of potassium ions (K⁺) into the forests
21 in the coniferous catchment was 50% of that in the deciduous catchment.

22 Stream discharge in the two catchments was measured from January 2006 to
23 September 2007 using parshall flumes (15 cm size) installed at the lower end of the

1 streams in both catchments (Figs. 1b and 1c). Water levels in the flumes were recorded
2 at 5-min intervals by a Trutrack ® automatic sensors (model number: WT-HR64K),
3 and the relationship between the water level and the stream discharge manually
4 measured at the flumes, the H-Q relation, was used to estimate the specific stream
5 discharge ($\text{m}^3\text{d}^{-1}\text{ha}^{-1}$ or $\text{L s}^{-1}\text{ha}^{-1}$) throughout the observation period.

6

7 *3.3. Stream water collection*

8 Quick-flow component in stream water is very difficult to quantify immediately in
9 field sites. Thus in our study, notwithstanding the time intervals after rainstorms, the
10 baseflow situation was qualitatively defined as when the hydrograph was on the falling
11 limb and stream discharge seemed to be small and constant even though the small
12 rainfall was supplied; implying when the quick-flow component seemed not to
13 contribute directly to stream discharge (i.e., no increase in stream discharge).

14 Stream water during no rainfall periods (baseflow) in both catchments was
15 manually and randomly collected in polyethylene bottles from January 2006 to August
16 2007 (total numbers of baseflow samples in both catchments were 54, but were 30 for
17 collection from October 2006 to August 2007). The minimum and maximum stream
18 discharge (i.e., baseflow discharge in this case) at collection were $0.02 \text{ L s}^{-1} \text{ ha}^{-1}$ (8
19 May 2006) and $3.19 \text{ L s}^{-1} \text{ ha}^{-1}$ (8 October 2006) in the coniferous catchment, and 0.53
20 $\text{L s}^{-1} \text{ ha}^{-1}$ (23 March 2006) and $5.06 \text{ L s}^{-1} \text{ ha}^{-1}$ (8 October 2006) in the deciduous
21 catchment, respectively.

22 Two rainstorms (Storm 1 and Storm 2, 26 to 27 September 2006 and 14 to 16 July
23 2007, respectively) caused increased stream water. Flow samples were collected during

1 these storms in polyethylene bottles by at 60 to 90 min intervals through an American
2 SIGMA ® water samplers (model number: 900) on the upstream side of the partial
3 flumes. The numbers of water samples in both catchments were 24 and 29 for Storm 1
4 and Storm 2, respectively. Storm 1 had a small-scale rainstorm with 40.4 mm of total
5 rainfall, maximum rainfall intensity of $1.9 \text{ mm } 5 \text{ min}^{-1}$, and 8.2 days yr^{-1} in rainfall
6 frequency for the same storm scale. In contrast, Storm 2 was a typhoon storm with
7 117.0 mm of total rainfall, maximum rainfall intensity of $1.6 \text{ mm } 5 \text{ min}^{-1}$, and 2.9 days
8 yr^{-1} in rainfall frequency for the same storm scale.

9 Glass bottles were also used for intermittent manual sampling of the stream water
10 during no rainfall periods and rainstorms to verify the effect of chemical adsorption
11 caused by the difference in collection materials. The concentration of major metals and
12 EEM spectrum of stream water was then compared with those of the water samples
13 collected simultaneously in the polyethylene bottles; they were below measurement
14 errors for total elements and ionic materials described in Sections 3.4.1. and 3.4.2. and
15 $< 0.13 \text{ QSU}$ for FAM. This finding implies that no influence to the concentration of
16 major metals and the EEM spectra of samples was brought by the difference in
17 collection materials.

18 Water samples collected in polyethylene bottles were transferred to glass bottles as
19 soon as possible. The samples during no rainfall periods were stored at $4 \text{ }^\circ\text{C}$ within one
20 day after collection, whereas the samples during rainstorms were stored within 3 days
21 after collection, because the water collection during rainstorms ran for 2 or 3 days. All
22 stream water samples were filtered within 5 days (120 hrs.) after collection. The
23 $0.45\text{-}\mu\text{m}$ mixed cellulose ester filters (Advantec Japan or Sartorius Mechatronics

1 Japan) were used in the filtration. Then analyses were conducted for total elements
2 (T-Na, T-Mg, T-Ca, and T-Si), free ions (Na^+ , Mg^{2+} , and Ca^{2+} ; K^+ was scarcely detected
3 in the stream water), and EEM spectra.

4

5 *3.4. Measurement of concentrations in water samples*

6 *3.4.1. Total elements (T-Na, T-Mg, T-Ca, and T-Si)*

7 At first, 5 mL of hydrochloric acid (HCl : 3 mol L^{-1}) was added to 40 mL of filtered
8 water to obtain the high sensitivity for the measurement of major metals by adjusting
9 the pH between 0.5 and 1.0, the approximate pH of standard solutions for the major
10 metals of an inductively coupled plasma atomic emission spectrometer (ICP) that we
11 used. A Shimadzu ® ICP (model number: ICPS-1000 IV, Detection limit is 0.01 mg
12 L^{-1}) was used to measure the concentration of total elements in the filtrate. The
13 injection volume of water sample was 30 mL. Standard solutions for Na, Mg, Ca, and
14 Si, of which the concentrations were 1, 5, and 25 mg L^{-1} , were used to calibrate the
15 results (measurement errors: $\leq \pm 0.01 \text{ mg L}^{-1}$ for Na-, $\leq \pm 0.04 \text{ mg L}^{-1}$ for Mg-, $\leq \pm 0.17$
16 mg L^{-1} for Ca-, and $\leq \pm 0.01 \text{ mg L}^{-1}$ for Si-analyses).

17

18 *3.4.2. Free ions (Na^+ , Mg^{2+} , and Ca^{2+})*

19 Free ions in the stream water filtrate without added HCl were measured with a
20 TOA-DKK ® ion chromatograph (IC; model number: IA-200 for anions and IA-300
21 for cations). The minimum injection volume of water sample into the instruments was
22 0.2 mL for anions and 0.02 mL for cations, measurement temperature was $40 \pm 4^\circ\text{C}$,
23 and detection limit is 0.01 mg L^{-1} . The eluant (mixed solution of phthalic acid, tris

1 hydroxymethyl aminomethane, and boric acid for anions; and methane-sulfonic acid
2 solution for cations) have a pH of 3.0. Single-point calibration by measurement of a
3 standard solution of each ion (10 mg L⁻¹ of Cl⁻, NO₃⁻, Na⁺, and Mg²⁺, and 20 mg L⁻¹ of
4 SO₄²⁻, K⁺, and Ca²⁺) was used (measurement errors: $\leq \pm 0.1$ mg L⁻¹). Bicarbonate
5 (HCO₃⁻) was measured by the titration for the stream water samples of 50 mL, which
6 was calculated from the alkalinity under pH=4.8 (measurement error: $\leq \pm 0.3$ mg L⁻¹).

7 The pH of the stream water during the observation period ranged between 6.0 and
8 8.0. Thus, the concentrations of ions measured by IC with an eluant pH of 3.0 may
9 have been somewhat higher than their natural concentrations in the stream water.
10 Although we didn't quantify the concentration of organic acid (OA), chromatic peaks
11 likely to be the formic acid and oxalic acid were detected mainly in water samples
12 from the coniferous catchment.

13

14 3.4.3. *Non-free ionic fraction (NIF)*

15 Strellis *et al.* (1996) indicated that laboratory standard calibration was the most
16 likely sources of large systematic errors in ICP and IC analyses. Thus, the
17 concentrations of the standard solutions, used for each calibration for ICP and IC, were
18 alternately measured with IC and ICP to identify and correct any instrumental
19 discrepancies between ICP and IC analyses. Each measured concentration, however,
20 ranged within the measurement error, implying that the measured concentrations were
21 mostly same as the originals of each standard solution and consequently the
22 instrumental discrepancies were negligible in our measurement.

23 The difference in concentration between total elements and free ions of metal

1 elements were equated with the concentration of non-free ionic fractions (NIF: all
2 materials that were not measurable with IC).

3

4 3.4.4. Fulvic acid-like materials (FAM)

5 The EEM spectra in the filtrate for the soil extracts and stream water samples
6 without added HCl were measured with a JASCO® 3-dimensional EEM spectrometer
7 (model number: FP-6600), at excitation wavelengths of 220–550 nm and emission
8 wavelengths of 250–600 nm (i.e., Ex/Em: 220-550/250-600) and a scanning speed of
9 2000 nm min⁻¹ (measurement error was $\leq \pm 0.08$ QSU). In our measurements (e.g., Fig.
10 3), a visible single peak in fluorescence intensity corresponding to FAM appeared in
11 the water samples during rainstorms at around Ex/Em: 340/440 (Coble *et al.*, 1993;
12 Suzuki *et al.*, 1998; Mostofa *et al.*, 2005; Fellman *et al.*, 2010), which could reflect the
13 terrestrially organic matter derived from vascular plants or soil organic matter in
14 wetlands and forested environment (Coble, 1996; Coble *et al.*, 1998). Although another
15 peak in fluorescence intensity appeared at around Ex/Em: 260/440 (Fig. 3c to 3f)
16 corresponding to fluorescence that resembles fulvic acid (Fellman *et al.*, 2010), we
17 adopted exclusively the peaks at around Ex/Em: 340/440 for our analyses. In contrast,
18 as shown in Fig. 3a and 3b, we did not measure FAM during no rainfall periods
19 because visible fluorescent peaks did not always appear, so that no data is presented.

20 Irrespective of stream discharge during rainstorms, some stream water samples (2
21 and 5 samples for Storm 1 in the coniferous and deciduous catchments, respectively,
22 and 10 samples for Storm 2 in both catchments) exhibited small peaks at around
23 Ex/Em: 270/320 (< 2.4 QSU and < 1.3 QSU in the coniferous and deciduous

1 catchments, respectively), possibly corresponding to peaks in amino acid-like (or
2 protein-like) materials (Coble *et al.*, 1993; Fellman *et al.*, 2010). As shown in Fig. 3,
3 the peaks indicating humic acid-like (e.g., at around Ex/Em: 480/520) and other
4 unknown materials were not detected visibly in our analyses.

5 To determine the fluorescence intensity of FAM (*F-FAM*), the peaks in fluorescence
6 intensity corresponding to FAM were normalized relative to the fluorescence intensity
7 of a 10 $\mu\text{g L}^{-1}$ quinine sulfate solution (= 10 QSU) at Ex/Em: 345/440; close to the
8 peak obtained for FAM (Ex/Em: 340/440). The solvent (sulfuric acid, 0.1 mol L^{-1} ;
9 Wako Pure Chemical Industries Japan) of this solution exhibited no fluorescence
10 spectrum. The fluorescence intensity of the quinine sulfate solution was verified every
11 140 min (for every batch of seven water samples) to determine any instrumental errors
12 which may have been caused by fluctuations in illumination.

13 In Terajima and Moriizumi (2013), the correlation lines of the *F-FAM* relative to
14 DOC concentration showed the linear correlations with different slopes for each water
15 sample at $\text{DOC} < 22 \text{ mg L}^{-1}$ and $F\text{-FAM} < 120 \text{ QSU}$ but became convex at $\text{DOC} > 22$
16 mg L^{-1} and $F\text{-FAM} > 120 \text{ QSU}$. This shows that attenuation of the FAM spectrum due
17 to high FAM concentration probably began at values $> 120 \text{ QSU}$. *F-FAM* of the soil
18 extracts from soil surfaces were greater than $> 120 \text{ QSU}$ (Fig. 4a). Thus the FAM
19 spectrum of these samples (without dilution) could have been attenuated and
20 consequently natural *F-FAM* of the soil extracts at soil surfaces may be greater than
21 the data shown in Fig. 4a. In contrast, because the *F-FAM* of all stream water filtrate
22 was $< 20 \text{ QSU}$, we did not dilute the water samples to avoid FAM spectrum
23 attenuation.

1 The *F*-FAM of water should usually be corrected for the inner filtering effect, for
2 example, on the basis of Lakowicz (1999). However, the correction coefficients of the
3 stream water filtrate, of which all *F*-FAM was below 20 QSU in our study, ranged
4 within the measurement error of the fluorescence intensity of 10 $\mu\text{g L}^{-1}$ quinine sulfate
5 solution. Accordingly, except for the soil extracts, the inner filter correction of the
6 stream water filtrate found to be negligible and thus we did not conduct the correction
7 for stream water.

8

9 *3.5. Statistical analysis*

10 To understand roughly the bonding possibility of NIF in the headwater streams, we
11 examined the relationship between stream discharge, the fluxes of Si, NIF, and FAM
12 by calculating their linear correlation coefficients (*r*-values with *p*- and *n*-values)
13 during no rainfall periods and two rainstorms. Then, total elements, free ions, NIF, and
14 Si (all units are in $[\text{mg L}^{-1}]$) were multiplied by the specific stream discharge at
15 corresponding water collection $[\text{L s}^{-1} \text{ ha}^{-1}]$ to obtain their specific fluxes $[\text{mg s}^{-1} \text{ ha}^{-1}]$.
16 In addition, to equivalently compare the relative differences in NIF transport affected
17 by the FAM amount in the stream water of the coniferous and deciduous catchments,
18 the *F*-FAM [QSU] (i.e., apparent concentration) was multiplied by the specific stream
19 discharge at corresponding water collection $[\text{L s}^{-1} \text{ ha}^{-1}]$ and we obtained the apparent
20 specific flux of FAM $[\text{QSU L s}^{-1} \text{ ha}^{-1}]$ in the stream water of both catchments.

21 Because the specific flux is obtained from multiplying the concentration of major
22 metals by specific stream discharge, the relationship between the specific flux and
23 specific stream discharge seems inevitably to show a good correlation. For example in

1 a headwater in Japan, however, the flux of wash load (i.e., multiplying the
2 concentration of wash loads by stream discharge) does not always necessarily represent
3 good correlations with stream discharge (Terajima *et al.*, 1997). Besides, the sources of
4 stream water, major metals, Si, NIF, and FAM are surmised to be spatially and
5 temporally different even in small headwater catchments: which seems too hard to
6 understand the relationship between the complicated hydrological processes and the
7 cause of changes in concentration. In addition, the specific flux is convenient both to
8 compare the chemical characteristics in various streams where catchment area and
9 stream discharge is usually different and to separate the flow component of stream
10 water as described in Section 3.6. Accordingly, in the strict sense, although the
11 relationships between specific stream discharge, the specific flux of major metals, Si,
12 NIF, and FAM do not necessary reveal the bonding forms of Na, Mg, and Ca in stream
13 water, the specific flux (involving the difference in stream discharge and catchment
14 area) can be one of the useful factors capable of understanding the transport dynamics
15 based on the mass balance of chemical materials.

16

17 *3.6. Separation of the stream water components and NIF*

18 Hirano *et al.* (2009) suggested that in the catchments we observed, the Si
19 concentration could be used as index for assessing the baseflow component of the
20 stream water. The baseflow component of the stream water derives from groundwater
21 or deep subsurface water, and as a result of reactions between the water and the soil or
22 basement rocks during the relatively long contact time between them, the Si
23 concentration in the baseflow is high (Wels *et al.*, 1991). Hirano *et al.* (2009) also

1 showed that, based on the mass conservation law and the Si flux, the baseflow
2 component of the stream water during rainstorms could be calculated as follows:

$$3 \quad Q_{\text{(baseflow in rainstorm)}} = Q_{\text{(rainstorm)}} (1 - Q_{\text{(quick-flow)}} / Q_{\text{(rainstorm)}}), \quad (1)$$

4 where $Q_{\text{(baseflow in rainstorm)}}$ is the baseflow component of the stream water during
5 rainstorms; $Q_{\text{(rainstorm)}}$ is the measured stream discharge during rainstorms, including
6 both baseflow and quick-flow components; and $Q_{\text{(quick-flow)}}$ is the quick-flow
7 component of the stream water calculated from the Si flux of near-surface water, as
8 represented by rainwater containing relatively low Si concentrations (a maximum of
9 0.27 mg L^{-1} in the presented case). Thus, in Eq. 1, $Q_{\text{(quick-flow)}} / Q_{\text{(rainstorm)}}$ is the ratio of
10 the quick-flow component to the total stream discharge during rainstorms, and $(1 - Q_{\text{(quick-flow)}} / Q_{\text{(rainstorm)}}$)
11 represents the ratio of the baseflow component to the total stream
12 discharge during rainstorms. Both these ratios can change over the course of a storm.

13 The NIF flux originating from the quick-flow component of the stream water
14 (ΔNIF) is calculated as follows:

$$15 \quad \Delta NIF = NIF_{\text{(rainstorm)}} - NIF_{\text{(baseflow in rainstorm)}}, \quad (2)$$

16 where $NIF_{\text{(rainstorm)}}$ is the measured NIF flux in the stream water during rainstorms,
17 including both the baseflow and quick-flow components; and $NIF_{\text{(baseflow in rainstorm)}}$ is
18 the NIF flux derived from the baseflow component during rainstorms, of which the
19 calculation will be presented in Section 5.3.1. (i.e., Eq. 5). In Eq. 2, ΔNIF would be
20 derived mostly from relatively shallow soils with high FAM (Fig. 4a) resulted from
21 shallow subsurface and near-surface water as biomatflow.

22 On the basis of Eqs. 1 and 2, with combining the other Eqs. noted in the sections of
23 5.2.1 and 5.3.1 (i.e., Eqs. 3 to 6), NIF dynamics in stream water and bonding

1 possibility of the major metals with FAM and/or nano-scale phyllosilicates such as clay
2 minerals will be discussed.

3

4 **4. Results**

5 *4.1. FAM distribution in soil*

6 The concentrations of water extractable FAM and DOC in soils in the coniferous
7 and deciduous catchments are shown in Fig. 4. Regardless of the slope position, FAM
8 (Fig. 4a) in the coniferous catchment occurred mainly in the shallow part of soil above
9 50 cm depth. In the deciduous catchment, higher FAM was measured at the mid and
10 lower slope positions than at the ridge top, even below 50 cm depth. The distribution
11 of DOC (Fig. 4b) was similar to that of FAM though the concentrations below 50 cm
12 depth ranged between a few mg L⁻¹ and 10 mg L⁻¹ especially in the coniferous
13 catchment.

14

15 *4.2. Major metals and Si in stream water during no rainfall periods*

16 We examined the changes in stream water chemistry and the proportion of NIF
17 relative to total elements in stream water (baseflow) during no rainfall periods from
18 2006 to 2007 (Fig. 5). The pH of the stream water ranged between 6.0 and 7.0 in both
19 catchments, however, that of the baseflow during the summer season tended to have
20 been 0.5 - 1.0 pH units higher than that during the winter season in both catchments.

21 Fig. 5 shows that, except for the concentrations of T-Ca and Ca²⁺ in the coniferous
22 catchment, which respectively ranged from 6 to 10 mg L⁻¹ and from 5 to 8 mg L⁻¹, the
23 concentrations of the total elements (T-Na, -Mg, and -Ca) and free ions (Na⁺, Mg²⁺,

1 and Ca^{2+}) were mostly below 5 mg L^{-1} in both catchments.

2 The concentrations of total elements were higher in the coniferous catchment than
3 in the deciduous catchment throughout the observation period. In addition, Fig. 6
4 shows that the flow chemistry in free ions showed the Ca-HCO_3 type in both
5 catchments. Although the concentration of free ions tended to rise through winter into
6 summer in both catchments, it was higher in the coniferous catchment than in the
7 deciduous catchment (i.e., mean concentrations were 1.4, 0.4, and 3.6 mg L^{-1} higher
8 for T-Na, T-Mg, and T-Ca, respectively; and 1.4, 0.4, and 3.6 mg L^{-1} higher for Na^+ ,
9 Mg^{2+} , and Ca^{2+} , respectively).

10 The mean Si concentration was 6.22 mg L^{-1} in the coniferous catchment and 4.88
11 mg L^{-1} in the deciduous catchment, 1.3 times higher on average in the coniferous
12 catchment than in the deciduous catchment. However, because the stream discharge
13 (i.e., baseflow in this case) in the coniferous catchment was about 0.7 - 0.8 times less
14 on average than that in the deciduous catchment, the specific fluxes of Si were
15 eventually similar in the two catchments throughout the observation period.

16 The proportion of NIF relative to total element ranged from 0.6% to 40.0% in the
17 coniferous catchment and from 7.3% to 58.0% in the deciduous catchments (refer to no
18 rainfall periods in Table 1a). They tended to become lower (7% to 19 % on average)
19 after May 2007 except during rainfall periods and NIF-Ca in the deciduous catchment
20 (Fig. 5), resulting from the rise of concentration of free ions relative to total elements
21 through winter into summer as shown in Fig. 6. In addition, the proportions of NIF in
22 the coniferous catchment (Table 1a) were 13.6% (in NIF-Na), 11.3% (in NIF-Mg), and
23 25.4% (in NIF-Ca) lower on average, with a significant difference under all $p < 0.0001$

1 and $n = 30$, than those in the deciduous catchment. Thus in the stream water during no
2 rainfall periods, although the concentrations of total elements and free ions were
3 mostly higher in the coniferous catchment than in the deciduous catchment, the
4 proportion of NIF in the stream water were less in the coniferous catchment than in the
5 deciduous catchment.

6

7 *4.3. Major metals, Si, and FAM in stream water during rainstorms*

8 We next examined the changes in stream water chemistry and the proportion of
9 NIF relative to total elements during rainstorms. The changes in pH of low and peak
10 stream discharge during the rainstorms in the two catchments were from 6.0 to 7.5. The
11 result in Storm 2 (a typhoon storm with total rainfall of 117 mm) is specifically shown
12 in Fig. 7. In the coniferous catchment, stream discharge initially peaked at 2.5 L s^{-1}
13 ha^{-1} at 00:00 LT on 15 July and peaked a second time at $8.5 \text{ L s}^{-1} \text{ ha}^{-1}$ at 12:30 LT on
14 15 July. In contrast, in the deciduous catchment, the stream discharge during the
15 morning of 15 July was about $1 \text{ L s}^{-1} \text{ ha}^{-1}$ without clear peak, but a peak of 8.0 L s^{-1}
16 ha^{-1} , similar in amount to the second peak in the coniferous catchment, occurred at
17 12:30 LT on 15 July.

18 T-Na and Na^+ concentrations in the coniferous catchment were relatively about 1.0
19 and 1.4 mg L^{-1} higher on average than those in the deciduous catchment. However,
20 T-Mg and Mg^{2+} concentrations were almost the same in both catchments: same on
21 average in T-Mg and 0.2 mg L^{-1} higher on average in Mg^{2+} in the coniferous catchment
22 than in the deciduous catchment. In contrast, T-Ca concentrations were similar in both
23 catchments (0.18 mg L^{-1} lower on average in the coniferous catchment than in the

1 deciduous catchment), whereas Ca^{2+} concentrations in the coniferous catchment were
2 about 1.7 mg L^{-1} higher on average than those in the deciduous catchment. These facts
3 provide that the NIF-Ca concentration (ratio) in the coniferous catchment was
4 consequently lower than that in the deciduous catchment.

5 The mean Si concentration in the coniferous catchment was 6.09 mg L^{-1} and that in
6 the deciduous catchment was 3.50 mg L^{-1} , 2.59 mg L^{-1} higher in the coniferous
7 catchment than in the deciduous catchment. This difference in concentrations resulted
8 in higher Si fluxes in the coniferous catchment than in the deciduous catchment during
9 Storm 2, because stream discharge was similar for the two catchments.

10 The maximum apparent concentrations of FAM in Storm 2 were 12 QSU in the
11 coniferous catchment and 15 QSU in the deciduous catchment, whereas they were 8.6
12 QSU and 8.7 QSU in Storm 1, respectively. Because the maximum stream discharge in
13 Storm 2 was similar in both catchments, the apparent flux of FAM in the coniferous
14 catchment was lower than that in the deciduous catchment.

15 The proportion of NIF relative to total elements during the rainstorms (refer to
16 Table 1b and 1c) ranged, in the coniferous catchment, from 5.9% to 41.8% in Storm 1
17 and from 1.9% to 72.4% in Storm 2. Contrastively, that in the deciduous catchment
18 ranged from 23.4% to 49.6% in Storm 1 and 24.5% to 60.5% in Storm 2. The mean
19 proportions of NIF in the coniferous catchment were 10.2% to 26.3% lower in Storm 1
20 and 12.2% to 19.1% lower in Storm 2 than those in the deciduous catchment, with a
21 significant difference under all $p < 0.0001$, $n = 22$ (Storm 1), $n = 29$ (Storm 2), also
22 similar to the results during no rainfall periods (Table 1a).

23 The changes in chemical concentrations of major ions in stream water during the

1 rainstorms (piper trilinear diagrams; Piper, 1944) are shown in Fig. 8. On the basis of
2 the combination of the rate of free ions expressed in Fig. 8, the flow modes
3 contributing stream water are categorized as following 4 types:

4 I : $\text{CaSO}_4\text{-CaCl}_2$ type (non-carbonate hardness) originated from surface flow to soil
5 water

6 II: $\text{Ca}(\text{HCO}_3)_2$ type (carbonate hardness) originated from soil water to shallow
7 groundwater

8 III: NaHCO_3 type (carbonate alkali) originated from stagnated deep groundwater

9 VI: $\text{NaSO}_4\text{-NaCl}$ type (non-carbonate alkali) originated from sea water or hot spring

10 In the coniferous catchment, soil water and shallow ground water (Type I to II)
11 seems to have been the main component of stream water during both rainstorms.
12 However, the combination of major ions in Storm 2 (Fig. 8b left) changed
13 discontinuously from the boundary of Type I and II to the edge of Type I during the
14 rising limb of the hydrograph. In contrast, in the deciduous catchment, the combination
15 distributed between the middle of Type I and Type II in both rainstorms and did not
16 change discontinuously as indicated in the coniferous catchment.

17

18 **5. Discussion**

19 *5.1. Mode of non-free ionic fraction in stream water*

20 We hardly detected K^+ in the stream water of either catchment not only during no
21 rainfall periods but also rainstorms, although it was detected in the shallow soil,
22 indicating that K^+ moved scarcely from the soil to the streams, possibly because most
23 K^+ was absorbed directly from the soils by the trees and other plants or was

1 irreversibly adsorbed by such clay minerals as vermiculite. Thus K is excludable from
2 the following discussion on the mode of NIF.

3 NIF-Na, -Mg, and -Ca in natural water can be classified into five chemical forms,
4 which are listed in Table 2. Representative candidate free polyatomic ions in the
5 aquatic environment are MgF^+ , NaSO_4^- , and MgPO_4^- , but these ions are mainly found
6 in seawater (Howard, 1998); these free polyatomic ions mainly dissociate in fresh
7 water or are absent in soils so that the Na and Mg occur as free ions in fresh water
8 streams.

9 Except for calcium phosphate which is insoluble in fresh water (Guardado *et al.*,
10 2007), the lowest fresh water solubility among inorganic compounds (hydrate, sulfate,
11 carbonate, nitrate, and phosphate) of Na, Mg, and Ca, at 20 °C and pH 7.0, is that of
12 NaHCO_3 (1.04 mol L^{-1}), Mg(OH)_2 ($1.65 \times 10^{-4} \text{ mol L}^{-1}$), and CaCO_3 ($1.50 \times 10^{-4} \text{ mol}$
13 L^{-1}). Therefore, in fresh water, the lowest saturated concentrations of Na^+ , Mg^{2+} , and
14 Ca^{2+} derived from the dissolution of NaHCO_3 , Mg(OH)_2 , and CaCO_3 are calculated as
15 about 23.9 g L^{-1} , 4.0 mg L^{-1} , and 6.0 mg L^{-1} , respectively. These calculated values,
16 except for Ca^{2+} in the coniferous catchment from April to June 2007 (7.5 mg L^{-1} at
17 most in Fig. 5), were enough higher (i.e., unsaturated in free ions) than the maximum
18 concentrations measured in the stream water. Moreover, the actual concentrations of
19 Na^+ , Mg^{2+} , and Ca^{2+} in the stream water at pH 6.0 to 8.0 could be lower than those
20 measured by ion-analyzer using an eluant with pH 3.0. Thus, the stream water were
21 hardly saturated by Na^+ , Mg^{2+} , and Ca^{2+} , and Na, Mg, and Ca derived from the
22 compounds (hydrate, sulfate, carbonate, nitrate, and phosphate) ought to occur mostly
23 as free ions in stream water with pH from 6.0 to 8.0. In addition, cations do not adsorb

1 onto Al- or Fe-hydroxides in freshwater environments with pH < 8.0, such as the pH of
2 the stream water in these headwater catchments, because they are electropositive
3 (AlOH_2^+ or FeOH_2^+).

4 Thus, in our study, except for calcium phosphate, the occurrence of NIF-Na, -Mg,
5 and -Ca in hydration and inorganic compounds scarcely needs consideration. Most
6 NIF-Na, -Mg, and -Ca, therefore, were probably bonded to organic materials or
7 nano-scale phyllosilicates such as clay minerals (e.g., Takahashi *et al.*, 1999). That is,
8 exploring the bonding of these materials is likely to be key to elucidate the transport of
9 Na, Mg, and Ca in the stream water of these headwater catchments.

11 5.2. NIF dynamics in stream water

12 5.2.1. Relationship between stream discharge, Si, and NIF during no rainfall periods

13 In contrast to the inverse relationship observed between stream discharge and the
14 Si concentration during Storm 2 (i.e., dilution of Si by quick-flow component with a
15 low concentration of Si; Fig. 7), the Si concentration during no rainfall periods (Fig. 5)
16 increased as the stream discharge (i.e., baseflow in this case) increased. This result
17 suggests that changes in the Si production are strongly controlled by the baseflow
18 component of stream water.

19 The correlation coefficients (r) and slopes of the relationships between stream
20 discharge and the Si flux and between the fluxes of Si and NIF during no rainfall
21 periods are shown in Table 3. In the correlation equations, the y-intercept was set to
22 zero, because discharge of Si does not occur without stream discharge, and no NIF
23 production occurs without stream discharge (i.e., without discharge of Si). Against the

1 wash load flux as mentioned in Section 3.5. (i.e., not always necessarily represent good
 2 correlations), the relationships are expressed as follows under the good correlation ($r =$
 3 0.99 , $p < 0.0001$, $n = 54$) and the different coefficient “ a ”:

$$4 \quad Si_{(baseflow)} = a Q_{(baseflow)}, \quad (3)$$

5 where $Si_{(baseflow)}$ is the Si flux in the stream water during the no rainfall periods, the
 6 coefficient a represents the average Si concentration, and $Q_{(baseflow)}$ is the stream
 7 discharge during no rainfall periods (i.e. baseflow). When the stream water was
 8 collected (and thus the Si concentration was measured), the maximum stream
 9 discharge was $3.19 \text{ L s}^{-1} \text{ ha}^{-1}$ and $5.06 \text{ L s}^{-1} \text{ ha}^{-1}$ in the coniferous and deciduous
 10 catchments, respectively. Thus, in the strict sense, when we calculate the baseflow
 11 component of the stream water during the rainstorms ($Q_{(baseflow \text{ in rainstorm})}$ in Eqs. 1, 5,
 12 and 6, Eq 3 is available to the baseflow component below $3.19 \text{ L s}^{-1} \text{ ha}^{-1}$ and 5.06 L s^{-1}
 13 ha^{-1} in the coniferous and deciduous catchments, respectively.

14 In contrast, the correlations between the fluxes of Si and NIF in the stream water
 15 during the no rainfall periods were also high ($r > 0.92$, $p < 0.0001$, $n = 30$) in both
 16 catchments (Table 3). The NIF flux during no rainfall periods ($NIF_{(baseflow)}$) can
 17 therefore be expressed by $Si_{(baseflow)}$ and Eq. 3 as follows:

$$18 \quad NIF_{(baseflow)} = b Si_{(baseflow)} = ab Q_{(baseflow)}. \quad (4)$$

19 Eq 4 indicates that the ratio of the NIF discharge to the Si discharge in stream water
 20 was approximately constant during the no rainfall periods. The values of the coefficient
 21 b are shown in Table 3. Even though Si is naturally included in nano-scale
 22 phyllosilicates, the relationship between Si and NIF expressed by Eq. 4 does not
 23 necessarily explain the transport form of the major metals bonded to nano-scale

1 phyllosilicates (e.g., substitution in lattices, invasion into the interlayer spaces, or ion
2 adsorption on the surfaces) because of occurring simple silicic acid ions. However, the
3 high correlation between Si and NIF during the no rainfall periods may have some
4 implications regarding the transport of Si and NIF, because during the rainstorms the Si
5 flux (Table 4) did not usually correlate strongly (i.e., $r < 0.80$) with NIF fluxes. In
6 particular, during Storm 1 (Table 4a), the correlations tended to be weak except for
7 NIF-Mg in the coniferous catchment, even when the baseflow component accounted
8 for most of the stream water (94% and 97% in the coniferous catchment and deciduous
9 catchment, respectively; Hirano *et al.*, 2009).

10

11 5.2.2. Relationship between stream discharge, Si, and NIF during rainstorms

12 The correlation coefficients (r) and slopes of the relationships between stream
13 discharge and the Si flux and between the fluxes of Si and NIF during rainstorms are
14 shown in Table 4.

15 The concentrations of Si in both catchments during the rainstorm decreased slightly
16 (a few mg L^{-1}) with an increase in stream discharge, indicating that the Si, originated in
17 the baseflow component in stream water, could have been diluted with the quick-flow
18 component containing the low Si concentration. This finding was consistent with those
19 from no rainfall periods, that is, more baseflow equates to higher Si concentration.

20 In both rainstorms, as well as the cases during the no rainfalls, the Si flux
21 correlated strongly ($r \geq 0.80$, $p < 0.0001$, $n = 24$ and 29 for the coniferous catchment
22 and deciduous catchment, respectively) with the stream discharge in both catchments.
23 The values of the coefficient a , however, were smaller in both catchments than its

1 value during the no rainfall periods (i.e., $a_{(\text{baseflow})} > a_{(\text{Storm 1})} > a_{(\text{Storm 2})}$). Therefore,
2 compared with the no rainfall periods, the rate of increase in the Si flux relative to that
3 in the stream water was lower during rainstorms, probably because the low Si
4 concentration of the quick-flow component of the stream water caused the Si
5 concentration in the total stream water to be relatively low. This finding suggests that,
6 as indicated in Section 5.2.1, the Si production during rainstorms is also strongly
7 controlled by Si in the baseflow component of the stream water.

8 The weak correlations ($r < 0.60$) between the fluxes of Si and NIF in some cases in
9 Table 4 suggest that, in contrast to no rainfall periods (Table 3), NIF was not
10 necessarily increased according to the increase in Si during rainstorms. Though the
11 ratio of NIF to Si reflects in part the different values of the b coefficient (Tables 3 and
12 4), the low correlations ($r < 0.60$, Table 4) may reflect either an increased or reduced
13 transport of NIF by the quick-flow component of the stream water in those cases. Thus,
14 in analyses of NIF transport during rainstorms, NIF should be classified according to
15 whether it derives from the baseflow or the quick-flow component of the stream water.
16 In contrast, as shown in the deciduous catchment in Table 4b, strong correlations ($r \geq$
17 0.80) may reveal that the flow components of the stream water did not implicate the
18 result in an increase in transport of Si and NIF.

19

20 5.3. Bonding possibility of Na, Mg, and Ca to FAM

21 5.3.1. Origin of FAM and NIF during rainstorms

22 Water extractable FAM tended to be stored in the shallower part of soil in the
23 coniferous catchment than in the deciduous catchment (Fig. 4 and Section 4.1.). In

1 contrast, soil porosity and saturation hydraulic conductivity are similar in both
2 catchments (Section 2.2.). These findings may imply that the hydrological process,
3 such as near-surface flow represented by biomatflow caused dominantly in the
4 coniferous catchment, may have controlled the downward movement of FAM.

5 In Storm 1, the apparent fluxes of FAM were mostly $\leq 2.0 \text{ QSU L s}^{-1} \text{ ha}^{-1}$ in both
6 catchments, and the maximum fluxes were $4.9 \text{ QSU L s}^{-1} \text{ ha}^{-1}$ and $6.4 \text{ QSU L s}^{-1} \text{ ha}^{-1}$
7 in the coniferous and deciduous catchments, respectively. In contrast, in Storm 2, the
8 apparent fluxes were $\geq 2.0 \text{ QSU L s}^{-1} \text{ ha}^{-1}$ in both catchments, and the maximum
9 fluxes were $53.3 \text{ QSU L s}^{-1} \text{ ha}^{-1}$ and $113.6 \text{ QSU L s}^{-1} \text{ ha}^{-1}$ in the coniferous and
10 deciduous catchments, respectively. Thus, the production of FAM was relatively small
11 during Storm 1, which was a small rainstorm that contributed only a small quick-flow
12 component to the total stream water (6% and 3% in the coniferous and deciduous
13 catchments, respectively; Hirano *et al.*, 2009) and consequently produced smaller
14 changes in the stream hydrograph than Storm 2. Thus, during rainstorms, most FAM
15 likely originated at relatively shallow depth in the soil as shown in Fig. 4a.

16 During the heavy rainstorm, both surface or near-surface water (i.e., quick-flow
17 component) and deep soil water was dominant in the coniferous catchment, whereas
18 both shallow and deep soil water was dominant in the deciduous catchment (Fig. 8).
19 Thus NIF and FAM production could be strongly controlled by the main flow
20 component of stream water, which was different between the coniferous and deciduous
21 catchment. Accordingly, when we analyze the relationship between FAM and NIF
22 during rainstorms, we should take into account whether the NIF in the stream water
23 derived primarily from the baseflow (deep soil) or quick-flow (near-surface or shallow

1 soil).

2 As indicated in Section 5.2.2, the Si production in the stream water during the
3 rainstorms seems to have been controlled mainly by Si included in the baseflow
4 component of the stream water (e.g., $a_{(\text{baseflow})} > a_{(\text{Storm 1})} > a_{(\text{Storm 2})}$ in Tables 3 and 4).
5 In addition, the maximum stream discharge during the no rainfall periods (i.e.,
6 maximum baseflow) were $3.19 \text{ L s}^{-1} \text{ ha}^{-1}$ and $5.06 \text{ L s}^{-1} \text{ ha}^{-1}$ in the coniferous and
7 deciduous catchments, respectively; corresponding to 0.6 and 1.3 times the maximum
8 baseflow component calculated for Storm 2 using Eq. 1. In other words, specifically in
9 the coniferous catchment, the maximum baseflow component during Storm 2 was more
10 than the baseflow during the no rainfall periods.

11 Accordingly, if we could extrapolate Eq. 4 for calculating the baseflow component
12 in the coniferous catchment during Storm 2, we can modify Eq. 4 for the two
13 catchments to express NIF originating from the baseflow component during the
14 rainstorms as follows:

$$15 \quad NIF_{(\text{baseflow in rainstorm})} = b Si_{(\text{baseflow in rainstorm})} = ab Q_{(\text{baseflow in rainstorm})}, \quad (5)$$

16 where $Si_{(\text{baseflow in rainstorm})}$ is the Si flux originating from the baseflow component of the
17 stream water during rainstorms. By combining Eqs. 1, 2, and 5, we obtain:

$$18 \quad \Delta NIF = NIF_{(\text{rainstorm})} - ab Q_{(\text{baseflow in rainstorm})} \\ 19 \quad = NIF_{(\text{rainstorm})} - ab Q_{(\text{rainstorm})} (1 - Q_{(\text{quick-flow})} / Q_{(\text{rainstorm})}). \quad (6)$$

20 Eq. 6 can thus be used to examine the relationship between NIF and FAM transported
21 in the stream water during rainstorms, which is mainly contained in the quick-flow
22 component passed through relatively shallow soil with a high FAM content (Fig. 4a).

23

1 5.3.2. Relationship between FAM, NIF, and Δ NIF during the rainstorms

2 The relationships between the fluxes of FAM, Si, NIF, and Δ NIF in the stream
3 water during rainstorms are shown in Table 5. In these correlations, the y-intercept (d)
4 is a non-zero value because NIF can be transported in the stream water during the no
5 rainfall periods, when the FAM flux is very small. In the case of Storm 1, we could not
6 determine whether the samples were collected on the rising or falling limb of the
7 hydrograph because the small rainfall amount produced only small changes in the
8 hydrograph, but in the case of Storm 2, we classified the samples according to when
9 they were collected. Thus, for the relatively weak correlations during Storm 2 ($r <$
10 0.70), we also calculated correlations for samples collected during the rising and
11 falling limbs of the hydrograph separately, and we show respectively those correlation
12 coefficients in parentheses in Table 5.

13 For Storm 1 (Table 5a), when the quick-flow component accounted for $\leq 6\%$ of
14 the stream water (Hirano *et al.*, 2009), we were unable to calculate correlation
15 coefficients for the FAM– Δ NIF-Ca relationship in the deciduous catchment, because
16 Δ NIF-Ca, calculated with Eq. 6, was generally 0 mg s^{-1} . Except for FAM versus
17 Δ NIF-Na in the deciduous catchment ($r = 0.86$, $p = 0.0009$, $n = 16$), the correlations
18 between FAM and Si, NIF, and Δ NIF were low ($r \leq 0.60$, $n = 22$ for the coniferous
19 catchment and $n = 16$ for the deciduous catchment). Thus, these low correlations likely
20 reflect the effect of the high proportion of baseflow component ($\geq 94\%$ of the stream
21 discharge; Hirano *et al.*, 2009) on the variation of FAM, derived mostly from the
22 shallow soil by the quick-flow component. These findings indicate that the transport of
23 NIF (or Δ NIF), that is, the bonding of Na, Mg, and Ca to FAM, hardly differed during

1 the small rainstorm from that during the no rainfall periods because of the relatively
2 small change in the quick-flow component of the stream water.

3 Takahashi *et al.* (1999) mentioned that the inorganic particles coated by humic
4 material (organic-inorganic complex) should have been regarded as important
5 particulate matter which shows different affinities to various ions from inorganic
6 particulates without the organic coating. In addition, they indicated that the colloidal
7 inorganic particles coated with humic substances were expected to be significantly
8 important in the environment. Thus, as well as the case during the no rainfalls, Na, Mg,
9 and Ca may have been bonded not only to FAM (Choppin and Shanbhag, 1981; Livens,
10 1991) but also to nano-scale phyllosilicates, organic acids of which the concentration
11 was implicitly indicated by our measurement in the ion-analyzer and the 3d-EEM,
12 other unknown materials such as citric, tartaric, lactic, and malic acids (Stevenson,
13 1994), or to else which is soluble in freshwater (Guardado *et al.*, 2007).

14 In Storm 2 (Table 5b), when the quick-flow component accounted for $\geq 30\%$ of the
15 stream discharge (Hirano *et al.*, 2009), the correlations between FAM and Δ NIF were
16 strong ($r \geq 0.81$, $p < 0.0001$, $n = 22$) in the coniferous catchment. This result supports
17 the interpretation that the quick-flow component during the large rainstorm in the
18 coniferous catchment, as shown at the edge of Type I in Fig. 8, may have transported
19 FAM and Δ NIF together from the relatively shallow soil layer. In contrast, in the
20 deciduous catchment, the strong correlations ($r \geq 0.83$, $p < 0.0001$, $n = 26$) of FAM
21 with both NIF and Δ NIF may have resulted from their derivation from soil water and
22 shallow groundwater (Fig. 8) that passed through the shallow to deep part of soil
23 containing higher FAM concentrations than those in the coniferous catchment (Fig.

1 4a).

2 Furthermore, FAM was also correlated more strongly with Si in the deciduous
3 catchment ($r = 0.87, p < 0.0001, n = 26$) than in the coniferous catchment ($r = 0.73, p <$
4 $0.0001, n = 26$). Together with the strong correlations between Si and NIF in the
5 deciduous catchment during Storm 2 (Table 4b), these findings may indicate the
6 interpretation that during heavy storms, Na, Mg, and Ca bond more likely to FAM and
7 also possibly to nano-scale phyllosilicates (organic–inorganic interactions as indicated
8 by Takahashi *et al.*, 1999) in the headwater stream of the deciduous catchment,
9 compared with the coniferous catchment. This thing may result from the interaction
10 between rainwater pathways in relatively deep part of soil compared to the
11 near-surface and FAM percolated into the deeper part of soil in the deciduous
12 catchment than in the coniferous catchment.

13

14 5.4. Effect of environmental factors on NIF production

15 Using a chemical model, Mantoura *et al.* (1978) showed that, in freshwater lakes,
16 compounds of Mg and Ca with sulfate ($-\text{SO}_4$) and carbonate ($-\text{CO}_3$) accounted for \leq
17 1% of their total concentrations, and it has since been assumed that most Na, Mg, and
18 Ca occurs as free ions in natural freshwater bodies. However, in our study (Figs. 5, 7,
19 and Table 1), NIF-Na, -Mg, and -Ca in stream water sometimes accounted for half or
20 more of the total concentrations of these elements. This was predominant in the
21 deciduous catchment, suggesting that NIF are not a negligible feature in freshwater
22 streams when the species of Na, Mg, and Ca are considered. Although the baseflow pH
23 was about 1.0 pH unit higher in the coniferous catchment than in the deciduous

1 catchment, the concentrations of free ions in the coniferous catchment were 1 to 4 mg
2 L⁻¹ higher than those in the deciduous catchment. In other words, increased production
3 of free ions due to the lower stream water pH never provides the lower concentrations
4 of free ions in the deciduous catchment (i.e., the higher proportion of NIF). In addition,
5 the flux of Na⁺, Mg²⁺, and Ca²⁺ to the forests were approximately the same in both
6 catchments (Section 3.2.). Therefore, the proportion of NIF must indicate that the
7 bonding of Na, Mg, and Ca between the soils and streams may account for the high
8 percentage of NIF in the stream water.

9 No human activities, such as thinning to reduce tree density or fertilizing, took
10 place in either catchment during the observation period. Furthermore, geographical,
11 topographical, geological, and climatic conditions were almost the same in both
12 catchments. Consequently, the differences in chemistry of Na, Mg, and Ca in the
13 stream water between the two catchments; specifically both in the higher free ion
14 concentration in the coniferous catchment than in the deciduous catchment (Fig. 6) and
15 in the proportion of NIF which was from 10% to 26% lower on average in the
16 coniferous catchment than in the deciduous catchment (Table 1); may primarily reflect
17 the effects of the different vegetation on the bonding of Na, Mg, and Ca.

18 In particular, Fig. 8 also emphasizes that the main flow pathways contributing to
19 stream water during Storm 2 could have been different in the coniferous and deciduous
20 catchments. The stream water (and thus FAM) in the coniferous catchment during
21 Storm 2 was likely originated in near-surface water and shallow groundwater, whereas
22 it was likely originated in soil water to shallow groundwater in the deciduous
23 catchment. The unique flow generation (biomatflow) from near-surface at the slope

1 during rainstorms in the coniferous catchment could be one of the causes to transport
2 much FAM to stream water and consequently to store little FAM in the deep part of
3 soil (Fig. 4a).

4

5 **6. Conclusions**

6 We investigated the transport of major metals by measuring the concentrations of Na,
7 Mg, Ca, Si, and fulvic acid-like materials in two mountainous headwater streams in
8 Japan, one surrounded by coniferous and the other by deciduous forest. The
9 conclusions are not necessarily being generalized because only data sets for two
10 rainstorms and two headwater catchments were prepared in the presented study.
11 However, together with the data during no rainstorms, the following outlines can be
12 mentioned for the transport and bonding possibility of major metals caused in the
13 headwater streams by differing vegetation:

14 (1) The proportion of non-free ionic fractions to total metal elements ranged from 0%
15 up to 70% in two streams during both no rainfall period and rainstorms and was
16 lower in the coniferous catchment than in the deciduous catchment, though most Na,
17 Mg, and Ca has previously been assumed as free ions in natural freshwater
18 environments.

19 (2) In the stream water during no rainfall periods, as well as during small rainstorms,
20 some non-free ionic fractions in two streams were probably transported in bonding to
21 unknown materials including nano-scale phyllosilicates (clay minerals), organic acids,
22 or else.

23 (3) During heavy rainstorms, non-free ionic fractions in two streams could have mostly

1 been in bonding to humic substances and possibly to nano-scale phyllosilicates;
2 which was dominantly enhanced in the deciduous catchment. This finding could have
3 been due to the difference in flow pathways contributing to stream water: both surface
4 or near-surface water (i.e., overland flow or biomatflow,) and shallow groundwater
5 was dominant in the coniferous catchment, whereas both soil water (i.e., subsurface
6 flow) and shallow groundwater was dominant in the deciduous catchment.

7 (4) Above findings imply that the environmental controls, such as implication among
8 the differences in rainstorm size (i.e., weather condition), the distribution and content
9 of humic substances in soils (i.e., vegetational condition), and the resultant rainwater
10 flow pathways contributing to stream water (i.e., hydrological condition), provided
11 the variable organic-inorganic interactions in the headwater streams where
12 geographical, topographic, geological, and climatic conditions are similar.

13 Headwaters are the most elementary topographic unit to examine catchment
14 hydrology. Consequently, the data on bonding of major metals, linked with the effect
15 of environmental controls in headwater streams, should be fundamental and useful to
16 understand the chemical species of major metals in the complicated and confusing
17 nature as stream catchments. It is hoped that the accumulation of data sets based on
18 field observations will improve the understanding of chemical species of major metals
19 which was believed to occur mostly as free ions in freshwater environments.

20

21 **Acknowledgement**

22 We thank the Rijin-kai Association, the owner of the Nariki catchment, for
23 permission to use the catchments for our observations. The present study was partially

1 funded by a grant for environmental research from the Sumitomo Foundation, Japan
2 (No. 083295).

3

4 **References**

- 5 Artinger, R., Bukau, G., Geyer, S., Fritz, P., Wolf, M., Kim, J.I., 2000.
6 Characterization of groundwater humic substances: Influence of sedimentary
7 organic carbon. *Appl. Geochem.* 15, 97-116.
- 8 Choppin, G.R., Shanbhag, P.M., 1981. Binding of calcium by humic acid. *J. Inorg.*
9 *Nucl. Chem.* 43, 921-922.
- 10 Coble, P., Schultz, C., Mopper, K., 1993. Fluorescence contouring analysis of DOC
11 intercalibration experiment samples: a comparison of techniques. *Mar. Chem.* 41,
12 173–178.
- 13 Coble, P., 1996. Characterization of marine and terrestrial DOM in seawater using
14 excitation-emission matrix spectroscopy. *Mar. Chem.* 51, 325-346.
- 15 Coble, P., Del Castillo, C., Avril, B., 1998. Distribution and optical properties of
16 CDOM in the Arabian Sea during the 1995 SW monsoon. *Deep-Sea Res. II* 45,
17 2195–2223.
- 18 Fellman, J.B., Hood, E., Robert G. M. Spencer, R.G.M., 2010. Fluorescence
19 spectroscopy opens new windows into dissolved organic matter dynamics in
20 freshwater ecosystems: A review. *Limnol. Oceanogr.* 55, 2452–2462.
- 21 Guardado, I., Urrutia, O., Garcia-Mina, J.M., 2007. Size distribution, complexing
22 capacity, and stability of phosphate-metal-humic complexes. *J. Agric. Food Chem.*
23 55, 408-413.

- 1 Hirano, T., Terajima, T., Nakamura, T., Aoki, F., Sakai, M., 2008. Surface and near-
2 surface runoff at slopes of the unmanaged coniferous forests and the natural
3 deciduous forests in Nariki nested catchment. *Trans. Jpn. Geomorph. Union.* 29,
4 255-280 (in Japanese with English abstract and captions).
- 5 Hirano, T., Terajima, T., Nakamura, T., Sakai, M., Aoki, F., Nanami, A., 2009. The
6 differences in the short-term runoff characteristics between the coniferous catchment
7 and the deciduous catchment, the effect of storm size on stormflow generation
8 processes of small forested catchments. *J. Jpn. Soc. Hydrol. Water Resour.* 22, 24-39
9 (in Japanese with English abstract and captions).
- 10 Howard, A.G., 1998. *Aquatic environmental chemistry, Oxford chemistry primers: 57,*
11 *p90, Oxford University Press.*
- 12 Inosato, H., Kanno, S., Shindo, S., Watanabe, K., 1980. *Local geology in Japan, Kanto*
13 *region. Asakura Publ. Co. Ltd., Tokyo (in Japanese).*
- 14 Lakowicz, J.R., 1999. *Principles of fluorescence spectroscopy, 2nd ed. Springer.*
- 15 Livens, F.R., 1991. Chemical reactions of metals with humic material. *Environ.*
16 *Pollution* 70, 183-208.
- 17 Majzik, A., Tombacz, E., 2007. Interaction between humic acid and montmorillonite in
18 the presence of calcium ions I. Interfacial and aqueous phase equilibria: adsorption
19 and complexation. *Organic Geochem.* 38, 1319-1329.
- 20 Malcolm, R.L., 1990. *Humic substances in soil and crop sciences: Selected readings*
21 *(eds. P. MacCarthy et al.), ASAI & SSSA, Madison, p13-35.*
- 22 Mantoura, R.F.C., Dickson, A., Riley, J.P., 1978. The complexation of metals with
23 humic materials in natural waters. *Estuarine Coastal Mar. Sci.* 6, 387-408.

- 1 McDowell W.H., 2003. Dissolved organic matter in soils—future directions and
2 unanswered questions. *Geoderma* 113, 179-186.
- 3 Mostofa, K., Honda, Y., Sakugawa, H., 2005. Dynamics and optical nature of
4 fluorescent dissolved organic matter in river waters in Hiroshima Prefecture, Japan.
5 *Geochem. J.* 39, 257–271.
- 6 Nogi, T., 2007. Temporal and spatial changes in rainfall amount and chemistry in an
7 unmanaged evergreen coniferous forest and natural deciduous broadleaf forest.
8 Graduation thesis of Chiba University, 29pp., in Japanese (unpublished).
- 9 Piper, A.M., 1944. A graphic procedure in the geochemical interpretation of
10 water-analyses. *Trans. Am. Geophys. Union* 25, 914-928.
- 11 Romkens, P.F., Bril, J., Salomons, W., 1996. Interaction between Ca²⁺ and dissolved
12 organic carbon: implication for metal mobilization. *Applied Geochem.* 11, 109-115.
- 13 Sidle, R.C., Hirano, T., Gomi, T., Terajima, T., 2007. Hortonian overland flow from
14 Japanese forest plantations – an aberration, the real thing, or something in between ?
15 *Hydrol. Process.* 21, 3237-3247.
- 16 Stevenson, F.J. 1994. *Humus Chemistry*. 2nd ed., 497pp., Wiley. New York.
- 17 Strellis, D.A., Hwang, H.H., Anderson, T.F., Landsberger, S. 1996. A comparative study
18 of IC, ICP-AES, and NAA measurements on chlorine, bromine, and sodium in
19 natural waters. *J. Radioanal. Nucl. Chem.* 211, 473-484.
- 20 Suzuki, Y., Nakaguchi, Y., Hiraki, K., Nagao, S., Kudo, M., Kimura, M., 1998.
21 Characteristics of the fluorescent substances in the Yodo River system by
22 three-dimensional excitation emission matrix spectroscopy. *Chikyu Kagaku* 32,
23 21–30.

- 1 Takahashi, Y., Minai, Y., Ambe, S., Makide, Y., Ambe, F., 1999. Comparison of
2 adsorption behavior of multiple inorganic ions on kaolinite and silica in the
3 presence of humic acid using the multi tracer technique. *Geochimica et*
4 *Cosmochimica Acta.* 63, 815-836.
- 5 Takahashi, Y., Kimura, T., Minai, Y., 2002. Direct observation of Cm (III) – fulvate
6 species on fulvic acid-montmorillonite hybrid by laser-induced fluorescence
7 spectroscopy. *Geochimica et Cosmochimica Acta.* 66, 1-12.
- 8 Terajima, T., Sakamoto, T., Nakai, Y., Kitamura, K., 1997. Suspended sediment
9 discharge in subsurface flow from the head hollow of a small forested watershed,
10 northern Japan. *Earth Sur. Proc. & Landf.* 22, 987-1000.
- 11 Terajima, T., Moriizumi, M., 2013. Temporal and spatial changes in dissolved organic
12 carbon concentration and fluorescence intensity of fulvic acid like materials in
13 mountainous headwater catchments. *J. Hydrol.* 479, 1-12.
- 14 Thurman, E.M., 1985. *Organic geochemistry of natural waters.* Martinus Nijhoff/Dr W.
15 Junk Publishers. Dordrecht, 497pp.
- 16 Tipping, E. 1993. Modeling ion binding by humic acids. *Physicochemical and*
17 *Engineering Aspects.* 73, 117-131.
- 18 Tokyo Prefectural Public Work Institute, 2002. *Geological map of the Okutama area.*
19 Tokyo Prefecture (in Japanese).
- 20 Wels, C., Cornett, R.I., Lazerte, B.D., 1991. Hydrograph separation: A comparison of
21 geochemical and isopycnic tracers. *J. Hydrol.* 122, 253-274.
- 22 Wilson, M.A., Tran, N.H., Milev, A.S., Kannangara, G.S.K., Volk, H., Lu, G.Q.M.,
23 2008. Nanomaterials in soils. *Geoderma* 146, 291-302.




Figure 1. Location and topographic maps of the Nariki catchment and headwater catchments. (a) Nariki catchment, (b) Coniferous catchment, (c) Deciduous catchment. Contour intervals of the Nariki catchment (a) and headwater catchments (b and c) are 10 and 5 m, respectively.

(a) Coniferous catchment

(b) Deciduous catchment

Tree canopy



Forest floor



Figure 2. Tree canopies and understory vegetation in the headwater catchments in the summer of 2006.

Coniferous catchment

Deciduous catchment




Figure 3 Typical EEM spectra for the different stream flow regimes. The arrows show the visible peaks in fluorescence intensity (FI) at around Ex/Em: 340/440 during the rainstorms. Contour intervals of FI are 1.0.




Figure 4. Distribution of water extractable fulvic acid-like materials (FAM) and dissolved organic carbon (DOC) in soils.





Figure 5. Concentrations of major metals (Na, Mg, and Ca) and silicon (Si), and the proportion of non-free ionic fractions (NIF) relative to total elements in stream water during no rainfall periods from 2006 to 2007. N.D. in the hyetograph shows no data

(a) Coniferous catchment



(b) Deciduous catchment




Figure 6. Seasonal changes in major ions in stream water during no rainfall periods, which were derived from the arithmetic mean of the concentration data during two months





Figure 7. Concentrations of major metals (Na, Mg, and Ca) and silicon (Si), peak fluorescence intensity (F -FAM) and the proportion of non-free ionic fractions (NIF) relative to total elements in stream water during a typhoon storm in 2007 (Storm 2).

(a) Storm 1 (26-27 Sep., 2006)



(b) Storm 2 (14-16 Jul., 2007)




Figure 8. Piper-trilinear diagrams of stream water chemistry (compositional changes in major ions) during the rainstorms.

- I : CaSO_4 - CaCl_2 type (non-carbonate hardness) originated from surface flow to soil water.
- II : $\text{Ca}(\text{HCO}_3)_2$ type (carbonate hardness) originated from soil water to shallow groundwater.
- III : NaHCO_3 type (carbonate alkali) originated from stagnated deep groundwater.
- IV : NaSO_4 - NaCl type (non-carbonate alkali) originated from sea water or hot spring.

Table 1. Proportion of non-free ionic fractions (NIF) relevant to total elements in the stream water (unit : %)

(a) No rainfall periods		Coniferous catchment	Deciduous catchment
NIF-Na	Range	4.4~25.6	7.3~50.3
	Average	13.0	26.6
NIF-Mg	Range	0.6~35.6	10.7~46.3
	Average	17.7	29.0
NIF-Ca	Range	6.2~40.0	32.5~58.0
	Average	19.6	45.0
(b) Storm 1		Coniferous catchment	Deciduous catchments
NIF-Na	Range	16.5~33.5	25.1~43.5
	Average	24.8	35.0
NIF-Mg	Range	25.0~41.8	33.9~48.5
	Average	31.8	42.0
NIF-Ca	Range	5.9~29.1	23.4~49.6
	Average	14.2	40.5
(c) Storm 2		Coniferous catchment	Deciduous catchment
NIF-Na	Range	3.3~38.2	24.5~51.3
	Average	16.4	33.9
NIF-Mg	Range	24.5~72.4	32.6~59.8
	Average	34.6	46.8
NIF-Ca	Range	1.9~72.1	34.3~60.5
	Average	28.0	47.1

Table 2. Non-free ionic forms of sodium (Na), magnesium (Mg), and calcium (Ca), and the possible bonding partner or ligand group in aquatic environment.

Non-free ionic forms of Na, Mg, and Ca	Possible bonding partner or ligand group
◆ Inorganic materials	
Hydration	⇒ H ₂ O, OH ⁻
Adsorption / Substitution in phyllosilicates	⇒ Si, Al
Inorganic compounds	⇒ F ⁻ , SO ₄ ²⁻ , CO ₃ ²⁻ , NO ₃ ⁻ , PO ₄ ³⁻ Al- or Fe- hydroxide, etc
◆ Organic materials	
Adsorption on humic substances	⇒ Carboxyl group : COO ⁻ Phenolic group : Ph-OH Amino group : -NH ₂ Imino group : -NH
Complex of organic acids	⇒ Oxalate : C ₂ O ₄ ²⁻ Formic acid : HCOO ⁻ Pyrrole (Chlorophyll) : C ₄ H ₅ N- etc: -NH ₂ , -NH

Table 3. Linear correlation coefficients (r) of the relationship between stream discharge (Q), the flux of Si, and non-free ionic fractions (NIF) during no rainfall periods. The y-intercept of the correlation line was on the origin because Si does not occur without stream discharge, also implying no NIF production without baseflow discharge. Thus, the coefficient a and b are relatively expressed as $y=ax$ where y is Si and x is Q and $y=bx$ where y is NIF and x is Si.

x	y	<u>Coniferous catchment</u>		<u>Deciduous catchment</u>	
		r	a or b	r	a or b
Q vs.	Si	0.99	6.84	0.99	4.93
	NIF-Na	0.93	0.10	0.93	0.21
Si vs.	NIF-Mg	0.93	0.05	0.92	0.08
	NIF-Ca	0.96	0.34	0.96	0.50

Table 4. Linear correlation coefficients (r) of the relationship between stream discharge (Q) and the flux of Si, and non-free ionic fractions (NIF) during the rainfalls. The coefficient a and b are same as those in Table 3. The r -values which exceeded 0.80 (strong correlation) are expressed by the bold style.

(a) Storm 1		<u>Coniferous catchment</u>		<u>Deciduous catchment</u>	
x	y	r	a or b	r	a or b
Q vs.	Si	0.90	6.01	0.80	3.89
	NIF-Na	0.79	0.20	0.22	0.25
Si vs.	NIF-Mg	0.89	0.08	0.69	0.10
	NIF-Ca	0.54	0.15	0.58	0.30

(b) Storm 2		<u>Coniferous catchment</u>		<u>Deciduous catchment</u>	
x	y	r	a or b	r	a or b
Q vs.	Si	0.99	5.82	0.99	2.96
	NIF-Na	0.40	0.06	0.97	0.25
Si vs.	NIF-Mg	0.71	0.08	0.92	0.28
	NIF-Ca	0.59	0.30	0.87	1.32

Table 5. Linear correlation coefficients (r) of the relationship between the fluxes of fulvic acid like materials (FAM), Si, non-free ionic fractions (NIF), and non-free ionic fractions originated in the baseflow component of stream water (Δ NIF) during the rainstorms; the coefficient c and constant d which are expressed as $y=cx+d$, where y is the Si or NIF and x is FAM. The r -values in the parentheses under $r < 0.70$ in Storm 2 show the linear correlation coefficients during the rising (left) and falling limb (right) of the hydrograph, respectively, and they which exceeded 0.80 (strong correlation) are expressed by the bold style. We were impossible to obtain the r -values between FAM and Δ NIF-Ca in the deciduous catchment in Storm 1, because Δ NIF-Ca, calculated from Equation 6, was mostly 0 mg s⁻¹.

(a) Storm 1		Coniferous catchment			Deciduous catchment		
x	y	r	c	d	r	c	d
FAM vs.	Si	0.26	0.11	3.35	0.51	0.09	2.38
	NIF-Na	0.31	0.04	0.60	0.44	0.04	0.56
	NIF-Mg	0.26	0.01	0.27	0.54	0.02	0.21
	NIF-Ca	0.14	0.02	0.48	0.60	0.11	0.52
	Δ NIF-Na	0.31	0.03	0.23	0.86	0.04	-0.03
	Δ NIF-Mg	0.26	0.01	0.25	0.57	0.01	0.01
	Δ NIF-Ca	0.13	0.02	0.35	—	—	—
(b) Storm 2		Coniferous catchment			Deciduous catchment		
x	y	r	c	d	r	c	d
FAM vs.	Si	0.73	0.89	6.69	0.87	0.26	2.69
	NIF-Na	0.20 (0.28, -0.52)	0.02	1.42	0.83	0.06	1.05
	NIF-Mg	0.65 (0.68, 0.62)	0.06	0.97	0.97	0.09	0.25
	NIF-Ca	0.50 (0.50, 0.52)	0.17	4.38	0.98	0.42	0.97
	Δ NIF-Na	0.88	0.15	-0.32	0.53 (0.92 , -0.31)	0.01	0.45
	Δ NIF-Mg	0.81	0.06	0.59	0.98	0.07	0.01
	Δ NIF-Ca	0.87	0.15	0.52	0.99	0.29	-0.44