

# Chloroplast-mediated regulation of CO<sub>2</sub>-concentrating mechanism by Ca<sup>2+</sup>-binding protein CAS in the green alga *Chlamydomonas reinhardtii*

Lianyong Wang<sup>a</sup>, Takashi Yamano<sup>a</sup>, Shunsuke Takane<sup>a</sup>, Yuki Niikawa<sup>a</sup>, Chihana Toyokawa<sup>a</sup>, Shin-ichiro Ozawa<sup>b</sup>, Ryutaro Tokutsu<sup>c</sup>, Yuichiro Takahashi<sup>b</sup>, Jun Minagawa<sup>c</sup>, Yu Kanesaki<sup>d</sup>, Hirofumi Yoshikawa<sup>e</sup>, and Hideya Fukuzawa<sup>a,1</sup>

<sup>a</sup>Graduate School of Biostudies, Kyoto University, Kyoto 606-8502, Japan; <sup>b</sup>Research Institute for Interdisciplinary Science, Okayama University, Okayama 700-8530, Japan; <sup>c</sup>Division of Environmental Photobiology, National Institute for Basic Biology, Okazaki 444-8585, Japan; <sup>d</sup>NODAI Genome Research Center, Tokyo University of Agriculture, Tokyo 156-8502, Japan; and <sup>e</sup>Department of Bioscience, Tokyo University of Agriculture, Tokyo 156-8502, Japan

Edited by Julian I. Schroeder, University of California, San Diego, La Jolla, CA, and approved September 20, 2016 (received for review April 25, 2016)

Aquatic photosynthetic organisms, including the green alga *Chlamydomonas reinhardtii*, induce a CO<sub>2</sub>-concentrating mechanism (CCM) to maintain photosynthetic activity in CO<sub>2</sub>-limiting conditions by sensing environmental CO<sub>2</sub> and light availability. Previously, a novel high-CO<sub>2</sub>-requiring mutant, H82, defective in the induction of the CCM, was isolated. A homolog of calcium (Ca<sup>2+</sup>)-binding protein CAS, originally found in *Arabidopsis thaliana*, was disrupted in H82 cells. Although *Arabidopsis* CAS is reported to be associated with stomatal closure or immune responses via a chloroplast-mediated retrograde signal, the relationship between a Ca<sup>2+</sup> signal and the CCM associated with the function of CAS in an aquatic environment is still unclear. In this study, the introduction of an intact CAS gene into H82 cells restored photosynthetic affinity for inorganic carbon, and RNA-seq analyses revealed that CAS could function in maintaining the expression levels of nuclear-encoded CO<sub>2</sub>-limiting-inducible genes, including the HCO<sub>3</sub><sup>-</sup> transporters high-light activated 3 (*HLA3*) and low-CO<sub>2</sub>-inducible gene A (*LCIA*). CAS changed its localization from dispersed across the thylakoid membrane in high-CO<sub>2</sub> conditions or in the dark to being associated with tubule-like structures in the pyrenoid in CO<sub>2</sub>-limiting conditions, along with a significant increase of the fluorescent signals of the Ca<sup>2+</sup> indicator in the pyrenoid. *Chlamydomonas* CAS had Ca<sup>2+</sup>-binding activity, and the perturbation of intracellular Ca<sup>2+</sup> homeostasis by a Ca<sup>2+</sup>-chelator or calmodulin antagonist impaired the accumulation of *HLA3* and *LCIA*. These results suggest that *Chlamydomonas* CAS is a Ca<sup>2+</sup>-mediated regulator of CCM-related genes via a retrograde signal from the pyrenoid in the chloroplast to the nucleus.

acclimation | bicarbonate transporter | calcium signaling | photosynthesis | pyrenoid

Carbon dioxide (CO<sub>2</sub>) is a key environmental signal for physiological responses in many organisms (1). For photosynthetic organisms, CO<sub>2</sub> is essential for survival. In vascular plants, guard cells in leaves control the opening and closure of stomata in response to environmental CO<sub>2</sub> concentrations, with these events controlled by protein kinase HT1 (2) and carbonic anhydrase (3). In aquatic conditions, the CO<sub>2</sub> diffusion rate is ~10,000-fold lower compared with that in air (4). Therefore, aquatic photosynthetic organisms, including microalgae, are frequently exposed to limiting-CO<sub>2</sub> stress. To acclimate to this stress, most microalgae possess a CO<sub>2</sub>-concentrating mechanism (CCM) to increase CO<sub>2</sub> concentrations around the CO<sub>2</sub>-fixation enzyme ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) and to maintain adequate photosynthetic efficiency.

The eukaryotic CCM has been studied in the model green alga *Chlamydomonas reinhardtii* (5). CCM1/CIA5 was identified as a zinc-finger-type regulatory factor for the induction of most limiting-CO<sub>2</sub>-induced genes, including *HLA3* (high-light activated 3), *LCIA* (low-CO<sub>2</sub>-inducible gene A), and *LCIB* (low-CO<sub>2</sub>-inducible gene B) (6–9). *HLA3* is an ATP-binding cassette transporter localized to the plasma membrane and associated with HCO<sub>3</sub><sup>-</sup> transport from the

outside of cells into the cytosol (10–13). *LCIA* is a possible anion channel localized to the chloroplast envelope and is associated with inorganic carbon (Ci) (CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup>) uptake into the chloroplast stroma in cooperation with *HLA3* (12, 14). *LCIB* is a chloroplast soluble protein whose localization is associated with distinct CO<sub>2</sub>-acclimation states, including high-CO<sub>2</sub> (HC) (5 to 0.5%), low-CO<sub>2</sub> (LC) (0.03 to 0.5%), and very-low-CO<sub>2</sub> (VLC) (<0.03%) (15). In HC and LC conditions, *LCIB* is dispersed throughout the chloroplast stroma and is essential for the survival in LC conditions (11, 16, 17). In contrast, in VLC conditions, *LCIB* is localized as a ring-like structure in the vicinity of the pyrenoid (14, 17), where Rubisco is enriched for CO<sub>2</sub>-fixation. Although the function of *LCIB* in each CO<sub>2</sub>-acclimation state remains to be elucidated, it is proposed that *LCIB* functions not only in LC conditions but also in VLC conditions for CO<sub>2</sub> uptake (14).

In addition to CO<sub>2</sub>, Ca<sup>2+</sup> also plays a role in the regulation of photosynthesis (18) and could mediate CO<sub>2</sub> signal transduction (19). As a molecular component related to the Ca<sup>2+</sup> signal, a thylakoid Ca<sup>2+</sup>-binding protein, CAS, has been shown to mediate the transient elevation of cytosolic Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>cyt</sub>), as well as stromal Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>stro</sub>), in guard cells

## Significance

Ca<sup>2+</sup> and CO<sub>2</sub> are fundamental biological signaling molecules in microbes, animals, and plants. Although Ca<sup>2+</sup> was proposed to act as a second messenger in CO<sub>2</sub> signaling in guard cells of terrestrial plants, the role of Ca<sup>2+</sup> in CO<sub>2</sub> signal transduction pathways in aquatic photosynthetic organisms remains largely unknown. We show here that a chloroplast Ca<sup>2+</sup>-binding protein, CAS, changes its localization in response to environmental CO<sub>2</sub> conditions and regulates the expression of nuclear-encoded limiting-CO<sub>2</sub>-induced genes, including two key bicarbonate transporters. These findings led us to propose a model for the participation of Ca<sup>2+</sup> signals in chloroplast-regulated CO<sub>2</sub> signal transduction of aquatic photosynthetic organisms and help us to further understand the role of Ca<sup>2+</sup> in CO<sub>2</sub> signal transduction in eukaryotes.

Author contributions: L.W., T.Y., and H.F. designed research; L.W., T.Y., S.T., Y.N., C.T., S.-i.O., R.T., Y.K., and H.F. performed research; T.Y., Y.T., J.M., H.Y., and H.F. contributed new reagents/analytic tools; L.W., T.Y., Y.N., and H.F. analyzed data; and L.W., T.Y., and H.F. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

Data deposition: The RNA-seq raw data (Tables S2 and S3) in this paper have been deposited in the DNA Data Bank of Japan (DDBJ) Sequence Read Archive (DRA) (accession no. DRA004677).

<sup>1</sup>To whom correspondence should be addressed. Email: fukuzawa@lif.kyoto-u.ac.jp.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1606519113/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1606519113/-DCSupplemental).

of *Arabidopsis thaliana* and to regulate plant immune responses and stomatal closure (20–22). In *Chlamydomonas*, an ortholog of CAS (Cre12.g497300) was also detected in the thylakoid membrane fraction (23, 24). It was proposed that CAS is required for photoacclimation by regulating induction of light-harvesting complex stress-related protein 3 (LHCSR3) (25) and for forming a super complex for photosynthetic cyclic electron flow (CEF) in anaerobic conditions (26). From *Chlamydomonas* insertion mutant library screening experiments, we previously isolated a mutant strain, H82, in which a hygromycin resistance gene cassette was inserted into *CAS* (27). H82 cells showed decreased Ci affinity and did not accumulate HLA3 or LCIA in LC conditions.


In this study, we show the suborganellar localization of CAS in the chloroplast in vivo and its  $\text{Ca}^{2+}$ -binding activity in vitro. Furthermore, using complemented strains of H82, the link between CAS and regulation of the CCM is elucidated by examining the expression patterns of HLA3 and LCIA in response to  $\text{CO}_2$  and  $\text{Ca}^{2+}$  changes. From these results, we propose the existence of chloroplast-mediated regulation of the CCM by  $\text{Ca}^{2+}$ -binding protein CAS in parallel with regulation by CCM1/CIA5.

## Results

**Complementation of H82 Mutant Phenotype by CAS Gene.** To examine whether disruption of *CAS* by insertion of a hygromycin resistance gene cassette is responsible for the phenotype of H82, an 8.2-kb PCR-amplified genomic DNA fragment containing the WT *CAS* gene with its own predicted promoter was cotransformed into H82 cells with a paromomycin resistance gene cassette, *aphVIII* (Fig. S1A). From 774 paromomycin-resistant transformants, two strains showing similar growth rates to that of WT cells in LC conditions were isolated and designated as C-1 and C-2 (Fig. S1B). In parallel, 12 strains from 92 paromomycin resistance transformants generated by transforming a plasmid pTY2b-CAS to H82 exhibited recovered growth rate (Fig. S1C) and accumulated CAS (Fig. S1D) in LC conditions. Then, C-1 was used for further analyses. The HC-requiring phenotype of H82 cells was dependent on light intensity (Fig. S1E and *SI Results and Discussion*), and the addition of excess  $\text{Ca}^{2+}$  did not recover the retarded growth rate of the H82 cells (Fig. S1B and *SI Results and Discussion*).

Because it was reported that CEF activity increases in LC conditions possibly for increased ATP demand for ABC-type bicarbonate transporter HLA3 in CCM (12, 28) and that CAS is associated with the CEF complex formation in anaerobic conditions (26), it was possible that CAS-dependent CEF might contribute to CCM. However, CEF activity was not affected by the lack of CAS protein in the LC conditions at 120  $\mu\text{mol photons}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  that we examined (Fig. S1F and *SI Results and Discussion*). This result coincides with the previous finding that down-regulation of CAS did not impair CEF activity in aerobic conditions (26).

To evaluate induction of the CCM in LC conditions, photosynthetic affinities for Ci were evaluated by measuring the rates of photosynthetic oxygen ( $\text{O}_2$ ) evolution of WT, H82, and C-1 cells. At pH 7.8 (ratio of  $\text{HCO}_3^-:\text{CO}_2 = 28:1$ ), the Ci concentration required for half maximal velocity [ $K_{0.5}$  (Ci)] in C-1 cells was  $58 \pm 8 \mu\text{M}$ , which was similar to that of  $50 \pm 8 \mu\text{M}$  in WT cells and  $\sim 19$  times lower than that of  $1,087 \pm 113 \mu\text{M}$  in H82 cells (Fig. 1A and Table S1). Even at pH 6.2 ( $\text{HCO}_3^-:\text{CO}_2 = 0.7:1$ ) and pH 7.0 ( $\text{HCO}_3^-:\text{CO}_2 = 4.3:1$ ), the respective  $K_{0.5}$  (Ci) values of C-1 were 6.5-fold and 19.7-fold lower than that of H82 cells (Fig. S1G and Table S1). Because maximum rates of photosynthesis ( $V_{\text{max}}$ ) of the three strains WT, H82, and C-1 were similar with each other at each pH condition (Table S1), the decreased Ci affinity in H82 cells could be partly explained by a defect in Ci uptake activity. The accumulation and fixation of [ $^{14}\text{C}$ ]-labeled Ci in H82 cells were 0.13 mM and 0.34 nmol per



**Fig. 1.** Characterization of CAS insertion mutant H82 and its complemented strain C-1. (A) Inorganic carbon (Ci) affinity of WT, H82, C-1, and transgenic H82 cells containing inducible genes of HLA3 and LCIA (AH-1) grown in medium with  $\text{NH}_4^+$  or  $\text{NO}_3^-$  as nitrogen sources in low- $\text{CO}_2$  (LC) conditions for 12 h. Photosynthetic  $\text{O}_2$ -evolving activity was measured in external dissolved Ci concentrations at pH 7.8, and the Ci concentration required for half maximal velocity [ $K_{0.5}$  (Ci)] was calculated. Data in all experiments are mean values  $\pm$  SD from three biological replicates. \* $P < 0.01$  by Student's *t* test. (B and C) Accumulation (B) and fixation (C) of Ci in WT, H82, and C-1 cells. Cells were grown in LC conditions for 12 h, and the intracellular Ci concentration and carbon fixation were measured using a silicone-oil layer method. SIS, sorbitol impermeable space. (D) Accumulation of CAS, LCIA, HLA3, and LCIB in WT, H82, and C-1 cells. Cells were grown in high- $\text{CO}_2$  (HC) or LC conditions for 12 h. Histone H3 was used as a loading control.

microliter of sorbitol impermeable space (SIS) after 80 s of illumination, respectively, which was lower than 0.22 mM and 1.34 nmol  $\mu\text{L SIS}^{-1}$  of C-1 cells (Fig. 1B and C).

In addition to recovery of the accumulation of CAS and photosynthetic affinities for Ci in C-1, LC-induced accumulation of HLA3 and LCIA was also restored (Fig. 1D). To further evaluate the contribution of the defect in the accumulation of HLA3 and LCIA to the decreased Ci affinity in H82 cells, strain AH-1 was generated, in which two recombinant plasmids, pTY2b-LCIA and pTY2b-HLA3 (12), were introduced into H82 cells and in which HLA3 and LCIA could be induced by switching the nitrogen source from  $\text{NH}_4^+$  to  $\text{NO}_3^-$  (Fig. S1H). Although the  $K_{0.5}$  (Ci) value of  $1,087 \pm 113 \mu\text{M}$  in AH-1 cells was similar to that of  $1,087 \pm 113 \mu\text{M}$  in H82 cells in  $\text{NH}_4^+$  conditions, that of AH-1 cells decreased to  $430 \mu\text{M}$  from  $793 \pm 91 \mu\text{M}$  in H82 cells by expression of both HLA3 and LCIA in  $\text{NO}_3^-$  conditions (Fig. 1A and Table S1). However, the  $K_{0.5}$  (Ci) value in AH-1 cells was still  $\sim 10$ -fold higher than those in WT and C-1 cells. These results suggested that decreased Ci affinity in H82 cells was partially caused by a defect in the accumulation of HLA3 and LCIA, but other additional factors could be responsible for the HC-requiring phenotype of H82 cells.


### CAS-Dependent Regulation of Nuclear-Encoded LC-Inducible Genes.

To evaluate the cause of decreased Ci affinity of H82 cells other than HLA3 and LCIA, genes whose expression was affected by

the *CAS* mutation were screened by RNA-seq analysis (Table S2). WT, H82, and C-1 cells were grown in HC or LC conditions for 0.3 and 2 h, and the respective transcriptome profiles were compared. After exposure to HC and LC conditions for 0.3 h, no gene other than *CAS* was affected by the mutation. In contrast, in LC conditions for 2 h, the expression levels of 44 genes in addition to *CAS* were significantly different [FDR (false discovery rate) < 0.05] in H82 cells compared with those in WT and C-1 cells (Table S3). Among them, the expression levels of 13 genes were decreased more than fourfold by the *CAS* mutation, including the following: *HLA3*; *LCIA*; *DNJ31* encoding putative DnaJ-like chaperonin; *CAH4* and *CAH5* encoding mitochondrial carbonic anhydrases; *PPP30* encoding type 2C protein phosphatase; chloroplast carrier protein 1 (*CCP1*) and *CCP2* encoding putative chloroplast envelope membrane proteins; low-CO<sub>2</sub>-inducible gene D (*LCID*); *LHCSR3.1* and *LHCSR3.2* encoding *LHCSR3*; and two unknown genes (*Cre12.g541550* and *Cre26.g756747*). Eventually, the transcriptional abundances of these 12 genes, except for gene *Cre12.g541550*, were decreased by the *CIA5* mutation (9).

In *Chlamydomonas*, accumulation of *LHCSR3*, which is essential for energy-dependent quenching (qE), is shown to be dependent on *CAS* in high-light condition (25). Because its function during LC acclimation related to *CAS* is unclear, we examined *LHCSR3* accumulation in LC conditions in a time-dependent manner (Fig. S1I). After shifting to LC conditions, significant accumulation of *LHCSR3* was detected within 2 h in WT cells, supporting the previous report that *LHCSR3* is induced by CO<sub>2</sub>-limiting stress (29, 30). Although the accumulation of *LHCSR3* in H82 cells was twofold lower than that in WT and C-1 cells in LC at 2 h or 4 h, similar accumulation levels of *LHCSR3* were detected in each strain at 12 h (Fig. S1J). These results suggested that *CAS* regulates *LHCSR3* accumulation at an early stage of CCM induction. In contrast, other CCM-related genes, including *HLA3* and *LCIA*, were transiently induced at the mRNA levels at 0.3 h in H82 cells, but their mRNA levels could not be maintained at the same levels as in WT and C-1 cells at 2 h in contrast to *LCIB* (Fig. S1K), suggesting that *CAS* could be required for maintaining the mRNA levels of *HLA3* and *LCIA* after the initial induction of these genes by CCM1/*CIA5*.

**Requirement of Ca<sup>2+</sup> for LC-Induced Accumulation of *HLA3* and *LCIA*.** As in the case of *AtCAS*, the N terminus of *CrCAS* also has Ca<sup>2+</sup>-binding activity (Fig. S2A–C and SI Results and Discussion). To know the link between Ca<sup>2+</sup> signal via *CAS* and regulation of HCO<sub>3</sub><sup>-</sup> transporters, the accumulation of *HLA3* and *LCIA* in WT cells grown in the presence of 1,2-bis(*o*-aminophenoxy)ethane-*N,N,N,N'*-tetraacetic acid (BAPTA), a membrane-impermeable Ca<sup>2+</sup>-specific chelator (31), was measured in LC conditions. The accumulation of *HLA3* and *LCIA* was dramatically decreased in the presence of 0.5 mM BAPTA (Fig. 2A) as in the case of *LHCSR3* (25, 30), and concomitantly *K*<sub>0.5</sub> (Ci) increased twofold (72 ± 9 μM to 144 ± 14 μM) (Fig. 2B and Table S1). Considering that the addition of 0.75 mM CaCl<sub>2</sub> rescued the accumulation of *HLA3* and *LCIA* and decreased Ci affinity, extracellular Ca<sup>2+</sup> is necessary for the accumulation of these HCO<sub>3</sub><sup>-</sup> transporters and for the photosynthetic Ci affinity. To further examine the regulation of these HCO<sub>3</sub><sup>-</sup> transporters by an intracellular Ca<sup>2+</sup> signal, the impact of *N*-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide hydrochloride (W-7), a membrane-permeable calmodulin antagonist, on the accumulation of *HLA3* and *LCIA* and photosynthetic Ci affinity was examined. In the presence of W-7, concentration-dependent effects of decreased accumulation of *HLA3* and *LCIA* were observed, as in the case of *LHCSR3* (Fig. S2D) (25, 30), and concomitantly *K*<sub>0.5</sub> (Ci) increased from 59 ± 18 μM (mock) to 235 ± 48 μM in the presence of 75 μM W-7 (Fig. S2E and Table S1).




**Fig. 2.** The effects of extracellular calcium (Ca<sup>2+</sup>) chelator 1,2-bis(*o*-aminophenoxy)ethane-*N,N,N,N'*-tetraacetic acid (BAPTA) on the accumulation of *HLA3*, *LCIA*, *LHCSR3*, *LCIB*, and *CAS* (A) and on inorganic carbon (Ci) affinity in low-CO<sub>2</sub> (LC) conditions at pH 7.8 (B). Histone H3 was used as a loading control. High-CO<sub>2</sub> (HC)-grown WT cells were centrifuged, resuspended by fresh high-salt medium supplemented with 20 mM 3-(*N*-morpholino)propanesulfonic acid (HSM) medium in the presence of the indicated chemicals and cultured in LC conditions for 2 h. For the culturing medium without any indications, HSM liquid medium contained 0.068 mM CaCl<sub>2</sub>. Data in all experiment indicate mean value ± SD from three biological replicates.

In contrast, addition of *N*-(6-aminohexyl)-1-naphthalenesulfonamide hydrochloride (W-5), a biologically inactive calmodulin antagonist, did not show any significant effects. As a control, *LCIB* was normally induced in the presence of both chemicals. These results suggested that a calmodulin-mediated Ca<sup>2+</sup> signal could play roles in the LC-induced accumulation of *HLA3* and *LCIA*.

**Relocation of Thylakoid Membrane-Localized *CAS* by LC in Light.** To examine the localization of *CAS*, total protein, soluble/insoluble, chloroplast envelope, and thylakoid membrane fractions were isolated and probed with antibodies against *CAS*, thylakoid membrane-localized D1, soluble protein *LCIB*, plasma membrane-localized H<sup>+</sup>-ATPase (12, 32), and chloroplast envelope-localized *LCIA* (12) (Fig. S3A–C). *CAS* was mainly detected in the insoluble and thylakoid membrane fractions where D1 was enriched, as shown in the previous proteome analyses (23, 24). Similar to the case of *AtCAS*, both the N and C terminus of *CrCAS* were exposed to the stromal side of thylakoid membrane (Fig. S3D and SI Results and Discussion). Next, to elucidate the detailed subcellular localization of *CAS* in vivo, an indirect immunofluorescence assay using an anti-*CAS* antibody was performed (Fig. 3A). In HC conditions with light illumination at 120 μmol photons·m<sup>-2</sup>·s<sup>-1</sup> (HC-light), fluorescent signals were observed as dispersed speckles in the chloroplast. In contrast, after shifting to LC conditions in light (LC-light), the fluorescent signals were aggregated in the pyrenoid after 2 h, and the aggregation was observed as tubule-like structures inside the pyrenoid at 12 h. These localization patterns were consistent with the results using a complemented strain FN-1 expressing FLAG-tag fused *CAS* and an anti-FLAG antibody (Fig. S3E–G). Furthermore, to observe *CAS* localization in living cells, we also generated a complemented strain, CL-1, expressing exogenous *CAS* tagged with Clover, a *Chlamydomonas*-adapted modified GFP (33) (Fig. S3E and H). Fluorescent signals of *CAS*-Clover were also distinctly aggregated and observed as a tubule-like structure in the pyrenoid in LC-light conditions at 12 h (Fig. S3I). The aggregated tubule-like signals were dispersed throughout the chloroplast when transferred to HC-light conditions (Fig. S3J and K).

Changing localization in the chloroplast in response to CO<sub>2</sub> was also shown in the case of *LCIB*, and light as well as CO<sub>2</sub> could affect its localization (17). Similarly, the aggregated fluorescent signals of *CAS* in LC-acclimated cells became diffuse






**Fig. 3.** Subcellular localization of CAS. (A) Localization of CAS in WT and H82 cells was assessed using an indirect immunofluorescence assay with an anti-CAS antibody. Cells grown in high- $\text{CO}_2$  (HC) conditions were shifted to low- $\text{CO}_2$  (LC) conditions for 2 h or 12 h in light, and then the LC-acclimated cells (12 h) were transferred from light to dark for 2 h in the LC condition. (B) The effect of dichlorophenyl-dimethylurea (DCMU) (10  $\mu\text{M}$ ), 2,5-dibromo-3-methyl-6-isopropylbenzoquinone (DBMIB) (10  $\mu\text{M}$ ), or cycloheximide (CHX) (10  $\mu\text{g}\cdot\text{mL}^{-1}$ ) on the localization of CAS in WT cells. The cells cultured in each condition used in A were also used for these drug treatments. The HC-grown cells were incubated under LC-light conditions in the presence of DCMU or DBMIB for 2 h. For CHX treatments, HC-grown cells were transferred to LC-light conditions in the presence of CHX for 12 h. The LC-light-acclimated cells were subjected to LC-dark conditions in the presence of CHX for 2 h. The LC-dark-acclimated cells were transferred to LC-light conditions with or without CHX for 2 h. (C) Localization of CAS in a *CCM1* insertion mutant, C16, and its parental 5D strain using an indirect immunofluorescence assay. Cells grown in HC conditions were shifted to LC-light conditions for 2 h or 12 h. For all panels, each image is placed with the flagella facing upward on the panel. Merged images show the fluorescence image superimposed with its differential interference contrast image. (Scale bars: 5  $\mu\text{m}$ .)

after transferring from LC-light to LC conditions in the dark (LC-dark) within 2 h (Fig. 3A and Fig. S3J). Considering the fact that the localization of CAS to the pyrenoid (as well as the accumulation of HLA3, LCIA, and LCIB) was inhibited in the presence of dichlorophenyl-dimethylurea (DCMU) and 2,5-dibromo-3-methyl-6-isopropylbenzoquinone (DBMIB) (Fig. 3B and Fig. S3I, K, and L) and that these inhibitors also suppress the CCM (34), the localization of CAS to the pyrenoid could be

important for regulation of the CCM in LC-light conditions where the CCM was active.

Next, to determine whether the aggregation of CAS in LC-light is associated with de novo protein synthesis, the effect of cycloheximide (CHX) was examined. By switching from HC-light to LC-light conditions or from LC-light to LC-dark conditions, a change in the localization of fluorescent signals derived from CAS was observed in the presence of CHX (Fig. 3B). These results suggested that the LC-induced CAS relocation was not associated with de novo protein synthesis, in contrast to the case of LCIB (35). Moreover, the addition of CHX inhibited the accumulation of LCIB but not CAS in LC conditions (Fig. S3M), and the addition of BAPTA and W-7 did not affect CAS localization (Fig. S3N). Because the accumulation of CAS and its relocation in response to  $\text{CO}_2$  were not impaired in strain C16, a *CCM1* insertion mutant (6) (Fig. 3C and Fig. S3O), relocation of CAS was regulated by external  $\text{CO}_2$  concentration irrespective of *CCM1*/*CIA5* function. Additionally, the impaired photosynthetic Ci affinity in H82 cells caused poor Ci consumption in the culture medium, affecting the localization of LCIB in LC conditions (Fig. S3P–S and SI Results and Discussion).

**Increased  $\text{Ca}^{2+}$  Concentrations in the Pyrenoid in LC-Light Conditions.** Because CAS had low  $\text{Ca}^{2+}$ -binding affinity (20), a high  $\text{Ca}^{2+}$  concentration should be required to bind  $\text{Ca}^{2+}$ . This aspect of CAS raised the possibility that the subcellular regions where free  $\text{Ca}^{2+}$  is enriched could be related to CAS localization. To test this hypothesis, we monitored the fluorescence of Calcium Green-1, AM, a  $\text{Ca}^{2+}$ -sensitive fluorescent dye, in WT and H82 cells (Fig. 4 and Fig. S4). In both HC and LC conditions, apparent fluorescent signals were detected in regions that overlapped with chlorophyll. Notably, distinct high levels of fluorescent signals were observed in the region of the pyrenoid in both WT and H82 cells, especially in LC-light conditions, thereby implying that free  $\text{Ca}^{2+}$  might be concentrated in the pyrenoid. These increased fluorescent signals in LC-light conditions were not impaired by the CAS mutation (Fig. 4 and Fig. S4A) as well as BAPTA and W-7 (Fig. S4B). In contrast, the fluorescent signals in the pyrenoid were decreased after shifting from LC-light to LC-dark conditions for 2 h. Additionally, the signal intensities of  $\text{Ca}^{2+}$  indicator in chloroplast were slightly stronger in H82 than in WT cells after switching to LC conditions for 12 h (Fig. 4 and Fig. S4A), suggesting a higher free  $\text{Ca}^{2+}$  concentration in the chloroplast of H82 cells. Considering that depletion of CAS protein does not affect the total cellular



**Fig. 4.** Calcium Green-1, AM fluorescence in WT and H82 cells. Cells were grown in high- $\text{CO}_2$  (HC) or low- $\text{CO}_2$  (LC) conditions for 12 h in light, or cells in LC-light were transferred to LC-dark conditions for 2 h, and then incubated with Calcium Green-1, AM (CG-1) at room temperature for 30 min. Fluorescence images derived from CG-1 treatments and chlorophyll (CHL) are shown. Each image is placed with the flagella facing upward on the panel. DIC, differential interference contrast image. (Scale bars: 5  $\mu\text{m}$ .)

Ca<sup>2+</sup> content (25), increased free Ca<sup>2+</sup> could be caused by the redistribution of internal Ca<sup>2+</sup>.

## Discussion

In this study, we identified a Ca<sup>2+</sup>-binding protein as a regulator of the CCM by characterization of a CAS insertion mutant, H82, and its complemented strain, C-1. RNA-seq analyses of these strains revealed that CAS is required for maintaining the expression of 13 nuclear-encoded LC-induced genes after induction (Table S3). Of those genes regulated by CAS, *HLA3* and *LCIA*, whose accumulation was decreased in H82 cells (Fig. 1D), are involved in HCO<sub>3</sub><sup>-</sup> uptake for operation of the CCM (11–14). However, simultaneous expression of *HLA3* and *LCIA* could only partially rescue the decreased Ci affinity of H82 cells (Fig. 1A), suggesting that some of the 13 CO<sub>2</sub>-limiting-inducible genes other than *HLA3* and *LCIA* could contribute to the operation of the CCM in *Chlamydomonas* cells.

Thylakoid-membrane-localized CAS changed its localization from dispersed in HC-light or LC-dark conditions, where the CCM was inactive, to an aggregated tubule-like structure in the pyrenoid in LC-light conditions, where the CCM is active (Fig. 3A and Fig. S3I), which means that the aggregation of CAS to the pyrenoid in response to the availability of environmental light and LC is important for regulation of the CCM. Considering that some of the thylakoid membrane penetrates the pyrenoid, which are termed pyrenoid tubules (36), it is possible that CAS could be localized in or along the pyrenoid tubules in LC-light conditions and could change its localization reversibly in response to CO<sub>2</sub> conditions. The change in CAS localization was independent of de novo protein synthesis (Fig. 3B), which was unlike the case of LCIB (35), suggesting that aggregated CAS is not newly synthesized and that CAS itself could move in response to light and CO<sub>2</sub> conditions. This distinct localization of CAS could be explained by the thylakoid membrane remodeling observed previously in varying light conditions (37) although the actual mechanism for the change in localization requires further analysis.

Previously, based on the fact that the accumulation of chloroplast envelope-localized *LCIA* is required for the expression of *HLA3*, unknown retrograde signals from the chloroplast to nuclear gene have been suggested to support the CCM (12). This study further revealed a regulatory pathway related to chloroplast-retrograde signaling from the thylakoid- and/or pyrenoid-tubule-localized CAS to nuclear genes because CAS was essential to maintain the expression of 13 genes possibly important for operation of the CCM, including both *HLA3* and *LCIA*. Additionally, the fact that accumulation of CCM1 was not inhibited by the loss of CAS protein (27), and vice versa (Fig. S3O), and that CAS could relocate to pyrenoid tubules in response to LC-light irrespective of CCM1 suggest that CCM1 and CAS could function in parallel to each other in the regulation of CCM, including HCO<sub>3</sub><sup>-</sup> transporters (Fig. S5 and SI Results and Discussion).

When chelating external Ca<sup>2+</sup> by application of BAPTA, the accumulation of *HLA3* and *LCIA* was decreased, and the addition of CaCl<sub>2</sub> restored accumulation (Fig. 2A). Because BAPTA cannot permeate the plasma membrane, it remains in the extracellular space (31) and prevents the elevation of both [Ca<sup>2+</sup>]<sub>cyt</sub> and [Ca<sup>2+</sup>]<sub>stro</sub> (22), meaning that the elevation of intracellular Ca<sup>2+</sup> is required for the accumulation of these two HCO<sub>3</sub><sup>-</sup> transporters. In terrestrial plants, the observation that CO<sub>2</sub>-induced changes in [Ca<sup>2+</sup>]<sub>cyt</sub> and stomatal closing were attenuated by chelating or without adding external Ca<sup>2+</sup> (19) leads to a hypothesis for the participation of [Ca<sup>2+</sup>]<sub>cyt</sub> in CO<sub>2</sub> signal transduction in guard cells. Considering that CAS also mediates transient elevation of [Ca<sup>2+</sup>]<sub>cyt</sub> and [Ca<sup>2+</sup>]<sub>stro</sub> in *Arabidopsis* (21, 22), it is possible that the regulation of these HCO<sub>3</sub><sup>-</sup> transporters by CAS could be through Ca<sup>2+</sup> signals resulting from the influx of [Ca<sup>2+</sup>]<sub>ext</sub> into *Chlamydomonas* cells.

Moreover, with the use of a Ca<sup>2+</sup> indicator, free Ca<sup>2+</sup> could be concentrated in the pyrenoid, especially in LC-light conditions (Fig. 4 and Fig. S4A). Because the Ca<sup>2+</sup>-binding characteristics of CAS involve low affinity and high capacity (20), the coexistence of CAS and higher concentrations of Ca<sup>2+</sup> in the pyrenoid could be important to activate CAS function. A rhodanese-like domain conserved in the C terminus of CAS is thought to exhibit a regulatory function rather than an enzymatic one (18), which might account for the consequent signal transduction followed by binding of Ca<sup>2+</sup> at its N terminus. Furthermore, because the chloroplastic Ca<sup>2+</sup> has been shown to be important for the chloroplast metabolism and the function of the thylakoid (18), where CAS protein is localized (Fig. S3B), it is possible that the absence of CAS could cause damage on thylakoid or pyrenoid structure in H82 cells in LC conditions. Of note, stronger fluorescent signals derived from a Ca<sup>2+</sup> indicator in the pyrenoid in LC-light conditions was also observed in H82 cells (Fig. 4 and Fig. S4A), which was not affected by perturbation of intracellular Ca<sup>2+</sup> homeostasis (Fig. S4B), suggesting that the change in Ca<sup>2+</sup> concentration is not directly regulated by CAS and could act upstream of the Ca<sup>2+</sup> signal for regulation of the CCM.

In addition to *HLA3* and *LCIA*, the mRNA abundance and accumulation of *LHCSR3* were significantly decreased by the impairment of CAS in LC conditions at 2 h (Table S3 and Fig. S1 I and J), supporting the previous finding that CAS knock-down *Chlamydomonas* strains showed decreased levels of *LHCSR3* accumulation (25). Considering that the accumulation of *HLA3*, *LCIA*, and *LHCSR3* was simultaneously decreased by the application of BAPTA or W-7 in LC at 2 h, the expression of these proteins could be regulated by CAS protein or Ca<sup>2+</sup> signal in a similar way.

In contrast to CCM1/*CIA5*, CAS is highly conserved in vascular plants, as well as green eukaryotic algae, over the course of evolution. Our findings showed that chloroplast-mediated retrograde signaling pathways via CAS were already developed in the green algae lineage, which could throw light on understanding the cross-talk between Ca<sup>2+</sup>- and CO<sub>2</sub>-dependent signal transduction pathways in photosynthetic organisms. Furthermore, considering that the induction of the CCM is dependent on limiting-CO<sub>2</sub> conditions, as well as on light intensity (29), the relationship between CO<sub>2</sub>-limiting stress and high-light stress could be clarified by further analyses of this CAS mutant. The trigger of relocation of CAS into the pyrenoid, as well as Ca<sup>2+</sup> in response to changes in the availability of CO<sub>2</sub> and light, could be a key regulatory factor for the CO<sub>2</sub>-sensing mechanism in photosynthetic eukaryotes.

## Materials and Methods

*C. reinhardtii* strain C-9 (mt<sup>-</sup>) was used as a WT line (6). *CCM1* insertion mutant C16, as well as its parental 5D strain (6), and a *PGRL1* insertion mutant (38) were characterized previously. For the complementation assay, the DNA genomic fragment was amplified using specific primers (Table S4). The genotypes of strains used in this study are listed in Table S5. In physiological assays, cells were cultured in Tris acetate-phosphate liquid medium for preculture and subcultured in modified high-salt medium (NH<sub>4</sub><sup>+</sup>) supplemented with 20 mM 3-(*N*-morpholino)propanesulfonic acid (Mops) to midlog phase (OD<sub>730</sub> 0.3 to 0.5) for photoautotrophic growth. For all culture conditions without other conditions specified, cells were cultured at 25 °C with illumination at 120 μmol photons·m<sup>-2</sup>·s<sup>-1</sup>.

Additional information is described in the SI Materials and Methods.

**ACKNOWLEDGMENTS.** We thank Gilles Peltier and Xenie Johnson (Commissariat à l'Énergie Atomique/Centre National de la Recherche Scientifique) for providing the *pgrl1* mutant and its complemented strain; Göran Samuelsson (Umeå University) for the anti-CAH3 antibody; and Michael Hippler (University of Münster) for the anti-LHCSR3 antibody. We also thank Kosuke Tsuda for technical assistance. This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grants 25120714 and 16H04805 (to H.F.) and 25840109 (to T.Y.); the Japan Science and Technology Agency (JST) Advanced Low Carbon Technology Research and Development Program (ALCA) (H.F.); JST Core Research for Evolutional Science and Technology (CREST) (Y.T. and S.-i.O.); and a Cooperative Research grant of the Genome Research for BioResource, NODAI Genome Research Center, Tokyo University of Agriculture.

- Cummins EP, Selfridge AC, Sporn PH, Sznajder JJ, Taylor CT (2014) Carbon dioxide-sensing in organisms and its implications for human disease. *Cell Mol Life Sci* 71(5): 831–845.
- Hashimoto M, et al. (2006) *Arabidopsis* HT1 kinase controls stomatal movements in response to CO<sub>2</sub>. *Nat Cell Biol* 8(4):391–397.
- Hu H, et al. (2010) Carbonic anhydrases are upstream regulators of CO<sub>2</sub>-controlled stomatal movements in guard cells. *Nat Cell Biol* 12(1):87–93, sup pp 1–18.
- Jones HG (1992) *Plants and Microclimate: A Quantitative Approach to Environmental Plant Physiology* (Cambridge Univ Press, Cambridge UK), 2nd Ed.
- Wang Y, Zhang J, Spalding MH (2015) The CO<sub>2</sub> concentrating mechanism and photosynthetic carbon assimilation in limiting CO<sub>2</sub>: How *Chlamydomonas* works against the gradient. *Plant J* 82(3):429–448.
- Fukuzawa H, et al. (2001) *Ccm1*, a regulatory gene controlling the induction of a carbon-concentrating mechanism in *Chlamydomonas reinhardtii* by sensing CO<sub>2</sub> availability. *Proc Natl Acad Sci USA* 98(9):5347–5352.
- Xiang Y, Zhang J, Weeks DP (2001) The *Cia5* gene controls formation of the carbon concentrating mechanism in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* 98(9):5341–5346.
- Miura K, et al. (2004) Expression profiling-based identification of CO<sub>2</sub>-responsive genes regulated by CCM1 controlling a carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant Physiol* 135(3):1595–1607.
- Fang W, et al. (2012) Transcriptome-wide changes in *Chlamydomonas reinhardtii* gene expression regulated by carbon dioxide and the CO<sub>2</sub>-concentrating mechanism regulator CIA5/CCM1. *Plant Cell* 24(5):1876–1893.
- Im CS, Grossman AR (2002) Identification and regulation of high light-induced genes in *Chlamydomonas reinhardtii*. *Plant J* 30(3):301–313.
- Duanmu D, Miller AR, Horken KM, Weeks DP, Spalding MH (2009) Knockdown of limiting-CO<sub>2</sub>-induced gene HLA3 decreases HCO<sub>3</sub><sup>-</sup> transport and photosynthetic Ci affinity in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* 106(14):5990–5995.
- Yamano T, Sato E, Iguchi H, Fukuda Y, Fukuzawa H (2015) Characterization of cooperative bicarbonate uptake into chloroplast stroma in the green alga *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* 112(23):7315–7320.
- Gao H, Wang Y, Fei X, Wright DA, Spalding MH (2015) Expression activation and functional analysis of HLA3, a putative inorganic carbon transporter in *Chlamydomonas reinhardtii*. *Plant J* 82(1):1–11.
- Wang Y, Spalding MH (2014) Acclimation to very low CO<sub>2</sub>: Contribution of limiting CO<sub>2</sub> inducible proteins, LCIB and LCIA, to inorganic carbon uptake in *Chlamydomonas reinhardtii*. *Plant Physiol* 166(4):2040–2050.
- Vance P, Spalding MH (2005) Growth, photosynthesis, and gene expression in *Chlamydomonas* over a range of CO<sub>2</sub> concentrations and CO<sub>2</sub>/O<sub>2</sub> ratios: CO<sub>2</sub> regulates multiple acclimation states. *Can J Bot* 83(7):796–809.
- Wang Y, Spalding MH (2006) An inorganic carbon transport system responsible for acclimation specific to air levels of CO<sub>2</sub> in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* 103(26):10110–10115.
- Yamano T, et al. (2010) Light and low-CO<sub>2</sub>-dependent LCIB-LCIC complex localization in the chloroplast supports the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant Cell Physiol* 51(9):1453–1468.
- Hochmal AK, Schulze S, Trompelt K, Hippler M (2015) Calcium-dependent regulation of photosynthesis. *Biochim Biophys Acta* 1847(9):993–1003.
- Kim TH, Böhmer M, Hu H, Nishimura N, Schroeder JI (2010) Guard cell signal transduction network: Advances in understanding abscisic acid, CO<sub>2</sub>, and Ca<sup>2+</sup> signaling. *Annu Rev Plant Biol* 61:561–591.
- Han S, Tang R, Anderson LK, Woerner TE, Pei ZM (2003) A cell surface receptor mediates extracellular Ca<sup>2+</sup> sensing in guard cells. *Nature* 425(6954):196–200.
- Nomura H, Komori T, Kobori M, Nakahira Y, Shiina T (2008) Evidence for chloroplast control of external Ca<sup>2+</sup>-induced cytosolic Ca<sup>2+</sup> transients and stomatal closure. *Plant J* 53(6):988–998.
- Nomura H, et al. (2012) Chloroplast-mediated activation of plant immune signalling in *Arabidopsis*. *Nat Commun* 3:926.
- Allmer J, Naumann B, Markert C, Zhang M, Hippler M (2006) Mass spectrometric genomic data mining: Novel insights into bioenergetic pathways in *Chlamydomonas reinhardtii*. *Proteomics* 6(23):6207–6220.
- Terashima M, Specht M, Naumann B, Hippler M (2010) Characterizing the anaerobic response of *Chlamydomonas reinhardtii* by quantitative proteomics. *Mol Cell Proteomics* 9(7):1514–1532.
- Petroustos D, et al. (2011) The chloroplast calcium sensor CAS is required for photoacclimation in *Chlamydomonas reinhardtii*. *Plant Cell* 23(8):2950–2963.
- Terashima M, et al. (2012) Calcium-dependent regulation of cyclic photosynthetic electron transfer by a CAS, ANR1, and PGRL1 complex. *Proc Natl Acad Sci USA* 109(43): 17717–17722.
- Wang L, Yamano T, Kajikawa M, Hirono M, Fukuzawa H (2014) Isolation and characterization of novel high-CO<sub>2</sub>-requiring mutants of *Chlamydomonas reinhardtii*. *Photosynth Res* 121(2-3):175–184.
- Lucker B, Kramer DM (2013) Regulation of cyclic electron flow in *Chlamydomonas reinhardtii* under fluctuating carbon availability. *Photosynth Res* 117(1-3):449–459.
- Yamano T, Miura K, Fukuzawa H (2008) Expression analysis of genes associated with the induction of the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant Physiol* 147(1):340–354.
- Maruyama S, Tokutsu R, Minagawa J (2014) Transcriptional regulation of the stress-responsive light harvesting complex genes in *Chlamydomonas reinhardtii*. *Plant Cell Physiol* 55(7):1304–1310.
- Polisenky DH, Braam J (1996) Cold-shock regulation of the *Arabidopsis* TCH genes and the effects of modulating intracellular calcium levels. *Plant Physiol* 111(4): 1271–1279.
- Campbell AM, Coble AJ, Cohen LD, Chng TH, Russo KM (2001) Identification and DNA sequence of a new H<sup>+</sup>-ATPase in the unicellular green alga *Chlamydomonas reinhardtii*. *J Phycol* 37(4):536–542.
- Lauersen KJ, Kruse O, Mussnug JH (2015) Targeted expression of nuclear transgenes in *Chlamydomonas reinhardtii* with a versatile, modular vector toolkit. *Appl Microbiol Biotechnol* 99(8):3491–3503.
- Badger MR, Kaplan A, Berry JA (1980) Internal inorganic carbon pool of *Chlamydomonas reinhardtii*-evidence for a carbon-dioxide concentrating mechanism. *Plant Physiol* 66(3):407–413.
- Yamano T, Asada A, Sato E, Fukuzawa H (2014) Isolation and characterization of mutants defective in the localization of LCIB, an essential factor for the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Photosynth Res* 121(2-3): 193–200.
- Engel BD, et al. (2015) Native architecture of the *Chlamydomonas* chloroplast revealed in situ cryo-electron tomography. *eLife* 4:e04889, and correction (2015) 4:e11383.
- Chuartzman SG, et al. (2008) Thylakoid membrane remodeling during state transitions in *Arabidopsis*. *Plant Cell* 20(4):1029–1039.
- Tolletter D, et al. (2011) Control of hydrogen photoproduction by the proton gradient generated by cyclic electron flow in *Chlamydomonas reinhardtii*. *Plant Cell* 23(7): 2619–2630.
- Dang KV, et al. (2014) Combined increases in mitochondrial cooperation and oxygen photoreduction compensate for deficiency in cyclic electron flow in *Chlamydomonas reinhardtii*. *Plant Cell* 26(7):3036–3050.
- Alric J, Laverne J, Rappaport F (2010) Redox and ATP control of photosynthetic cyclic electron flow in *Chlamydomonas reinhardtii* (l) aerobic conditions. *Biochim Biophys Acta* 1797(1):44–51.
- Karlsson J, et al. (1998) A novel alpha-type carbonic anhydrase associated with the thylakoid membrane in *Chlamydomonas reinhardtii* is required for growth at ambient CO<sub>2</sub>. *EMBO J* 17(5):1208–1216.
- Duanmu D, Wang Y, Spalding MH (2009) Thylakoid lumen carbonic anhydrase (CAH3) mutation suppresses air-Dier phenotype of LCIB mutant in *Chlamydomonas reinhardtii*. *Plant Physiol* 149(2):929–937.
- Fujiwara T, et al. (2015) A nitrogen source-dependent inducible and repressible gene expression system in the red alga *Cyanidioschyzon merolae*. *Front Plant Sci* 6:657.
- Trapnell C, et al. (2012) Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* 7(3):562–578.