TITLE:

Shoulder horizontal abduction stretching effectively increases shear elastic modulus of pectoralis minor muscle

AUTHOR(S):

Umehara, Jun; Nakamura, Masatoshi; Fujita, Kosuke; Kusano, Ken; Nishishita, Satoru; Araki, Kojiro; Tanaka, Hiroki; Yanase, Ko; Ichihashi, Noriaki

CITATION:

Umehara, Jun ...[et al]. Shoulder horizontal abduction stretching effectively increases shear elastic modulus of pectoralis minor muscle. Journal of Shoulder and Elbow Surgery 2017, 26(7): 1159-1165

ISSUE DATE:

2017-07

URL:

<http://hdl.handle.net/2433/231253>

RIGHT:

© 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>; The full-text file will be made open to the public on 01 July 2018 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

How do you stretch pectoralis minor?

1 Shoulder horizontal abduction stretching effectively increases shear elastic modulus of
2 pectoralis minor muscle

3

4 Jun Umehara, M.S.^{1,2)*}, Masatoshi Nakamura, Ph.D.³⁾, Kosuke Fujita, M.S.⁴⁾, Ken Kusano,
5 B.S.¹⁾, Satoru Nishishita, M.S.¹⁾, Kojiro Araki, M.S.⁵⁾, Hiroki Tanaka, M.S.²⁾, Ko Yanase, M.S.¹⁾,
6 Noriaki Ichihashi, Ph.D.¹⁾

7

8 1) Human Health Sciences, Graduate School of Medicine, Kyoto University, Kyoto, Japan

9 2) Rehabilitation Unit, Kyoto University Hospital, Kyoto, Japan

10 3) Institute for Human Movement and Medical Sciences, Niigata University of Health and
11 Welfare, Niigata, Japan

12 4) Rehabilitation group, Department of Medical Technique, Nagoya University, Nagoya, Japan

13 5) Department of Rehabilitation, Sapporo Tokushukai Hospital, Sapporo, Japan

14

15 ***Corresponding author**

16 Jun Umehara (✉)

17 Human Health Sciences, Graduate School of Medicine, Kyoto University,

18 53 Shogoin-Kawahawa-cho, Kyoto 606-8507, Japan

19 Phone: +81-75-751-3935; Fax: +81-75-751-3909

How do you stretch pectoralis minor?

20 E-mail: umehara.jun.77z@st.kyoto-u.ac.jp

21

22 **Financial biases:** None

23

24 **Ethics committee approval**

25 The study design was approved by the ethic committee of Kyoto University Graduate School

26 and the Faculty of Medicine (R0314).

27

28 **Acknowledgement**

29 This study was supported by the Grant – in – Aid for Scientific Research (B) 15H03043.

30

How do you stretch pectoralis minor?

31 Abstract

32 Background

33 The stretching maneuver for pectoralis minor muscle, which is shoulder horizontal abduction
34 or scapular retraction, is performed in a clinical and sport setting because the tightness of this
35 muscle may contribute to scapular dyskinesis. The effective stretching maneuver for pectoralis
36 minor muscle is unclear in vivo. The purpose of this study was to verify the effective stretching
37 maneuver for pectoralis minor muscle in vivo using ultrasonic shear wave elastography.

38 Methods

39 Eighteen healthy men participated in this study. Elongation of the pectoralis minor muscle was
40 measured for three stretching maneuvers (shoulder flexion, shoulder horizontal abduction, and
41 scapular retraction) at three shoulder elevation angles (30°, 90°, and 150°). The shear elastic
42 modulus used as the index of muscle elongation was computed using ultrasonic shear wave
43 elastography for the above nine stretching maneuver-angle combinations.

44 Results

45 The shear elastic modulus was highest in horizontal abduction 150° followed by horizontal
46 abduction 90°, horizontal abduction 30°, scapular retraction 30°, scapular retraction 90°,
47 scapular retraction 150°, flex 150°, flex90° and flex30°. The shear elastic modulus of horizontal
48 abduction 90° and horizontal abduction 150° were significantly higher than other stretching
49 maneuvers. There was no significant difference between horizontal abduction 90° and
50 horizontal abduction 150°.

How do you stretch pectoralis minor?

51 **Conclusions**

52 This study determined that shoulder horizontal abduction at elevations of 90° and 150° were
53 the most effective stretching maneuvers for pectoralis minor muscle in vivo.

54

55 **Level of evidence:** Basic Science Study.

56

57 **Keyword**

58 Stretching;

59 Pectoralis minor muscle;

60 Ultrasonic shear wave elastography;

61 Shear elastic modulus;

62 Shoulder horizontal abduction;

63 Rehabilitation

64

How do you stretch pectoralis minor?

65 Introduction

66 In shoulder rehabilitation, clinical evaluation, and intervention for scapular dyskinesis are
67 important because of its relation to various shoulder injuries, such as subacromial
68 impingement^{4,12,17,19,23}, rotator cuff tear^{14,24}, unstable shoulder²⁰ and frozen shoulder¹⁰. The
69 literature suggests that scapular dyskinesis may be caused by multiple factors such as bone,
70 joint, neurological, or soft tissue mechanisms¹³. In soft tissue mechanisms, the tightness of the
71 pectoralis minor muscle (PMi) is one of the factors inducing scapular dyskinesis, which can be
72 examined and treated by a therapist⁸. Previous studies have reported that the tightness of the
73 PMi is related to posture, including scapular internal rotation in the resting position³ and
74 decreases in scapular external rotation and posterior tilt during arm elevation⁵. These changes,
75 which is the scapular internal rotation and anterior tilt, are similar to the change in scapular
76 motion found in many shoulder injuries^{19,20} and it is also possible that there might be a
77 relationship between shoulder injury and the tightness of the PMi. Therefore, the flexibility of
78 the PMi is important for preventing and improving scapular dyskinesis.

79 Stretching interventions are recommended to increase and improve muscle flexibility
80 and stretching of the PMi is frequently used in rehabilitation programs^{1,18,21}. Therefore, some
81 studies have investigated stretching maneuver of the PMi. In a previous study, Borstad and
82 Ludwig⁶ compared the length of the PMi during three stretching maneuver using an
83 electromagnetic motion capture system with skin surface markers in healthy adults. This study
84 concluded that the most effective PMi stretching maneuver was a unilateral corner self-stretch

How do you stretch pectoralis minor?

85 similar to horizontal abduction of shoulder joint. On the other hand, Muraki et al.²⁵ directly
 86 measured the length of the PMi during three passive shoulder motions and three stretching
 87 techniques using displacement sensors in fresh cadavers. They advocated that scapular
 88 retraction made the greatest change in PMi length. The contradiction of these two studies most
 89 likely resulted from differences between the subjects (living body vs. cadaver) or measurement
 90 methods. In addition, it is unknown whether or not the results of these previous studies apply
 91 to live people for effective stretching positions of PMi, because Borstad and Ludewig⁶ did not
 92 measure the tension of the PMi during stretching, but instead measured the distance between
 93 the coracoid process and the fourth rib/sternum junction, and Muraki et al.²⁵ used cadavers in
 94 their study. Therefore, an investigation of effective in vivo stretching maneuver of the PMi
 95 determined by measuring muscle tension during stretching is needed.

96 A new ultrasound-based technology called ultrasonic shear wave elastography (SWE)
 97 was developed, allowing reliable and non-invasive measurement of soft tissue viscoelastic
 98 properties². SWE monitors the propagation of shear waves generated in tissue using acoustic
 99 radiation forces and is able to evaluate the shear elastic modulus of individual muscles²⁶.
 100 Because of the strong liner relationship, identified by prior studies, between passive muscle
 101 tension calculated by traditional method and shear elastic modulus measured by SWE in
 102 vitro^{9,15}, SWE has been used in many stretching studies of skeletal muscle^{9,15,27,28}. In addition,
 103 our previous studies indicate an increase in shear elastic modulus with muscle elongation during

How do you stretch pectoralis minor?

104 stretching^{27,28}. Therefore, SWE has proven to be a valid technology for noninvasively
105 investigating muscle elongation in vivo.

106 Regarding stretching maneuver of the PMi, a unilateral corner self-stretch or scapular
107 retraction at a 30° shoulder flexion angle have been recommended by Borstad and Ludwig⁶ or
108 Muraki et al.²⁵, respectively. Muraki et al.²⁵ also described that the PMi can be stretched by
109 150° passive shoulder flexion and scaption as well as scapular retraction or shoulder horizontal
110 abduction. Thus, we hypothesized that shoulder horizontal abduction or scapular retraction with
111 the shoulder in elevated positions is an effective maneuver for stretching the PMi. The objective
112 of this study was to quantitatively verify the effective stretching maneuver of the PMi using the
113 shear elastic modulus measured by SWE in vivo.
114

How do you stretch pectoralis minor?

115 **Materials and Methods**

116 **Participants**

117 Eighteen men (age, 26.2 ± 4.0 years; height, 171.1 ± 5.0 cm; weight, 67.4 ± 7.8 kg) with no
118 orthopaedic or nervous system abnormalities in their upper limbs participated in this study.
119 Participants were recruited from the students of our institution. The subject orally confirmed
120 that they complied with the following exclusion criteria: females, athletes, or those who perform
121 any extensive exercise, and subjects having a history of orthopaedic disease or neuropathy in
122 their upper limbs. The sample size was calculated by the G*power software (Version 3.1.,
123 Heinrich Heine University, Dusseldorf, Germany) for a one-way analysis of variance (ANOVA)
124 with repeated measures (effect size = 0.25, α error = 0.05, power = 0.8), which showed that 17
125 subjects were required. This study protocol conformed to the Helsinki Declaration and was
126 approved by the Ethics Review Board of our institution.

127

128 **Experimental procedures**

129 This study was an experimental study, with randomized allocation of stretching intervention for
130 each subject using a random number table. Healthy male subjects were randomly recruited.
131 After the aim and procedures were explained to all subjects, the subjects underwent nine
132 stretching maneuvers performed by one researcher. The outcome was measured and analyzed
133 by another researcher.

134 All procedures were performed by the same two investigators who both had physical

How do you stretch pectoralis minor?

therapist licenses: one investigator measured the shear elastic modulus using SWE, while the other performed the stretching maneuver. The non-dominant upper limb was chosen for intervention. Each subject lay on their side on a bed with their non-intervention arm under their head, their trunk parallel to the long axis of the bed, and both the hip and knee flexed 45°. The relax position (Rest) was defined as that the shoulder in 0° flexion, 0° abduction, the elbow fully extended, and the palm of the hand parallel the bed. In this study, three stretching maneuvers (flexion, horizontal abduction, and scapular retraction) were investigated at the three shoulder elevation angles (30°, 90°, and 150°) for a total of nine stretching positions, on the basis of previous studies^{6,25}. For passive shoulder motion, the interventional shoulder of the subject was passively flexed at 30° (Flex30), 90° (Flex90), and 150° (Flex150) by the investigator (Figure 1). For shoulder horizontal abduction, the interventional shoulder was passively horizontally abducted as much as possible at shoulder elevation angles of 30° (Hab30), 90° (Hab90), and 150° (Hab150) while the shoulder was maximally externally rotated and the elbow was flexed 90° by the investigator (Figure 2). For scapular retraction, the interventional fully flexed elbow was maximally pressed along the longitudinal axis of the humerus, and the interventional scapular was passively maximally retracted at the shoulder for flexion angles of 30° (Retract30), 90° (Retract90), and 150° (Retraction150) by the investigator (Figure 3). The subjects were stretched until reaching a point of discomfort (but not pain), as verbally acknowledged by the subjects. During all stretching and measurement acquisition, subjects

How do you stretch pectoralis minor?

154 were instructed to relax as much as possible.

155

156 **Instrumentation**

157 In this study, the shear elastic modulus measured by SWE (Aixplorer, SuperSonic Imagine,
158 Aix-en-Provence, France) with an ultrasound transducer (15-4 MHz linear probe) was defined
159 as the indicator of muscle elongation of the PMi. The shear elastic modulus was calculated from
160 the shear wave propagation speed generated by the transducer². The shear elastic modulus (G)
161 was calculated from the shear wave propagation speed (V) using the following formula:

162 $G = \rho V^2$

163 Where ρ is the muscle density, assumed to be 1,000 kg/m³. A previous study showed that there
164 is a significant correlation between the shear elastic modulus, which was measured by SWE,
165 and the muscle elongation, which was measured by a traditional tension test^{9,15}.

166 The shear elastic modulus was measured in all measurement positions using SWE. The
167 measurement place was defined as the midpoint between the coracoid process and the fourth
168 rib/sternum junction, identified in the ultrasonic image. The probe was placed in parallel with
169 muscle fascicle of the PMi. The region of interest (ROI) was established near the center point
170 of the muscle belly in the ultrasound image. The shear elastic modulus was measured three
171 times at each measurement site and the mean value was used for analysis. All analyses were
172 performed by the researcher who was blinded to the stretching positions by anonymizing all

How do you stretch pectoralis minor?

ultrasonic images. The subjects were instructed to hold their breath during measurement of the shear elastic modulus to prevent PMi elongation due to the movement of the rib cage.

Five healthy men (age, 25.8 ± 3.7 years; height, 172.8 ± 5.0 cm; weight, 65.8 ± 4.6 kg) were used to evaluate the reliability of the ultrasound measurement. The measurement was acquired for three passive shoulder motions and three stretching maneuvers. The reliability of the shear elastic modulus measurement was confirmed using intraclass correlation coefficients (1,3) ($ICC_{1,3}$) with a 95% confidence interval (95% CI). $ICC_{1,3}$ values, which represent intra-observer reliability in the intra-day, were calculated from the shear elastic modulus. $ICC_{1,3}$ values fell within a range of 0.90 – 0.99 for all measurements (Table I). A previous study that investigated the reliability coefficient suggested that a range of 0.81 – 1.00 was almost perfect¹⁶. Therefore, the measured values of the shear elastic modulus in our study were considered reproducible since the $ICC_{1,3}$ observed was almost perfect, according to this previous study.

Data analysis

Statistical analysis was performed using IBM SPSS Statistics 22 (International Business Machines Corporation). To find whether the PMi was elongated in the nine stretching positions, differences in the shear elastic modulus between Rest and each stretching position were assessed using a paired Student's t-test with Bonferroni revision. Additionally, when the shear elastic modulus was found to be significantly different from that at Rest, a one-way ANOVA

How do you stretch pectoralis minor?

192 with repeated measures was used to determine the effect of passive motion or stretching
193 maneuver on the shear elastic moduli amongst them. If a significant main effect was found,
194 then a Bonferroni multiple comparison for the post hoc test was performed. A confidence level
195 of 0.05 was used in all statistical tests.
196

How do you stretch pectoralis minor?

197 **Result**

198 The shear elastic modulus for each measurement is shown in Table II. The shear elastic modulus
199 was highest at Hab150, followed by Hab90, Hab30, Retraction30, Retraction90, Retraction150,
200 Flex150, Flex90, and Flex30. The shear elastic moduli of these positions, except Flex30 and
201 Flex90, were significantly higher than the elastic modulus at Rest ($P < .05$ or $P < .01$, Table II).
202 For the measurement positions where the shear elastic modulus was significantly higher than
203 that at Rest, a one-way ANOVA with repeated measures was used to indicate a significant main
204 effect ($P < .001$, $F = 29.0$). For the positions showing significantly higher shear elastic moduli
205 than the elastic modulus at Rest, a Bonferroni multiple comparison for the post-hoc test was
206 performed, indicating that the shear elastic moduli of Hab90 and Hab150 were significantly
207 higher than those of the other positions. However, there was no significant difference between
208 Hab90 and Hab150. In addition, although the shear elastic modulus of Hab30 was significantly
209 higher than those of Flex150 and Retraction150 ($P < .001$, respectively), there were no
210 significant differences among the others (Figure 4).

211

How do you stretch pectoralis minor?

212 Discussion

213 This is the first to determine effective stretching maneuver of the PMi using shear elastic
214 modulus values measured by SWE, which quantitatively reflects the grade of muscle elongation
215 during stretching in vivo. The main finding of this study was that maximal horizontal abduction
216 of the shoulder at an elevation angle of 90° and 150° effectively elongates the PMi muscle.

217 We hypothesized that the PMi could be elongated effectively by shoulder horizontal
218 abduction or scapular retraction at elevated shoulder positions i.e., Hab150 or Retraction150.
219 In this study, our results showed that the shear elastic modulus at all measurement positions
220 was higher than that at Rest, except for Flex30 and Flex90. Furthermore, the shear elastic
221 moduli of Hab90 and Hab150 were significantly greater than those of all measurement positions
222 whose shear elastic modulus was greater than at Rest. These results suggest that the most
223 effective stretching maneuvers of the PMi are Hab90 and Hab150, which is partly consistent
224 with our hypothesis.

225 Borstad and Ludewig⁶ compared the mean length change from the coracoid process of
226 the scapula to the fourth rib/sternum junction for three pectoralis minor stretches; unilateral
227 corner self-stretch, sitting manual stretch, and spine manual stretch. They concluded that
228 unilateral corner self-stretch, in which a subject abducts the humerus to 90° with the palm on a
229 wall and then rotates their trunk away from the elevated arm to increase the shoulder horizontal
230 abduction, lengthened the pectoralis minor muscle most effectively. Our results, that the shear
231 elastic moduli of Hab90 and Hab150 were significantly higher than those of other measurement

How do you stretch pectoralis minor?

positions, were consistent with the findings of the previous study by Borstad and Ludewig⁶. On the other hand, Muraki et al. directly measured PMi lengthening during three passive shoulder motions and three stretching techniques using fresh cadaveric transthoracic specimens. Their study concluded that scapular retraction at an angle of 30° flexion, in which the examiner exerted a posterosuperior pressure to the elbow along the longitudinal axis of the humerus, resulted in the greatest change in PMi length measured by displacement sensor²⁵, this is inconsistent with our results. This contradiction probably originates from the difference in methodology. The horizontal abduction of the shoulder might stretch the pectoralis major muscle and the clavipectoral fascia, which may directly impact the ability of the pectoralis minor muscle to receive stretch influence. Removing these tissues overlying the PMi in order to expose this muscle, as stated in Murakis' study, could be the reason for the contradiction. In addition, there was a glaring difference in the nature of the study medium, i.e., live tissue versus cadaver tissue, which likely contributed to this inconsistency. It is possible that the difference in viscoelasticity and other material properties of the shoulder joint between a living body and a cadaver affected the elongation of the PMi¹¹. Contrary to these previous studies, our study examined the applicability of various stretching maneuver of the PMi in live people using the shear elastic modulus values measured by SWE.

The shear elastic moduli of Hab30, Hab90, and Hab150 were significantly higher than the elastic modulus of Flex150. These results indicate that shoulder horizontal abduction is a

How do you stretch pectoralis minor?

251 more effective means of stretching the PMi than shoulder flexion, and that scapular motion is
252 probably responsible for this difference. From an anatomical perspective, the external rotation
253 and posterior tilt of the scapula stretches the PMi⁶ because this muscle originates on the third,
254 fourth, and fifth ribs and runs superolaterally, inserting at the coracoid process of the scapula.
255 Previous studies measuring scapular motion reported that during shoulder flexion the scapula
256 externally rotates, upwardly rotate, and tilts posteriorly²², and that the scapula externally rotates
257 and tilts posteriorly during shoulder horizontal abduction⁷. Comparing the scapular motion of
258 shoulder flexion and that of shoulder horizontal abduction in these previous studies, the scapular
259 external rotation during shoulder horizontal abduction was greater than that during shoulder
260 flexion. Thus, the results of the current study indicating that the shear elastic moduli of Hab30,
261 Hab90, and Hab150 were significantly higher than the elastic modulus of Flex150 suggests that
262 scapular external rotation contributes more to stretching the PMi than scapular posterior tilt.
263 Furthermore, the scapular motion also relates to the fact that the shear elastic moduli of Hab90
264 and Hab150 were found to be significantly higher than those of Retraction30, Retraction90, and
265 Retraction150. The PMi could be more stretched by shoulder horizontal abduction than scapular
266 retraction because the scapular external rotation of shoulder horizontal abduction is greater than
267 that of scapular retraction. However, there was no study investigating scapular motion during
268 scapular retraction. Further research is required to elucidate scapular motion during scapular
269 retraction using electromagnetic sensors or optoelectronic markers.

How do you stretch pectoralis minor?

270 When the shear elastic modulus at Rest was compared to the elastic moduli of other
271 measurement positions, it was found to be significantly lower than all except the Flex30 and
272 Flex90 positions. Considering these results, although Hab90 and Hab150 were the most
273 effective for stretching the PMi, all measurement positions, except Flex30 and Flex90,
274 effectively stretch the PMi. In the clinical setting, patients requiring stretching of the PMi
275 frequently have a limited range of shoulder motion, and a shoulder instability. Therefore, Hab30
276 or Retraction30 might be better suited for these patients. Further research is required to
277 investigate the effects of stretching intervention of the PMi in the patients with a limited range
278 of motion and a shoulder instability.

279 Our determination of horizontal abduction of the shoulder at elevation angles of 90°
280 and 150° as effective means of stretching the PMi may be beneficial in clinical and athletic
281 settings. However, this study should be interpreted with note of the following. First, the
282 participants in this study were healthy young men as prescribed by the exclusion criteria.
283 Therefore, it is unknown whether similar effects can always be expected in patients with
284 impingement syndrome or frozen shoulder. Second, we could not measure the scapular motion
285 during the stretching maneuvers. Further research investigating scapular motion during
286 stretching is required to identify any potential relationship between scapular motion and
287 elongation of the PMi. Third, the shear elastic modulus of the lateral fiber groups of the PMi
288 was measured in this study; thus, similar behavior cannot always be expected in the medial

How do you stretch pectoralis minor?

289 fiber groups of the PMi. However, we presume that there are few differences between the shear
290 elastic moduli of the lateral and medial fiber groups of the PMi because Muraki et al.²⁵ reported
291 that there was no difference of lengthening of the PMi. Thus, the shear elastic modulus of the
292 PMi measured in the current study might represent that of the whole PMi muscle.
293

How do you stretch pectoralis minor?

294 **Conclusions**

295 We quantitatively investigated effective stretching maneuver of the pectoralis minor muscle
296 using shear elastic modulus values obtained by ultrasonic shear wave elastography. Our results
297 showed that shoulder horizontal abduction at shoulder elevation angles of 90° and 150°
298 effectively elongated the pectoralis minor muscle. The stretching maneuver of the pectoralis
299 minor muscle proposed in this study may be useful for clinical application.

300

How do you stretch pectoralis minor?

301 Reference

- 302 1. Bang MD, Deyle GD. Comparison of supervised exercise with and without manual
303 physical therapy for patients with shoulder impingement syndrome. J. Orthop. Sports
304 Phys. Ther. 2000 Mar;30(3):126–37. doi:10.2519/jospt.2000.30.3.126
- 305 2. Bercoff J, Tanter M, Fink M. Supersonic shear imaging: A new technique for soft
306 tissue elasticity mapping. IEEE Trans. Ultrason. Ferroelectr. Freq. 2004
307 Apr;51(4):396–409. doi:10.1109/TUFFC.2004.1295425
- 308 3. Borstad JD. Resting position variables at the shoulder: evidence to support a posture-
309 impairment association. Phys. Ther. 2006 Apr;86(4):549–57.
- 310 4. Borstad JD, Ludewig PM. Comparison of scapular kinematics between elevation and
311 lowering of the arm in the scapular plane. Clin. Biomech. 2002;17:650–659.
312 doi:10.1016/S0268-0033(02)00136-5
- 313 5. Borstad JD, Ludewig PM. The effect of long versus short pectoralis minor resting
314 length on scapular kinematics in healthy individuals. J. Orthop. Sports Phys. Ther.
315 2005;35(4):227–238. doi:10.2519/jospt.2005.1669
- 316 6. Borstad JD, Ludewig PM. Comparison of three stretches for the pectoralis minor
317 muscle. J. Shoulder Elbow Surg. 2006 Jan;15(3):324–30. doi:10.1016/j.jse.2005.08.011
- 318 7. Bourne D a., Choo AMT, Regan WD, MacIntyre DL, Oxland TR. Three-dimensional
319 rotation of the scapula during functional movements: An in vivo study in healthy
320 volunteers. J. Shoulder Elb. Surg. 2007;16(2):150–162. doi:10.1016/j.jse.2006.06.011

How do you stretch pectoralis minor?

- 321 8. Cools AMJ, Struyf F, De Mey K, Maenhout A, Castelein B, Cagnie B. Rehabilitation
322 of scapular dyskinesis: from the office worker to the elite overhead athlete. Br. J.
323 Sports Med. 2014;48(8):692–7. doi:10.1136/bjsports-2013-092148
- 324 9. Eby SF, Song P, Chen S, Chen Q, Greenleaf JF, An K-N. Validation of shear wave
325 elastography in skeletal muscle. J. Biomech. 2013 Sep 27;46(14):2381–7.
326 doi:10.1016/j.jbiomech.2013.07.033
- 327 10. Fayad F, Roby-Brami A, Yazbeck C, Hanneton S, Lefevre-Colau M-M, Gautheron V,
328 et al. Three-dimensional scapular kinematics and scapulohumeral rhythm in patients
329 with glenohumeral osteoarthritis or frozen shoulder. J. Biomech. 2008;41(2):326–332.
330 doi:10.1016/j.jbiomech.2007.09.004
- 331 11. Gottsauner-Wolf F, Grabowski JJ, Chao EYS, An KN. Effects of freeze/thaw
332 conditioning on the tensile properties and failure mode of bone-muscle-bone units: A
333 biomechanical and histological study in dogs. J. Orthop. Res. 1995;13(1):90–95.
334 doi:10.1002/jor.1100130114
- 335 12. Hébert LJ, Moffet H, McFadyen BJ, Dionne CE. Scapular behavior in shoulder
336 impingement syndrome. Arch. Phys. Med. Rehabil. [Internet]. 2002;83(1):60–69.
337 doi:10.1053/apmr.2002.27471
- 338 13. Kibler W Ben, Ludewig PM, McClure PW, Michener L a, Bak K, Sciascia AD.
339 Clinical implications of scapular dyskinesis in shoulder injury: the 2013 consensus

How do you stretch pectoralis minor?

- 340 statement from the “Scapular Summit”. Br. J. Sports Med. [Internet]. 2013;47(14):877–
341 85. doi:10.1136/bjsports-2013-092425
- 342 14. Kijima T, Matsuki K, Ochiai N, Yamaguchi T, Sasaki Y, Hashimoto E, et al. In vivo 3-
343 dimensional analysis of scapular and glenohumeral kinematics: comparison of
344 symptomatic or asymptomatic shoulders with rotator cuff tears and healthy shoulders.
345 J. Shoulder Elb. Surg. 2015;24(11):1817–1826. doi:10.1016/j.jse.2015.06.003
- 346 15. Koo TK, Guo J-Y, Cohen JH, Parker KJ. Relationship between shear elastic modulus
347 and passive muscle force: an ex-vivo study. J. Biomech. 2013 Aug 9;46(12):2053–9.
348 doi:10.1016/j.jbiomech.2013.05.016
- 349 16. Landis JR, Koch GG. The measurement of observer agreement for categorical data.
350 Biometrics. 1977 Mar;33(1):159–74.
- 351 17. Lawrence RL, Braman JP, Laprade RF, Ludewig PM. Comparison of 3-Dimensional
352 Shoulder Complex Kinematics in Individuals With and Without Shoulder Pain, Part 1:
353 Sternoclavicular, Acromioclavicular, and Scapulothoracic Joints. J. Orthop. Sport.
354 Phys. Ther. 2014;44(9):636-A8. doi:10.2519/jospt.2014.5339
- 355 18. Ludewig PM, Borstad JD. Effects of a home exercise programme on shoulder pain and
356 functional status in construction workers. Occup Environ Med. 2003;60:841–849.
- 357 19. Ludewig PM, Cook TM. Alterations in shoulder kinematics and associated muscle
358 activity in people with symptoms of shoulder impingement. Phys. Ther. 2000

How do you stretch pectoralis minor?

- 359 Mar;80(3):276–91.
- 360 20. Matias R, Pascoal AG. The unstable shoulder in arm elevation: A three-dimensional
361 and electromyographic study in subjects with glenohumeral instability. Clin. Biomech.
362 2006;21:S52–S58. doi:10.1016/j.clinbiomech.2005.09.014
- 363 21. McClure PW, Bialker J, Neff N, Williams G, Karduna A. Shoulder function and 3-
364 dimensional kinematics in people with shoulder impingement syndrome before and
365 after a 6-week exercise program. Phys. Ther. 2004 Sep;84(9):832–48.
- 366 22. McClure PW, Michener L a., Sennett BJ, Karduna AR. Direct 3-dimensional
367 measurement of scapular kinematics during dynamic movements in vivo. J. Shoulder
368 Elb. Surg. 2001;10(3):269–277. doi:10.1067/mse.2001.112954
- 369 23. McClure PW, Michener L a, Karduna AR. Shoulder function and 3-dimensional
370 scapular kinematics in people with and without shoulder impingement syndrome. Phys.
371 Ther. 2006;86(8):1075–1090.
- 372 24. Mell AG, LaScalza S, Guffey P, Ray J, Maciejewski M, Carpenter JE, et al. Effect of
373 rotator cuff pathology on shoulder rhythm. J. Shoulder Elb. Surg. 2005;14(1):S58–S64.
374 doi:10.1016/j.jse.2004.09.018
- 375 25. Muraki T, Aoki M, Izumi T, Fujii M, Hidaka E, Miyamoto S. Lengthening of the
376 Pectoralis Minor Muscle During Passive Shoulder Motions and Stretching Techniques:
377 A Cadaveric Biomechanical Study. Phys. Ther. 2009 Apr 1;89(4):333–341.

How do you stretch pectoralis minor?

- 378 doi:10.2522/ptj.20080248
- 379 26. Shiina T, Nightingale KR, Palmeri ML, Hall TJ, Bamber JC, Barr RG, et al. WFUMB
- 380 Guidelines and Recommendations for Clinical Use of Ultrasound Elastography: Part 1:
- 381 Basic Principles and Terminology. *Ultrasound Med. Biol.* 2015 May;41(5):1126–1147.
- 382 doi:10.1016/j.ultrasmedbio.2015.03.009
- 383 27. Umegaki H, Ikezoe T, Nakamura M, Nishishita S, Kobayashi T, Fujita K, et al. The
- 384 effect of hip rotation on shear elastic modulus of the medial and lateral hamstrings
- 385 during stretching. *Man. Ther.* 2014 Aug 13;7–10. doi:10.1016/j.math.2014.07.016
- 386 28. Umehara J, Ikezoe T, Nishishita S, Nakamura M, Umegaki H, Kobayashi T, et al.
- 387 Effect of hip and knee position on tensor fasciae latae elongation during stretching: An
- 388 ultrasonic shear wave elastography study. *Clin. Biomech.* 2015 Sep;30(10):1056–1059.
- 389 doi:10.1016/j.clinbiomech.2015.09.007
- 390
- 391

How do you stretch pectoralis minor?

Figure 1. Passive shoulder flexion; left figure is Flex30, centre figure is Flex90, and right figure is Flex150.

How do you stretch pectoralis minor?

Figure 2. Shoulder horizontal abduction stretching; A is Hab30, B is Hab90, and C is Hab150.

How do you stretch pectoralis minor?

398

399 **Figure 3.** Scapular retraction stretching; A is Retraction30, B is Retraction90, and C is

400 Retraction150.

401

How do you stretch pectoralis minor?

Figure 4. Multiple comparisons of shear elastic modulus; *: $p < .01$ Hab150 is significantly higher than others except for Hab90, †: $p < .01$ Hab90 is significantly higher than others except for Hab150, ‡: $p < .001$ Hab30 is significantly higher than Flex150 and Retraction150.

How do you stretch pectoralis minor?

407 **Table I** Reliability of shear elastic modulus.

Measurement position	ICC _{1,3}	95% CI
Rest	0.99	0.94–1.00
Flex30	0.95	0.79–1.00
Flex90	0.94	0.74–0.99
Flex150	0.97	0.86–1.00
Hab30	0.90	0.55–0.99
Hab90	0.95	0.77–0.99
Hab150	0.99	0.94–1.00
Retraction30	0.95	0.78–0.99
Retraction90	0.95	0.76–0.99
Retraction150	0.93	0.69–0.99

408 ICC: intraclass correlation coefficient, 95% CI: 95% confidence interval

409

How do you stretch pectoralis minor?

410 **Table II** Shear elastic modulus of pectoralis minor muscle in measurement positions.

Measurement position	shear elastic modulus mean value $\pm$ SD (kPa)	95% CI	Comparison with Rest (P value)
Rest	12.5 $\pm$ 2.6	11.3–13.7	
Flex30	12.8 $\pm$ 4.3	10.8–14.8	.99
Flex90	10.3 $\pm$ 3.0	9.0–11.7	.54
Flex150	18.0 $\pm$ 5.8	15.3–20.7	.02
Hab30	31.1 $\pm$ 7.0	27.9–34.4	<.001
Hab90	56.7 $\pm$ 19.1	47.9–65.6	<.001
Hab150	58.7 $\pm$ 20.0	49.5–68.0	<.001
Retraction30	27.8 $\pm$ 13.4	21.6–34.0	.003
Retraction90	24.0 $\pm$ 12.2	18.4–29.7	.007
Retraction150	19.3 $\pm$ 7.7	15.7–22.8	.02

411 SD: standard deviation, 95% CI: 95% confidence interval

412