TITLE:

Acute effects of stretching on passive properties of human gastrocnemius muscle-tendon unit: Analysis of differences between hold-relax and static stretching

AUTHOR(S):

Nakamura, Masatoshi; Ikezoe, Tome; Tokugawa, Takahiro; Ichihashi, Noriaki

CITATION:

Nakamura, Masatoshi ...[et al]. Acute effects of stretching on passive properties of human gastrocnemius muscle-tendon unit: Analysis of differences between hold-relax and static stretching. Journal of Sport Rehabilitation 2015, 24(3): 286-292

ISSUE DATE:

2015-08-01

URL:

<http://hdl.handle.net/2433/231321>

RIGHT:

Accepted author manuscript version reprinted, by permission, from Journal of Sport Rehabilitation, 2015, 24 (3): 286-292, <https://doi.org/10.1123/jsr.2014-0164>. © Human Kinetics, Inc. The full-text file will be made open to the public on 01 August 2015 in accordance with publisher's 'Terms and Conditions for Self-Archiving'. This is not the published version. Please cite only the published version. この論文は出版社版ではありません。引用の際には出版社版をご確認ください。

- 1 **Acute effects of stretching on passive properties of human gastrocnemius muscle–tendon unit:**
- 2 **Analysis of differences between hold–relax and static stretching**
- 3

4 **Abstract**

5 **Context:** Hold–relax stretching (HRS) and static stretching (SS) are commonly used to
6 increase the range of motion (ROM) of a joint and to decrease muscle stiffness. However,
7 whether there are differences between acute effects of HRS and SS on end ROM, passive
8 torque, and muscle stiffness is unclear. In addition, any differences between the mechanisms
9 by which HRS and SS lead to an increase in end ROM are also unclear.

10 **Objective:** The purpose of this study was to compare the acute effects of HRS and SS on the
11 passive properties of the gastrocnemius muscle–tendon unit (MTU), end ROM, passive
12 torque, and muscle stiffness in vivo and to investigate the factors involved in increasing end
13 ROM.

14 **Design:** A cross-over experimental design.

15 **Participants:** Thirty healthy men with no history of neuromuscular disease or
16 musculoskeletal injury involving the lower limbs (21.7 ± 1.2 years).

17 **Intervention:** Both HRS and SS of 30 s was repeated four times, lasting a total of 2 min.

18 **Main Outcome Measures:** End ROM, passive torque, and muscle stiffness were measured
19 during passive ankle dorsiflexion using a dynamometer and ultrasonography before and
20 immediately after HRS and SS.

21 **Results:** The results showed that end ROM and passive torque at the end ROM significantly
22 increased immediately after both HRS and SS, whereas muscle stiffness significantly

23 decreased. In addition, the percent change in passive torque at end ROM upon use of the HRS
24 technique was significantly higher than that after use of the SS technique. However, the
25 percent change in muscle stiffness following SS was significantly higher than that with HRS.
26 **Conclusion:** These results suggest that both HRS and SS can effectively decrease muscle
27 stiffness of the gastrocnemius MTU. In addition, these results suggest that HRS induces a
28 change in the passive torque at end ROM, i.e., sensory perception, rather than changing
29 muscle stiffness.

30 Keywords: muscle stiffness, ultrasonography, proprioceptive neuromuscular facilitation
31 stretching, passive torque, stretch tolerance.

32

33 INTRODUCTION

34 Stretching exercises are commonly used in clinical and athletic settings and can be classified
35 as static stretching (SS), dynamic stretching, ballistic stretching, and proprioceptive
36 neuromuscular facilitation (PNF) stretching. PNF stretching and SS are the most popular type
37 of exercises¹. SS is a stretching technique in which the target muscle is elongated and held at
38 the lengthened position for a certain period of time. Many studies have reported that joint
39 range of motion (ROM) increases immediately after SS²⁻⁴.

40

41 PNF stretching techniques, which involve hold-relax stretching (HRS), are based on
42 the work of Voss et al⁵. **HRS is a stretching technique comprising a combination of SS and**
43 **isometric contraction of the agonist muscle performed in the elongated position.** The target
44 muscle is stretched for a certain period. An isometric contraction at the lengthened position is
45 then performed, followed by another set of SS^{6,7}. Similar to SS, many studies have reported
46 that ROM increases immediately after HRS⁶⁻⁸. **In addition, recent studies regarding acute**
47 **effects have reported that HRS is more effective than SS for increasing ROM⁹⁻¹¹.** Many of the
48 previous studies have used ROM as an outcome measurement of flexibility for stretching
49 exercises.

50

51 However, measurement of ROM is also influenced by psychological factors and

stretch tolerance, such as pain and stretch tolerance, in addition to the viscoelasticity of muscles, tendons, ligaments, and joint capsules^{12, 13}. Therefore, an alternative approach is to measure the passive torque during passive stretching. The overall stiffness of the muscle–tendon unit (MTU) can be estimated by calculating the relationship between passive torque and joint angle¹⁴. Recent studies have shown that gastrocnemius muscle stiffness can be assessed by measuring the displacement of the myotendinous junction (MTJ) during passive ankle dorsiflexion using a dynamometer and ultrasonography and that muscle stiffness decreases after 3–5 min of SS¹⁵⁻¹⁸.

In addition, the passive torque at end ROM was defined as a stretch tolerance^{10, 13, 18, 19}, and an increase in the passive torque at end ROM was defined as a modification of stretch tolerance. Many studies have reported that the passive torque at end ROM increases after SS^{10, 18, 20} and HRS^{6, 8-11}, which suggests that a change in stretch tolerance occurs after stretching.

Previous studies have compared the acute effects of HRS and SS on end ROM and stretch tolerance. However, to the best of our knowledge, there has been no study examining the acute effect of HRS on muscle stiffness and comparing the acute effects on muscle stiffness between HRS and SS. Because decreased muscle stiffness can lead to improvements

71 in athletic performance or prevention of injury²¹⁻²³, a clear understanding of the differences in
72 the effects of HRS and SS on muscle stiffness is important in clinical and athletic settings.

73 In addition, any differences between the mechanisms by which HRS and SS lead to
74 an increase in end ROM are also unclear. Therefore, this study aimed to compare the acute
75 effects of HRS and SS on passive properties of gastrocnemius MTU in vivo and to
76 investigate the factors involved in increasing end ROM. We hypothesized that both HRS and
77 SS could increase end ROM, but that the underlying mechanisms for these effects may differ,
78 with the effect of SS influenced by the decrease in muscle stiffness and that of HRS
79 influenced by the modified stretch tolerance.

80

81 **METHODS**

82 **Study Design**

83 A cross-over experimental design was used to compare the acute effects of HRS and SS on
84 passive properties of gastrocnemius MTU in vivo and to investigate the factors related to
85 increasing end ROM. All participants visited the laboratory on two occasions separated by at
86 least 1 week but no more than 2 weeks to take into account the influence of the measurements
87 and the minimize the carry over effect. Each participant performed HRS once and SS once
88 but in a random order. The subjects were instructed not to begin any other stretching program
89 during the experimental period. All measurements were performed prior to (PRE) and

90 immediately after (POST) HRS and SS. The subjects were familiarized with the procedure
91 and were instructed to remain relaxed throughout the measurement period.

92

93 **Participants**

94 Thirty men volunteered for this study (age, 21.7 ± 1.2 years; height, 170.0 ± 5.3 cm; body
95 mass, 62.4 ± 7.8 kg). Participants with a history of neuromuscular disease or musculoskeletal
96 injury involving the lower limbs were excluded. All participants were fully informed of the
97 procedures and purpose of the study and gave their written informed consent. This study was
98 approved by the ethics committee.

99

100 **Procedures**

101 **Measurements of end ROM and passive torque**

102 The subjects were instructed to lie in the prone position on a dynamometer table (MYORET
103 RZ-450, Kawasaki Heavy Industries, Kobe, Japan) with their hips securely held in place with
104 an adjustable lap belt (Fig 1). The knee of the dominant leg was kept in full extension, and
105 the foot of the same leg was attached securely to the dynamometer footplate with adjustable
106 lap belts. The ankle was passively dorsiflexed at a constant velocity of $5^\circ/\text{s}$, starting from 30°
107 plantarflexion to end dorsiflexion ROM. In this study, end dorsiflexion ROM was defined as
108 the angle achieved by the joint when the point of discomfort, but not pain was reached¹⁶⁻¹⁸.

109 Passive plantarflexion torque was measured using a dynamometer in a similar manner as end
110 ROM was measured. Passive torque at end ROM was defined as a stretch tolerance, and an
111 increase in the passive torque at end ROM was defined as a modification of stretch tolerance
112 during stretching^{10, 13, 18, 19}.

113

114 **Measurement of muscle stiffness**

115 B-mode ultrasonography (Famio Cube SSA-520A; Toshiba Medical Systems Corporation,
116 Tochigi, Japan) was used to determine the displacement of MTJ of medial gastrocnemius
117 (MG) during passive ankle dorsiflexion. MTJ was identified and visualized as a continuous
118 sagittal plane on the ultrasound image using an 8-MHz linear-array probe. An acoustically
119 reflective marker was placed on the skin under the ultrasound probe to confirm that the probe
120 did not move during the measurements¹⁶. We defined MTJ displacement as the distance
121 between MTJ and an acoustically reflective marker. A custom-made fixation device was used
122 to secure the probe to the skin. Ultrasound images of MTJ were quantified using an open-
123 source digital measurement software (Image J, National Institutes of Health, Bethesda,
124 Maryland, USA). To accurately measure MTJ, it was identified at the inner-most edges of the
125 fascia surrounding the muscle where it fuses with the tendon. MTJ displacement was
126 measured between 0° and 30° of ankle dorsiflexion. According to a previous study²⁴,
127 movement of the dynamometer was stopped at 0° and 30°, and at these angles, the

measurement of MTJ by ultrasonography and that of passive torque by dynamometer were synchronously performed. Muscle stiffness was defined as the value obtained by dividing the change in passive torque (between 0° and 30°) by the MTJ displacement¹⁷.

Surface electromyography

Electromyography (EMG; TeleMyo2400; Noraxon USA, Inc., Scottsdale, AZ, USA) was used to confirm that the subjects were relaxed and to ensure that muscles were inactive during passive dorsiflexion. Surface electrodes (Blue Sensor M, Ambu, Denmark) with a 2.0-cm interelectrode distance were placed on certain portions of muscle bellies of the following muscles: MG, lateral gastrocnemius (LG). The EMG sampling rate was 1500 Hz.

EMG activity was recorded from MG and LG while the subjects were performing an isometric maximum voluntary contraction (MVC). MVC of MG and LG were obtained during maximal isometric plantar flexion the ankle at 0° using the dynamometer. Strong verbal encouragement was provided during the contraction to promote maximal effort. EMG activity was calculated using root mean square (RMS), and full wave rectification was performed using an RMS smoothing algorithm with a window interval of 50 ms. EMG activity recorded during the measurements was expressed as a percentage of MVC.

147 HRS and SS

148 Both HRS and SS were performed using the dynamometer in the prone position with the knee
149 extended, similar to the measurements of end ROM and passive torque. In HRS, initially, the
150 ankle was passively dorsiflexed at a constant velocity of 5°/s, starting from 30° plantar
151 flexion to end dorsiflexion ROM, and was held at the end angle for 15 s. The subjects were
152 then instructed to perform MVC of the plantar flexors for 5 s in the same position. **After this**
153 **contraction, the ankle was held at the end angle for an additional 10 s. After each 30-s HRS**
154 **stretch, the ankle was returned to 30° plantar flexion. This application of HRS for 30 s was**
155 **repeated four times, for a total time of 2 min.**

156

157 During SS, the ankle was passively dorsiflexed, starting from 30° plantar flexion to end
158 dorsiflexion ROM and was held at the end angle for 30 s. This SS technique of 30 s was
159 repeated four times, lasting a total of 2 min. In both HRS and SS techniques, the angles of
160 stretching were the same for each stretching. **In addition, for both the HRS and SS techniques,**
161 **we used constant angle stretching, which is routinely used to stretch MTU^{17, 18, 25}, to**
162 **standardize stretching intensity, which is routinely used to stretch MTU^{17, 18, 25}. We previously**
163 **confirmed that the SS protocol with stretching for more than 2 min significantly decreases**
164 **muscle stiffness²⁶. Therefore, we adopted the SS and HRS protocols with 2-min stretching**
165 **durations in total.**

166

167 **Reliability of the measurements**

168 All measurements were repeated twice on different days to assess test–retest reliability (10

169 healthy men; age, 21.8 ± 1.2 years; height, 172.0 ± 3.4 cm; body mass, 63.2 ± 8.9 kg). The

170 measurements were performed with at least a 1-week interval, but not longer than a 2-week

171 interval, between the two tests.

172

173 **Statistical analyses**

174 SPSS (version 17.0; SPSS Japan INC., Tokyo, Japan) was used for statistical analyses. For all

175 variables, significance of differences between HRS and SS techniques at PRE was assessed

176 using the unpaired t-test. Significant differences between PRE and POST were determined for

177 both the HRS and SS techniques using the paired t-test with Bonferroni correction. In178 addition, the percent change between the PRE and POST conditions was calculated to clarify179 the differences in the effects of HRS and SS on passive properties (end ROM, muscle180 stiffness, and stretch tolerance): percent change = (PRE value – POST value) / (PRE value) ×181 100. Because the Shapiro–Wilk tests showed that the percent change was not normally182 distributed, differences in the rates of change between the HRS and SS groups were

183 determined using the Mann–Whitney U test.

184 Differences were considered statistically significant at an alpha level of $P < 0.05$.

185

186 The reliability of end ROM, passive torque at end ROM, and muscle stiffness
187 measurements were examined using the intra-class correlation coefficient (ICC). On the basis
188 of the reliability coefficients, the standard error of measurement (SEM) was calculated (SEM
189 = SD $\sqrt{1 - \text{ICC}}$ for each measurement. Descriptive data are shown as mean $\pm$ standard
190 deviation (SD).

191

192 **RESULTS**

193 Reliability assessments for end ROM, passive torque at end ROM, and muscle
194 stiffness are shown in Table 1. ICC (1, 1) was between 0.891 and 0.957 ($P < 0.01$), and SEM
195 was between 0.4 and 1.4.

196 Changes in variables between PRE and POST in HRS and SS techniques are shown in
197 Table 2. There were no significant differences in any variable between HRS and SS
198 techniques at PRE. In both techniques, POST values of end ROM and passive torque at end
199 ROM were significantly higher than PRE values. Furthermore, for both techniques the POST
200 value of muscle stiffness was significantly lower than the PRE value.

201 A comparison of the percent changes between PRE and POST conditions induced by
202 the HRS and SS techniques is shown in Table 3. There was no significant difference in the
203 percent change in the end ROM between the HRS and SS techniques. The percent change in

204 muscle stiffness in the SS technique was significantly higher than that in the HRS technique,
205 whereas the percent change in passive torque at the end ROM in the HRS technique was
206 significantly higher than that in the SS technique. There were no significant differences in the
207 PRE and POST EMG activities of MG and LG, which were <2.0% MVC during all the
208 measurements in both techniques (table 2).

209

210 **DISCUSSION**

211 This study investigated the acute effects of 2 min of HRS and SS on passive properties of
212 gastrocnemius MTU. The major findings of this study was that the percent change in passive
213 torque at end ROM in HRS technique was significantly higher than that in SS technique,
214 whereas that in muscle stiffness in SS was significantly higher than that in HRS. These
215 results suggest that HRS affects the stretch tolerance, rather than muscle stiffness, in contrast
216 to SS, which is consistent with our hypothesis. To the best of our knowledge, this is the first
217 report to compare the acute effects of HRS and SS on passive properties of gastrocnemius
218 MTU, including end ROM, muscle stiffness, and stretch tolerance.

219

220 Our results showed that POST value of end ROM was significantly higher than PRE
221 value in both techniques. In addition, POST value of muscle stiffness was significantly lower
222 than PRE values in both techniques. These results suggest that an increase in end ROM is a

223 reflection of a decrease in muscle stiffness, which is consistent with the results of other
224 studies examining the acute effects of SS¹⁵⁻¹⁸. The previous study¹⁸ suggested that the effect
225 of stretching on end ROM may be related to decreases in muscle stiffness and a modification
226 in stretch tolerance. We measured passive torque at end ROM, as a stretch tolerance, and
227 POST value of passive torque at end ROM was significantly higher than PRE value in both
228 techniques. This result suggests that an increase in end ROM is also related to a modification
229 in stretch tolerance, which is consistent with the results of previous studies examining the
230 acute effects of SS^{10, 18, 20} and HRS^{6, 8-11}. Therefore, we concluded that decreased muscle
231 stiffness and a modification in stretch tolerance occurring after 2 min of HRS and SS could
232 contribute to an increase in end ROM¹⁸.

233

234 However, Magnusson et al.¹⁰ reported that 90 s of HRS increased ROM without a
235 corresponding decrease in passive torque, suggesting that the increase in end ROM was
236 because of a modification in stretch tolerance rather than changes in the passive properties of
237 MTU¹⁰. In addition, many other studies^{6, 8, 9, 11} have concluded that the increase in end ROM
238 is predominantly because of a modification in stretch tolerance using HRS durations less than
239 90 s. These results do not agree with our results showing that the decrease in muscle stiffness
240 also contributes toward increasing end ROM. We consider that this discrepancy may be
241 because of differences in HRS duration. In this study, HRS duration was 2 min (four

242 repetitions of a 30-s HRS technique), which was comparatively longer than previous studies⁶,
243 ^{8, 9, 11}. In addition, this discrepancy may be due to differences in the target muscle for HRS,
244 which was the gastrocnemius MTU in our study and the hamstring MTU in the previous
245 study¹⁰. Furthermore, the method used for HRS also may have contributed to the difference in
246 the results of our study and the previous study¹⁰. Specifically, the contraction duration in our
247 study was 20 s (4 times × 5 s), whereas that in the previous study¹⁰ was 6 s. Therefore, in
248 addition to a modification in stretch tolerance, muscle stiffness may also change after HRS
249 technique in this study.

250

251 The decrease in muscle stiffness after SS may be associated with a change in the
252 properties of intramuscular connective tissues rather than muscle fiber lengthening^{16, 17}.
253 Although the detailed mechanism underlying the decrease in muscle stiffness after HRS is
254 not known, the acute effects of HRS on the properties of intramuscular connective tissue,
255 such as endomysium, perimysium, and epimysium, may also contribute to the decrease in
256 muscle stiffness after HRS and SS

257

258 In this study, our results showed that there was no significant difference in the change
259 in end ROM between HRS and SS techniques, which is consistent with previous studies^{27, 28}.
260 In contrast, previous studies^{9-11, 29, 30} have reported that the acute effect of HRS on end ROM

261 was greater than that of SS, which is inconsistent with our results. This discrepancy may be
262 because of differences in target muscles. We examined the effects of HRS and SS on
263 gastrocnemius MTU, whereas previous studies⁹⁻¹¹ examined these effects on hamstring MTU.
264 Further study is required to clarify differences in effects between stretching maneuvers on
265 various target muscles.

266

267 The second major finding of this study was that there was a greater increase in passive
268 torque at end ROM in HRS technique compared with SS technique, whereas there was a
269 greater decrease in muscle stiffness in SS technique compared with HRS technique. We
270 consider that the decrease in muscle stiffness in SS technique was greater than that in HRS
271 technique because of the stretching duration. The subjects were instructed to perform MVC
272 of the plantar flexors for 5 s between stretching maneuvers in HRS technique. Therefore, the
273 target muscle was elongated for a total of 100 s in HRS technique, whereas the muscle was
274 elongated for a total of 120 s in SS technique. The lengthening deformation of intramuscular
275 connective tissue (e.g., endomysium, perimysium, and epimysium) may also contribute to the
276 decrease in muscle stiffness^{16, 17}. In HRS technique, shortened muscle fiber during a
277 voluntary isometric contraction leads to deflection (“slack” in intramuscular connective
278 tissue), which may hamper the decrease in muscle stiffness³¹. Therefore, our findings suggest
279 that the effect on muscle stiffness in HRS is lower than that in SS because of differences in

280 stretch duration between HRS and SS. In addition, previous studies^{32, 33} reported that the
281 tendon stiffness decreases after an isometric contraction. Therefore, there was the possibility
282 that the decrease in tendon stiffness during HRS technique was greater compared with SS
283 technique. Because we did not measure the tendon stiffness in both techniques, further study
284 is needed to clarify the effects of HRS and SS techniques on this outcome. Decreased muscle
285 stiffness can be beneficial in improving athletic performance or preventing injury²¹⁻²³.
286 Therefore, because SS technique might be more beneficial in in improving athletic
287 performance or preventing injury, further study is needed to clarify the effects of HRS and SS
288 not only on passive properties, such as end ROM and muscle stiffness, but also on improving
289 performance and preventing injury.

290

291 Our results showed that there was a greater increase in passive torque at end ROM in
292 HRS technique compared with SS technique. Regarding the mechanism of a modification in
293 stretch tolerance, afferent input from muscles and joints during stretching may inhibit signals
294 from nociceptive fibers, which may increase pain thresholds^{8, 10, 11}. In addition, it is possible
295 that sensory afferents affect interneuron release of enkephalins, which could help reduce
296 transmission of nociception in the dorsal horn during stretching, thereby increasing the pain
297 threshold. The analgesic effects achieved by increasing the pain threshold may have altered
298 the stretch tolerance. Our results also suggest that HRS could increase pain thresholds, i.e., a

299 modification in stretch tolerance, to greater levels than those achievable with SS. It is
300 possible that the greater modification in stretch tolerance may be because of a voluntary
301 contraction in HRS technique^{8, 10}. Compared with SS, HRS technique, which places stronger
302 loads on MTU by a voluntary contraction, may increase pain thresholds. With respect to the
303 contraction intensity in HRS, a previous study³⁵ suggested that max isometric contractions
304 may not be required for firing sensory afferents or for inducing the anti-nociceptive signals.
305 Therefore, further study is required to more closely examine contraction intensity.

306

307 This study had some limitations. First, the examiner performing the measurements
308 was not blinded to the groups. Second, we examined only the acute effects of SS and HRS on
309 the passive properties. Thus, we did not examine the prolonged effects after more than a few
310 days or the effects of a stretching training program that lasts several weeks. Therefore, the
311 results may not apply to long-term stretching programs.

312

313 **Conclusion**

314 Our results suggest that both HRS and SS can increase end ROM, which may be
315 because of the decreases in the muscle stiffness and modified stretch tolerance during the
316 stretch application. In addition, compared to SS, HRS may have a greater effect on the
317 alteration of stretch tolerance rather than the decrease in muscle stiffness.

318

319 **Acknowledgment**

320 This work was supported by a Grant-in-Aid for Japan Society for the Promotion of Science

321 (JSPS) Fellows.

322

323 References

- 324 **1.** Haff GG. Roundtable Discussion: Flexibility Training. *Strength and Conditioning*
325 *Journal*. 2006;28(2):64-85.

- 326 **2.** O'Sullivan K, Murray E, Sainsbury D. The effect of warm-up, static stretching and
327 dynamic stretching on hamstring flexibility in previously injured subjects. *BMC*
328 *Musculoskelet Disord*. 2009;10:37.

- 329 **3.** Hubley CL, Kozey JW, Stanish WD. The effects of static stretching exercises and
330 stationary cycling on range of motion at the hip joint*. *J Orthop Sports Phys Ther*.
331 1984;6(2):104-109.

- 332 **4.** Ryan ED, Beck TW, Herda TJ, et al. Do practical durations of stretching alter muscle
333 strength? A dose-response study. *Med Sci Sports Exerc*. 2008;40(8):1529-1537.

- 334 **5.** Voss DE, Ionta MK, Myers BJ, eds. *Proprioceptive neuromuscular facilitation:*
335 *Patterns and techniques*. 3rd ed: Lipponcott Williams & Wilkins; 1985.

- 336 **6.** Youdas JW, Haefflinger KM, Kreun MK, Holloway AM, Kramer CM, Hollman JH.
337 The efficacy of two modified proprioceptive neuromuscular facilitation stretching
338 techniques in subjects with reduced hamstring muscle length. *Physiother Theory*
339 *Pract*. 2010;26(4):240-250.

- 340 **7.** Spernoga SG, Uhl TL, Arnold BL, Gansneder BM. Duration of Maintained Hamstring
341 Flexibility After a One-Time, Modified Hold-Relax Stretching Protocol. *J Athl Train*.

- 342 2001;36(1):44-48.
- 343 **8.** Azevedo DC, Melo RM, Alves Correa RV, Chalmers G. Uninvolved versus target
344 muscle contraction during contract: relax proprioceptive neuromuscular facilitation
345 stretching. *Phys Ther Sport*. 2011;12(3):117-121.
- 346 **9.** Mitchell UH, Myrer JW, Hopkins JT, Hunter I, Feland JB, Hilton SC. Acute stretch
347 perception alteration contributes to the success of the PNF "contract-relax" stretch. *J*
348 *Sport Rehabil*. 2007;16(2):85-92.
- 349 **10.** Magnusson SP, Simonsen EB, Aagaard P, Dyhre-Poulsen P, McHugh MP, Kjaer M.
350 Mechanical and physical responses to stretching with and without preisometric
351 contraction in human skeletal muscle. *Arch Phys Med Rehabil*. 1996;77(4):373-378.
- 352 **11.** O'Hora J, Cartwright A, Wade CD, Hough AD, Shum GL. Efficacy of static stretching
353 and proprioceptive neuromuscular facilitation stretch on hamstrings length after a
354 single session. *J Strength Cond Res*. 2011;25(6):1586-1591.
- 355 **12.** Sale D, Quinlan J, Marsh E, McComas AJ, Belanger AY. Influence of joint position on
356 ankle plantarflexion in humans. *J Appl Physiol*. 1982;52(6):1636-1642.
- 357 **13.** Weppler CH, Magnusson SP. Increasing muscle extensibility: a matter of increasing
358 length or modifying sensation? *Physical therapy*. 2010;90(3):438-449.
- 359 **14.** Magnusson SP, Simonsen EB, Aagaard P, Kjaer M. Biomechanical responses to
360 repeated stretches in human hamstring muscle in vivo. *Am J Sports Med*.

- 361 1996;24(5):622-628.
- 362 **15.** Kay AD, Blazeovich AJ. Moderate-duration static stretch reduces active and passive
- 363 plantar flexor moment but not Achilles tendon stiffness or active muscle length. *J*
- 364 *Appl Physiol.* 2009;106(4):1249-1256.
- 365 **16.** Morse CI, Degens H, Seynnes OR, Maganaris CN, Jones DA. The acute effect of
- 366 stretching on the passive stiffness of the human gastrocnemius muscle tendon unit. *J*
- 367 *Physiol.* 2008;586(1):97-106.
- 368 **17.** Nakamura M, Ikezoe T, Takeno Y, Ichihashi N. Acute and prolonged effect of static
- 369 stretching on the passive stiffness of the human gastrocnemius muscle tendon unit in
- 370 vivo. *J Orthop Res.* 2011;29(11):1759-1763.
- 371 **18.** Mizuno T, Matsumoto M, Umemura Y. Viscoelasticity of the muscle-tendon unit is
- 372 returned more rapidly than range of motion after stretching. *Scand J Med Sci Sports.*
- 373 2013;23(1):23-30.
- 374 **19.** Gajdosik RL, Allred JD, Gabbert HL, Sonsteng BA. A stretching program increases
- 375 the dynamic passive length and passive resistive properties of the calf muscle-tendon
- 376 unit of unconditioned younger women. *Eur J Appl Physiol.* 2007;99(4):449-454.
- 377 **20.** Halbertsma JP, van Bolhuis AI, Goeken LN. Sport stretching: effect on passive
- 378 muscle stiffness of short hamstrings. *Arch Phys Med Rehabil.* 1996;77(7):688-692.
- 379 **21.** Worrell TW, Smith TL, Winegardner J. Effect of hamstring stretching on hamstring

- 380 muscle performance. *J Orthop Sports Phys Ther.* 1994;20(3):154-159.
- 381 **22.** Shellock FG, Prentice WE. Warming-up and stretching for improved physical
382 performance and prevention of sports-related injuries. *Sports Med.* 1985;2(4):267-278.
- 383 **23.** Bixler B, Jones RL. High-school football injuries: effects of a post-halftime warm-up
384 and stretching routine. *Fam Pract Res J.* 1992;12(2):131-139.
- 385 **24.** Nakamura M, Ikezoe T, Takeno Y, Ichihashi N. Effects of a 4-week static stretch
386 training program on passive stiffness of human gastrocnemius muscle-tendon unit in
387 vivo. *Eur J Appl Physiol.* 2012;112(7):2749-2755.
- 388 **25.** Kay AD, Blazeovich AJ. Reductions in active plantarflexor moment are significantly
389 correlated with static stretch duration. *Eur J Sport Sci.* 2008;8:41-46.
- 390 **26.** Nakamura M, Ikezoe T, Takeno Y, Ichihashi N. Time course of changes in passive
391 properties of the gastrocnemius muscle-tendon unit during 5 min of static stretching.
392 *Man Ther.* 2013;18(3):211-215.
- 393 **27.** Puentedura EJ, Huijbregts PA, Celeste S, et al. Immediate effects of quantified
394 hamstring stretching: hold-relax proprioceptive neuromuscular facilitation versus
395 static stretching. *Physical therapy in sport : official journal of the Association of*
396 *Chartered Physiotherapists in Sports Medicine.* 2011;12(3):122-126.
- 397 **28.** Condon SM, Hutton RS. Soleus muscle electromyographic activity and ankle
398 dorsiflexion range of motion during four stretching procedures. *Physical therapy.*

- 399 1987;67(1):24-30.
- 400 **29.** Maddigan ME, Peach AA, Behm DG. A comparison of assisted and unassisted
401 proprioceptive neuromuscular facilitation techniques and static stretching. *Journal of*
402 *strength and conditioning research / National Strength & Conditioning Association.*
403 2012;26(5):1238-1244.
- 404 **30.** Miyahara Y, Naito H, Ogura Y, Katamoto S, Aoki J. Effects of proprioceptive
405 neuromuscular facilitation stretching and static stretching on maximal voluntary
406 contraction. *Journal of strength and conditioning research / National Strength &*
407 *Conditioning Association.* 2013;27(1):195-201.
- 408 **31.** MacIntosh BR, MacNaughton MB. The length dependence of muscle active force:
409 considerations for parallel elastic properties. *J Appl Physiol.* 2005;98(5):1666-1673.
- 410 **32.** Kay AD, Blazeovich AJ. Isometric contractions reduce plantar flexor moment, Achilles
411 tendon stiffness, and neuromuscular activity but remove the subsequent effects of
412 stretch. *J Appl Physiol.* 2009;107(4):1181-1189.
- 413 **33.** Kubo K, Kanehisa H, Kawakami Y, Fukunaga T. Effects of repeated muscle
414 contractions on the tendon structures in humans. *Eur J Appl Physiol.* 2001;84(1-
415 2):162-166.
- 416

417 Figure 1 Experimental set-up of passive dorsiflexion test

418

419 The ankle of the dominant leg was attached securely to the dynamometer footplate by

420 adjustable lap belts to prevent the heel that moving away from the footplate.

421

422 **Table 1 Reliability assessment for end ROM, passive torque at end ROM, muscle**

423 **stiffness**

	Test 1	Test 2	ICC (1, 1) (95% CI)	Difference ^a (95% CI)	<u>SEM</u>
End ROM	32.1 ± 2.8	32.6 ± 2.5	0.891 (0.642–0.971)	0.5 ± 1.2 (–0.3–1.3)	<u>1.4</u>
passive torque at end ROM	43.2 ± 9.7	42.1 ± 7.9	0.893 (0.648–0.972)	–1.1 ± 4.4 (–4.3–2.0)	<u>0.9</u>
muscle stiffness	38.1 ± 5.2	36.5 ± 4.5	0.957 (0.846–0.989)	–1.6 ± 4.5 (–4.9–1.6)	<u>0.4</u>

424 Data are means ± standard deviation

425 a: Difference = Test 2 value – Test 1 value

426

427 **Table 2 Changes between PRE and POST values in both HRS and SS techniques**

Outcome	Technique	PRE	POST	P value	Difference ^a (95% CI)
End ROM (°)	HRS	33.7 ± 2.7	39.5 ± 3.1**	<u>P < 0.025</u>	<u>5.8±1.5 (5.2-6.3)</u>
	SS	33.8 ± 3.0	39.1 ± 3.2**	<u>P < 0.025</u>	<u>5.3±1.3 (4.8-5.8)</u>
Passive torque at end ROM (Nm)	HRS	41.7 ± 10.7	47.2 ± 10.5**	<u>P < 0.025</u>	<u>5.6±3.5 (4.2-6.9)</u>
	SS	41.5 ± 9.0	43.5 ± 9.9**	<u>P < 0.025</u>	<u>2.0±2.3 (1.1-2.8)</u>
Muscle stiffness (Nm/cm)	HRS	36.1 ± 13.3	31.6 ± 10.2**	<u>P < 0.025</u>	<u>-4.5±4.7 (-6.3- -2.7)</u>
	SS	37.3 ± 17.1	25.3 ± 12.1**	<u>P < 0.025</u>	<u>-12.0±9.2 (-15.4- -8.5)</u>
MG EMG activity (%MVC)	HRS	1.6 ± 0.7	1.6 ± 0.7	P = 0.45	<u>0.0±0.4 (-0.1-0.1)</u>
	SS	1.7 ± 0.7	1.8 ± 0.7	P = 0.62	<u>0.1±0.4 (-0.1-0.2)</u>
LG EMG activity (%MVC)	HRS	1.5 ± 1.0	1.4 ± 0.9	P = 0.94	<u>-0.1±0.4 (-0.2-0.1)</u>
	SS	1.6 ± 0.9	1.7 ± 0.9	P = 0.41	<u>0.0±0.5 (-0.1-0.2)</u>
TA EMG activity (%MVC)	HRS	1.2 ± 0.8	1.1 ± 0.8	P = 0.78	<u>0.0±0.5 (-0.2-0.2)</u>
	SS	1.3 ± 1.0	1.2 ± 0.8	P = 0.20	<u>-0.1±0.6 (-0.3-0.1)</u>

428 **: P < 0.05; significant difference in change between PRE and POST

429 Data are means ± standard deviation

430 a: Difference = POST value – PRE value

431

432 **Table 3 Comparisons of the rates of change between HRS and SS techniques**

433

Rate of change (%)	HRS <u>technique</u>	SS <u>technique</u>	P value
End ROM	17.2 ± 4.4	15.8 ± 4.2	P = 0.26
Passive torque at the end ROM	14.5 ± 11.8	4.7 ± 6.3	P < 0.01
Muscle stiffness	-10.9 ± 10.2	-30.4 ± 14.3	P < 0.01

434

435