TITLE:

High-resolution suborganellar localization of Ca²⁺-binding protein CAS, a novel regulator of CO₂-concentrating mechanism

AUTHOR(S):

Yamano, Takashi; Toyokawa, Chihana; Fukuzawa, Hideya

CITATION:

Yamano, Takashi ...[et al]. High-resolution suborganellar localization of Ca²⁺-binding protein CAS, a novel regulator of CO₂-concentrating mechanism. *Protoplasma* 2018, 255(4): 1015-1022

ISSUE DATE:

2018-07

URL:

<http://hdl.handle.net/2433/231982>

RIGHT:

This is a post-peer-review, pre-copyedit version of an article published in 'Protoplasma'. The final authenticated version is available online at: <https://doi.org/10.1007/s00709-018-1208-2>; The full-text file will be made open to the public on 25 January 2019 in accordance with publisher's 'Terms and Conditions for Self-Archiving'. ; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

[Click here to view linked References](#)

1 **Title:** High-resolution suborganellar localization of Ca²⁺-binding protein CAS, a novel
2 regulator of CO₂-concentrating mechanism

3
4
5 **Authors:** Takashi Yamano, Chihana Toyokawa, and Hideya Fukuzawa

6
7
8
9 **Affiliation:** Graduate School of Biostudies, Kyoto University, Kyoto, 606-8502, Japan

10
11
12
13 **Corresponding author:**

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

9 Hideya Fukuzawa

10 Graduate School of Biostudies, Kyoto University, Kyoto, 606-8502, Japan

11 Phone: +81-75-753-4298

12 FAX: +81-75-753-9228

13 E-mail: fukuzawa@lif.kyoto-u.ac.jp

14 Abstract

15 Many aquatic algae induce a CO₂-concentrating mechanism (CCM) associated with active
16 inorganic carbon transport to maintain high photosynthetic affinity using dissolved inorganic
17 carbon even in low-CO₂ (LC) conditions. In the green alga *Chlamydomonas reinhardtii*, a
18 Ca²⁺-binding protein CAS was identified as a novel factor regulating the expression of CCM-
19 related proteins including bicarbonate transporters. Although previous studies revealed that
20 CAS associates with the thylakoid membrane and changes its localization in response to CO₂
21 and light availability, its detailed localization in the chloroplast has not been examined *in*
22 *vivo*. In this study, high-resolution fluorescence images of CAS fused with a
23 *Chlamydomonas*-adapted fluorescence protein, Clover, were obtained by using a sensitive
24 hybrid detector and an image deconvolution method. In high-CO₂ (5% v/v) conditions, the
25 fluorescence signals of Clover displayed a mesh-like structure in the chloroplast and part of
26 the signals discontinuously overlapped with chlorophyll autofluorescence. The fluorescence
27 signals gathered inside the pyrenoid as a distinct wheel-like structure at 2 h after transfer to
28 LC-light condition, and then localized to the center of the pyrenoid at 12 h. These results
29 suggest that CAS could move in the chloroplast along the thylakoid membrane in response to
30 lowering CO₂ and gather inside the pyrenoid during the operation of the CCM.

31

32 **Key words:** Bicarbonate transporter, Ca²⁺-binding protein, *Chlamydomonas*, CO₂-
33 concentrating mechanism, Pyrenoid

34

35 Introduction

36 Photosynthetic organisms can sense and respond to changes of several environmental factors,
37 such as light, CO₂, temperature, and various nutrient availabilities, to optimize and/or
38 maintain their photosynthetic activity. Among these stresses, the shortage of CO₂ supply
39 impacts many physiological aspects of plants, especially photosynthetic efficiency due to the
40 low affinity of ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) against CO₂. In
41 aquatic environments, CO₂-limiting stress is caused not only by the low catalytic activity of
42 Rubisco but also by the 10,000-fold slower diffusion rate of CO₂ in aquatic conditions than
43 that in atmospheric conditions (Jones 1992). To acclimate to this stress, many aquatic
44 organisms possess a CO₂-concentrating mechanism (CCM), which involves the active
45 transport of inorganic carbon (Ci; CO₂ and HCO₃⁻) and enzymatic conversion between CO₂
46 and HCO₃⁻ to maintain the Ci pool and concentrate CO₂ in the vicinity of Rubisco (Fukuzawa
47 et al. 2012; Wang et al. 2015).

48 Molecular aspects of the eukaryotic CCM have been mainly studied using the eukaryotic
49 green alga *Chlamydomonas reinhardtii* as a model. So far, it was reported that high-light
50 activated 3 (HLA3) and low-CO₂ (LC)-inducible protein A (LCIA) are associated with the
51 HCO₃⁻ transport system, which facilitate HCO₃⁻ uptake from outside of cells to the
52 chloroplast stroma across physiological barriers such as the plasma membrane and
53 chloroplast envelope (Gao et al. 2015; Yamano et al. 2015). HLA3 belongs to a multidrug-
54 resistance-related protein subfamily of the ATP-binding cassette transporter superfamily (Im
55 and Grossman 2001) and localizes to the plasma membrane (Yamano et al. 2015). LCIA
56 belongs to a formate-nitrite transporter family (Mariscal et al. 2006), in which proteins form a
57 pentameric aquaporin-like channel rather than an active transporter (Wang et al. 2009), and
58 localizes to the chloroplast envelope (Wang and Spalding 2014; Yamano et al. 2015).

59 Because simultaneous knockdown or knockout of *HLA3* and *LCIA* causes a dramatic
60 decrease in photosynthetic Ci uptake, and simultaneous overexpression of these genes raised
61 photosynthetic Ci affinity and internal Ci accumulation, HLA3 and LCIA are cooperatively
62 associated with HCO₃⁻ transport to increase the Ci pool in the chloroplast stroma (Duanmu et
63 al. 2009; Wang and Spalding 2014; Gao et al. 2015; Yamano et al. 2015). These LC-
64 inducible proteins are regulated by zinc-containing regulatory protein CCM1/CIA5
65 (Fukuzawa et al. 2001; Xiang et al. 2001; Miura et al. 2004). Recently, we isolated a novel
66 high-CO₂ (HC)-requiring mutant H82 (Wang et al. 2014) and revealed that a Ca²⁺-binding
67 protein, CAS, is also essential for the operation of the CCM by regulating the expression of
68 *HLA3* and *LCIA* (Wang et al. 2016). CAS was initially identified in *Arabidopsis thaliana*

(Han et al. 2003) and was detected in the thylakoid membrane fraction (Nomura et al. 2008). By proteomic analysis and indirect immunofluorescence assays, *Chlamydomonas* CAS was also localized to the thylakoid membrane and especially inside the pyrenoid, which is a prominent structure in the chloroplast of the cells cultured in LC conditions (Wang et al. 2016). In many algae, the pyrenoid develops as a spherical proteinaceous structure surrounded with starch sheathes in the chloroplast. Some of the thylakoid membrane penetrates into the pyrenoid, termed pyrenoid tubules (Ohad et al. 1967), and multiple parallel minitubules are bundled within the pyrenoid tubule (Engel et al. 2015). Recently, *Chlamydomonas* CAS was also detected in the protein fraction of purified eyespot (Trippens et al. 2017). Considering that *Arabidopsis* CAS regulates nuclear-encoded genes related to plant immune responses (Nomura et al. 2012) and that *Chlamydomonas* CAS also regulates nuclear-encoded genes related to the CCM such as *HLA3* and *LCIA* (Wang et al. 2016), CAS-mediated retrograde signaling systems from the chloroplast to the nucleus appear to be conserved during the evolution of the plant lineage. However, the actual function of CAS and its detailed subcellular localization *in vivo* remain to be determined.

In this study, by combination of sensitive hybrid detector system, optimization of imaging parameters, and image deconvolution technique, we revealed distinct localization patterns of CAS in HC and LC conditions at high resolution. This result could help in understanding the function of CAS associated with the retrograde signal regulating stress-responsive genes.

Materials and Methods

Cell culture and growth conditions

Chlamydomonas reinhardtii strain C-9 (photosynthetically WT strain originally provided by the IAM Culture Collection held at Tokyo University, and available from the *Chlamydomonas* Resource Center as strain CC-5098), and transgenic lines were cultured in Tris-acetate-phosphate (TAP) medium for maintenance. For physiological and biochemical experiments, a 5 mL volume of cells were grown in liquid TAP medium for pre-cultivation, and diluted with modified high-salt medium supplemented with 20 mM 3-(N-Morpholino)propanesulfonic acid (HSM) to an OD₇₃₀ of ~0.05. Then, the cells were grown under HC (5% v/v) conditions at 120 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ until midlog phase with OD₇₃₀ of 0.3 to 0.5. For LC induction, HC-acclimated cells were centrifuged at 600 $\times$ g, and pellets were resuspended in 50 mL of fresh HSM medium, and cultured in LC (0.04% v/v) conditions at 120 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ for indicated time periods.

Plasmid construction and transformation

The genomic sequence of *CAS* was amplified by PCR with PrimeSTAR GXL (Takara Bio) using genomic DNA extracted from strain C-9 as a template with forward primer TP-clover-F (5'-TTTGCAGGATGCATATGCAGCTTGCTAACGCTCCT-3') and reverse primer gCAS-clover-R (5'-CGATGACGTCAGATCTCGAGCGGGGGGCGGGCAG-3'). The PCR products were purified and cloned into pOptimized Clover vector (Lauersen et al. 2015) digesting with *Nde*I and *Bg*/II using a SLiCE cloning method (Motohashi 2015). For the introduction of a flexible amino acid linker between CAS and Clover, two synthetic oligo nucleotides, gCAS_clover_linker-F (5'-CCCCCGCTCGAGATCTGGCGGCGCGGCCGCGGGCAGATCTGACGTCATCG-3') and gCAS_clover_linker-R (5'-CGATGACGTCAGATCTGCCCCGCGGCCGCGGCCGCGGCCGAGATCTCGAGCGGGGG-3') was annealed and then cloned into the above plasmid digesting with *Bg*/II using a SLiCE cloning method (18-bp nucleotide sequences encoding flexible linker are shown by underlines). This expression plasmid of CAS-Clover was transformed into the H82 mutant (Wang et al. 2014) by electroporation using a NEPA-21 electroporator (NEPAGENE), as described previously (Yamano et al. 2013). The transformants were incubated at 25°C for 24 h with gentle shaking and illumination of less than 1.5 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and spread over TAP plates containing 30 $\mu\text{g mL}^{-1}$ hygromycin.

Immunoblotting analyses

Extracted total proteins suspended in SDS loading buffer containing 50 mM Tris HCl (pH 8.0), 25% (vol/vol) glycerol, 2% (wt/vol) SDS, and 0.1 M DTT were incubated at 37°C for 30 min and subsequently centrifuged at 13,000 $\times g$ for 5 min. The supernatant was loaded onto an SDS-polyacrylamide gel electrophoresis (SDS/PAGE) gel for the separation of proteins. Next, proteins were transferred to polyvinylidene fluoride (Pall Life Science) membranes using a semidry blotting system. Membranes were blocked with 5% (wt/vol) skim milk powder (Wako) in phosphate-buffered saline (PBS). Blocked membranes were washed with PBS containing 0.1% (vol/vol) Tween 20 (PBS-T) and treated with anti-CAS (1:5,000 dilution) or anti-Histone H3 (1:20,000 dilution) antibodies. To recognize the primary antibody, a horseradish peroxidase-conjugated goat anti-rabbit IgG antibody (Life Technologies) was used as a secondary antibody in a dilution of 1:10,000. After washing with PBS-T, immunoreactive signals were detected using Luminata Crescendo Western HRP

substrate (Merck Millipore) and images were obtained using ImageQuant LAS-4010 (GE Healthcare).

Photosynthetic oxygen evolution

For evaluating the affinity for Ci , the rate of dissolved Ci -dependent photosynthetic O_2 evolution was measured. Cells harvested after growth in HC and LC conditions were suspended in Ci -depleted Hepes- NaOH buffer (pH 7.8) at $10 \mu\text{g mL}^{-1}$ chlorophyll. Photosynthetic O_2 evolution was measured by applying a Clark-type O_2 electrode (Hansatech Instruments), as described previously (Yamano et al. 2008).

Capture of high-resolution fluorescence images

To reduce *Chlamydomonas* cell movement, $2.5 \mu\text{L}$ cells were placed between a coverslip and a thin agarose pad (Skinner et al. 2013), and then 16-bit digital fluorescence images were acquired with oil immersion objective lens (HC PL APO $63\times/1.40$; Leica) using an inverted laser-scanning confocal fluorescence microscope TCS SP8 (Leica) equipped with a sensitive hybrid detector (HyD). CAS-Clover was excited at 488 nm and emission was detected at 500–520 nm. Image scanning was performed with pinhole size of 0.6 Airy units, with z-stack distance of the scan at 150 nm, at a pixel size of 25 nm, and with a line scan speed of 200 Hz. Huygens Essential software (Scientific Volume Imaging B.V.) was used for data processing. Deconvolution of confocal datasets was performed using the point-spread function (PSF) theoretically calculated from the microscopic parameters attached to the data and classic maximum likelihood estimation (CMLE) algorithm (settings: maximum iterations: 100; signal-to-noise: 20; quality criterion: 0.05).

Results

Isolation of transgenic lines expressing CAS-Clover

To examine the subcellular localization of CAS *in vivo*, we generated transgenic lines expressing CAS fused with Clover (CAS-Clover), a *Chlamydomonas*-adapted modified green fluorescence protein (Lauersen et al. 2015). We modified the expression plasmid of CAS-Clover used previously (Wang et al. 2016) by introducing a flexible amino acid linker (Gly-Gly-Ala-Ala-Ala-Gly) between CAS and Clover to minimize interference by the protein fusion (Fig. 1a). This plasmid was used to transform the H82 mutant, from which 960 transformants showing paromomycin resistance were obtained, and nine transformants designated as CL-1–CL-9 showing fluorescence signals derived from CAS-Clover inside the

pyrenoid were screened. By immunoblotting analysis using an anti-CAS antibody, a band of approximately 63 kDa corresponding to the predicted size of the CAS-Clover fusion protein was detected (Fig. 1b). Among these transformants, strain CL-2 showed the strongest fluorescence signal and was selected for further analyses. The values of maximum O₂-evolving activity ($V_{\max}$) and $K_{0.5}$ (Ci), the Ci concentration required for half $V_{\max}$, of CL-2 were similar to those of wild-type (WT) cells (Fig. 1c), indicating that decreased photosynthetic Ci-affinity of H82 was complemented by expressing the CAS-Clover.

High-resolution suborganellar localization of CAS-Clover *in vivo*

High-resolution fluorescence images of the CL-2 cells expressing CAS-Clover were obtained using the combination of a sensitive hybrid detector and an image deconvolution technique. In HC conditions, the fluorescence signals were distributed across the entire chloroplast and several punctuate spots were also observed (Fig. 2a). By defocusing of confocal images, fluorescence signals displayed a mesh-like structure, and part of the signals discontinuously overlapped with chlorophyll autofluorescence (Fig. 2b). Considering that CAS was detected in the fraction enriched with the thylakoid membrane (Wang et al. 2016), CAS could be not uniformly but discontinuously distributed on the thylakoid membrane in HC conditions.

Next, when the cells were shifted from HC to LC conditions, the fluorescence signals were detected inside the pyrenoid as a distinct wheel-like structure at 2 h (Fig. 3a–c). When we shifted the focus along the z-axis direction, a strong fluorescent spot was also observed in the lateral region of the chloroplast, which overlapped with the region of eyespot observed in a differential interference contrast image (Fig. 3a). Although the autofluorescence signals of the eyespot were detected in the WT cells, their signal intensities were significantly weaker than that of CL-2 cells with the same microscopic conditions (Fig. 3b), indicating that the fluorescence signals of the eyespot region in CL-2 cells were mostly derived from CAS-Clover. By defocusing of confocal images in the pyrenoid region, the wheel-like structure consisting of several fibers were clearly observed (Fig. 3c). Inside the developed pyrenoid, chlorophyll autofluorescence were hardly detected (Fig. 3a), which was consistent with a previous report (Uniacke and Zerges 2007). This is possibly because the mean diameter of the pyrenoid tubule is very thin at 107 ± 26 nm (Engel et al. 2015), or the amount of chlorophyll could be much decreased in the pyrenoid tubules. By enhancing the contrast of fluorescence, thin fibers were observed, which could be derived from the structure of the pyrenoid tubules (Fig. 3e). In LC conditions after 12 h, the wheel-like structure had almost disappeared, and CAS-Clover was localized to the center of the pyrenoid (Fig. 3f).

Considering that part of the thylakoid membrane, termed the pyrenoid tubules, penetrates into the pyrenoid and fuses at the center of the pyrenoid, forming a knotted core (Engel et al. 2015; Meyer et al. 2016) and that relocation of CAS was not associated with *de novo* protein synthesis (Wang et al. 2016), dispersed CAS-Clover in the chloroplast in HC conditions could move and gather into the pyrenoid along the thylakoid membranes during CCM induction.

Discussion

In this study, we determined suborganellar localization of CAS based on fluorescence images of functional CAS-Clover *in vivo* at high resolution. CAS showed distinct different localization patterns between HC and LC conditions. Dispersed localization of CAS-Clover in HC conditions changed to a wheel-like structure in LC conditions at 2 h and aggregated inside the pyrenoid at 12 h. In particular, this wheel-like localization of CAS-Clover was clearly observed for the first time in this study, strengthening the hypothesis that CAS gathers inside the pyrenoid along the pyrenoid tubules during the operation of the CCM (Wang et al. 2016). Although the relocation of CAS in the chloroplast and its importance for regulation of the CCM has been proposed, it remains unclear how CAS moves along thylakoid membranes.

One possible mechanism is posttranslational modification. Other CCM-related proteins, such as LCIB and CAH3, also change their localization in response to CO₂ availability and undergo phosphorylation when CO₂ availability is limiting (Blanco-Rivero et al. 2012; Yamano et al. 2010). LCIB is an indispensable factor in the CCM and is observed as dispersed speckles in the chloroplast in HC conditions, but changes its localization as a ring-like structure in the vicinity of the pyrenoid in the LC-adapted cells (Yamano et al. 2010), which is distinctly different from the CAS localization pattern. Because *de novo* protein synthesis inhibits the relocation of LCIB (Yamano et al. 2014), but does not affect that of CAS (Wang et al. 2016), the regulatory mechanism of relocation could be different between these proteins. An α -type carbonic anhydrase, CAH3, is shown to be associated with dehydration of HCO₃⁻ to CO₂ within the lumen of pyrenoid tubules (Karlsson et al. 1998). Although CAH3 is associated with the donor side of PSII in the stroma of thylakoid membranes in HC conditions, CAH3 is partly concentrated in the pyrenoid tubules, which does not contain PSII, to provide CO₂ to Rubisco in LC conditions (Blanco-Rivero et al. 2012). Moreover, LCI5/EPYC1 was the first reported protein phosphorylated in response to CO₂-limiting conditions (Turkina et al. 2006). LCI5/EPYC1 is colocalized with Rubisco in

the pyrenoid matrix and assists in the formation of the pyrenoid and the packing of Rubisco in the pyrenoid in LC conditions by linking with Rubisco (Mackinder et al. 2016). In *Arabidopsis*, it is reported that a light-dependent thylakoid protein kinase STN8 phosphorylates a stroma-exposed Thr380 residue of CAS (flanking sequence is SGTKFLP and phosphorylated Threonine is underlined; Vainonen et al. 2008), which is also conserved as Thr370 (flanking sequence is TSTRRLP and putative phosphorylated Threonine is underlined) in *Chlamydomonas* CAS. Based on these results, phosphorylation could be an important factor to regulate the relocation and/or function of CCM-related proteins. Identifying kinases, phosphorylation sites, and obtaining high-resolution images of these proteins could lead to a better understanding of the regulatory mechanism of suborganellar protein relocation.

Another possible mechanism is the structural dynamics of thylakoid membranes. CAS has a hydrophobic sequence that separates the protein sequence into an N-terminus with a Ca²⁺-binding region and a C-terminus with a rhodanese-like domain, and it is thought that CAS anchors to the thylakoid membrane via the hydrophobic sequence (Wang et al. 2016). A recent study revealed that both the structural stability and flexibility of thylakoid membranes is essential for dynamic protein reorganization (Iwai et al. 2014). It is possible that CAS also moves along with the membrane dynamics, although directional movement of the thylakoid membrane from dispersed chloroplast region into the pyrenoid and *vice versa* is unknown.

Recently, CAS was detected in a purified fraction of the *Chlamydomonas* eyespot and also involved in regulating the positive phototactic response under continuous illumination (Trippens et al. 2017). Consistent with this result, we first observed that the fluorescence signal of CAS-Clover overlapped with the eyespot *in vivo*. Ca²⁺ influx through the channel rhodopsins in the eyespot region play an important role for the regulation of phototactic behavior, but the primary Ca²⁺ sensing mechanism is unknown. Using our knockout mutant H82, the regulatory roles of CAS associated with the positive phototactic response could be more clarified.

It has become clear that the pyrenoid is important not only for CO₂ fixation but also for the regulation of the CCM (Meyer et al. 2017; Mitchell et al. 2017). So far, hundreds of proteins with unknown function have been identified in the purified pyrenoid (Mackinder et al. 2016), and there could be other CCM-related proteins that could relocate in the chloroplast in response to the CO₂ availability as reported previously (Yamano et al. 2010). Further screening of mutants showing aberrant localization patterns of these proteins could lead to understanding the regulatory mechanism of suborganellar relocation in response to

environmental stresses (Yamano et al. 2014). Obtaining high-resolution images described in this study could be useful for observing the suborganellar localization of proteins, especially for ones localized in small compartments such as the pyrenoid in a single cell.

Acknowledgments

This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grants 16H04805 (to H.F.) and 16K07399 (to T.Y.) and the Japan Science and Technology Agency Advanced Low Carbon Technology Research and Development Program (to H.F.).

References

- Blanco-Rivero A, Shutova T, Roman MJ, Villarejo A, Martinez F (2012)** Phosphorylation controls the localization and activation of the luminal carbonic anhydrase in *Chlamydomonas reinhardtii*. PLoS One 7:e49063 doi:10.1371/journal.pone.0049063
- Duanmu D, Miller AR, Horken KM, Weeks DP, Spalding MH (2009)** Knockdown of limiting-CO₂-induced gene HLA3 decreases HCO₃⁻ transport and photosynthetic Ci affinity in *Chlamydomonas reinhardtii*. Proc Natl Acad Sci USA 106:5990–5995
- Engel BD, Schaffer M, Cuellar LK, Villa E, Plitzko JM, Baumeister W (2015)** Native architecture of the *Chlamydomonas* chloroplast revealed by *in situ* cryo-electron tomography. eLife 4:e04889 doi:10.7554/eLife.04889
- Fukuzawa H, Miura K, Ishizaki K, Kucho K, Saito T, Kohinata T, Ohyama K (2001)** *CcmI*, a regulatory gene controlling the induction of a carbon-concentrating mechanism in *Chlamydomonas reinhardtii* by sensing CO₂ availability. Proc Natl Acad Sci USA 98:5347–5352
- Fukuzawa, H, Ogawa T, Kaplan A (2012)** The uptake of CO₂ by cyanobacteria and microalgae. In: Eaton-Rye JJ, Tripathy BC, Sharkey TD (eds) Photosynthesis "plastid biology, energy conversion and carbon assimilation." Springer, Advances in Photosynthesis and Respiration 34:625–650
- Gao H, Wang Y, Fei X, Wright DA, Spalding MH (2015)** Expression activation and functional analysis of HLA3, a putative inorganic carbon transporter in *Chlamydomonas reinhardtii*. Plant J 82:1–11
- Han S, Tang R, Anderson LK, Woerner TE, Pei ZM. (2003)** A cell surface receptor mediates extracellular Ca²⁺ sensing in guard cells. Nature 425:196–200

- 305 **Im CS, Grossman AR** (2002) Identification and regulation of high light-induced genes in
306 *Chlamydomonas reinhardtii*. Plant J 30:301-313
- 307 **Iwai M, Yokono M, Nakano A** (2014) Visualizing structural dynamics of thylakoid
308 membranes. Sci Rep 4:3768 doi:10.1038/srep03768
- 309 **Jones HG** (1992) Plants and microclimate: A quantitative approach to environmental plant
310 physiology, 2nd ed. Cambridge University Press: Cambridge, UK.
- 311 **Karlsson J, Clarke AK, Chen ZY et al** (1998) A novel alpha-type carbonic anhydrase
312 associated with the thylakoid membrane in *Chlamydomonas reinhardtii* is required for
313 growth at ambient CO₂. EMBO J 17:1208-1216
- 314 **Lauersen KJ, Kruse O, Mussnug JH** (2015) Targeted expression of nuclear transgenes in
315 *Chlamydomonas reinhardtii* with a versatile, modular vector toolkit. Appl Microbiol
316 Biotechnol 99:3491–3503
- 317 **Mackinder LCM, Meyer M, Mettler-Altmann T et al** (2016) A repeat protein links
318 Rubisco to form the eukaryotic carbon-concentrating organelle. Proc Natl Acad Sci
319 USA 113:5958–5963
- 320 **Mariscal V, Moulin P, Orsel M, Miller AJ, Fernandez E, Galvan A** (2006) Differential
321 regulation of the *Chlamydomonas* *Nar1* gene family by carbon and nitrogen. Protist
322 157:421–433
- 323 **Meyer MT, McCormick AJ, Griffiths H** (2016) Will an algal CO₂-concentrating
324 mechanism work in higher plants? Curr Opin Plant Biol 31:181–188
- 325 **Meyer MT, Whittaker C, Griffiths H** (2017) The algal pyrenoid: key unanswered
326 questions. J Exp Bot 68:3739-3749
- 327 **Mitchell MC, Metodieva G, Metodiev MV, Griffiths H, Meyer MT** (2017) Pyrenoid loss
328 impairs carbon-concentrating mechanism induction and alters primary metabolism in
329 *Chlamydomonas reinhardtii*. J Exp Bot 68:3891-3902
- 330 **Miura K, Yamano T, Yoshioka S et al** (2004) Expression profiling-based identification of
331 CO₂-responsive genes regulated by CCM1 controlling a carbon-concentrating
332 mechanism in *Chlamydomonas reinhardtii*. Plant Physiol 135:1595–1607
- 333 **Motohashi K** (2015) A simple and efficient seamless DNA cloning method using SLiCE
334 from *Escherichia coli* laboratory strains and its application to SLiP site-directed
335 mutagenesis. BMC Biotechnol 15:47 doi:10.1186/s12896-015-0162-8
- 336 **Nomura H, Komori T, Kobori M, Nakahira Y, Shiina T** (2008) Evidence for chloroplast
337 control of external Ca²⁺-induced cytosolic Ca²⁺ transients and stomata closure. Plant J
338 53:988–998

- 339 **Nomura H, Komori T, Uemura S et al** (2012) Chloroplast-mediated activation of plant
immune signaling in *Arabidopsis*. Nat Commun 3:926 doi:10.1038/ncomms1926
- 340
- 341 **Ohad I, Siekevitz P, Palade GE** (1967) Biogenesis of chloroplast membranes. I. Plastid
dedifferentiation in a dark-grown algal mutant (*Chlamydomonas reinhardtii*). J Cell Biol
35:521–552
- 342
- 343
- 344 **Skinner SO, Sepúlveda LA, Xu H, Golding I** (2013) Measuring mRNA copy number in
individual *Escherichia coli* cells using single-molecule fluorescent *in situ* hybridization.
Nat Protoc 8:1100–1113
- 345
- 346
- 347 **Trippens J, Reißweber T, Kreimer G** (2017) The chloroplast calcium sensor protein
CAS affects phototactic behaviour in *Chlamydomonas reinhardtii* (Chlorophyceae) at
low light intensities. Phycologia 56:261–270
- 348
- 349
- 350 **Turkina MV, Blanco-Rivero A, Vainonen JP, Vener AV, Villarejo A** (2006) CO₂
limitation induces specific redox-dependent protein phosphorylation in *Chlamydomonas*
reinhardtii. Proteomics 6:2693–2704
- 351
- 352
- 353 **Uniacke J and Zerges W** (2007) Photosystem II assembly and repair are differentially
localized in *Chlamydomonas*. Plant Cell 19:3640–3654
- 354
- 355 **Vainonen JP, Sakuragi Y, Stael S et al** (2008) Light regulation of CaS, a novel
phosphoprotein in the thylakoid membrane of *Arabidopsis thaliana*. FEBS J 275:1767–
1777
- 356
- 357
- 358 **Wang L, Yamano T, Kajikawa M, Hirono M, Fukuzawa H** (2014) Isolation and
characterization of novel high-CO₂-requiring mutants of *Chlamydomonas reinhardtii*.
Photosynth Res 121:175–184
- 359
- 360
- 361 **Wang L, Yamano T, Takane S et al** (2016) Chloroplast-mediated regulation of CO₂-
concentrating mechanism by Ca²⁺-binding protein CAS in the green alga
Chlamydomonas reinhardtii. Proc Natl Acad Sci USA 113:12586–12591
- 362
- 363
- 364 **Wang Y, Huang Y, Wang J et al** (2009) Structure of the formate transporter FocA reveals a
pentameric aquaporin-like channel. Nature 462:467–472
- 365
- 366 **Wang Y, Spalding MH** (2014) Acclimation to very low CO₂: Contribution of limiting CO₂
inducible proteins, LCIB and LCIA, to inorganic carbon uptake in *Chlamydomonas*
reinhardtii. Plant Physiol 166:2040–2050
- 367
- 368
- 369 **Wang Y, Stessman DJ, Spalding MH** (2015) The CO₂ concentrating mechanism and
photosynthetic carbon assimilation in limiting CO₂: how *Chlamydomonas* works against
the gradient. Plant J 82:429–448
- 370
- 371

- 372 **Xiang Y, Zhang J, Weeks DP** (2001) The *Cia5* gene controls formation of the carbon
373 concentrating mechanism in *Chlamydomonas reinhardtii*. Proc Natl Acad Sci USA
374 98:5341–5346
- 375 **Yamano T, Miura K, Fukuzawa H** (2008) Expression analysis of genes associated with the
376 induction of the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. Plant
377 Physiol 147:340–354
- 378 **Yamano T, Tsujikawa T, Hatano K, Ozawa S, Takahashi Y, Fukuzawa H** (2010) Light
379 and low-CO₂-dependent LCIB-LCIC complex localization in the chloroplast supports
380 the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. Plant Cell Physiol
381 51:1453–1468
- 382 **Yamano T, Iguchi H, Fukuzawa H** (2013) Rapid transformation of *Chlamydomonas*
383 *reinhardtii* without cell-wall removal. J Biosci Bioeng 115:691–694
- 384 **Yamano T, Asada A, Sato E, Fukuzawa H** (2014) Isolation and characterization of mutants
385 defective in the localization of LCIB, an essential factor for the carbon-concentrating
386 mechanism in *Chlamydomonas reinhardtii*. Photosynth Res 121:193–200
- 387 **Yamano T, Sato E, Iguchi H, Fukuda Y, Fukuzawa H** (2015) Characterization of the
388 cooperative bicarbonate uptake into chloroplast stroma in the green alga
389 *Chlamydomonas reinhardtii*. Proc Natl Acad Sci USA 112:7315–7320
- 390

Figure legends

Fig. 1 Isolation of transgenic lines expressing CAS-Clover. (a) A schematic illustration of the expression plasmid of CAS-Clover. The plasmid was constructed based on the pOptimized Clover vector (Lauersen et al. 2015). Translation start (ATG) and stop (TAA) sites are shown. The genomic sequence of *CAS* is placed at the downstream of $P_{A/R}$, *HSP70A/RBCS2* tandem promoter, which is followed by first intron of *RBCS2*. The nucleotide acid sequence GGCGGCGCGGCCGCGGGC encoding the amino acid sequence Gly-Gly-Ala-Ala-Ala-Gly represents a synthetic flexible linker between CAS and Clover. The expression of CAS-Clover is terminated by the T_{RBCS2} , 3'-untranslated region of *RBCS2*. Restriction enzyme sites for cloning of *CAS* (*NdeI* and *BglII*) and for insertion check of the flexible linker (*NotI*) are shown. (b) Accumulation of CAS and CAS-Clover fusion protein in wild-type (WT), H82, and transformants (CL strains). Cells were grown in low-CO₂ (LC) conditions for 12 h. Histone H3 was used as a loading control. (c) Maximum photosynthetic O₂-evolving activity (V_{max} ; left) and inorganic carbon (Ci) affinity (right) of WT, H82, and CL-2 cells grown in LC conditions for 12 h. Photosynthetic O₂-evolving activity was measured in externally dissolved Ci concentrations at pH 7.8, and the $K_{0.5}$ (Ci), the Ci concentrations required for half V_{max} , were calculated. Data in all experiments are mean values $\pm$ standard deviation from three biological replicates. * $P < 0.001$ by Student's t test.

Fig. 2 Fluorescence signals derived from CAS-Clover in high-CO₂ (HC) conditions. (a) CL-2 cells were adapted to HC conditions. Defocused images +1.0 μ m from the focal plane are shown in the bottom row. Each image is placed with the flagella facing upward on the panel. DIC, differential interference contrast image. Scale bar, 2 μ m. (b) Enlarged fluorescence images of the white boxed area in (a) obtained by defocusing the sample from -0.6 to +0.8 μ m from the focal plane. Scale bar, 400 nm.

Fig. 3 Fluorescence signals derived from CAS-Clover in low-CO₂ (LC) conditions. (a) CL-2 cells grown in high-CO₂ conditions were transferred to LC conditions for 2 h. Defocused images +1.0 μ m from the focal plane are shown in the bottom row. Each image is placed with the flagella facing upward on the panel. White arrowheads indicate the eyespot region. DIC, differential interference contrast image. Scale bar, 2 μ m. (b) Autofluorescence image of wild-type (WT) cells grown in LC conditions for 2 h. White arrowheads indicate the eyespot region. Scale bar, 2 μ m. (c) Enlarged fluorescence images of the pyrenoid region by defocusing the sample from -0.6 to +0.8 μ m from the focal plane. Scale bar, 400 nm. (d)

425 Different images of CAS-Clover merged with chlorophyll grown in LC conditions for 2 h. (e)

426 High-contrast image of chlorophyll autofluorescence in the pyrenoid region. Scale bar, 400

427 nm. (f) Fluorescence images derived from CAS-Clover in LC conditions for 12 h. Scale bar,

428 2 μ m.

[Click here to view linked References](#)

1 **Title:** High-resolution suborganellar localization of Ca²⁺-binding protein CAS, a novel
2 regulator of CO₂-concentrating mechanism

3
4
5 **Authors:** Takashi Yamano, Chihana Toyokawa, and Hideya Fukuzawa
6

7
8
9 **Affiliation:** Graduate School of Biostudies, Kyoto University, Kyoto, 606-8502, Japan
10

11
12
13 **Corresponding author:**

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

9 Hideya Fukuzawa
10 Graduate School of Biostudies, Kyoto University, Kyoto, 606-8502, Japan

11 Phone: +81-75-753-4298

12 FAX: +81-75-753-9228

13 E-mail: fukuzawa@lif.kyoto-u.ac.jp

14 Abstract

15 Many aquatic algae induce a CO₂-concentrating mechanism (CCM) associated with active
16 inorganic carbon transport to maintain high photosynthetic affinity using dissolved inorganic
17 carbon even in low-CO₂ (LC) conditions. In the green alga *Chlamydomonas reinhardtii*, a
18 Ca²⁺-binding protein CAS was identified as a novel factor regulating the expression of CCM-
19 related proteins including bicarbonate transporters. Although previous studies revealed that
20 CAS associates with the thylakoid membrane and changes its localization in response to CO₂
21 and light availability, its detailed localization in the chloroplast has not been examined *in vivo*.
22 In this study, high-resolution fluorescence images of CAS fused with a *Chlamydomonas*-
23 adapted fluorescence protein, Clover, were obtained by using a sensitive hybrid detector and
24 an image deconvolution method. In high-CO₂ (5% v/v) conditions, the fluorescence signals
25 of Clover displayed a mesh-like structure in the chloroplast and part of the signals
26 discontinuously overlapped with chlorophyll autofluorescence. The fluorescence signals
27 gathered inside the pyrenoid as a distinct wheel-like structure at 2 h after transfer to LC-light
28 condition, and then localized to the center of the pyrenoid at 12 h. These results suggest that
29 CAS could move in the chloroplast along the thylakoid membrane in response to lowering
30 CO₂ and gather inside the pyrenoid ~~along the pyrenoid tubules, penetrated thylakoid~~
31 ~~membrane into pyrenoid,~~ during the operation of the CCM.

32
33 **Key words:** Bicarbonate transporter, Ca²⁺-binding protein, *Chlamydomonas*, CO₂-
34 concentrating mechanism, Pyrenoid

36 Introduction

37 Photosynthetic organisms can sense and respond to changes of several environmental factors,
38 such as light, CO₂, temperature, and various nutrient availabilities, to optimize and/or
39 maintain their photosynthetic activity. Among these stresses, the shortage of CO₂ supply
40 impacts many physiological aspects of plants, especially photosynthetic efficiency due to the
41 low affinity of ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) against CO₂. In
42 aquatic environments, CO₂-limiting stress is caused not only by the low catalytic activity of
43 Rubisco but also by the 10,000-fold slower diffusion rate of CO₂ in aquatic conditions than
44 that in atmospheric conditions (Jones 1992). To acclimate to this stress, many aquatic
45 organisms possess a CO₂-concentrating mechanism (CCM), which involves the active
46 transport of inorganic carbon (Ci; CO₂ and HCO₃⁻) and enzymatic conversion between CO₂
47 and HCO₃⁻ to maintain the Ci pool and concentrate CO₂ in the vicinity of Rubisco (Fukuzawa
48 et al. 2012; Wang et al. 2015).

49 Molecular aspects of the eukaryotic CCM have been mainly studied using the eukaryotic
50 green alga *Chlamydomonas reinhardtii* as a model. So far, it was reported that high-light
51 activated 3 (HLA3) and low-CO₂ (LC)-inducible protein A (LCIA) are associated with the
52 HCO₃⁻ transport system, which facilitate HCO₃⁻ uptake from outside of cells to the
53 chloroplast stroma across physiological barriers such as the plasma membrane and
54 chloroplast envelope (Gao et al. 2015; Yamano et al. 2015). HLA3 belongs to a multidrug-
55 resistance-related protein subfamily of the ATP-binding cassette transporter superfamily (Im
56 and Grossman 2001) and localizes to the plasma membrane (Yamano et al. 2015). LCIA
57 belongs to a formate-nitrite transporter family (Mariscal et al. 2006), in which proteins form a
58 pentameric aquaporin-like channel rather than an active transporter (Wang et al. 2009), and
59 localizes to the chloroplast envelope (Wang and Spalding 2014; Yamano et al. 2015).

60 Because simultaneous knockdown or knockout of *HLA3* and *LCIA* causes a dramatic
61 decrease in photosynthetic Ci uptake, and simultaneous overexpression of these genes raised
62 photosynthetic Ci affinity and internal Ci accumulation, HLA3 and LCIA are cooperatively
63 associated with HCO₃⁻ transport to increase the Ci pool in the chloroplast stroma (Duanmu et
64 al. 2009; Wang and Spalding 2014; Gao et al. 2015; Yamano et al. 2015). These LC-
65 inducible proteins are regulated by zinc-containing regulatory protein CCM1/CIA5
66 (Fukuzawa et al. 2001; Xiang et al. 2001; Miura et al. 2004). Recently, we isolated a novel
67 high-CO₂ (HC)-requiring mutant H82 (Wang et al. 2014) and revealed that a Ca²⁺-binding
68 protein, CAS, is also essential for the operation of the CCM by regulating the expression of
69 *HLA3* and *LCIA* (Wang et al. 2016). CAS was initially identified in *Arabidopsis thaliana*

(Han et al. 2003) and was detected in the thylakoid membrane fraction (Nomura et al. 2008). By proteomic analysis and indirect immunofluorescence assays, *Chlamydomonas* CAS was also localized to the thylakoid membrane and especially inside the pyrenoid, which is a prominent structure in the chloroplast of the cells cultured in LC conditions (Wang et al. 2016). In many algae, the pyrenoid develops as a spherical proteinaceous structure surrounded with starch sheathes in the chloroplast. Some of the thylakoid membrane penetrates into the pyrenoid, termed pyrenoid tubules (Ohad et al. 1967), and multiple parallel minitubules are bundled within the pyrenoid tubule (Engel et al. 2015). Recently, *Chlamydomonas* CAS was also detected in the protein fraction of purified eyespot (Trippens et al. 2017). Considering that *Arabidopsis* CAS regulates nuclear-encoded genes related to plant immune responses (Nomura et al. 2012) and that *Chlamydomonas* CAS also regulates nuclear-encoded genes related to the CCM such as *HLA3* and *LCIA* (Wang et al. 2016), CAS-mediated retrograde signaling systems from the chloroplast to the nucleus appear to be conserved during the evolution of the plant lineage. However, the actual function of CAS and its detailed subcellular localization *in vivo* remain to be determined.

In this study, by combination of sensitive hybrid detector system, optimization of imaging parameters, and image deconvolution technique, we revealed distinct localization patterns of CAS in HC and LC conditions at high resolution. This result could help in understanding the function of CAS associated with the retrograde signal regulating stress-responsive genes.

Materials and Methods

Cell culture and growth conditions

Chlamydomonas reinhardtii strain C-9 (photosynthetically WT strain originally provided by the IAM Culture Collection held at Tokyo University, and available from the *Chlamydomonas* Resource Center as strain CC-5098), and transgenic lines were cultured in Tris-acetate-phosphate (TAP) medium for maintenance. For physiological and biochemical experiments, a 5 mL volume of cells were grown in liquid TAP medium for pre-cultivation, and diluted with modified high-salt medium supplemented with 20 mM 3-(N-Morpholino)propanesulfonic acid (pH 7.0) to an OD₇₃₀ of ~0.05 ~~for photoautotrophic growth~~. Then, the cells were grown under HC (5% v/v) conditions at 120 μmol photons m⁻² s⁻¹ until midlog phase with OD₇₃₀ of 0.3 to 0.5. For LC induction, HC-acclimated cells were centrifuged at 600 × g, and pellets were resuspended in 50 mL of fresh HSM medium, and cultured in LC (0.04% v/v) conditions at 120 μmol photons m⁻² s⁻¹ for indicated time

~~periods. For all culture conditions, cells were cultured at 25°C with illumination at 120 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$.~~

Plasmid construction and transformation

The genomic sequence of *CAS* was amplified by PCR with PrimeSTAR GXL (Takara Bio) using genomic DNA extracted from strain C-9 as a template with forward primer TP-clover-F (5'-TTTGCAGGATGCATATGCAGCTTGCTAACGCTCCT-3') and reverse primer gCAS-clover-R (5'-CGATGACGTCAGATCTCGAGCGGGGGGCGGGCAG-3'). The PCR products were purified and cloned into pOptimized Clover vector (Lauersen et al. 2015) digesting with *Nde*I and *Bgl*II using a SLiCE cloning method (Motohashi 2015). For the introduction of a flexible amino acid linker between CAS and Clover, two synthetic oligo nucleotides, gCAS_clover_linker-F (5'-CCCCGCTCGAGATCTGGCGGCGCGGCCGCGGGCAGATCTGACGTCATCG-3') and gCAS_clover_linker-R (5'-CGATGACGTCAGATCTGCCCGCGGCCGCGCCGCGCCAGATCTCGAGCGGGGG-3') was annealed and then cloned into the above plasmid digesting with *Bgl*II using a SLiCE cloning method (18-bp nucleotide sequences encoding flexible linker are shown by underlines). This expression plasmid of CAS-Clover was transformed into the H82 mutant (Wang et al. 2014) by electroporation using a NEPA-21 electroporator (NEPAGENE), as described previously (Yamano et al. 2013). The transformants were incubated at 25°C for 24 h with gentle shaking and illumination of less than 1.5 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and spread over TAP plates containing 30 $\mu\text{g mL}^{-1}$ hygromycin.

Immunoblotting analyses

Extracted total proteins suspended in SDS loading buffer containing 50 mM Tris HCl (pH 8.0), 25% (vol/vol) glycerol, 2% (wt/vol) SDS, and 0.1 M DTT were incubated at 37°C for 30 min and subsequently centrifuged at 13,000 $\times g$ for 5 min. The supernatant was loaded onto an SDS-polyacrylamide gel electrophoresis (SDS/PAGE) gel for the separation of proteins. Next, proteins were transferred to polyvinylidene fluoride (Pall Life Science) membranes using a semidry blotting system. Membranes were blocked with 5% (wt/vol) skim milk powder (Wako) in phosphate-buffered saline (PBS). Blocked membranes were washed with PBS containing 0.1% (vol/vol) Tween 20 (PBS-T) and treated with anti-CAS (1:5,000 dilution) or anti-Histone H3 (1:20,000 dilution) antibodies. To recognize the primary antibody, a horseradish peroxidase-conjugated goat anti-rabbit IgG antibody (Life

Technologies) was used as a secondary antibody in a dilution of 1:10,000. After washing with PBS-T, immunoreactive signals were detected using Luminata Crescendo Western HRP substrate (Merck Millipore) and images were obtained using ImageQuant LAS-4010 (GE Healthcare).

Photosynthetic oxygen evolution

For evaluating the affinity for C_i , the rate of dissolved C_i -dependent photosynthetic O_2 evolution was measured. Cells harvested after growth in HC and LC conditions were suspended in C_i -depleted Hepes-NaOH buffer (pH 7.8) at $10 \mu\text{g mL}^{-1}$ chlorophyll. Photosynthetic O_2 evolution was measured by applying a Clark-type O_2 electrode (Hansatech Instruments), as described previously (Yamano et al. 2008).

Capture of high-resolution fluorescence images

To reduce *Chlamydomonas* cell movement, $2.5 \mu\text{L}$ cells were placed between a coverslip and a thin agarose pad (Skinner et al. 2013), and then 16-bit digital fluorescence images were acquired with oil immersion objective lens (HC PL APO $63\times/1.40$; Leica) using an inverted laser-scanning confocal fluorescence microscope TCS SP8 (Leica) equipped with a sensitive hybrid detector (HyD). CAS-Clover was excited at 488 nm and emission was detected at 500–520 nm. Image scanning was performed with pinhole size of 0.6 Airy units, with z-stack distance of the scan at 150 nm, at a pixel size of 25 nm, and with a line scan speed of 200 Hz. Huygens Essential software (Scientific Volume Imaging B.V.) was used for data processing. Deconvolution of confocal datasets was performed using the point-spread function (PSF) theoretically calculated from the microscopic parameters attached to the data and classic maximum likelihood estimation (CMLE) algorithm (settings: maximum iterations: 100; signal-to-noise: 20; quality criterion: 0.05).

Results

Isolation of transgenic lines expressing CAS-Clover

To examine the subcellular localization of CAS *in vivo*, we generated transgenic lines expressing CAS fused with Clover (CAS-Clover), a *Chlamydomonas*-adapted modified green fluorescence protein (Lauersen et al. 2015). We modified the expression plasmid of CAS-Clover used previously (Wang et al. 2016) by introducing a flexible amino acid linker (Gly-Gly-Ala-Ala-Ala-Gly) between CAS and Clover to minimize interference by the protein fusion (Fig. 1a). This plasmid was used to transform the H82 mutant, from which 960

transformants showing paromomycin resistance were obtained, and nine transformants designated as CL-1–CL-9 showing fluorescence signals derived from CAS-Clover inside the pyrenoid were screened. By immunoblotting analysis using an anti-CAS antibody, a band of approximately 63 kDa corresponding to the predicted size of the CAS-Clover fusion protein was detected (Fig. 1b). Among these transformants, strain CL-2 showed the strongest fluorescence signal and was selected for further analyses. The values of maximum O_2 -evolving activity (V_{max}) and $K_{0.5}$ (Ci), the Ci concentration required for half V_{max} , of CL-2 were similar to those of wild-type (WT) cells (Fig. 1c), indicating that decreased photosynthetic Ci-affinity of H82 was complemented by expressing the CAS-Clover.

High-resolution suborganellar localization of CAS-Clover *in vivo*

High-resolution fluorescence images of the CL-2 cells expressing CAS-Clover were obtained using the combination of a sensitive hybrid detector and an image deconvolution technique. In HC conditions, the fluorescence signals were distributed across the entire chloroplast and several punctuate spots were also observed (Fig. 2a). By defocusing of confocal images, fluorescence signals displayed a mesh-like structure, and part of the signals discontinuously overlapped with chlorophyll autofluorescence (Fig. 2b). Considering that CAS was detected in the fraction enriched with the thylakoid membrane (Wang et al. 2016), CAS could be not uniformly but discontinuously distributed on the thylakoid membrane in HC conditions.

Next, when the cells were shifted from HC to LC conditions, the fluorescence signals were detected inside the pyrenoid as a distinct wheel-like structure at 2 h (Fig. 3a–c). When we shifted the focus along the z-axis direction, a strong fluorescent spot was also observed in the lateral region of the chloroplast, which overlapped with the region of eyespot observed in a differential interference contrast image (Fig. 3a). Although the autofluorescence signals of the eyespot were detected in the WT cells, their signal intensities were significantly weaker than that of CL-2 cells with the same microscopic conditions (Fig. 3b), indicating that the fluorescence signals of the eyespot region in CL-2 cells were mostly derived from CAS-Clover. By defocusing of confocal images in the pyrenoid region, the wheel-like structure consisting of several fibers were clearly observed (Fig. 3**cb**). Inside the developed pyrenoid, chlorophyll autofluorescence were hardly detected (Fig. 3a), which was consistent with a previous report (Uniacke and Zerges 2007). This is possibly because the mean diameter of the pyrenoid tubule is very thin at 107 ± 26 nm (Engel et al. 2015), or the amount of chlorophyll could be much decreased in the pyrenoid tubules. By enhancing the contrast of fluorescence, thin fibers were observed, which could be derived from the structure of the

pyrenoid tubules (Fig. 3^{ed}). In LC conditions after 12 h, the wheel-like structure had almost disappeared, and CAS-Clover was localized to the center of the pyrenoid (Fig. 3^{fe}). Considering that part of the thylakoid membrane, termed the pyrenoid tubules, penetrates into the pyrenoid and fuses at the center of the pyrenoid, forming a knotted core (Engel et al. 2015; Meyer et al. 2016) and that relocation of CAS was not associated with *de novo* protein synthesis (Wang et al. 2016), dispersed CAS-Clover in the chloroplast in HC conditions could move and gather into the pyrenoid along the thylakoid membranes during CCM induction.

Discussion

In this study, we determined suborganellar localization of CAS based on fluorescence images of functional CAS-Clover *in vivo* at high resolution. CAS showed distinct different localization patterns between HC and LC conditions. Dispersed localization of CAS-Clover in HC conditions changed to a wheel-like structure in LC conditions at 2 h and aggregated inside the pyrenoid at 12 h. In particular, this wheel-like localization of CAS-Clover was clearly observed for the first time in this study, strengthening the hypothesis that CAS gathers inside the pyrenoid along the pyrenoid tubules during the operation of the CCM (Wang et al. 2016). Although the relocation of CAS in the chloroplast and its importance for regulation of the CCM has been proposed, it remains unclear how CAS moves along thylakoid membranes.

One possible mechanism is posttranslational modification. Other CCM-related proteins, such as LCIB and CAH3, also change their localization in response to CO₂ availability and undergo phosphorylation when CO₂ availability is limiting (Blanco-Rivero et al. 2012; Yamano et al. 2010). LCIB is an indispensable factor in the CCM and is observed as dispersed speckles in the chloroplast in HC conditions, but changes its localization as a ring-like structure in the vicinity of the pyrenoid in the LC-adapted cells (Yamano et al. 2010), which is distinctly different from the CAS localization pattern. Because *de novo* protein synthesis inhibits the relocation of LCIB (Yamano et al. 2014), but does not affect that of CAS (Wang et al. 2016), the regulatory mechanism of relocation could be different between these proteins. An α -type carbonic anhydrase, CAH3, is shown to be associated with dehydration of HCO₃⁻ to CO₂ within the lumen of pyrenoid tubules (Karlsson et al. 1998). Although CAH3 is associated with the donor side of PSII in the stroma of thylakoid membranes in HC conditions, CAH3 is partly concentrated in the pyrenoid tubules, which does not contain PSII, to provide CO₂ to Rubisco in LC conditions (Blanco-Rivero et al. 2012). Moreover, LCI5/EPYC1 was the first reported protein phosphorylated in response to

CO₂-limiting conditions (Turkina et al. 2006). LCI5/EPYC1 is colocalized with Rubisco in the pyrenoid matrix and assists in the formation of the pyrenoid and the packing of Rubisco in the pyrenoid in LC conditions by linking with Rubisco (Mackinder et al. 2016). In *Arabidopsis*, it is reported that a light-dependent thylakoid protein kinase STN8 phosphorylates a stroma-exposed Thr380 residue of CAS (flanking sequence is SGTKFLP and phosphorylated Threonine is underlined; Vainonen et al. 2008), which is also conserved as Thr370 (flanking sequence is TSTRRLP and putative phosphorylated Threonine is underlined) in *Chlamydomonas* CAS. Based on these results, phosphorylation could be an important factor to regulate the relocation and/or function of CCM-related proteins. Identifying kinases, phosphorylation sites, and obtaining high-resolution images of these proteins could lead to a better understanding of the regulatory mechanism of suborganellar protein relocation.

Another possible mechanism is the structural dynamics of thylakoid membranes. CAS has a hydrophobic sequence that separates the protein sequence into an N-terminus with a Ca²⁺-binding region and a C-terminus with a rhodanese-like domain, and it is thought that CAS anchors to the thylakoid membrane via the hydrophobic sequence (Wang et al. 2016). A recent study revealed that both the structural stability and flexibility of thylakoid membranes is essential for dynamic protein reorganization (Iwai et al. 2014). It is possible that CAS also moves along with the membrane dynamics, although directional movement of the thylakoid membrane from dispersed chloroplast region into the pyrenoid and *vice versa* is unknown.

Recently, CAS was detected in a purified fraction of the *Chlamydomonas* eyespot and also involved in regulating the positive phototactic response under continuous illumination (Trippens et al. 2017). Consistent with this result, we first observed that the fluorescence signal of CAS-Clover overlapped with the eyespot *in vivo*. Ca²⁺ influx through the channel rhodopsins in the eyespot region play an important role for the regulation of phototactic behavior, but the primary Ca²⁺ sensing mechanism is unknown. Using our knockout mutant H82, the regulatory roles of CAS associated with the positive phototactic response could be more clarified.

It has become clear that the pyrenoid is important not only for CO₂ fixation but also for the regulation of the CCM (Meyer et al. 2017; Mitchell et al. 2017). So far, hundreds of proteins with unknown function have been identified in the purified pyrenoid (Mackinder et al. 2016), and there could be other CCM-related proteins that could relocate in the chloroplast in response to the CO₂ availability as reported previously (Yamano et al. 2010). Further screening of mutants showing aberrant localization patterns of these proteins could lead to

improved understand the regulatory mechanism of suborganellar relocation in response to environmental stresses (Yamano et al. 2014). Obtaining high-resolution images described in this study could be useful for observing the suborganellar localization of proteins, especially for ones localized in small compartments such as the pyrenoid in a single cell.

Acknowledgments

This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grants 16H04805 (to H.F.) and [16K0739925840109](#) (to T.Y.) and the Japan Science and Technology Agency Advanced Low Carbon Technology Research and Development Program (to H.F.).

References

- Blanco-Rivero A, Shutova T, Roman MJ, Villarejo A, Martinez F (2012)** Phosphorylation controls the localization and activation of the luminal carbonic anhydrase in *Chlamydomonas reinhardtii*. PLoS One 7:e49063 doi:10.1371/journal.pone.0049063
- Duanmu D, Miller AR, Horken KM, Weeks DP, Spalding MH (2009)** Knockdown of limiting-CO₂-induced gene HLA3 decreases HCO₃⁻ transport and photosynthetic Ci affinity in *Chlamydomonas reinhardtii*. Proc Natl Acad Sci USA 106:5990–5995
- Engel BD, Schaffer M, Cuellar LK, Villa E, Plitzko JM, Baumeister W (2015)** Native architecture of the *Chlamydomonas* chloroplast revealed by *in situ* cryo-electron tomography. eLife 4:e04889 doi:10.7554/eLife.04889
- Fukuzawa H, Miura K, Ishizaki K, Kucho K, Saito T, Kohinata T, Ohyama K (2001)** *CcmI*, a regulatory gene controlling the induction of a carbon-concentrating mechanism in *Chlamydomonas reinhardtii* by sensing CO₂ availability. Proc Natl Acad Sci USA 98:5347–5352
- Fukuzawa, H, Ogawa T, Kaplan A (2012)** The uptake of CO₂ by cyanobacteria and microalgae. In: Eaton-Rye JJ, Tripathy BC, Sharkey TD (eds) Photosynthesis "plastid biology, energy conversion and carbon assimilation." Springer, Advances in Photosynthesis and Respiration 34:625–650
- Gao H, Wang Y, Fei X, Wright DA, Spalding MH (2015)** Expression activation and functional analysis of HLA3, a putative inorganic carbon transporter in *Chlamydomonas reinhardtii*. Plant J 82:1–11

- 307 **Han S, Tang R, Anderson LK, Woerner TE, Pei ZM.** (2003) A cell surface receptor
308 mediates extracellular Ca²⁺ sensing in guard cells. *Nature* 425:196–200
- 309 **Im CS, Grossman AR** (2002) Identification and regulation of high light-induced genes in
310 *Chlamydomonas reinhardtii*. *Plant J* 30:301–313
- 311 **Iwai M, Yokono M, Nakano A** (2014) Visualizing structural dynamics of thylakoid
312 membranes. *Sci Rep* 4:3768 doi:10.1038/srep03768
- 313 **Jones HG** (1992) *Plants and microclimate: A quantitative approach to environmental plant*
314 *physiology*, 2nd ed. Cambridge University Press: Cambridge, UK.
- 315 **Karlsson J, Clarke AK, Chen ZY et al** (1998) A novel alpha-type carbonic anhydrase
316 associated with the thylakoid membrane in *Chlamydomonas reinhardtii* is required for
317 growth at ambient CO₂. *EMBO J* 17:1208–1216
- 318 **Lauersen KJ, Kruse O, Mussnug JH** (2015) Targeted expression of nuclear transgenes in
319 *Chlamydomonas reinhardtii* with a versatile, modular vector toolkit. *Appl Microbiol*
320 *Biotechnol* 99:3491–3503
- 321 **Mackinder LCM, Meyer M, Mettler-Altmann T et al** (2016) A repeat protein links
322 Rubisco to form the eukaryotic carbon-concentrating organelle. *Proc Natl Acad Sci*
323 *USA* 113:5958–5963
- 324 **Mariscal V, Moulin P, Orsel M, Miller AJ, Fernandez E, Galvan A** (2006) Differential
325 regulation of the *Chlamydomonas* *Nar1* gene family by carbon and nitrogen. *Protist*
326 157:421–433
- 327 **Meyer MT, McCormick AJ, Griffiths H** (2016) Will an algal CO₂-concentrating
328 mechanism work in higher plants? *Curr Opin Plant Biol* 31:181–188
- 329 **Meyer MT, Whittaker C, Griffiths H** (2017) The algal pyrenoid: key unanswered
330 questions. *J Exp Bot* 68:3739–3749
- 331 **Mitchell MC, Metodieva G, Metodiev MV, Griffiths H, Meyer MT** (2017) Pyrenoid loss
332 impairs carbon-concentrating mechanism induction and alters primary metabolism in
333 *Chlamydomonas reinhardtii*. *J Exp Bot* 68:3891–3902
- 334 **Miura K, Yamano T, Yoshioka S et al** (2004) Expression profiling-based identification of
335 CO₂-responsive genes regulated by CCM1 controlling a carbon-concentrating
336 mechanism in *Chlamydomonas reinhardtii*. *Plant Physiol* 135:1595–1607
- 337 **Motohashi K** (2015) A simple and efficient seamless DNA cloning method using SLiCE
338 from *Escherichia coli* laboratory strains and its application to SLiP site-directed
339 mutagenesis. *BMC Biotechnol* 15:47 doi:10.1186/s12896-015-0162-8

- 340 **Nomura H, Komori T, Kobori M, Nakahira Y, Shiina T** (2008) Evidence for chloroplast
- 341 control of external Ca^{2+} -induced cytosolic Ca^{2+} transients and stomata closure. *Plant J*
- 342 53:988–998
- 343 **Nomura H, Komori T, Uemura S et al** (2012) Chloroplast-mediated activation of plant
- 344 immune signaling in *Arabidopsis*. *Nat Commun* 3:926 doi:10.1038/ncomms1926
- 345 **Ohad I, Siekevitz P, Palade GE** (1967) Biogenesis of chloroplast membranes. I. Plastid
- 346 dedifferentiation in a dark-grown algal mutant (*Chlamydomonas reinhardtii*). *J Cell Biol*
- 347 35:521–552
- 348 **Skinner SO, Sepúlveda LA, Xu H, Golding I** (2013) Measuring mRNA copy number in
- 349 individual *Escherichia coli* cells using single-molecule fluorescent *in situ* hybridization.
- 350 *Nat Protoc* 8:1100–1113
- 351 **Trippens J, Reißweber T, Kreimer G** (2017) The chloroplast calcium sensor protein
- 352 CAS affects phototactic behaviour in *Chlamydomonas reinhardtii* (Chlorophyceae) at
- 353 low light intensities. *Phycologia* 56:261–270
- 354 **Turkina MV, Blanco-Rivero A, Vainonen JP, Vener AV, Villarejo A** (2006) CO_2
- 355 limitation induces specific redox-dependent protein phosphorylation in *Chlamydomonas*
- 356 *reinhardtii*. *Proteomics* 6:2693–2704
- 357 **Uniacke J and Zerges W** (2007) Photosystem II assembly and repair are differentially
- 358 localized in *Chlamydomonas*. *Plant Cell* 19:3640–3654
- 359 **Vainonen JP, Sakuragi Y, Stael S et al** (2008) Light regulation of CaS, a novel
- 360 phosphoprotein in the thylakoid membrane of *Arabidopsis thaliana*. *FEBS J* 275:1767–
- 361 1777
- 362 **Wang L, Yamano T, Kajikawa M, Hirono M, Fukuzawa H** (2014) Isolation and
- 363 characterization of novel high- CO_2 -requiring mutants of *Chlamydomonas reinhardtii*.
- 364 *Photosynth Res* 121:175–184
- 365 **Wang L, Yamano T, Takane S et al** (2016) Chloroplast-mediated regulation of CO_2 -
- 366 concentrating mechanism by Ca^{2+} -binding protein CAS in the green alga
- 367 *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* 113:12586–12591
- 368 **Wang Y, Huang Y, Wang J et al** (2009) Structure of the formate transporter FocA reveals a
- 369 pentameric aquaporin-like channel. *Nature* 462:467–472
- 370 **Wang Y, Spalding MH** (2014) Acclimation to very low CO_2 : Contribution of limiting CO_2
- 371 inducible proteins, LCIB and LCIA, to inorganic carbon uptake in *Chlamydomonas*
- 372 *reinhardtii*. *Plant Physiol* 166:2040–2050

- 373 **Wang Y, Stessman DJ, Spalding MH** (2015) The CO₂ concentrating mechanism and
374 photosynthetic carbon assimilation in limiting CO₂: how *Chlamydomonas* works against
375 the gradient. *Plant J* 82:429–448
- 376 **Xiang Y, Zhang J, Weeks DP** (2001) The *Cia5* gene controls formation of the carbon
377 concentrating mechanism in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA*
378 98:5341–5346
- 379 **Yamano T, Miura K, Fukuzawa H** (2008) Expression analysis of genes associated with the
380 induction of the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant*
381 *Physiol* 147:340–354
- 382 **Yamano T, Tsujikawa T, Hatano K, Ozawa S, Takahashi Y, Fukuzawa H** (2010) Light
383 and low-CO₂-dependent LCIB-LCIC complex localization in the chloroplast supports
384 the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant Cell Physiol*
385 51:1453–1468
- 386 **Yamano T, Iguchi H, Fukuzawa H** (2013) Rapid transformation of *Chlamydomonas*
387 *reinhardtii* without cell-wall removal. *J Biosci Bioeng* 115:691–694
- 388 **Yamano T, Asada A, Sato E, Fukuzawa H** (2014) Isolation and characterization of mutants
389 defective in the localization of LCIB, an essential factor for the carbon-concentrating
390 mechanism in *Chlamydomonas reinhardtii*. *Photosynth Res* 121:193–200
- 391 **Yamano T, Sato E, Iguchi H, Fukuda Y, Fukuzawa H** (2015) Characterization of the
392 cooperative bicarbonate uptake into chloroplast stroma in the green alga
393 *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* 112:7315–7320

Figure legends

Fig. 1 Isolation of transgenic lines expressing CAS-Clover. (a) A schematic illustration of the expression plasmid of CAS-Clover. The plasmid was constructed based on the pOptimized Clover vector (Lauersen et al. 2015). Translation start (ATG) and stop (TAA) sites are shown. The genomic sequence of *CAS* is placed at the downstream of *P_{A/R}*, *HSP70A/RBCS2* tandem promoter, which is followed by first intron of *RBCS2*. The nucleotide acid sequence GGCGGCGCGGCCGCGGGC encoding the amino acid sequence Gly-Gly-Ala-Ala-Ala-Gly represents a synthetic flexible linker between CAS and Clover. The expression of CAS-Clover is terminated by the *T_{RBCS2}*, 3'-untranslated region of *RBCS2*. Restriction enzyme sites for cloning of *CAS* (*NdeI* and *BglII*) and for insertion check of the flexible linker (*NotI*) are shown. (b) Accumulation of CAS and CAS-Clover fusion protein in wild-type (WT), H82, and transformants (CL strains). Cells were grown in low-CO₂ (LC) conditions for 12 h. Histone H3 was used as a loading control. (c) Maximum photosynthetic O₂-evolving activity ($V_{\max}$; left) and inorganic carbon (Ci) affinity (right) of WT, H82, and CL-2 cells grown in LC conditions for 12 h. Photosynthetic O₂-evolving activity was measured in externally dissolved Ci concentrations at pH 7.8, and the $K_{0.5}$ (Ci), the Ci concentrations required for half $V_{\max}$, were calculated. Data in all experiments are mean values $\pm$ standard deviation from three biological replicates. * $P < 0.001$ by Student's t test.

Fig. 2 Fluorescence signals derived from CAS-Clover in high-CO₂ (HC) conditions. (a) CL-2 cells were adapted to HC conditions. Defocused images +1.0 μm from the focal plane are shown in the bottom row. Each image is placed with the flagella facing upward on the panel. DIC, differential interference contrast image. Scale bar, 2 μm . (b) Enlarged fluorescence images of the white boxed area in (a) obtained by defocusing the sample from -0.6 to +0.8 μm from the focal plane. Scale bar, 400 nm.

Fig. 3 Fluorescence signals derived from CAS-Clover in low-CO₂ (LC) conditions. (a) CL-2 cells grown in high-CO₂ conditions were transferred to LC conditions for 2 h. Defocused images +1.0 μm from the focal plane are shown in the bottom row. Each image is placed with the flagella facing upward on the panel. White arrowheads indicate the eyespot. DIC, differential interference contrast image. Scale bar, 2 μm . (b) Autofluorescence image of wild-type (WT) cells grown in LC conditions for 2 h. White arrowheads indicate the eyespot region. Scale bar, 2 μm . (c) Enlarged fluorescence images of the pyrenoid region by defocusing the sample from -0.6 to +0.8 μm from the focal plane. Scale bar, 400 nm. (d)

429 Different images of CAS-Clover merged with chlorophyll grown in LC conditions for 2 h.
430 (~~de~~) High-contrast image of chlorophyll autofluorescence in the pyrenoid region. Scale bar,
431 400 nm. (~~ef~~) Fluorescence images derived from CAS-Clover in LC conditions for 12 h. Scale
432 bar, 2 μ m.

Conflict of Interest:

The authors declare that they have no conflict of interest.

Figure2

[Click here to download Figure Fig.2.tif](#)

Figure3

[Click here to download Figure Fig.3.tif](#)