TITLE:

Technique of Dorsal Transversely Oriented Transposition Flap for Web Reconstruction in Toe Syndactyly Surgery

AUTHOR(S):

Saito, Susumu; Suzuki, Yoshihisa; Suzuki, Shigehiko

CITATION:

Saito, Susumu ...[et al]. Technique of Dorsal Transversely Oriented Transposition Flap for Web Reconstruction in Toe Syndactyly Surgery. Journal of Foot and Ankle Surgery 2015, 54(6): 1119-1123

ISSUE DATE:

2015-11

URL:

<http://hdl.handle.net/2433/232632>

RIGHT:

© 2015. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>; The full-text file will be made open to the public on 01 November 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

Title

A Technique of Dorsal Transversely-oriented Transposition Flap for Web Reconstruction in Toe Syndactyly Surgeries

Abstract

In toe syndactyly, dorsal, or planter commissural flap, in combination with skin grafts, ensures an acceptable result; however, parallel unsightly scars in the longitudinal direction on the dorsum of the toes sometimes fail to satisfy patients' and/or parents of patients' aesthetic expectations. To address this issue, we developed a novel technique using a transversely-oriented transposition flap for web reconstruction, which can spare the dorsal interdigital skin maximally to shift the dorsal scars planterly to be inconspicuous. The design of the flap is simple and uncomplicated surgically. Moreover, the donor site morbidity is minimal, owing to the good healing potential of transverse scars. This technique could be an alternative in web reconstruction of toe syndactylies, especially in cases of high cosmetic priority.

Keywords

flap, polysyndactyly, surgery, syndactyly, toe

1 **Introduction**

2 The aim of syndactyly surgery is to reconstruct not only functionally-independent but
3 also cosmetically-pleasing digits. In syndactyly of the toes, surgeries are occasionally
4 indicated to meet patients' and/or parents of patients' cosmetic requests. As in hand
5 syndactyly, reconstructive procedures using a dorsal, or planter commissural flap, in
6 combination with skin grafts, have been employed; however, unsightly longitudinal
7 scars on the dorsum of the toes sometimes fail to satisfy their expectations (Fig. 1).
8 Also, the symmetrical appearance of the dorsal scars remains a reminder that the
9 toes were artificially separated. In an attempt to address these issues, we developed
10 a novel design technique, using a dorsal transversally-oriented transposition flap for
11 web reconstruction. Our technique is described below.

13 **Surgical Technique**

14 The technique is employed for web creation in toe syndactyly and polysyndactyly of
15 the toes. Surgeries are performed under general anesthesia and tourniquet control. A
16 transversely-oriented flap is designed dorsally on the base of one of the involved toes
17 (Fig.1). Its pedicle, measuring approximately 5 mm in width (the width of the flap can
18 be adjusted according to the size of the toes and skin tension), is placed at the
19 proximal third to quarter interdigital space of the proximal phalanx. Two parallel

1 incisions, constituting both edges of the flap, are made to run transversely and slightly
2 curve distalward to join at the opposite aspect of the toe (Fig. 2(B), Fig. 3(B)). A dorsal
3 zigzag incision is then made distally to create multiple triangular interdigitating flaps.
4 On the planter side, a small triangular flap is designed at the base and a zigzag
5 incision is extended distally in the same manner as on the dorsal side. With care not
6 to injure the underlying adipose layer, the transversely-oriented flap is raised (Fig.
7 2(C)) and redirected 90 degrees to be inset into the commissure (Fig. 2(D), Fig. 3(C)).
8 The V-shaped distal tip is trimmed into a rectangular configuration (Fig. 3(D)). After
9 tight closure of the donor site using absorbable dermal sutures, the lateral aspects of
10 the toes are resurfaced with the interdigitating flaps. If skin defects remain, they are
11 covered with full-thickness skin grafts harvested from the submalleolar region. A bulky
12 dressing is applied and the extremity is held in cast immobilization. Sutures are
13 removed 3 weeks postoperatively. A sponge-pad (Reston™, 3M Critical & Chronic
14 Care Solutions) compression on the commissure is maintained for 3 to 6 months (Fig.
15 4).

16

17 Discussion

18 During a period of 6 years, we operated on 11 webs of 8 patients, 5 boys and 3 girls,
19 with congenital syndactylous webs of the foot, using the technique. In the 11, 9 were

1 corrections of simple syndactyly of the toes, and 2 of postaxial polysyndactyly of the
2 toes. Except for the case of a girl of the age of 13, all the cases were in the age range
3 of 1 to 2 years when surgery was performed. In most of the syndactyly cases, the total
4 surface areas for skin grafts were less than 1.0 cm², especially for defects of the most
5 proximal portion of the separated surfaces. In polysyndactyly cases, skin grafts were
6 not needed. Neither flap loss nor delay in wound healing was observed in any one
7 case. All the transverse scars of the donor sites became inconspicuous and
8 physiological configuration of the web was reconstructed (Fig. 5(B), Fig. 6(B)). None
9 of the cases showed a substantial scar contracture or web creep during the mean
10 follow-ups of 26 months (range, 12-45).

11 Most published papers regarding toe syndactyly and polysyndactyly of the toes were
12 originated from East Asian countries^{1-3, 6-9}. Recently, European surgeons, Marsh and
13 Floyd, reported their surgical experiences of toe syndactyly, and investigated patients'
14 satisfaction⁴. The investigation revealed that the patients' concerns about their
15 conditions were successfully reduced after operation, suggesting that if psychological
16 problems of patients with toe syndactyly were highlighted, surgical indication would
17 increase.

18 Originally, corrections of syndactylous conditions of the toes were performed in the
19 almost same maneuvers as for hand syndactyly, such as a dorsal rectangular flap, or

1 bi-planer rectangular or triangular flaps for web reconstruction¹⁻⁵, in combination with
2 skin grafts for lateral resurfacings. Since the 1990s, two technical variations with
3 different surgical concepts have been advocated: one is an open technique^{6, 7}, and
4 the other is a graft-less method^{8, 9}. In the open technique, the residual skin defects of
5 the lateral aspects of the separated toes are intentionally left open to be secondarily
6 healed. This technique shortens the operation time and eliminates the need for
7 postoperative cares for grafts; however, the risk of web creep can be heightened due
8 to wound constriction. Kawabata et al. documented a series of cases treated with the
9 open technique, where a dorsal rectangular flap was made more proximal than usual
10 to compensate for possible web creep⁶. Hirosaka et al. also reported their
11 experiences with the open technique. They mentioned duration of the lateral
12 epithelization averaged 5 weeks⁷. On the other hand, the graft-less method includes
13 variations of island flap techniques, where the newly-created commissure is
14 reconstructed by island flaps, and the lateral raw aspects of the toes are closed
15 primarily with interdigital skin flaps^{8, 9}. Ito and Arai described a graft-less technique
16 using a planter pentagonal island flap⁸. This flap technique required wide
17 subcutaneous dissections for distalward mobilization. Their surgical indication was
18 limited to cases which had sufficient interdigital skin to close the lateral aspects
19 directly. Moreover, they suggested that especially in syndactyly cases, removal of

1 interdigital fat tissue was always necessitated to lessen tension in the closure. Lim et
2 al. used a dorsal pentagonal island flap for commissural reconstruction⁹. Their
3 technique successfully achieved functional web creation; however, dorsal longitudinal
4 scars continuing from the base of the web to the dorsum of the metatarsal had a
5 tendency to develop scar hypertrophy or keloid formation.

6 Considering these technical issues, we developed a surgical concept, which satisfies
7 the functional and aesthetic aspects as followed: a smooth transition from the dorsum
8 of the foot to the commissure to create a physiological slope of the web, no linear
9 scars on the lateral aspects to avoid web creep, no dorsal longitudinal scars to avoid
10 scar hypertrophy, simple design and easy flap elevation. Since 2008, acknowledging
11 the good healing potential of transverse scars, we switched the axis of the traditional
12 rectangular flap from longitudinal to transverse direction. Lateral resurfacing was
13 performed using both interdigitating flaps and skin grafts. By sparing the dorsal
14 interdigital skin maximally, both the dorsal scars and grafted sites were shifted more
15 planterly to be inconspicuous. Furthermore, the asymmetrical appearance of dorsal
16 scars does not reveal the anomalous condition, thereby protecting patients' privacy.

17 A technical concern regarding our method is that the tapered shape of the flap might
18 pose the potential risk of distal necrosis. We raised the flap with care to preserve the
19 vascular networks in the subcutaneous layer, and held it in place with a bulky dressing

and cast immobilization as in skin grafting. Additionally, the most distal tip of the flap is always trimmed. These procedures minimized the possibility of flap failures. Our surgical concept does not include graft-less reconstruction. In order to eliminate the possibility of future lateral pain caused by the de-fattening maneuver used in the direct lateral closures, we chose not to utilize both the open and the graft-less method. A minimized surface area of skin graft might contribute to lessen the risk of web creep.

The limitation of our study is 2-fold. Firstly, is a lack of experiences of complex, or multiple syndactylies of the toes. Secondly, is a limited term of follow-ups. Further data on variations of conditions with long-term follow-ups will clarify the applicability of the technique.

In conclusion, the design of the transversely-oriented dorsal transposition flap is simple and uncomplicated surgically. Using this technique, the dorsal interdigital skin can be spared maximally, and the donor site morbidity is minimal, owing to the good healing potential of transverse scars. This technique could be an alternative in web reconstruction of toe syndactylies, especially in cases of high cosmetic priority.

Acknowledgement/s

None

References

1. Nogami H. Polydactyly and polysyndactyly of the fifth toe. Clin Orthop Relat Res. 204:261-265, 1986.
2. Onishi K, Nakajima T, Yoneda K, Aoki T. Operation for polysyndactyly of the fifth toe using Z-plasty. Br J Plast Surg. 45:554-556, 1992.
3. Hayashi A, Yanai A, Komuro Y, Nishida M. A New Surgical Technique for Polysyndactyly of the Toes without Skin Graft. Plast Reconstr Surg. 114:433-438, 2004.
4. Marsh DJ, Floyd D. Toe syndactyly revisited. J Plast Reconstr Aesthet Surg. 64:535-540, 2011.
5. Park S, Eguchi T, Tokioka K, Minegishi M. Reconstruction of incomplete syndactyly of the toes using both dorsal and plantar flaps. Plast Reconstr Surg. 98:534-537, 1996
6. Kawabata H1, Ariga K, Shibata T, Matsui Y Open treatment of syndactyly of the foot. Scand J Plast Reconstr Surg Hand Surg. 37:150-154, 2003.
7. Hikosaka M1, Ogata H, Nakajima T, Kobayashi H, Hattori N, Onishi F, Tamada I. Advantages of open treatment for syndactyly of the foot: defining its indications. Scand J Plast Reconstr Surg Hand Surg. 43:148-152, 2009.
8. Itoh Y, Arai K. A new operation for syndactyly and polysyndactyly of the foot

without skin grafts. *Br J Plast Surg.* 48:306-311, 1995.

9. Lim YJ, Teoh LC, Lee EH. Reconstruction of syndactyly and polysyndactyly of the toes with a dorsal pentagonal island flap: a technique that allows primary skin closure without the use of skin grafting. *J Foot Ankle Surg.* 46:86-92, 2007.

Figure Legends

Fig. 1. Illustrations showing the surgical design of the transversely-oriented dorsal transposition flap. Pairs of dot indicate the dorsal peak of the webs.

Fig. 2. A series of photographs showing the process of web reconstruction by the transversely-oriented transposition flap for toe syndactyly. (A) Dorsal view of 1-year-old boy's incomplete syndactyly of the second web of the left foot. (B) The flap, 5 mm in width, is designed on the dorsum of the second toe. (C) Flap elevation. (D) The flap is redirected 90° to be inset into the commissure. The both edges of the donor wound are approximated.

Fig. 3. A series of photographs showing the process of web reconstruction by the transversely-oriented transposition flap for postaxial polysyndactyly of the toes. (A) Dorsal view showing a 1-year-old girl's postaxial polysyndactyly of the right foot. (B) After resection of the fifth toe, the transversely-oriented flap is designed on the dorsum of the fourth toe. (C) The flap is redirected 90° to be inset into the

1 commissure. The both edges of the donor wound are approximated. (D) After
2 trimming of the distal end of the flap, the flap is inset into the commissure. Lateral
3 resurfacings of the both toes are completed without skin grafting.

4 **Fig.4.** Postoperative sponge-pad compression applying onto the newly-created
5 commissure. (A) Toe syndactyly. (B) Polysyndactyly of the lessor toes.

6 **Fig. 5.** Postoperative photographs of toe syndactyly (the same case in Fig. 2) at
7 follow-up of 12 months. (A) The transverse scar at the donor site on the base of the
8 second toe is significantly inconspicuous. (B) The commissure shows no contracture.

9 **Fig. 6.** Postoperative photographs of polysyndactyly of the toe (the same case in Fig.
10 3). (A) The transverse scar at the donor site on the base of the forth toe is significantly
11 inconspicuous. (B) Natural configuration of the slope of the web is created.

12

13

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6