

Title	New Marine Antifouling Compounds from the Red Alga <i>Laurencia</i> sp.
Author(s)	Oguri, Yuko; Watanabe, Mami; Ishikawa, Takafumi; Kamada, Takashi; Vairappan, Charles; Matsuura, Hiroshi; Kaneko, Kensuke; Ishii, Takahiro; Suzuki, Minoru; Yoshimura, Erina; Nogata, Yasuyuki; Okino, Tatsufumi
Citation	Marine Drugs (2017), 15(9)
Issue Date	2017-8-28
URL	http://hdl.handle.net/2433/233623
Right	© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. (CC BY 4.0).
Type	Journal Article
Textversion	publisher

Article

New Marine Antifouling Compounds from the Red Alga *Laurencia* sp.

Yuko Oguri ¹, Mami Watanabe ¹, Takafumi Ishikawa ¹, Takashi Kamada ², Charles S. Vairappan ², Hiroshi Matsuura ³, Kensuke Kaneko ⁴, Takahiro Ishii ⁵, Minoru Suzuki ⁶, Erina Yoshimura ⁷, Yasuyuki Nogata ⁸ and Tatsufumi Okino ^{1,9,*} 

- ¹ Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810, Japan; oguri-yuu@awi.co.jp (Y.O.); watamatsu@ees.hokudai.ac.jp (M.W.); tishikawa947@gmail.com (T.I.)
- ² Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Kota Kinabalu, Sabah 88400, Malaysia; takashi.kamada@ums.edu.my (T.K.); csv@mes.edu.my (C.S.V.)
- ³ National Institute of Technology, Asahikawa College, Asahikawa 071-8142, Japan; matsuura@asahikawa-nct.ac.jp
- ⁴ Graduate School of Pharmaceutical Sciences, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan; kaneko.kensuke.6a@kyoto-u.ac.jp
- ⁵ Faculty of Agriculture, University of the Ryukyus, Senbaru 1, Nishihara, Okinawa 903-0213, Japan; ishiit@agr.u-ryukyu.ac.jp
- ⁶ Coastal Branch of Natural History Museum and Institute, Chiba, 123 Yoshio, Katsuura, Chiba 299-5242, Japan; reiboku1965@yahoo.co.jp
- ⁷ CERES, Inc., 1-4-5 Midori, Abiko, Chiba 270-1153, Japan; erinamusi@yahoo.co.jp
- ⁸ Environmental Science Research Laboratory, Central Research Institute of Electric Power Industry, 1646 Abiko, Abiko, Chiba 270-1194, Japan; noga@criepi.denken.or.jp
- ⁹ Faculty of Environmental Earth Science, Hokkaido University, Sapporo 060-0810, Japan
- * Correspondence: okino@ees.hokudai.ac.jp; Tel.: +81-11-706-4519

Received: 31 July 2017; Accepted: 23 August 2017; Published: 28 August 2017

Abstract: Six new compounds, omaezol, intricatriol, hachijojimallenes A and B, debromoaplysinol, and 11,12-dihydro-3-hydroxyretinol have been isolated from four collections of *Laurencia* sp. These structures were determined by MS and NMR analyses. Their antifouling activities were evaluated together with eight previously known compounds isolated from the same samples. In particular, omaezol and hachijojimallene A showed potent activities ($EC_{50} = 0.15\text{--}0.23 \mu\text{g/mL}$) against larvae of the barnacle *Amphibalanus amphitrite*.

Keywords: antifouling; biofouling; barnacle; terpenoid; acetogenin; *Laurencia*; Rhodophyta

1. Introduction

Biofouling is a major cause of increased fuel consumption of ships and facilitates the spread of invasive marine organisms [1]. After many years of utilizing toxic antifoulants to control biofouling, the IMO (International Maritime Organization) banned the use of organotin compounds in 2008, paving the way for the development of environmentally friendly fouling-resistant coatings [2]. In addition, effective and new fouling treatment is needed to control and curb the introduction of non-indigenous species (NIS), that are introduced into local waters partly via ballast water uptake and discharge in vessels. The ballast water management convention will enter into force in September 2017. However, to effectively control NIS introduction, an effective antifouling treatment approach is still needed [3]. Search for natural antifouling compounds to solve biofouling has been “a work in progress” for almost four decades. Lately, a number of natural antifouling products have been reported and some of them have shown potent activities [4,5]. Furthermore, recent efforts to identify molecular mechanisms of antifouling compounds improve the possibility of their industrial importance [6]. As expected,

industrial utilization of natural products as antifouling agents would warrant systematic synthetic efforts, and these have been implemented [5]. The total synthesis of 10-isocyano-4-cadinene [7] and dolastatin 16 [8] were achieved by our group [9,10]. The red alga *Laurencia* sp. is a rich source of biologically active secondary metabolites [11,12] and produces antifouling compounds such as elatol [13], omaezallene [14], and 2,10-dibromo-3-chloro-7-chamigrene [15]. As a continuous effort, our group has been searching for novel and potent antifouling compounds from *Laurencia* sp. by using barnacle larvae and epiphytic diatom assays. Hence, here we report the structures (Figure 1) and activities of new antifouling compounds from *Laurencia* sp.




Figure 1. Structures of new compounds isolated from *Laurencia* sp.

2. Results

2.1. Omaezol (1) and Intricatiol (2)

Laurencia sp., which was collected in Omaezaki, Japan, was extracted with MeOH. The extract yielded omaezol (1), intricatiol (2), in addition to the previously isolated omaezallenes and intricatetraol [14]. The molecular formula of 1 was determined to be $C_{20}H_{35}BrO_2$ (m/z 368.1707, calcd. for $C_{20}H_{33}BrO$, 368.1709 [$M - H_2O$] $^+$) by HR-EIMS (High Resolution-Electron Ionization Mass Spectrometry), suggesting three degrees of unsaturation. The existence of a hydroxy group was revealed by an IR absorption at 3386 cm^{-1} . ^{13}C NMR data (Table 1) showed the presence of one double bond, suggesting a bicyclic structure. COSY correlations connected C-1 to C-3. HMBC peaks from H-19 to C-1, 5, 9, 10 and from H-20 to C-3, 4, 5 concluded a cyclohexane ring (Figure S1). Based on ^{13}C NMR chemical shifts, a bromine atom is attached to C-1 (δ 67.6) and a hydroxy group is connected to C-4 (δ 71.4). A side chain was determined by COSY correlation (H-11/H-18 and H-12/H-13) and HMBC peaks (H-18/C-7, 12 and H-14/C-13, 16, 17). Another hydroxy group is attached to C-7 (δ 74.6). Overlapping of NMR chemical shifts (C-6; δ 32.1, C-8; δ 31.8, C-9; δ 38.7, C-10; δ 39.4, H-6; δ 1.48, H-8; δ 1.49) hampered the positioning of the two remaining methylenes. However, HSQC-TOCSY peaks from H-8 to C-9 and H-6 to C-5 concluded the planar structure of 1. NOESY correlations determined relative configurations of the *trans*-decalin (Figure 2).




Figure 2. Key NOESY correlations observed in 2D NMR of compound 1.

The molecular formula of **2** was determined to be $C_{30}H_{52}Br_2Cl_2O_6$ (m/z 757.1584, calcd. for $C_{30}H_{53}Br_2Cl_2O_6$, 757.1581 $[M + H]^+$) by HR-FABMS (High Resolution-Fast Atom Bombardment Mass Spectrometry). The 1H and ^{13}C NMR spectra of **2** (Table 1) were similar to intricatetraol [16,17], but were not symmetric. The 2D NMR interpretations (Figure S2) showed that both compounds had the same carbon skeleton. However, ^{13}C chemical shift changes were observed (C-7: δ 84.1 in intricatetraol, δ 113.4 in **2**, and C-11: δ 77.5 in intricatetraol, δ 84.7 in **2**). C-7 in **2** was proposed to be an acetal carbon and C-11 was suggested to be a part of an ether instead of a secondary alcohol to make another five-membered ring. In conclusion, the planar structure of **2** was determined and its configuration was assumed to be the same as intricatetraol [16,17], which was extracted from the same sample.

Table 1. 1H and ^{13}C NMR spectroscopic data for compounds 1 and 2 in $CDCl_3$.

Carbon Number	Compound 1		Compound 2		Carbon Number	Compound 2	
	^{13}C	1H	^{13}C	1H		^{13}C	1H
1	67.6	3.91	27.8	1.689	24	27.5	1.695
2	30.2	2.44, 2.01	72.0		23	72.0	
3	41.9	1.55, 1.70	67.1	4.07	22	67.1	4.08
4	71.4		28.7	1.85, 2.61	21	28.9	1.75, 2.45
5	48.2	1.10	36.8	1.72, 1.89	20	36.9	1.47, 1.72
6	32.1	1.48, 2.00	72.2		19	73.3	
7	74.6		113.4		18	84.3	3.85
8	31.8	1.49, 1.90	27.9	1.60, 1.70	17	26.4	1.93
9	38.7	1.05, 1.76	32.8	1.91	16	32.7	1.57, 2.19
10	39.4		86.7		15	85.6	
11	33.6	1.67	84.7	3.61	14	76.0	3.57
12	30.2	1.52	29.6	1.55, 2.40	13	29.4	1.74
13	26.5	1.12			25	33.1	1.785
14	124.3	5.10			30	32.9	1.793
15	131.7				26	21.2	1.28
16	25.5	1.70			29	24.5	1.26
17	17.6	1.62			27	17.6	1.45
18	12.3	0.90			28	23.0	1.19
19	14.4	1.26					
20	29.7	1.18					

2.2. Hachijojimallenes A (3) and B (4)

Hachijojimallenes A (**3**) and B (**4**) were isolated together with *N*-methyl-2,3,6-tribromoindole [18] and pinnaterpene C [19] from a red alga, *Laurencia* sp., which was collected in Hachijojima Island, Japan. Based on HR-EIMS data, the molecular formula of **3** was determined to be $C_{15}H_{19}Br_3O_3$

(m/z 466.8851, calcd. for $C_{15}H_{18}Br_3O_2$, 466.8852 $[M - OH]^+$). The existence of a bromoallene moiety was suggested by an IR absorption at 1957 cm^{-1} , ^{13}C NMR chemical shifts (δ 74.1, 99.4, and 201.9), and 1H NMR chemical shifts (δ 5.45 and 6.08) (Table 2). COSY correlations established a carbon skeleton C-3—C-8—C-13—C-10 and an ethyl group. ^{13}C chemical shifts revealed that C-4, 6, and 7 were oxymethines and C-10 and 12 were brominated. These carbons and an acetal carbon (δ 102.1) were assembled by HMBC data to conclude the planar structure of **3** (Figure S3). Relative configurations were determined by NOEs (H-6/H-10, H-10/H-11 α , H-11 α /H-12, H-7/H-12, and H-8/H-13).

The molecular formula of **4** was determined to be $C_{15}H_{18}Br_2O_3$ (m/z 403.9627, calcd. for $C_{15}H_{18}Br_2O_3$, 403.9617 $[M]^+$) by HR-EIMS, suggesting the existence of another ring. The striking difference in its ^{13}C NMR (Table 2) was observed in C-9 (**3**; δ 102.1, **4**; δ 119.7) and C-12 (**3**; δ 51.3, **4**; δ 75.6). In addition, HMBC was observed from H-12 to C-9. In conclusion, the hemiacetal in **3** was replaced by acetal in **4** to form an ether between C-9 and C-12, which was debrominated at C-12. NOEs of **4** supported its relative configurations, similar to those of **3**.

Table 2. 1H and ^{13}C NMR spectroscopic data for compounds **3–5** in $CDCl_3$.

Carbon Number	Compound 3		Compound 4		Compound 5	
	^{13}C	1H	^{13}C	1H	^{13}C	1H
1	74.1	6.08	74.0	3.09	58.7	
2	201.9		201.9		103.4	
3	99.4	5.45	100.0	5.48	42.4	2.59
4	73.8	4.73	74.1	4.72	31.7	1.26, 1.74
5	39.3	1.81, 2.26	39.6	1.82, 2.35	43.0	1.74, 1.92
6	79.4	4.56	88.6	5.11	131.6	
7	81.4	4.61	82.8	4.77	159.4	
8	50.5	2.77	51.0	3.19	110.0	6.72
9	102.1		119.7		138.8	
10	53.6	4.07	42.8	4.13	122.1	6.74
11	44.3	2.50, 2.76	40.6	1.74, 2.70	122.4	6.92
12	51.3	3.79	75.6	4.03	21.5	2.33
13	44.7	1.97	50.2	2.13	24.2	1.31
14	24.7	1.47, 2.11	23.3	1.44, 1.60	203.5	9.78
15	11.6	1.03	12.5	0.99	13.1	1.04
9-OH		3.65				

2.3. Debromoaplysinal (**5**)

Re-investigation of the red alga *L. okamurae*, collected at Oshoro Bay, Hokkaido, Japan, yielded debromoaplysinal (**5**). The molecular formula of **5** was determined to be $C_{15}H_{18}O_2$ (m/z 231.1379, calcd. for $C_{15}H_{19}O_2$, 231.1380 $[M + H]^+$) by ESI-TOFMS. Its 1D NMR spectra (Table 2) revealed the existence of an aldehyde and a trisubstituted benzene. COSY and HMBC correlations (Figure S4) clarified its planar structure, which is similar to aplysinal and debromoaplysinal. In fact, chemical shifts in the trisubstituted benzene of **5** are similar to those of debromoaplysinal, and those in the cyclopentane of **5** are similar to those of aplysinal. NOESY peaks between H-13 and H-14 suggested *cis* configuration of Me-13 and the aldehyde. Irradiation of H-3 by DPGSE 1D NOE showed the enhancement of H-13. In conclusion, H-3, Me-13, and the aldehyde are located on the same face.

2.4. 11,12-Dihydro-3-hydroxyretinol (**6**)

Re-investigation of the red alga *L. nipponica* collected at Muroran, Hokkaido, Japan, yielded 11,12-dihydro-3-hydroxyretinol (**6**), along with three known halogenated chamigrene-type sesquiterpenoids [20,21]. The molecular formula of **6** was deduced to be $C_{20}H_{32}O_2$ (m/z 304.2400, calcd. for $C_{20}H_{32}O_2$, 304.2397 $[M]^+$) by HR-EIMS, accounting for five degrees of unsaturation. The 1H - and ^{13}C -NMR spectroscopic data (Table 3) as well as the HSQC experiment of **6** showed the presence

of eight sp^2 carbons, three vinyl methyls, a pair of geminal methyls, an oxymethylene, an oxymethine, four methylenes, and a quaternary carbon. These signals explained four degrees of unsaturation, implying that one ring was present in **6**. COSY and HMBC data (Figure S5) clearly indicated the gross structure of **6**. The *E* configurations of double bonds were deduced from the 1H - 1H coupling constants ($^3J_{7-8}$ 15.9 Hz) and ^{13}C -NMR chemical shifts of vinyl methyls (C-19; δ 13.2, and C-20; δ 17.2). Unfortunately, the configuration at C-3 was not determined due to its limited amount.

Table 3. 1H and ^{13}C NMR spectroscopic data for compound **6** in $CDCl_3$.

Carbon Number	^{13}C	1H	Carbon Number	^{13}C	1H
1	37.9		11	27.5	2.29
2	49.1	1.47, 1.76	12	40.0	2.12
3	65.9	4.00	13	140.0	
4	43.2	2.02, 2.37	14	124.2	5.44
5	125.9		15	60.1	4.18
6	138.2		16	22.4	1.71
7	124.0	5.92	17	31.0	1.06
8	139.0	6.00	18	29.5	1.06
9	134.6		19	13.2	1.79
10	131.4	5.40	20	17.2	1.71

2.5. Antifouling Activity

Most of the compounds obtained in this study were tested for antifouling activity against larvae of the barnacle *Amphibalanus amphitrite* (Table 4). The antifouling activity of **1** was potent ($EC_{50} = 0.23 \mu\text{g/mL}$) and its toxicity was low ($LC_{50} = 3.7 \mu\text{g/mL}$), while **2** was not potent ($EC_{50} > 10 \mu\text{g/mL}$). The antifouling activity of **3** was potent ($EC_{50} = 0.15 \mu\text{g/mL}$) and its toxicity was low ($LC_{50} = 9.8 \mu\text{g/mL}$), while **4** was a little less potent ($EC_{50} = 0.31 \mu\text{g/mL}$, $LC_{50} = 6.8 \mu\text{g/mL}$). The antifouling activity of the known compound, pinnaterpene C (**7**, $EC_{50} = 0.82 \mu\text{g/mL}$, $LC_{50} > 10 \mu\text{g/mL}$), was similar to that of **4**, while another known compound, *N*-methyl-2,3,6-tribromoindole, showed weak activity ($EC_{50} = 4.3 \mu\text{g/mL}$, not toxic at $10 \mu\text{g/mL}$). Compound **5** also showed moderate antifouling activity ($EC_{50} = 1.0 \mu\text{g/mL}$) and did not kill any larvae at $10 \mu\text{g/mL}$. Some known compounds (Figure 3) isolated from *L. okamuræ* were tested. Three aromatic sesquiterpenoids [22] showed antifouling activities (laurinterol **8**: $EC_{50} = 0.65 \mu\text{g/mL}$, $LC_{50} = 5.8 \mu\text{g/mL}$; isolaurinterol **9**: $EC_{50} = 0.34 \mu\text{g/mL}$, $LC_{50} > 10 \mu\text{g/mL}$; debromolaurinterol **10**: $EC_{50} = 0.5 \mu\text{g/mL}$, $LC_{50} > 10 \mu\text{g/mL}$), while α -bromocuparene (**11**) was inactive. Three sesquiterpenoids (Figure 3) isolated from *L. nipponica* collected in Muroran showed antifouling activities (prepacifenol **12**: $EC_{50} = 0.63 \mu\text{g/mL}$, $LC_{50} > 10 \mu\text{g/mL}$; pacifenol **13**: $EC_{50} = 2.36 \mu\text{g/mL}$, $LC_{50} > 10 \mu\text{g/mL}$; 2,10-dibromo-3-chloro-9-hydroxy- α -chamigrene **14** $EC_{50} = 2.51 \mu\text{g/mL}$, $LC_{50} > 10 \mu\text{g/mL}$). In addition, anti-diatom activities against the epiphytic marine diatoms, *Nitzschia* sp. and *Cylindrotheca closterium*, were tested for three chamigrenes and **6**. Compounds **6**, **12**, and **14** showed inhibition against both diatoms at $5\text{--}10 \mu\text{g/cm}^2$.




Figure 3. Structures of tested known compounds isolated from *Laurencia* sp.

Table 4. Antifouling activities (μM) against barnacle larvae.

Compound	EC ₅₀	LC ₅₀
1	0.59	9.6
2	—	—
3	0.31	20
4	0.76	17
5	4.3	—
7	1.6	—
8	2.2	20
9	1.2	—
10	2.3	—
11	—	—
12	1.5	—
13	5.5	—
14	6.1	—
CuSO ₄	0.72	1.4

—: EC₅₀ or LC₅₀ is more than 10 $\mu\text{g}/\text{mL}$.

3. Discussion

From a chemical point of view, **1** is a rare halo-diterpenoid, prenylated selinene-type compound like anhydroaplysiadiol [23]. Oxasqualenoids such as **2** have been proposed to be synthesized by the epoxide-opening cascades. These reactions could be catalyzed by vanadium-dependent bromoperoxidases [24]. However, we do not know how different cyclization patterns are controlled to produce **2** and the similar compound intricatetraol [16,17]. Although most of C₁₅ acetogenins from *Laurencia* contain only ether rings, **3** and **4** contain a carbocyclic ring, similar to lembynes [25]. Compound **5** is one of common laurane-type sesquiterpenes. Retinols occur in nature only in animals and in the limited green algae *Caulerpa* sp. [26] This is the first record of a retinane-type diterpene

(6) isolated from *Laurencia*. Although the genus of *Laurencia* is one of the most studied organisms in marine natural product chemistry [11,12], recent isolation techniques yielded six new compounds in this study. Twelve compounds showed antifouling activities against the barnacle larvae. Notably, the activities of 1 and 3 were equivalent to CuSO_4 ($\text{EC}_{50} = 0.18 \mu\text{g/mL}$). Three compounds showed antifouling activities against diatoms. However, most of compounds isolated from *Laurencia* sp. so far have not been tested for antifouling activity. Our results suggest that *Laurencia* is a potential source of antifouling compounds.

4. Materials and Methods

4.1. General Procedures

IR spectra were measured on a JASCO IR-700 spectrophotometer (JASCO, Tokyo, Japan). ^1H NMR and ^{13}C NMR spectra were recorded in CDCl_3 by using JEOL JNM-ECA600 (JEOL, Tokyo, Japan), JEOL JNM-EX400 (JEOL, Tokyo, Japan), or BRUKER ASX300 spectrometer (Bruker BioSpin, Faellanden, Switzerland), unless otherwise stated. EI-MS were obtained on a JEOL JMS-FABmate spectrometer (JEOL, Tokyo, Japan). FAB-MS were obtained on a JEOL JMS-HX110 spectrometer (JEOL, Tokyo, Japan). ESI-MS were obtained on a JEOL JMS-700TZ (JEOL, Tokyo, Japan) or BRUKER DALTONICS micro TOF-HS focus spectrometer (Bruker Daltonics, Bremen, Germany). Optical rotations were recorded on a HORIBA SEPA-300 polarimeter (Horiba, Kyoto, Japan).

4.2. Plant Material

Algal samples of *Laurencia* sp. were collected at Omaezaki, Shizuoka Prefecture, and Hachijojima Island, Tokyo, Japan. *L. okamurae* was collected at Oshoro Bay and *L. nipponica* was collected at Muroran, Hokkaido, Japan. The voucher specimens are deposited in the Herbarium of Graduate School of Science, Hokkaido University.

4.3. Omaezol and Intricatriol

The dried algal sample (250 g) was extracted and separated as described in a previous paper [12]. Omaezol (1, 3.2 mg) and intricatriol (2, 17.0 mg) were isolated by HPLC (YMC-Pack Pro C18 (YMC, Kyoto, Japan) with CH_3CN and H_2O) from the omaezallene containing silica-gel fraction.

1: $[\alpha]_{\text{D}}^{23} -67.4$ (c 0.17, CHCl_3); IR (neat), ν_{max} 3428, 1259, 1185, 1088, 1065, 954, 801 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Table 1.

2: $[\alpha]_{\text{D}}^{23} -45.2$ (c 0.39, CHCl_3); IR (neat), ν_{max} 3414, 2972, 1451, 1369, 1214, 1101, 756 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Table 1.

4.4. Hachijojimallenes A and B

The air-dried algae (80 g) were soaked in MeOH (500 mL) for seven days. The MeOH solution was concentrated in vacuo, and the residue was partitioned between EtOAc and H_2O . The EtOAc layer was then washed with water, dried over anhydrous Na_2SO_4 , and evaporated in vacuo to leave a dark green oily substance (1.23 g). The extract (700 mg) was fractionated by Si gel CC with a step gradient (hexane and EtOAc). The fraction (41 mg) eluted with hexane-EtOAc (9:1) was further subjected to PTLC (preparative thin-layer chromatography) with hexane to give 1-methyl-2,3,6-tribromoindole (4.0 mg), which was determined on the basis of ^1H NMR and MS data. The fraction (411 mg) eluted with hexane-EtOAc (7:3) was further chromatographed on a Si gel column with hexane-EtOAc (3:1) (each eluate, 10 mL) to give 10 fractions. The fourth fraction (79.1 mg) was then separated by PTLC with hexane-EtOAc (3:1) to afford hachijojimallene B (4) (3.5 mg). The combined fifth and sixth fractions (74.9 mg) were further separated by repeated PTLC with hexane-EtOAc (1:1) to give hachijojimallene A (3) (40.0 mg). The seventh fraction (43.8 mg) was subjected to PTLC with hexane-EtOAc (3:1) followed by HPLC (Develosil-ODS-T-5 with CH_3CN and H_2O) to give pinnaterpene C (7, 15.0 mg).

3: $[\alpha]_D^{24} -88.2$ (*c* 0.13, CHCl₃); IR (neat), ν_{\max} 3490, 1957, 1258, 1184, 1165, 1143, 1120, 1038, 959, 788, 756 cm⁻¹; ¹H NMR and ¹³C NMR, see Table 1.

4: $[\alpha]_D^{24} -151$ (*c* 0.71, CHCl₃); IR (neat), ν_{\max} 1950, 1243, 1181, 1127, 1080, 1045, 1022, 970, 855, 821 cm⁻¹; ¹H NMR and ¹³C NMR, see Table 1.

4.5. Debromoaplysinal

The air-dried algae (100 g) were soaked in MeOH (3 L) for seven days. The MeOH solution was concentrated in vacuo, and the residue was partitioned between EtOAc and H₂O. The EtOAc layer was then washed with water, dried over anhydrous Na₂SO₄, and evaporated in vacuo to obtain the crude extract (4.3 g). The extract (2.1 g) was fractionated by Si gel column chromatography with a step gradient (hexane and EtOAc). The fraction (1.4 g) eluted with hexane-EtOAc (3:1) was further subjected to two sets of Si gel column chromatography and PTLC with toluene to give debromoaplysinal (**5**) (0.9 mg). The other known compounds (laurinterol **8**, 30.6 mg; isolaurinterol **9**, 12.4 mg; debromolaurinterol **10**, 11.0 mg; α -bromocuparene **11**, 1.1 mg) were isolated from the same fraction (hexane-EtOAc 3:1) in a similar way.

5: $[\alpha]_D^{24} -91.9$ (*c* 0.08, CHCl₃); IR (neat), ν_{\max} 2927, 2856, 1731, 1593, 1499, 1456, 1377, 1266, 1146, 1117, 971, 805 cm⁻¹; ¹H NMR and ¹³C NMR, see Table 1.

4.6. 11,12-Dihydro-3-hydroxyretinol

Fresh algal sample (400 g wet weight) was extracted in MeOH at room temperature for three days. The resulting MeOH extract was concentrated in vacuo and partitioned between diethyl ether and H₂O. The diethyl ether fraction (340 mg) was subjected to Si gel CC eluting with a gradient of hexane and EtOAc in an increasing polarity. Fraction 4, eluted with hexane-EtOAc (3:1), was subjected to PTLC to yield pacifenol (**12**, 4.1 mg) and prepacifenol (**13**, 6.3 mg). Fraction 5, eluted with hexane-EtOAc (1:1), was subjected to PTLC with CHCl₃-MeOH (97:3) to yield 2,10-dibromo-3-chloro-9-hydroxy- α -chamigrene (**14**, 7.3 mg). Fraction 7, eluted with hexane-EtOAc (1:3), gave **6** (1.3 mg) after purification with PTLC using CHCl₃-MeOH (95:5).

6: $[\alpha]_D^{27} -61.9$ (*c* 0.07, CHCl₃); ¹H NMR and ¹³C NMR, see Table 1.

4.7. Antifouling Assay

An antifouling assay against larvae of the barnacle *Amphibalanus amphitrite* was conducted according to the previous literature [9]. The antifouling assay against diatoms *Nitzschia* sp. and *Cylindrotheca closterium* was conducted as follows. The diatoms were maintained in test tubes under 16 h light and 8 h dark cycle conditions at 25 °C in modified Jorgensen's medium. Pure compounds were applied to a sample zone (16 mm in diameter) of cellulose TLC aluminum sheet (58 × 58 mm), and then each treated sheet was placed in Petri dish (15 × 90 mm in diameter). After that, 25-mL aliquots of modified Jorgensen's medium were introduced into Petri dishes with treated sheets, and inoculated with 1 mL of cultivated diatom suspension in equivalent cell densities. Petri dishes were sealed with Parafilm to ensure a closed system, and were incubated under the same conditions. Cell growth was estimated daily, and the adhesive condition was evaluated and compared to those of the control after seven days of incubation.

Supplementary Materials: The following are available online at www.mdpi.com/1660-3397/15/9/267/s1, Figure S1: Key 2D NMR correlations of compound **1**; Figure S2: Key 2D NMR correlations of compound **2**; Figure S3: Key 2D NMR correlations of compound **3**; Figure S4: Key 2D NMR correlations of compound **5**; Figure S5: Key 2D NMR correlations of compound **6**; Figure S6: ¹H NMR spectrum of omaezol (**1**) in CDCl₃; Figure S7: ¹³C NMR spectrum of omaezol (**1**) in CDCl₃; Figure S8: ¹H NMR spectrum of intricatriol (**2**) in CDCl₃; Figure S9: ¹³C NMR spectrum of intricatriol (**2**) in CDCl₃; Figure S10: ¹H-¹H COSY spectrum of intricatriol (**2**) in CDCl₃; Figure S11: HSQC spectrum of intricatriol (**2**) in CDCl₃; Figure S12: HMBC spectrum of intricatriol (**2**) in CDCl₃; Figure S13: NOESY spectrum of intricatriol (**2**) in CDCl₃; Figure S14: ¹H NMR spectrum of hachijojimallene

A (3) in CDCl₃; Figure S15: ¹³C NMR spectrum of hachijojimallene A (3) in CDCl₃; Figure S16: ¹H-¹H COSY spectrum of hachijojimallene A (3) in CDCl₃; Figure S17: HMQC spectrum of hachijojimallene A (3) in CDCl₃; Figure S18: HMBC spectrum of hachijojimallene A (3) in CDCl₃. Figure S19: NOESY spectrum of hachijojimallene A (3) in CDCl₃; Figure S20: ¹H NMR spectrum of hachijojimallene B (4) in CDCl₃; Figure S21: ¹³C NMR spectrum of hachijojimallene B (4) in CDCl₃. Figure S22: HMQC spectrum of hachijojimallene B (4) in CDCl₃; Figure S23: HMBC spectrum of hachijojimallene B (4) in CDCl₃; Figure S24: ¹H NMR spectrum of debromoaplysinal (5) in CDCl₃; Figure S25: ¹³C NMR spectrum of debromoaplysinal (5) in CDCl₃; Figure S26: DPGFSE 1D NOE spectrum of debromoaplysinal (5) in CDCl₃; Figure S27: ¹H NMR spectrum of 11,12-dihydro-3-hydroxyretinol (6) in CDCl₃; Figure S28: ¹³C NMR spectrum of 11,12-dihydro-3-hydroxyretinol (6) in CDCl₃.

Acknowledgments: This work was supported by JSPS KAKENHI Grant Number 16H04975 and JST. The authors would like to thank Tomohiko Kawamura (University of Tokyo) for providing the diatoms.

Author Contributions: T.O. conceived and designed the experiments; Y.O., M.W., H.M., C.S.V., and M.S. performed the experiments and analyzed the data on omaezol, intricatriol, and hachijojimallenes; T.I. and K.K. contributed the debromoaplysinal experiments; T.K. and T.I. contributed the 11,12-dihydro-3-hydroxyretinol experiments; E.Y. and Y.N. contributed the larval assays; T.O. wrote the paper.

Conflicts of Interest: The authors declare no conflict of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

References

1. Davidson, I.; Scianni, C.; Hewitt, C.; Everett, R.; Holm, E.; Tamburri, M.; Ruiz, G. Assessing the drivers of ship biofouling management—Aligning industry and biosecurity goals. *Biofouling* **2016**, *32*, 411–428. [[CrossRef](#)]
2. Callow, J.A.; Callow, M.E. Trends in the development of environmentally friendly fouling-resistant marine coatings. *Nat. Commun.* **2011**, *2*, 244. [[CrossRef](#)] [[PubMed](#)]
3. Fernandes, J.A.; Santos, L.; Vance, T.; Fileman, T.; Smith, D.; Bishop, J.D.D.; Viard, F.; Queirós, A.M.; Merino, G.; Vuisman, E.; et al. Costs and benefits to European shipping of ballast-water and hull-fouling treatment: Impacts of native and non-indigenous species. *Mar. Policy* **2016**, *64*, 148–155. [[CrossRef](#)]
4. Almeida, J.R.; Vasconcelos, V. Natural antifouling compounds: Effectiveness in preventing invertebrate settlement and adhesion. *Biotechnol. Adv.* **2015**, *33*, 343–357. [[CrossRef](#)]
5. Qian, P.; Li, Z.; Xu, Y.; Fusetani, N. Marine natural products and their synthetic analogs as antifouling compounds: 2009–2014. *Biofouling* **2015**, *31*, 101–122. [[CrossRef](#)] [[PubMed](#)]
6. Qian, P.; Chen, L.; Xu, Y. Molecular mechanisms of antifouling compounds. *Biofouling* **2013**, *29*, 381–400. [[CrossRef](#)] [[PubMed](#)]
7. Okino, T.; Yoshimura, E.; Hirota, H.; Fusetani, N. New antifouling sesquiterpenes from four nudibranchs of the family Phyllidiidae. *Tetrahedron* **1996**, *52*, 9447–9454. [[CrossRef](#)]
8. Tan, L.T.; Goh, B.P.L.; Tripathi, A.; Lim, M.G.; Dickinson, G.H.; Lee, S.S.C.; Teo, S.L.M. Natural antifoulants from the marine cyanobacterium *Lyngbya majuscula*. *Biofouling* **2010**, *26*, 685–695. [[CrossRef](#)] [[PubMed](#)]
9. Casalme, L.O.; Yamauchi, A.; Sato, A.; Petitbois, J.G.; Nogata, Y.; Yoshimura, E.; Okino, T.; Umezawa, T.; Matsuda, F. Total synthesis and biological activity of dolastatin 16. *Org. Biomol. Chem.* **2017**, *15*, 1140–1150. [[CrossRef](#)] [[PubMed](#)]
10. Nishikawa, K.; Nakahara, H.; Shirokura, Y.; Nogata, Y.; Yoshimura, E.; Umezawa, T.; Okino, T.; Matsuda, F. Total synthesis of 10-isocyano-4-cadinene and its stereoisomers and evaluations of antifouling activities. *J. Org. Chem.* **2011**, *76*, 6558–6573. [[CrossRef](#)] [[PubMed](#)]
11. Suzuki, M.; Vairappan, C.S. Halogenated secondary metabolites from Japanese species of the red algal genus *Laurencia* (Rhodomelaceae, Ceramiales). *Curr. Top. Phytochem.* **2005**, *7*, 1–34.
12. Wang, B.G.; Gloer, J.B.; Ji, N.Y.; Zhao, J.C. Halogenated organic molecules of Rhodomelaceae origin: Chemistry and biology. *Chem. Rev.* **2013**, *113*, 3632–3685. [[CrossRef](#)] [[PubMed](#)]
13. König, G.M.; Wright, A.D. *Laurencia rigida*: Chemical investigation of its antifouling dichloromethane extract. *J. Nat. Prod.* **1997**, *60*, 967–970. [[CrossRef](#)]
14. Umezawa, T.; Oguri, Y.; Matsuura, H.; Yamazaki, S.; Suzuki, M.; Yoshimura, E.; Furuta, T.; Nogata, Y.; Serisawa, Y.; Matsuyama-Serisawa, K.; et al. Omaezallene from red alga *Laurencia* sp.: Structure elucidation, total synthesis, and antifouling activity. *Angew. Chem. Int. Ed.* **2014**, *53*, 3909–3912. [[CrossRef](#)] [[PubMed](#)]

15. Al-Lihaibi, S.S.; Abdel-Lateff, A.; Alarif, W.M.; Nogata, Y.; Ayyad, S.N.; Okino, T. Potent antifouling metabolites from Red Sea organisms. *Asian J. Chem.* **2015**, *27*, 2252–2256. [[CrossRef](#)]
16. Suzuki, M.; Matsuo, Y.; Takeda, S.; Suzuki, T. Intricatetraol, a halogenated triterpene alcohol from the red alga *Laurencia intricata*. *Phytochemistry* **1993**, *33*, 651–656. [[CrossRef](#)]
17. Morimoto, Y.; Okita, T.; Takaishi, M.; Tanaka, T. Total synthesis and determination of the absolute configuration of (+)-intricatetraol. *Angew. Chem. Int. Ed.* **2007**, *46*, 1132–1135. [[CrossRef](#)]
18. Carter, G.T.; Rimehart, K.L., Jr.; Li, L.H.; Kuentzel, S.L.; Connor, J.L. Brominated indoles from *Laurencia brongniartii*. *Tetrahedron Lett.* **1978**, *19*, 4479–4482. [[CrossRef](#)]
19. Fukuzawa, A.; Kumagai, Y.; Masamune, T.; Furusaki, A.; Matsumoto, T.; Katayama, C. Pinnaterpenes A, B, and C, new dibromoditerpenes from the red alga *Laurencia pinnata* Yamada. *Chem. Lett.* **1982**, *11*, 1389–1392. [[CrossRef](#)]
20. Suzuki, M.; Kurosawa, E.; Furusaki, A. The structure and absolute stereochemistry of a halogenated chamigrene derivative from the red alga *Laurencia* species. *Bull. Chem. Soc. Jpn.* **1988**, *61*, 3371–3373. [[CrossRef](#)]
21. Abe, T.; Masuda, M.; Suzuki, T.; Suzuki, M. Chemical races in the red alga *Laurencia nipponica* (Rhodomelaeae, Ceramiales). *Phycol. Res.* **1999**, *47*, 87–95. [[CrossRef](#)]
22. Suzuki, M.; Kurosawa, E. Halogenated and non-halogenated aromatic sesquiterpenes from the red algae *Laurencia okamurae* Yamada. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 3352–3354. [[CrossRef](#)]
23. Takahashi, Y.; Suzuki, M.; Abe, T.; Masuda, M. Anhydroaplysiadiol from *Laurencia japonensis*. *Phytochemistry* **1998**, *48*, 987–990. [[CrossRef](#)]
24. Kaneko, K.; Washio, K.; Umezawa, T.; Matsuda, F.; Morikawa, M.; Okino, T. cDNA cloning and characterization of vanadium-dependent bromoperoxidases from the red alga *Laurencia nipponica*. *Biosci. Biotechnol. Biochem.* **2014**, *78*, 1310–1319. [[CrossRef](#)]
25. Vairappan, C.S.; Daitoh, M.; Suzuki, M.; Abe, T.; Masuda, M. Antibacterial halogenated metabolites from the Malaysian *Laurencia* species. *Phytochemistry* **2001**, *58*, 291–297. [[CrossRef](#)]
26. Blackman, A.J.; Wells, R.J. Caulerpol, a diterpene alcohol, related to vitamin A, from *Caulerpa brownii* (algae). *Tetrahedron Lett.* **1976**, *17*, 2729–2730. [[CrossRef](#)]



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).