TITLE:

Comparison of methyl eugenol metabolites, mitochondrial COI, and rDNA sequences of *Bactrocera philippinensis* (Diptera: Tephritidae) with those of three other major pest species within the dorsalis complex

AUTHOR(S):

Tan, Keng Hong; Wee, Suk-Ling; Ono, Hajime; Nishida, Ritsuo

CITATION:

Tan, Keng Hong ...[et al]. Comparison of methyl eugenol metabolites, mitochondrial COI, and rDNA sequences of *Bactrocera philippinensis* (Diptera: Tephritidae) with those of three other major pest species within the dorsalis complex. *Applied Entomology and Zoology* 2013, 48(3): 275-282

ISSUE DATE:

2013-08

URL:

<http://hdl.handle.net/2433/235534>

RIGHT:

This is a post-peer-review, pre-copyedit version of an article published in *Applied Entomology and Zoology*. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s13355-013-0183-5>; この論文は出版社版でありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

**Comparison of methyl eugenol metabolites, mitochondrial COI and
rDNA sequences of *Bactrocera philippinensis* (Diptera: Tephritidae)
with three other major pest species within the *dorsalis* complex**

Keng Hong Tan¹, Suk-Ling Wee², Hajime Ono³ and Ritsuo Nishida³

¹ Tan Hak Heng, 20, Jalan Tan Jit Seng, 11200, Penang, Malaysia

² School of Environmental and Natural Resource Sciences, Faculty of Science and
Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor Darul Ehsan,
Malaysia

³ Laboratory of Chemical Ecology, Graduate School of Agriculture, Kyoto University,
Kyoto, 606-8502, Japan

Hajime Ono

Email: onoono@kais.kyoto-u.ac.jp

Abstract

Males of the oriental fruit fly, *Bactrocera dorsalis* (Hendel), and some of its sibling species show a strong affinity to methyl eugenol (ME). Methyl eugenol ingested by male flies is biotransformed in the crop to two ME-metabolites that eventually accumulate in the rectal gland, which is known to serve as a reservoir for *B. dorsalis* sex pheromones. Upon ME-feeding, males of laboratory and wild *B. philippinensis* Drew and Hancock selectively accumulated two metabolites, 2-allyl-4,5-dimethoxyphenol and (*E*)-coniferyl alcohol, in the rectal gland, as was seen in *B. dorsalis sensu stricto*, *B. invadens* Drew, Tsuruta and White and *B. papayae* Drew and Hancock. Phylogenetic analysis of COI and rDNA sequence data of these four taxa also revealed a tight relationship among *B. philippinensis*, *B. dorsalis s.s.*, *B. invadens* and *B. papayae* (all four are members of the *dorsalis*-species complex). This result corroborates the pheromone analysis. The usefulness of pheromonal analysis as a chemotaxonomy tool to complement molecular and other analyses in the differentiation of closely related sibling species within the *Bactrocera dorsalis* complex, where using morphological characters had been inadequate, is highlighted.

Key words: *Bactrocera philippinensis* – *Bactrocera dorsalis* species complex – pheromone – methyl eugenol – mitochondrial DNA

Introduction

Currently, there are 75 species within the *Bactrocera dorsalis* Hendel species complex (Diptera: Tephritidae) (Clarke et al. 2005), an increase of over twenty species from the 52 species revised by Drew and Hancock (1994). Of these, 26 species are known to be responsive to the male attractant methyl eugenol (ME), including *B. dorsalis* s.s., *B. invadens* Drew, Tsuruta and White, *B. papayae* Drew and Hancock (synonym *B. dorsalis sensu lato* – see discussion), and *B. philippinensis* Drew and Hancock; all of which are considered serious and highly invasive fruit pests (Clarke et al. 2005; Tan et al. 2011).

Bactrocera philippinensis is recorded as an endemic and notable pest species in the Philippines and has a distinct geographical range from *B. dorsalis* s.s. (Clarke et al, 2005). In Palau, it was first recorded and mistakenly identified as *B. dorsalis* in September, 1996 but confirmed as *B. philippinensis* five years later (http://www.spc.int/Pacifly/Species_profiles/B_philippinensis.htm).

Females of *B. dorsalis* from different localities/countries have significantly different average aculeus length in descending order India>Thailand>Hawaii>Taiwan (Mahmood 1999). Many females from these places, except Taiwan, have aculeus length that overlaps with that of *B. papayae* and *B. philippinensis* (Mahmood 2004). This together with other morphometric data led Mahmood (2004) to conclude that “no tenable method has been found to separate specimens of *B. dorsalis* from *B. papayae*

and *B. philippinensis* using external morphological characters”.

It was reported that females of *B. philippinensis* may be differentiated from *B. papayae* by having shorter length of scales found on the distal end of the eversible membrane of the ovipositor (Drew and Hancock, 1994). Nevertheless, recent work via electron scanning microscopy by Mahmood (2004) showed no difference in the length of scales on the distal end of the eversible membrane of the ovipositor between *B. papayae* and *B. philippinensis*. These inconsistencies show that the morphological characters and their morphometrics may be more consistent with population-level, rather than species-level, variations. Furthermore, phylogenetic studies have shown that *B. philippinensis* is monophyletic with respect to *B. dorsalis* s.s. and *B. papayae* (Armstrong and Cameron 2000; Muraji and Nakahara 2001; Zhang et al. 2010; Krosch 2012a, b). Therefore, it is of great importance and urgency to understand the phylogenetic relationship using mitochondrial genes sequences in conjunction with other characters such as sex pheromone profiles, which would provide a distinctive feature on diversification among the sibling species (Tan et al. 2011).

Like other notorious *Bactrocera* pest species, males of *B. philippinensis* are strongly attracted to and compulsively feed on ME, which is found naturally in over 450 plant species from 80 families spanning across 38 orders (Tan and Nishida 2012). Consumption of ME significantly improves male mating performance and competitiveness of *B. dorsalis* s.s. (Shelly and Dewire 1994, Shelly 2000; Tan and Nishida 1996; Shelly and Nishida 2004; Orankanok et al. 2011), *B. papayae* (Tan and Nishida 1996, 1998), *B. carambolae* (Wee et al. 2007). ME acts as a sex pheromone

precursor in *B. dorsalis* s.s. and *B. papayae* in which ME is biotransformed to (*E*)-coniferyl alcohol (E-CF) and 2-allyl-4,5-dimethoxyphenol (DMP) (Fig. 1) (Nishida et al. 1988; Tan and Nishida 1996, 1998). The two volatile phenylpropanoids, temporarily stored in the rectal gland and emitted at dusk prior to courtship, were shown to act as sex pheromone that attracted conspecific females of *B. dorsalis* s.s. (Nishida et al. 2000) and *B. papayae* (Tan and Nishida 1996, 1998; Hee and Tan 1998; Khoo et al. 2000). The same ME metabolites, E-CF and DMP, were also detected in the male rectal gland of *B. invadens*, a highly invasive species in the African continent (Tan et al. 2011).

After consuming ME, irradiated males of *B. philippinensis* showed a mating advantage compared with males never exposed to the attractant (Shelly et al. 1996). Since the rectal/sex pheromone volatiles have not been identified in this species, we aim to determine the fate of ME after consumption by males and to examine the phylogenetic relationship of *B. philippinensis* in relation to three other putative species, *B. dorsalis* s.s., *B. invadens* and *B. papayae*, within the *B. dorsalis* complex, by using the mitochondrial COI and rDNA sequences. Lately, a study showed that *B. dorsalis* s.s., *B. papayae* and *B. philippinensis* represent one biological species based on independent data sets on mitochondrial DNA and wing shape (Schutze et al. 2012).

Materials and methods

Insects

Bactrocera philippinensis, originally from Guimaras, Philippines, and positively identified by R.A.I. Drew in 2009, was cultured in the FAO/IAEA Agriculture and Biotechnology Laboratory, Seibersdorf, Austria. Due to strict quarantine restrictions, experiments were conducted exclusively on sexually mature males (21-25 day-old) at the Seibersdorf laboratory.

Chemical analysis

Gas chromatography-mass spectrometry (GC-MS) was performed with an Agilent 5975 inert XL MSD mass spectrometer (electron impact ionization at 70 eV) linked to an Agilent 6890 gas chromatograph equipped with a HP-5MS column (28 m × 0.25 mm, 0.25 µm film thickness), using helium as a carrier gas and programmed from 60°C (1 min holding) to 280°C at a rate of 10°C/min. The GC quantification was carried out with an HP 5890 series II plus using a HP-1 column (30 m × 0.25 mm, 0.25 µm film thickness). The oven temperature was programmed from 60°C (2 min holding) to 240°C at a rate of 10°C/min using 1-hexadecanol (Wako Pure Chemical Industries, Japan) as an internal standard. The carrier gas was helium; and detection was by a flame ionization detector (FID).

Feeding test and rectal sample preparation

Males of *B. philippinensis* (21-25 day old; approximately the 50th generation) were allowed to feed on methyl eugenol (Aldrich Chemicals Co., USA) impregnated into small filter paper discs (Advantec, antibiotic test disc, thick, 8 mm diameter) (10

131 $\mu\text{l}/\text{disc}$) *ad libitum* for *ca.* 15 min and were then kept in a cylindrical plastic rearing
132 cage (15 cm diameter x 20 cm height) with a sufficient amount of adult food
133 (sugar-yeast hydrolysate mixture) and water at 24-30°C under ambient light conditions.

134 Rectal glands were dissected from the males at 6, 12, 24 and 48 hours following ME
135 feeding. Each extracted gland was individually preserved in 250 μl of redistilled
136 ethanol in a 1 ml screw-capped glass vial, and then, kept in a freezer at -20°C until
137 chemical analysis. The significant differences of the contents between DMP and E-CF
138 were analyzed by Wilcoxon signed-ranks test.

140 Field trapping of wild male flies

141 Wild males were captured using ME-baited sticky traps at Castillejos, Zambales,
142 Philippines (Luzon Island) in 2007. The whole fly was carefully extracted from the
143 sticky trap and preserved individually in a vial containing redistilled ethanol (*ca* 250 μl).
144 Each of the specimens was subjected to morphological identification as *B.*
145 *philippinensis* before their rectal glands were extracted individually. A total of 18 males
146 were examined chemically.

148 Identification and quantification of ME-metabolites in male rectal gland

149 Chemical identification was based on comparison of their retention time and mass
150 fragment pattern with those of authentic chemicals. Quantification of the
151 ME-metabolites was conducted as previously described by Tan et al. (2011).

153 Molecular cloning and sequence analysis

154 DNA extraction from individual adults, PCR amplification, and sequence analysis were
155 performed as described by Tan et al. (2011). PCR amplifications were performed for the
156 mitochondrial cytochrome oxidase subunit I (COI) gene and rDNA containing a part of
157 the 16S rRNA gene, the tRNA^{val} gene, and a part of the 12S rRNA gene. The forward (f)
158 and reverse (r) primer pairs were as follows: for COI, f
159 (5'-ATTTATAATGTAATTGTAACAGC-3') and r
160 (5'-GAAGTATTTAARTTTCGRTCTG-3'); and for rDNA, f
161 (5'-TTCAGTGGGCAGGTCAGACT-3') and r (5'-
162 ATATGCACACATCGCCCGTC-3'). The PCR products were cloned into the pGEM-T
163 Easy vector (Promega, WI, USA) and sequences of the clones were determined using
164 T7 and SP6 universal primers. The DNA sequence data have been deposited in the
165 DDBJ/EMBL-Bank/GenBank as listed in Table 1.

167 Phylogenetic analysis

168 The DNA sequences were determined for *B. dorsalis* s.s., *B. invadens*, *B. papayae*, *B.*
169 *philippinensis* (these species are taxonomically real, yet biologically dubious, entities)
170 and another sibling species of the *dorsalis* complex, *B. carambolae*, which has a
171 distinctive sex pheromone profile compared with the four species investigated; and two
172 unrelated ME-responsive species, *B. correcta* and *B. zonata* were included in the
173 phylogenetic analysis to provide an insight into the relationship between sex pheromone
174 diversification and genetic distance amongst the fruit fly species. The corresponding

genes of the complete mitochondrial genome of *B. cucurbitae*, a non-ME responsive *Bactrocera* species but a cue-lure- and raspberry ketone-responsive pest (JN635562), were used as out groups.

Sequence alignment was performed using the program Clustal W 1.8.3. The phylogenetic trees were generated by aligning the determined sequences excluding the primer regions together with those of Tokushima et al. (2010) and Tan et al. (2011) using the corresponding genes of *B. cucurbitae* as an out group. The maximum likelihood (ML) method was conducted by MEGA 5 (Tamura et al. 2011). Parameters were based on the General Time Reversible model with among-site rate heterogeneity according to a Gamma distributed with Invariant sites (G+I).

Results

Identification and accumulation of methyl eugenol metabolites in the rectal glands

Two ME-derived phenylpropanoids, DMP and E-CF (Fig. 1), were detected from rectal glands of male *B. philippinensis* following ME feeding. The identity of the metabolites were confirmed by GC-MS data as follows.

DMP (2-allyl-4,5-dimethoxyphenol). GC: RI (relative retention index on HP-5MS):

1630. MS: m/z (%) 194 (100, M^+), 179 (87), 163 (10), 151 (16), 136 (7), 133 (8), 123 (32), 119 (9), 105 (7), 95 (10), 91 (23), 79 (11), 77 (16), 69 (27), 53 (11).

E-CF ((*E*)-Coniferyl alcohol). GC: RI 1748. MS m/z (%): 180 (71, M^+), 162 (10), 147 (13), 137 (100), 131 (15), 124 (52), 119 (25), 109 (13), 103 (18), 91 (39), 77 (19), 65

(14), 55 (12).

Figure 2 shows the quantities of DMP and E-CF present in the rectal gland of male *B. philippinensis* at 6, 12, 24 and 48 h after feeding on ME. Both phenylpropanoids increased in quantity with time and peaked at 24 h, and thereafter a decrease was observed at 48 h post feeding. The total quantity of the two compounds was approximately 20 µg/gland at 24 h post ME feeding. The individual contents of DMP and E-CF in the rectal gland of the males were variable at any given time after feeding on ME. While no significant difference in contents between DMP and E-CF was detected at 6h and 48h post ME feeding, the contents of E-CF were significantly higher than those of DMP at 12h and 24h post ME feeding (Wilcoxon signed-ranks test, 12h: $Z = -2.24$, $N = 8$, $P = 0.025$; 24h: $Z = -1.99$, $N = 6$, $P = 0.046$).

Phenylpropanoid contents in the wild males

Some of the wild *B. philippinensis* male flies captured by ME-baited sticky traps in Luzon, Philippines, lacked both DMP and E-CF or either one of the metabolites. Hence, it contributed to the high variation in total contents per fly [Mean content $\pm$ S.E.: DMP 4.7 ± 6.7 µg/gland; E-CF, 1.6 ± 3.5 µg/gland]. The ratio of DMP and E-CF detected in the rectal glands varied significantly for wild males with a bias towards DMP, while laboratory-raised males have less variation (Fig. 3).

Phylogenetic analyses of *B. philippinensis* in relation to other *Bactrocera* species

The fragments of COI and rDNA were amplified by PCR, and the length of

the resulting nucleotide was 497 bp and from 884 to 888 bp, respectively. Both COI and rDNA analyses show that a lineage comprising of *B. philippinensis*, *B. dorsalis* s.s., *B. invadens*, *B. papayae* and *B. carambolae* was quite distinct from the two individual lineages derived from *B. zonata* and *B. correcta* (Fig. 4). These phylogenetic trees based on the mitochondrial genes clearly show that *B. philippinensis* belongs to the same clade as the *B. dorsalis* s.s. as well as *B. invadens*, *B. papayae* and *B. carambolae*.

Discussion

After consuming ME, two rectal (sex pheromone) volatile components, DMP and E-CF, were detected in *B. philippinensis* males. These phenylpropanoid components were also previously detected in the other three closely related species of the *B. dorsalis* complex after pharmacophagy of ME, namely *B. dorsalis* s.s. (Nishida et al., 2000) and *B. papayae* (Tan and Nishida 1996), and *B. invadens* (Tan et al. 2011). But it was found to differ partially from its sibling species, *B. carambolae* (Wee and Tan 2007), and totally from other ME-responsive species that are not within the *dorsalis* complex – namely *B. correcta* and *B. zonata* (Tokushima et al. 2010, Tan et al. 2011). Insect sex pheromone is often highly species-specific and often serves as an important chemical cue, although not solely responsible, for mate recognition and helps to delimit gene flow among different populations through different blend, ratios and isomers, as in the European ermine moth, *Yponomeuta* spp. (Löfstedt et al. 1991). Therefore, the presence of identical sex pheromone profiles of DMP and E-CF among the four species strongly

indicates that the species belong to the same biological species because they possess the same ‘chemical’ language for sexual communication. It is speculated that if their movements had not been restricted by international trade and quarantine restrictions, they would have been able to ‘meet’ and mate amongst themselves without difficulty. Previous mating study conducted in the Philippines also showed that there is high reproductive compatibility between *B. philippinensis* and *B. dorsalis* s.s. (Medina et al. 1988). Additionally, mating compatibility studies conducted by A. Jessup, in the FAO/IAEA Agriculture and Biotechnology Laboratory, Seibersdorf, Austria, have shown that *B. invadens* and *B. philippinensis* were able to interbreed readily with *B. dorsalis* s.s. and reproduced “highly fertile F₁ and F₂ hybrids” (A. Jessup, IAEA — personal communication 2010, 2011).

Similarly, the ability to interbreed between *B. dorsalis* s.s. and *B. papayae* and yield viable offspring up to F₃ (Tan 2003) indicated that the two sibling species are not distinct biological species. This is further supported by the genetic evidence that one of three actin gene alleles in *B. dorsalis* s.s. and *B. papayae* – allele BdorA1 and allele BpapA2, respectively, have identical DNA sequence (Naeole and Haymer 2003). Hence, *B. papayae* has been referred to as *B. dorsalis sensu lato*. Recent population genetic analysis also corroborated these species as one and the same biological species (Schutze et al. 2012; Krosch et al. 2012a).

Similar trend of variations in the ratio of sex pheromone components was also reported in *B. papayae* males whereby laboratory raised *B. papayae* males produced higher amount of E-CF than DMP and the vice versa was seen in the wild conspecific

263 males (Wee and Tan 2001). Prolonged inbreeding in enclosures may induce insects to
264 adapt to laboratory conditions that may affect changes in the pheromone titers
265 (Giebultowicz et al. 1992; Raina et al. 1989). It was also suggested that the production
266 and ratio of DMP to E-CF may be indirectly related to the contemporary needs of
267 individual flies, as DMP was demonstrated to be a much weaker sex attractant than
268 E-CF, for *B. papayae* females (Hee and Tan, 1998). In addition, DMP was observed to
269 deter birds better than ME, followed by E-CF (Nishida and Fukami, 1990). Thus, it was
270 hypothesized that the E-CF as a sex pheromone component is more crucial than
271 allomone component (DMP) in the laboratory where the need to deter predators is
272 greatly reduced hence the consistence in higher E-CF to DMP ratio. Additionally, the
273 variations may indicate that the ratio of the sex pheromone components may not be a
274 good indicator for distinguishing tephritid species as the ratio may be dependent on
275 natural sources (often unknown) of the chemicals which the males feed on, as found in
276 certain *Bulbophyllum* orchid floral fragrance or secretion that contains either one or
277 both of the sex pheromone components besides ME (Tan et al. 2002, 2006).

278 The wild *B. philippinensis* males that did not possess ME-metabolites may
279 indicate that they had either no opportunity to feed on natural sources of ME or likely
280 just attained sexual maturity to be attracted for the first time to ME to initiate
281 pharmacophagy. This phenomenon was also shown in wild *B. papayae* (Nishida et al.
282 1988; Tan and Nishida 1996, 1998; Wee and Tan 2001). Further work is needed to
283 clarify the nature of pheromone acquisition, sequestration, emission and behavioral
284 effects on the courtship sequences in *B. philippinensis*, particularly under its natural

environment, in comparison with those studies conducted for other sibling species.

Phylogenetic analyses on *B. philippinensis* from Guimaras and Luzon, Philippines — based on the comparison of nucleotide sequences in the mitochondrial genes, COI and rDNA — showed that the species belongs to the same species clade as *B. carambolae*, *B. dorsalis* s.s., *B. invadens* and *B. papayae* but differs from the *B. zonata* species complex which is consistent with previous rDNA analysis by Muraji and Nakahara (2001) and Krosch et al. (2012a). Similar phylogenetic relationship based on the two mitochondrial genes was obtained for seven pest sibling species, namely *B. carambolae*, *B. dorsalis* s.s., *B. kandiensis*, *B. musae* (Tryon), *B. occipitalis*, *B. papayae* and *B. philippinensis*, which appeared to have a common ancestor, within the *B. dorsalis* complex (Zhang et al. 2010). This further shows that *B. philippinensis* is indeed very closely related to the other ME-responsive sibling species within the *B. dorsalis* species complex. Furthermore, a recent phylogenetic analysis using fruit flies collected in the Ryukyu Islands, Taiwan, and the Philippines, and from fruits imported from the Philippines and China intercepted at Narita International Airport showed fruit flies possessing one of five banding patterns were grouped as *B. philippinensis* based on the sequence of 1,138 bp mitochondrial DNA fragment from 16S to 12S rDNA (Muraji et al. 2008). This led to an inference that the fruit flies collected in Sakishima region, Japan, might have flown in directly from the Philippines.

In conclusion, the above evidence in this paper – chemical ecology and chemotaxonomy as well as DNA analyses together with other evidence from different perspectives, i.e., mating compatibility and hybrid viability, as well as molecular

analyses, strongly indicate that *B. philippinensis* and its three very close sibling species, namely – *B. dorsalis* s.s., *B. invadens* and *B. papayae*, belong to the same biological species. Therefore, based on the integrative and comprehensive approach, these four species, currently discriminated by indistinct and variable morphological characters that are unreliable for differentiating the sibling species (Mahmood 2004), should be recognized taxonomically as either different populations or strains of a single biological species, *B. dorsalis*. More importantly, they should not remain as the status quo of four distinct putative species – currently based solely on indistinct morphological characters, which have caused much confusion worldwide in the identification of the sibling species in quarantine surveys and interceptions, as well as untold economic losses due to quarantine restrictions in international trade, especially for third world or developing countries.

Acknowledgements

The authors would like to thank Andrew Jessup and Mark Schutze (FAO/IAEA Agriculture and Biotechnology Laboratory, Seibersdorf, Austria) for the supply of sexually mature live male flies of *B. philippinensis* and specimens of *B. invadens* in alcohol, W. Orankanok and S. Chinvinijkul (Department of Agricultural Extension, Bangkok) for *B. correcta* and *B. dorsalis*, S. Permalloo (Entomology Division, Ministry of AgroIndustry and Fisheries, Mauritius) for *B. zonata*, and Hiroshi Enomoto for the wild *B. philippinensis* flies from Luzon island. Two of us (Tan and Wee) also express our sincere gratitude to Jorge Hendrichs of International Atomic Energy Agency, Vienna

for sponsoring our trip to Vienna during which feeding and extraction experiments were conducted. This work was partly supported by International Atomic Energy Agency-Coordinated Research Projects Contract numbers 16066 & 16160, and the Grant-in-Aid for Scientific Research from JSPS (No. 23380035) from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and JST in Research for Promoting Technological Seeds (under R. Nishida).

References

- Armstrong KF, Cameron CM (2000) Species identification of tephritids across a broad taxonomic range using ribosomal DNA. In: Tan KH (ed) Area-wide Control of Fruit Flies and other Insect Pests. Penerbit USM, Penang, Malaysia, pp 203-710
- Clarke AR, Armstrong KF, Carmichael AE, Milne JR, Raghu S, Roderick GK, Yeates DK (2005) Invasive phytophagous pests arising through a recent tropical evolutionary radiation: The *Bactrocera dorsalis* complex of fruit flies. Annu Rev Entomol 50: 293-319
- Drew RAI, Hancock DL (1994) The *Bactrocera dorsalis* complex of fruit flies (Diptera: Tephritidae: Dacinae) in Asia. Bull Entomol Res Suppl. 2: 1-68
- Giebultowicz JM, Webb RE, Raina AK, Ridgway RL (1992) Effects of temperature and age on daily changes in pheromone titer in laboratory reared and wild gypsy moth (Lepidoptera: Lymantriidae). Environ Entomol 21: 822-826

- 1
- 2
- 3 350 Hee AKW, Tan KH (1998) Attraction of female and male *Bactrocera papayae* to
- 4
- 5 351 conspecific males fed with methyl eugenol and attraction of females to male sex
- 6
- 7
- 8 352 pheromone components. J Chem Ecol 24: 753-764
- 9
- 10
- 11 353 Khoo CCH, Yuen KH, Tan KH (2000) Attraction of female *Bactrocera papayae* to sex
- 12
- 13 354 pheromone components with two different release devices. J Chem Ecol 26:
- 14
- 15 355 2487-2496
- 16
- 17
- 18 356 Krosch MN, Schutze MK, Armstrong KF, Graham GC, Yeates DK, Clarke AR (2012a)
- 19
- 20
- 21 357 A molecular phylogeny for the Tribe Dacini (Diptera: Tephritidae): systematic and
- 22
- 23 358 biogeographic implications. Mol Phylogenet Evol 64: 513-523
- 24
- 25
- 26 359 Krosch MN, Schutze MK, Armstrong KF, Boontop Y, Boykin LM, Chapman TA,
- 27
- 28 360 Englezou A, Cameron SL, Clarke AR (2012b) Piecing together an integrative
- 29
- 30 361 taxonomic puzzle: microsatellite, wing shape and aedeagus length analyses of
- 31
- 32 362 *Bactrocera dorsalis* s.l. (Diptera: Tephritidae) find no evidence of multiple lineages
- 33
- 34 363 in a proposed contact zone along the Thai/Malay Peninsula. Syst Entomol 38: 2-13
- 35
- 36
- 37 364 Löfstedt C, Herrebout WM, Menken SBJ (1991) Sex pheromones and their potential
- 38
- 39 365 role in the evolution of reproductive isolation in small ermine moths
- 40
- 41 366 (Yponomeutidae). Chemoecology 2: 20-28.
- 42
- 43
- 44 367 Mahmood K (1999) Intraspecific variations in two pest species of the Oriental fruit fly
- 45
- 46 368 *Bactrocera dorsalis* (Hendel) (Tephritidae: Diptera) complex. Pak J Zool. 31:
- 47
- 48 369 315-321
- 49
- 50
- 51 370 Mahmood K (2004) Identification of pest species in Oriental fruit fly, *Bactrocera*
- 52
- 53 371 *dorsalis* (Hendel) (Diptera: Tephritidae) species complex. Pak J Zool 36: 219-230
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1
- 2
- 3 372 Medina FIS, Carillo PAV, Gregorio JS, Aguilar CP (1998) The mating compatibility
- 4
- 5 373 between *Bactrocera philippinensis* and *Bactrocera dorsalis*. Abstracts of 5th
- 6
- 7
- 8 374 international symposium on fruit flies of economic importance, June 1-5, 1998
- 9
- 10
- 11 375 Penang, Malaysia, p 155
- 12
- 13 376 Muraji M, Nakahara S (2001) Phylogenetic relationships among fruit flies, *Bactrocera*
- 14
- 15
- 16 377 (Diptera: Tephritidae), based on the mitochondrial rDNA sequences. Insect Mol Biol
- 17
- 18
- 19 378 10: 549–559
- 20
- 21 379 Muraji M, Nakahara S, Ishida T, Minoura K, Miyazaki I, Kohama T (2008) The
- 22
- 23
- 24 380 Philippines is a possible source of the *Bactrocera dorsalis* complex species
- 25
- 26
- 27 381 (Diptera:Tephritidae) occasionally collected in the Ryukyu Islands of Japan;
- 28
- 29 382 analyses of mitochondrial DNA. Appl Entomol Zool 43: 609-615
- 30
- 31
- 32 383 Naeole CKM, Haymer DS (2003) Use of oligonucleotide arrays for molecular
- 33
- 34
- 35 384 taxonomic studies of closely related species in the Oriental fruit fly (*Bactrocera*
- 36
- 37 385 *dorsalis*) complex. Mol Ecol Notes 3: 662–665
- 38
- 39
- 40 386 Nishida R, Tan KH, Serit M, Lajis NH, Sukari AM, Takahashi S, Fukami H (1988)
- 41
- 42
- 43 387 Accumulation of phenylpropanoids in the rectal glands of males of the Oriental fruit
- 44
- 45 388 fly, *Dacus dorsalis*. Experientia 44: 534-536
- 46
- 47
- 48 389 Nishida R, Fukami H (1990) Sequestration of distasteful compounds by some
- 49
- 50
- 51 390 pharmacophagous insects. J Chem Ecol 16: 151-164
- 52
- 53 391 Nishida R, Shelly TE, Kaneshiro KY, Tan KH (2000) Roles of semiochemicals in
- 54
- 55
- 56 392 mating systems: a comparison between oriental fruit fly and medfly. In: Tan KH (ed)
- 57
- 58
- 59 393 Area-wide control of fruit flies and other insect pests. Penerbit USM, Penang,
- 60
- 61
- 62
- 63
- 64
- 65

- 1
- 2
- 3 394 Malaysia, pp 631–637
- 4
- 5
- 6 395 Orankanok W, Chinvinijkul S, Sawatwangkhong A, Pinkaew S, Orankanok S (2011)
- 7
- 8 396 Methyl eugenol and pre-release diet improve mating performance of young
- 9
- 10
- 11 397 *Bactrocera dorsalis* and *Bactrocera correcta* males. J Appl Entomol doi:
- 12
- 13 398 10.1111/j.1439-0418.2011.01677.x
- 14
- 15
- 16 399 Raina AK, Stadelbacher EA, Ridgway RL (1989) Comparison of sex pheromone
- 17
- 18 400 composition and pheromone-mediated male behavior of laboratory-reared and wild
- 19
- 20
- 21 401 *Heliothis zea* (Lepidoptera: Noctuidae). J Chem. Ecol 15: 1259-1265
- 22
- 23
- 24 402 Schutze MK, Krosch MN, Armstrong KF, Chapman TA, Englezou A, Chomič A,
- 25
- 26 403 Cameron SL, Hailstones D, Clarke AR (2012) Population structure of *Bactrocera*
- 27
- 28 404 *dorsalis* s.s., *B. papayae* and *B. philippinensis* (Diptera: Tephritidae) in southeast
- 29
- 30 405 Asia: evidence for a single species hypothesis using mitochondrial DNA and
- 31
- 32 406 wing-shape data. BMC Evol Biol 12:130
- 33
- 34
- 35 407 Shelly TE (2000) Flower-feeding affects mating performance in male Oriental fruit flies
- 36
- 37 408 *Bactrocera dorsalis*. Ecol Entomol 25: 109-114
- 38
- 39
- 40 409 Shelly TE, Dewire AM (1994) Chemically mediated mating success in male Oriental
- 41
- 42 410 fruit flies (Diptera: Tephritidae). Ann Entomol Soc Am 87: 375-382
- 43
- 44
- 45 411 Shelly TE, Nishida R (2004) Larval and adult feeding on methyl eugenol and the mating
- 46
- 47 412 success of male oriental fruit flies, *Bactrocera dorsalis* (Hendel) (Diptera:
- 48
- 49 413 Tephritidae). Entomol Exp Appl 112:155-158
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1
- 2
- 3 414 Shelly T, Resilva S, Reyes M, Bignayan H (1996) Methyl eugenol and mating
- 4
- 5 415 competitiveness of irradiated male *Bactrocera philippinensis* (Diptera: Tephritidae).
- 6
- 7
- 8 416 Fla Entomol 79: 481-488
- 9
- 10
- 11 417 Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5:
- 12
- 13 418 Molecular evolutionary genetics analysis using maximum likelihood, evolutionary
- 14
- 15 419 distance, and maximum parsimony methods. Mol Biol Evol 28: 2731-2739
- 16
- 17
- 18 420 Tan KH (2003) Interbreeding and DNA analysis of sibling species within the
- 19
- 20 421 *Bactrocera dorsalis* complex. In: Recent trends on sterile insect technique and
- 21
- 22 422 area-wide integrated pest management – economic feasibility, control projects,
- 23
- 24 423 farmer organization and *Bactrocera dorsalis* complex control study. Research
- 25
- 26 424 Institute for Subtropics, Okinawa, Japan, pp 113-122
- 27
- 28
- 29 425 Tan KH, Nishida R (1996) Sex pheromone and mating competition after methyl eugenol
- 30
- 31 426 consumption in the *Bactrocera dorsalis* complex. In: McPherson BA, Steck, GJ (eds)
- 32
- 33 427 Fruit fly pests: a world assessment of their biology and management St. Lucid Press,
- 34
- 35 428 Florida, USA, pp 147-153
- 36
- 37
- 38 429 Tan KH, Nishida R (1998) Ecological significance of male attractant in the defence and
- 39
- 40 430 mating strategies of the fruit fly, *Bactrocera papayae*. Entomol Exp Appl 89:
- 41
- 42 431 155-158
- 43
- 44
- 45 432 Tan KH, Nishida R (2012) Methyl eugenol: Its occurrence, distribution, and role in
- 46
- 47 433 nature, especially in relation to insect behavior and pollination. J Insect Sci 12 (56):
- 48
- 49 434 1-74
- 50
- 51 435 Tan KH, Nishida R, Toong YC (2002) Floral synomone of a wild orchid, *Bulbophyllum*
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1
- 2
- 3 436 *cheiri*, lures *Bactrocera* fruit flies for pollination. J Chem Ecol 28:1161-1172
- 4
- 5
- 6 437 Tan KH, Tan LT, Nishida R (2006) Floral phenylpropanoid cocktail and architecture of
- 7
- 8 438 *Bulbophyllum vinaceum* orchid in attracting fruit flies for pollination. J Chem Ecol
- 9
- 10
- 11 439 32: 2429-2441
- 12
- 13
- 14 440 Tan KH, Tokushima I, Ono H, Nishida R (2011) Comparison of phenylpropanoid
- 15
- 16 441 volatiles in male rectal pheromone gland after methyl eugenol consumption, and
- 17
- 18 442 molecular phylogenetic relationship of four global pest fruit fly species: *Bactrocera*
- 19
- 20
- 21 443 *invadens*, *B. dorsalis*, *B. correcta* and *B. zonata*. Chemoecology 21: 25–33
- 22
- 23
- 24 444 Tokushima I, Orankanok W, Tan KH, Ono H, Nishida R (2010) Accumulation of
- 25
- 26 445 phenylpropanoid and sesquiterpenoid volatiles in male rectal pheromonal glands of
- 27
- 28 446 the guava fruit fly, *Bactrocera correcta*. J Chem Ecol 36: 1327–1334
- 29
- 30
- 31
- 32 447 Wee SL, Tan KH (2001) Allomonal and hepatotoxic effects following methyl eugenol
- 33
- 34 448 consumption in *Bactrocera papayae* male against *Gekko monarchus*. J Chem Ecol
- 35
- 36 449 27: 953-964
- 37
- 38
- 39
- 40 450 Wee SL, Tan KH (2007) Temporal accumulation of phenylpropanoids in male fruit flies,
- 41
- 42 451 *Bactrocera dorsalis* and *B. carambolae* (Diptera: Tephritidae) following methyl
- 43
- 44 452 eugenol consumption. Chemoecology 17: 81-85
- 45
- 46
- 47
- 48 453 Wee, SL, Tan, KH, Nishida R (2007) Pharmacophagy of methyl eugenol by males
- 49
- 50 454 enhances sexual selection of *Bactrocera carambolae*. J Chem Ecol 33: 1272-1282
- 51
- 52
- 53 455 Zhang B, Liu YH, Wu WX, Wang ZL (2010) Molecular phylogeny of *Bactrocera*
- 54
- 55 456 species (Diptera: Tephritidae: Dacini) inferred from mitochondrial sequences of 16S
- 56
- 57 457 rDNA and COI sequences. Fla Entomol 93: 369-377
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

Legends

Fig. 1 Chemical structures of methyl eugenol (ME) and its metabolites, 2-allyl-4,5-dimethoxyphenol (DMP) and *E*-coniferyl alcohol (E-CF) sequestered by *Bactrocera philippinensis* males post feeding on ME.

Fig. 2 Contents (mean $\pm$ S.E.) of 2-allyl-4,5-dimethoxyphenol (DMP: white) and *E*-coniferyl alcohol (E-CF: gray) sequestered by *Bactrocera philippinensis* males 6, 12, 24 and 48 h post feeding on methyl eugenol ($N = 8$ for 6, 12 and 48 h; $N = 6$ for 24 h).

Fig. 3 Contents of 2-allyl-4,5-dimethoxyphenol (DMP) and *E*-coniferyl alcohol (E-CF) accumulated in the individual rectal glands of wild *Bactrocera philippinensis* males captured in a field site in Luzon Island, Philippines (black) ($N = 18$) and laboratory males artificially fed on methyl eugenol (48 hr post treatment) (circle) ($N = 8$).

Fig. 4 Phylogenetic trees constructed by maximum likelihood (ML) analysis. The values shown at the nodes of the branches are bootstrap values (%) from 1000 replicate bootstrap samplings. Branch length is proportional to ML estimated genetic distances. The scale bars indicate the number of nucleotide substitutions per site. GenBank accession numbers are listed in Table 1. (A) Phylogenetic tree

1
2
3 480 deduced from COI sequences. (B) Phylogenetic tree deduced from rDNA
4
5
6 481 sequences.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. List of the Genbank accession numbers of *Bactrocera* species analysed

Accession number	Gene	Species	Geographic origin (year)	References
AB720885	COI	<i>B. philippinensis</i>	IAEA (Vienna) –A, origin: Guimaras, PH (2010)	This study
AB720891	rDNA	<i>B. philippinensis</i>	IAEA (Vienna) –A, origin: Guimaras, PH (2010)	This study
AB720886	COI	<i>B. philippinensis</i>	IAEA (Vienna) –B, origin: Guimaras, PH (2010)	This study
AB720892	rDNA	<i>B. philippinensis</i>	IAEA (Vienna) –B, origin: Guimaras, PH (2010)	This study
AB720882	COI	<i>B. philippinensis</i>	Wild-A, Luzon, PH (2007)	This study
AB720893	rDNA	<i>B. philippinensis</i>	Wild-A, Luzon, PH (2007)	This study
AB720883	COI	<i>B. philippinensis</i>	Wild-B, Luzon, PH (2007)	This study
AB720894	rDNA	<i>B. philippinensis</i>	Wild-B, Luzon, PH (2007)	This study
AB720884	COI	<i>B. philippinensis</i>	Wild-C, Luzon, PH (2007)	This study
AB720895	rDNA	<i>B. philippinensis</i>	Wild-C, Luzon, PH (2007)	This study
AB568103	COI	<i>B. dorsalis s.s.</i>	Laboratory, Bangkok, TH (2009)	Tan et al. (2011)
AB568107	rDNA	<i>B. dorsalis s.s.</i>	Laboratory, Bangkok, TH (2009)	Tan et al. (2011)
AB720887	COI	<i>B. dorsalis s.s.</i>	Wild, Oahu, US (2002)	This study
AB720896	rDNA	<i>B. dorsalis s.s.</i>	Wild, Oahu, US (2002)	This study
AB568104	COI	<i>B. invadens</i>	IAEA (Vienna), origin: KE (2009)	Tan et al. (2011)
AB568108	rDNA	<i>B. invadens</i>	IAEA (Vienna), origin: KE (2009)	Tan et al. (2011)
AB720888	COI	<i>B. papayae</i>	Wild, Penang, MY (2007)	This study
AB720897	rDNA	<i>B. papayae</i>	Wild, Penang, MY (2007)	This study
AB720889	COI	<i>B. carambolae</i>	Wild, Selangor, MY (2008)	This study
AB720898	rDNA	<i>B. carambolae</i>	Wild, Selangor, MY (2008)	This study

AB720890	COI	<i>B. tryoni</i>	IAEA (Vienna), origin: Queensland, AU (2007)	This study
AB720899	rDNA	<i>B. tryoni</i>	IAEA (Vienna), origin: Queensland, AU (2007)	This study
AB720881	COI	<i>B. correcta</i>	Wild, Bangkok, TH (2007)	This study
AB569585	rDNA	<i>B. correcta</i>	Wild, Bangkok, TH (2007)	Tokushima et al. (2010)
AB721013	COI	<i>B. correcta</i>	Wild, Guava orchard ^a , TH (2009)	This study
AB569586	rDNA	<i>B. correcta</i>	Wild, Guava orchard ^a , TH (2009)	Tokushima et al. (2010)
AB568102	COI	<i>B. correcta</i>	Laboratory, Bangkok, TH (2009)	Tan et al. (2011)
AB568106	rDNA	<i>B. correcta</i>	Laboratory, Bangkok, TH (2009)	Tan et al. (2011)
AB568105	COI	<i>B. zonata</i>	Laboratory, MU (2009)	Tan et al. (2011)
AB568109	rDNA	<i>B. zonata</i>	Laboratory, MU (2009)	Tan et al. (2011)

^a 14°07'55.559" North; 100°48'28.762" East

Fig. 1

Methyl eugenol (ME)

DMP

E-CF

Fig. 2

Fig. 3

A

B