TITLE:

Characteristics of carbonaceous aerosols emitted from peatland fire in Riau, Sumatra, Indonesia

AUTHOR(S):

Fujii, Yusuke; Iriana, Windy; Oda, Masafumi;
Puriwigati, Astiti; Tohno, Sususmu; Lestari, Puji;
Mizohata, Akira; Huboyo, Haryono Setiyo

CITATION:

Fujii, Yusuke ...[et al]. Characteristics of carbonaceous aerosols emitted from peatland fire in Riau, Sumatra, Indonesia. *Atmospheric Environment* 2014, 87: 164-169

ISSUE DATE:

2014-04

URL:

<http://hdl.handle.net/2433/237621>

RIGHT:

© 2014. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>; The full-text file will be made open to the public on 1 April 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Characteristics of carbonaceous aerosols emitted from peatland fire in Riau, Sumatra, Indonesia

Yusuke Fujii^{1*}, Windy Iriana², Masafumi Oda¹, Astiti Puriwigati²,
Susumu Tohno¹, Puji Lestari², Akira Mizohata³, Haryono Setiyo Huboyo¹

¹ Department of Socio-Environmental Energy Science, Graduate School of Energy Science, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan.

² Department of Environmental Engineering, Bandung Institute of Technology, JL. Ganesha No. 10, Bandung 40132, Indonesia.

³ Radiation Research Center, Osaka Prefecture University, Sakai, Osaka 599-8531, Japan.

Abstract

Biomass burning is a significant source of fine particulate matter (PM_{2.5}). Forest, bush, and peat fires in Kalimantan and Sumatra, Indonesia are major sources of transboundary haze pollution in Southeast Asia. However, limited data exist regarding the chemical characteristics of aerosols at sources. We conducted intensive field studies in Riau Province, Sumatra, Indonesia, during the peatland fire and non-burning seasons in 2012. We characterized PM_{2.5} carbonaceous aerosols emitted from peatland fire based on ground-based source-dominated sampling. PM_{2.5} aerosols were collected with two mini-volume samplers using Teflon and quartz fiber filters. Background aerosols were also sampled during the transition period between the non-burning and fire seasons. We analyzed the carbonaceous content (organic carbon (OC) and elemental carbon (EC)) by a thermal optical reflectance utilizing the IMPROVE_A protocol and the major organic components of the aerosols by a gas chromatography/mass spectrometry. PM_{2.5} aerosols emitted from peatland fire were observed in high concentrations of $7120 \pm 3620 \mu\text{g}/\text{m}^3$ and were primarily composed of OC ($71.0 \pm 5.11\%$ of PM_{2.5} mass). Levoglucosan exhibited the highest total ion current and was present at concentrations of $464 \pm 183 \mu\text{g}/\text{m}^3$. The OC/EC ratios (36.4 ± 9.08), abundances of eight thermally-derived carbon fractions, OC/Levoglucosan ratios (10.6 ± 1.96), and Levoglucosan/Mannosan ratios (10.6 ± 2.03) represent a signature profile that is inherent in peatland fire. These data will be useful in identifying contributions from single or multiple species in atmospheric aerosol samples collected from peatland fires.

Keywords: Carbonaceous aerosol; PM_{2.5}; Levoglucosan; Mannosan; Biomass Burning

* Corresponding author. Tel: +81-75-753-5618; Fax: +81-75-753-5619

E-mail address: fujii.yusuke.86n@st.kyoto-u.ac.jp

35 **Highlights**

36

- 37 ➤ PM_{2.5} aerosols emitted from peatland fire in Indonesia were characterized.
- 38 ➤ PM_{2.5} aerosols emitted from peatland fire were primarily composed of OC.
- 39 ➤ We found some source indicators that were inherent in peatland fire.

40

1. Introduction

Peatland is organic soil that has formed for over thousands of years from decomposed vegetation and other life forms, and peat deposits can extend up to seven meters in thickness (Wulandari, 2002). In Southeast Asia, smoke originating from peatland fires in Kalimantan and Sumatra in Indonesia is a major cause of transboundary haze pollution.

In general, approximately 80–90% of the smoke particles produced by biomass burning is in the $PM_{2.5}$ size range, and these particles are primarily composed of organic carbon, which constitutes 50–60% of the total particle mass (Phuleria *et al.*, 2005; Schollnberger *et al.*, 2002; Reid *et al.*, 2005). $PM_{2.5}$ aerosols present a high risk of deposition in the alveoli of lungs and are associated with a greater general health risk than coarse aerosols (Federal Register, 2006; Lippmann, 1998).

According to version 3 of the Global Fire Emissions Database (GFED), average $PM_{2.5}$ emissions from fire (including deforestation, savanna, forest, agricultural waste, and peat fires) from 1997 to 2010 in Indonesia are 2.9 Tg/year, accounting for 9.2% of global fire $PM_{2.5}$ emissions and 62% of Southeast Asian fire emissions. In Indonesia, peatland fire is a dominant source of $PM_{2.5}$ emissions, accounting for 55% of all fire sources. Thus, it is a significant emission source of $PM_{2.5}$ aerosols in Indonesia.

Many peatland fires occur on Sumatra and Kalimantan islands during the dry season, emitting

60 gases and smoke aerosols that cause atmospheric pollution (haze) and adversely affect the health
61 of people living in surrounding areas. For example, an unprecedented Indonesian fire episode
62 occurred in 1997–1998 due to the El Niño–Southern Oscillation (ENSO) event at that time.
63 Extensive forest fires, including peatland fires, resulted in the development of a smog blanket that
64 covered an immense area of 4 million km² in Southeast Asia. This smog affected the livelihoods
65 and health of 75 million people in six countries and completely toppled their lives. Moreover, the
66 smog resulted in the closure of airports; it was cited as the possible cause of an air crash near
67 Medan, Sumatra, and a tanker collision in the Straits of Malacca (Stolle and Tomich, 1999). Air
68 pollution attained previously unknown levels in East Kalimantan, Singapore, and Kuala Lumpur,
69 with daily average particulate matter reaching extremely hazardous levels (4000 µg/m³; Heil *et*
70 *al.*, 1998). Although ENSO events considerably contribute to the occurrence of Indonesian fires,
71 pollution from smoke haze is a recurrent problem in Indonesia and neighboring countries, even in
72 non-ENSO years (Tacconi, 2003).

73 Riau Province in Sumatra is one of the primary hotspots for peatland fire during the dry season,
74 and the smoke aerosols generated there cause haze in Riau and in neighboring countries such as
75 Malaysia and Singapore (Harahap, 2012; Hong, 2012). However, limited data exist regarding the
76 chemical characteristics of these smoke aerosols (Othman and Latif, 2013; See *et al*, 2007) and
77 the effects of aerosols from peatland fires on the atmospheric environment and human health. To

investigate these effects, the chemical characterization of fresh smoke aerosols from peatland fire is necessary. In this study, the carbonaceous species of PM_{2.5} aerosols emitted from peatland fire were characterized by directly sampling PM_{2.5} aerosols at fire hotspots in Riau Province. Moreover, we determined source indicators of carbonaceous species of smoke from peatland fires for source apportionment. These data can help in identifying single or multiple species in atmospheric aerosol samples that contribute to peatland fires.

2. Materials and methods

2.1. Sampling locations

The sampling locations in this study are illustrated in Fig. 1. The burning site and background site were located at Sepahat Village and Sukajadi Village, respectively, in Bengkalis Regency, Riau Province. The burning site was surrounded by peatland and forest, and the background site was located ~50 km away from the burning site and housing estates. Bengkalis Regency lies on the east coast of Sumatra Island and consists of several islands. This district covers an area of 1,204,423 km², in which nearly 85% of the land exhibits low topography and is covered with tropical forests, with an average elevation of only 2.0–6.1 m above sea level. Most of the soils are organosols or peat soils; that is, they contain abundant organic substances. The temperature in a Bengkalis is strongly influenced by the tropical marine climate and is typically 26–32°C. The

rainy season generally lasts from September to January, with an average rainfall of 809–4078 mm/year, and the dry season usually spans February to August. Fig. 2 illustrates monthly hotspot counts in Riau in 2011 and 2012 based on Indofire datasets, confirming that many hotspots are detected in the area during the dry season.

2.2. Sample collection and analysis

In 2012, background and peatland fire PM_{2.5} aerosol samples were collected on May 16–17 and June 13–17, respectively. Two mini-volume samplers (MiniVol™ TAS, Airmetrics) were utilized to continuously collect PM_{2.5} aerosols on Teflon and quartz fiber filters for 24 h (background) and 2.5–5.2 h (peatland fire), respectively, at a flow rate of 5 L/min. PM_{2.5} aerosols were collected at several burning sites, about 1.5 m away from peatland fire hotspots and a background site on 7 and 4 occasions, respectively. Wind speeds in the peatland fire samplings ranged from 0.610 to 6.21 m/s (average: 2.50 m/s) and aerosols were collected in smoke plumes. Quartz fiber filters were heat-treated at 900°C for 4 h before sampling to remove any absorbed organic materials.

Filter samples were analyzed to determine PM_{2.5} mass concentrations, carbonaceous content (organic carbon (OC) and elemental carbon (EC)), and the quantity of key biomarkers.

The Teflon filter was weighed using a microbalance (ME5-F, Sartorius) with a sensitivity of ±1 µg in a stable environment of 25.7 ± 0.432°C (average ± standard deviation) and 26.0 ± 0.607

119 RH% before and after sampling, respectively, to determine PM_{2.5} mass concentrations.

120 The carbonaceous contents of the aerosols collected in the quartz fiber filters were quantified
121 using a DRI Model 2001 OC/EC Carbon Analyzer, which employs thermal optical reflectance
122 following the IMPROVE_A protocol. The IMPROVE_A temperature defines temperature
123 plateaus for thermally-derived carbon fractions as follows: 140°C for OC₁, 280°C for OC₂,
124 480°C for OC₃, and 580°C for OC₄ in helium (He) carrier gas; 580°C for EC₁, 740°C for EC₂,
125 and 840°C for EC₃ in a mixture of 98% He and 2% oxygen (O₂) carrier gas (Chow *et al.*, 2007).

126 OC, EC, and total carbon (TC) were calculated from the eight carbon fractions as follows:

$$127 \quad 128 \quad OC = OC_1 + OC_2 + OC_3 + OC_4 + OP \quad (1)$$

$$129 \quad 130 \quad EC = EC_1 + EC_2 + EC_3 - OP \quad (2)$$

$$131 \quad 132 \quad TC = OC + EC \quad (3)$$

133
134 where OP (the amount of pyrolyzed OC) is defined as the carbon content measured after the
135 introduction of O₂ until reflectance returns to its initial value at the start of analysis.

136 Two key biomarkers, levoglucosan and mannosan, obtained from the quartz fiber filters were
137 quantified by gas chromatography/mass spectrometry (GC/MS). Organic compound speciation
138 was basically accomplished, following the procedures of Fabbri *et al.*, 2009 and Pashynska *et al.*,
139 2002. Aliquots from the quartz fiber filter were spiked with the internal standard of methyl
140 β-L-arabinopyranoside before extraction. Each spiked filter was extracted by ultrasonic agitation

for 3 × 20 min periods using 3 mL of a dichloromethane/methanol mixture (3/1, v/v) (dichloromethane: Wako, purity >99.5%; methanol: Wako, purity >99.7%). The combined extracts were reduced to approximately 100 µL using a rotary evaporator (250 hPa, 40°C). Subsequently, the concentrated extract was filtered through a Teflon syringe filter (pore size 0.1 µm) and dried completely under a nitrogen stream. Prior to analysis, the total extracts were converted to trimethylsilyl derivatives by reaction with 150 µL of N,O-bis-(trimethylsilyl)-trifluoroacetamide with 1% trimethylchlorosilane and 90 µL of pyridine for 3 h at 70°C.

The derivatized samples were analyzed on a Shimadzu GC/MS system (GCMS-QP2010-Plus, Shimadzu) equipped with an Rtx-5Sil MS column (with selectivity similar to that of a mixture of 5% diphenyl and 95% dimethyl polysiloxane, 30 m × 0.25 mm I.D., 0.25 µm, RESTEK). Helium (purity 99.9995%) with an average velocity of 35.3 cm/s was used as the carrier gas. The GC oven temperature program was as follows: isothermal at 80°C for 5 min, 80–180°C at 3°C/min, then 180–300°C at 20°C/min, and maintained at 300°C for 5 min. The injection port and transfer line were maintained at 300°C. The data for quantitative analysis were acquired in the electron impact mode (70 eV).

3. Results and discussion

3.1. $PM_{2.5}$ mass concentration

Background and peatland fire $PM_{2.5}$ concentrations determined by gravimetric analysis were $23.9 \pm 2.53 \mu\text{g}/\text{m}^3$ and $7120 \pm 3620 \mu\text{g}/\text{m}^3$, respectively; that is, peatland fire $PM_{2.5}$ concentrations were ~300 times higher than background concentrations. These extremely high $PM_{2.5}$ concentrations present a great risk to human health, particularly in the case of fire fighters. Since the high concentration is due to the sampling at 1.5 m away from the fire, a different value would be obtained in the different sampling distance.

3.2. OC and EC

The average OC concentrations from peatland fire and background were $4970 \pm 3620 \mu\text{g}/\text{m}^3$ and $9.42 \pm 1.97 \mu\text{g}/\text{m}^3$, respectively, and average EC concentrations were $133 \pm 46.6 \mu\text{g}/\text{m}^3$ and $3.23 \pm 0.795 \mu\text{g}/\text{m}^3$, respectively. In terms of the carbonaceous fractions in $PM_{2.5}$ aerosols, $PM_{2.5}$ aerosols emitted from peatland fire were composed of $71.0 \pm 5.11\%$ OC and $2.05 \pm 0.509\%$ EC. Conversely, background $PM_{2.5}$ aerosols were composed of $39.1 \pm 5.08\%$ OC and $13.5 \pm 3.04\%$ EC. The remaining undetermined mass can be attributed to the various functional groups in organic aerosols, inorganic ions, metals, soil and particle bound water.

OC to EC mass ratios (OC/EC) provide some indication of the origins of carbonaceous $PM_{2.5}$ (Cao *et al.*, 2005; Chow *et al.*, 1996; Gray *et al.*, 1986; Turpin and Huntzicker, 1991). In this study, the average peatland fire and background OC/EC ratios were 36.4 ± 9.08 and 2.99 ± 0.738 ,

respectively. Few differences in the OC/EC ratio were observed, regardless of PM_{2.5} mass concentrations, and the coefficient of variance for samples from peatland fire was 24.9%. See *et al.*, 2007 reported that the average OC/EC ratio was 2.42 for PM_{2.5} aerosols collected in the open field within a 100 m circumference from the boundary of the Indonesian peatland fires in 2005. This value is much smaller than our result of 36.4. The discrepancy is due to the different measurement techniques for the OC-EC split. See *et al.*, 2007 adopted thermal method whose temperature of OC-EC split was 350°C. The method would cause significant overestimation of EC concentration in aerosol (Gelencsér, 2004). On the other hand, we utilized thermal-optical reflectance method with pyrolysis correction. OC/EC ratios from our peatland fire and other vegetative burning sources are illustrated in Fig. 3. OC/EC ratios ranged from 4.34 to 79.7; this discrepancy can be partly attributed to variations in the type and moisture content of the burning material.

Abundances of eight thermally-derived carbon fractions differ by carbon sources (Cao *et al.*, 2005; Chow *et al.*, 2004; Watson *et al.*, 1994). Fig. 4 illustrates the abundances of eight thermally-derived carbon fractions at both the peatland fire and background sites by mass percentage of total carbon. Distinct differences in carbon fractions were observed between the two sites. OC₁ accounted for $31.7 \pm 2.21\%$ of TC in peatland fire samples but only $0.710 \pm 0.721\%$ of TC in background samples. OC₂ accounted for $47.5 \pm 0.948\%$ of TC in peatland fire

samples and $13.9 \pm 1.64\%$ of TC in background samples. OC₃ accounted for $7.44 \pm 1.57\%$ of TC in peatland fire samples and $40.7 \pm 2.62\%$ of TC in background samples. OC₄ accounted for $1.22 \pm 0.412\%$ of TC in peatland fire samples and $15.7 \pm 1.27\%$ of TC in background samples. OP accounted for $9.28 \pm 1.39\%$ of TC in peatland fire samples and $3.95 \pm 4.13\%$ of TC in background samples. Lower EC fractions were observed in the peatland fire than the background samples. Thus, it was found that TC in PM_{2.5} aerosols derived from peatland fires was primarily composed of OC₁ and OC₂.

Fig. 5 illustrates the average percentages of eight fractions in PM_{2.5} for the peatland fire samples and those from other sources. The vegetation burning profiles used as reference were obtained by ground-based source-dominated sampling of the plumes of small controlled burns of wood debris at the Big Bend National Park in the US (see Chow *et al.*, 2004). The carbon fraction abundances differ by emission source: OC₁ is enriched ($23.9 \pm 12.4\%$) in the vegetative burning profile; OC₂ is enriched ($34.7 \pm 2.16\%$) in the peatland fire profile; OC₃ is enriched ($37.6 \pm 15.9\%$) in the cooking profile; EC₂ is enriched ($23.3 \pm 10.4\%$) in the motor vehicle profile. In terms of the coefficient of variance, the carbon fraction obtained from peatland fire is more consistent than any source presented in Fig. 5. Thus, the peatland profile will be useful for source discrimination.

3.3. Key biomarkers

220 The organic compounds produced by peatland fires were analyzed by GC/MS, and an example
221 of typical GC/MS total ion current (TIC) tracers for the total extract of peatland fire samples is
222 presented in Fig. 6. Levoglucosan clearly exhibits the highest TIC peak in peatland fire samples,
223 although mannosan and palmitic acid were also detected. Palmitic acid is one of the most basic
224 units of plant fats, oils, and phospholipids (Simoneit, 2002). The major tracers in smoke emitted
225 from biomass burning are the thermal degradation (pyrolysis) products of the biopolymers of
226 cellulose and lignin in woody tissue. The thermal degradation of cellulose (also hemicelluloses)
227 yields dehydromonosaccharide derivatives, which are predominant compounds in the smoke
228 (Simoneit *et al.*, 1999). The major compound is levoglucosan, with minor and variable amounts
229 of galactosan and mannosan. These compounds cannot be formed by hydrolysis or by the
230 microbial alteration of carbohydrates; thus, they are specific to burning (Simoneit *et al.*, 1999). In
231 addition, levoglucosan is considerably resistant to atmospheric degradation (Fraser and
232 Lakshmanan, 2000). Therefore, levoglucosan and mannosan are regarded as key biomarkers in
233 peatland fire aerosols like other biomass burnings.

234 The average levoglucosan concentrations from peatland fire and background samples were 464
235 $\pm 183 \mu\text{g}/\text{m}^3$ and $0.278 \pm 0.155 \mu\text{g}/\text{m}^3$, respectively and the average mannosan concentration
236 were $47.5 \pm 25.6 \mu\text{g}/\text{m}^3$ and $0.0190 \pm 0.0108 \mu\text{g}/\text{m}^3$, respectively. Levoglucosan and mannosan in
237 background samples may have originated from wood burning for cooking, agricultural use,

among others. However, concentrations of levoglucosan and mannosan emitted in association with peatland fire events were much higher than their corresponding background concentrations.

OC/Levoglucosan ratios vary to some extent for different burning conditions and wood types (Pio, *et al.*, 2008). In this study, the average OC/Levoglucosan ratio in peatland fire was $10.6 \pm 1.96 \mu\text{gC}/\mu\text{g}$. For comparison of our peatland fire samples with biomass burning samples, the average OC/Levoglucosan ratios for various source samples are presented in Fig. 7, which shows significant differences in OC/Levoglucosan ratios between the peatland fire and other emission sources. Thus, this ratio may be useful for source discrimination for peatland fire. However, because levoglucosan is emitted from other biomass burning sources and there are other types of biomass burning in Riau, Sumatra, Indonesia, selection of only OC/Levoglucosan ratio for source apportionment of peatland fire can lead to overestimation of the contribution of peatland fire. Hence, other indicators for peatland fire are needed.

The relative amounts of the individual anhydrosaccharides in biomass smoke aerosols can be used for further source assignment of specific biofuels (Fabbri *et al.*, 2009; Alves *et al.*, 2010). Here, we investigated the Levoglucosan/Mannosan ratios and compared those ratios from peatland fire and other biomass burning sources. In this study, the average Levoglucosan/Mannosan ratios in peatland fire and background were 10.6 ± 2.03 and 14.5 ± 3.39 , respectively. For comparison of our peatland fire samples with other biomass burning samples,

the average Levoglucosan/Mannosan ratios for various source samples are presented in Table 1, which shows the significant differences in Levoglucosan/Mannosan ratios between the peatland fire and other emission sources. Thus, this ratio should be more useful for source discrimination for peatland fire than OC/Levoglucosan ratio.

4. Conclusions

PM_{2.5} carbonaceous aerosols were collected at a peatland fire hotspots and a background site on 7 and 4 occasions, respectively. PM_{2.5} aerosols emitted from peatland fire were observed in very high concentrations ($7120 \pm 3620 \mu\text{g}/\text{m}^3$) and were primarily composed of OC ($71.0 \pm 5.11\%$ of PM_{2.5} mass).

The OC/EC ratios (36.4 ± 9.08 for peatland fire), abundances of eight thermally-derived carbon fractions, OC/Levoglucosan ratios (10.6 ± 1.96 for peatland fire), and Levoglucosan/Mannosan ratios (10.6 ± 2.03 for peatland fire) observed here represent a signature profile that seems to be inherent in peatland fire emissions. Therefore, it is suggested that such profiles are useful for peatland fire source discrimination.

In future studies, additional chemical speciation for fresh and aged smoke from peatland fires will be required to estimate atmospheric environmental impacts and/or adverse health effects (e.g., humic-like substances, polycyclic aromatic hydrocarbons).

276

277

278 **Acknowledgement**

279

280 This study was partially supported by funds from the Grants-in Aid for Innovative Areas under
281 Grant No. 2012005 from the Japanese Ministry of Education, Culture, Sports, Science and
282 Technology (MEXT).

283

284

285 **References**

286

287 Alves, C.A., Gonçalves, C., Evtyugina, M., Pio, C.A., Mirante, F., Puxbaum, H., 2010.

288 Particulate Organic Compounds Emitted from Experimental Wildland Fires in a Mediterranean
289 Ecosystem. *Atmos. Environ.* 44: 2750–2759.

290 Cao, J.J., Wu, F., Chow, J.C., Lee, S.C., Li, Y., Chen, S.W., An, Z.S., Fung, K.K., Watson, J.G.,

291 Zhu, C.S. and Liu, S.X., 2005. Characterization and Source Apportionment of Atmospheric

292 Organic and Elemental Carbon during Fall and Winter of 2003 in Xi'an, China. *Atmos. Chem.*

293 *Phys.* 5: 3127–3137.

294 Chow, J.C., Watson, J.G., Lu, Z., Lowenthal, D.H., Frazier, C.A., Solomon, P.A., Thuillier, R.H.

295 and Magliano, K.L., 1996. Descriptive Analysis of PM_{2.5} and PM₁₀ at Regionally

296 Representative Locations during SJVAQS/AUSPEX. *Atmos. Environ.* 30: 2079–2112.

- 297 Chow, J.C., Watson, J.G., Chen, L.-W.A., Chang, M.C.O., Robinson, N.F., Trimble, D. and Kohl,
298 S., 2007. The IMPROVE_A Temperature Protocol for Thermal/Optical Carbon Analysis:
299 Maintaining Consistency with a Long-Term Database. *J. Air Waste Manage. Assoc.* 57: 1014–
300 10023.
- 301 Chow, J.C., Watson, J.G., Kuhns, H.D., Etyemezian, V., Lowenthal, D.H., Crow, D.J., Kohl, S.D.,
302 Engelbrecht, J.P. and Green, M.C., 2004. Source Profiles for Industrial, Mobile, and Area
303 Sources in Big Bend Regional Aerosol Visibility and Observational (BRAVO) Study.
304 *Chemosphere* 54: 185–208.
- 305 Fabbri, D., Torri, C., Simoneit, B.R.T., Marynowski, L., Rushdi, A.I., Fabiańska, M.J., 2009.
306 Levoglucosan and Other Cellulose and Lignin Markers in Emissions from Burning of Miocene
307 Lignites. *Atmos. Environ.* 43: 2286–2295.
- 308 Federal Register, 2006. National Ambient Air Quality Standards for Particulate Matter: Final
309 Rule. 40 CFR Parts 50, 53, and 58, volume 62 (138), October 17, 2006, U.S. EPA, Office of
310 Air and Radiation, Office of Air Quality Planning and Standards, Research Triangle Park, NC.
- 311 Fraser, M.P. and Lakshmanan, K., 2000. Using Levoglucosan as a Molecular Marker for the
312 Long-range Transport of Biomass Combustion Aerosols. *Environ. Sci. Technol.* 34, 4560–4564.
- 313 Gelencsér, A., 2004. Carbonaceous Aerosol, Springer, p.32.
- 314 GFED. [online] [Accessed 17 July 2013]. Available: <http://www.globalfiredata.org/>.

- 315 Gray, H.A., Cass, G.R., Huntzicker, J.J., Heyerdahl, E.K. and Rau, J.A., 1986. Characteristics of
316 Atmospheric Organic and Elemental Carbon Particle Concentrations in Los Angeles. *Environ.*
317 *Sci. Technol.* 20: 580–589.
- 318 Harahap, R., 2012. As dry season returns to Riau, so do fires and choking haze. The Jakarta Post,
319 June 13. [online] [Accessed 17 July 2013]. Available:
320 [http://www.thejakartapost.com/news/2012/06/13/as-dry-season-returns-riau-so-do-fires-and-ch](http://www.thejakartapost.com/news/2012/06/13/as-dry-season-returns-riau-so-do-fires-and-choking-haze.html)
321 [oking-haze.html](http://www.thejakartapost.com/news/2012/06/13/as-dry-season-returns-riau-so-do-fires-and-choking-haze.html).
- 322 Heil, A., Stolle, F., Mahmud, M. and Effendi, E., 1998. Air Pollution from Large Scale Forest and
323 Land Fires in Indonesia 1997–1998: Development and Impacts. The First Synthesis Working
324 Group Meeting on Land Use Planning and Management to Reduce Transboundary Pollution
325 from Fires in Southeast Asia. Bongor, Indonesia, 18–19 August 1998.
- 326 Hong C., 2012. ‘Indonesian’-made haze covers parts of KL, surrounding areas. The Straits Times,
327 June 16. [online] [Accessed 17 July 2013]. Available:
328 [http://www.thejakartapost.com/news/2012/06/16/indonesian-made-haze-covers-parts-kl-surrou](http://www.thejakartapost.com/news/2012/06/16/indonesian-made-haze-covers-parts-kl-surrounding-areas.html)
329 [nding-areas.html](http://www.thejakartapost.com/news/2012/06/16/indonesian-made-haze-covers-parts-kl-surrounding-areas.html).
- 330 Indofire. [online] [Accessed 17 July 2013]. Available: <http://www.indofire.org/indofire/hotspot>.
- 331 Lee, S., Baumann, K., Schauer, J.J., Sheesley, R.J., Naeher, L.P., Simone, M., Blake, D.R.,
332 Edgerton, E.S., Russell, A.G. and Clements, M., 2005. Gaseous and Particulate Emissions from

- 333 Prescribed Burning in Georgia. *Environ. Sci. Technol.* 39: 9049-9056.
- 334 Lippmann M., 1998. In The 1997 US EPA Standards for Particulate Matter and Ozone. *Issues*
- 335 *Environ. Sci. Tech.* 10, Hester, R.E. and Harrison, R.M. (Eds.), Royal Society of Chemistry:
- 336 75–99.
- 337 Lubis, A.M. and Harahap, R., 2013. Only rain can help S'pore now. The Jakarta Post, June 21.
- 338 [online] [Accessed 17 July 2013]. Available:
- 339 <http://m.thejakartapost.com/news/2013/06/21/only-rain-can-help-s-pore-now.html>.
- 340 Othman, M. and Latif, M.T., 2013. Dust and Gas Emissions from Small-Scale Peat Combustion.
- 341 *Aerosol Air Qual. Res.* 13: 1045–1059.
- 342 Phuleria, H.C., Fine, P.M., Zhu, Y., and Sioutas, C., 2005. Air Quality Impacts of the October
- 343 2003 Southern California Wildfires. *J. Geophys. Res.* 110, D07S20, doi:
- 344 10.1029/2004JD004626.
- 345 Pio, C.A., Legrand, M., Alves, C.A., Oliveria, T., Afonso, J., Caseiro, A., Puxbaum, H.,
- 346 Sanchez-Ochoa, A. and Gelensér, A., 2008. Chemical Composition of Atmospheric Aerosols
- 347 during the 2003 Summer Intense Forest Fire Period. *Atmos. Environ.* 42: 7530–7543.
- 348 Reid, J.S., Koppmann, R., Eck, T.F. and Eleuterio, D.P., 2005. A Review of Biomass Burning
- 349 Emissions Part II: Intensive Physical Properties of Biomass Burning Particles. *Atmos. Chem.*
- 350 *Phys.* 5: 799–825.

- 351 See, S.W., Balasubramanian, R., Rianawati, E., Karthikeyan, S. and Streets, D.G., 2007.
- 352 Characterization and Source Apportionment of Particulate Matter $\leq 2.5 \mu\text{m}$ in Sumatra,
- 353 Indonesia, during a Recent Peat Fire Episode. *Environ. Sci. Technol.* 41: 3488–3494.
- 354 Sheesley R.J. and Schauer, J.J., 2003. Characterization of Organic Aerosols Emitted from the
- 355 Combustion of Biomass Indigenous to South Asia. *J. Geophys. Res.* 108, D9, 4285, doi:
- 356 10.1029/2002JD002981.
- 357 Simoneit, B.R.T., 2002. Biomass Burning – A Review of Organic Tracers for Smoke from
- 358 Incomplete Combustion. *Appl. Geochem* 17: 129–162.
- 359 Simoneit, B.R.T., Schauer, J.J., Nolte, C.J., Oros, D.R., Elias, V.O., Fraser, M.P., Rogge, W.F. and
- 360 Cass, G.R., 1999. Levoglucosan, A Tracer for Cellulose in Biomass Burning and Atmospheric
- 361 Particles. *Atmos. Environ.* 33: 173–182.
- 362 Stolle, F. and Tomich, T.P., 1999. The 1997-1998 Fire Event in Indonesia. *Nat. Resour.* 35: 22–30.
- 363 Tacconi, L., 2003. Fires in Indonesia: Causes, Costs and Policy Implications. CIFOR Occasional
- 364 Paper 38.
- 365 Turpin, B.J. and Huntzicker, J.J., 1991. Secondary Formation of Organic Aerosol in the Los
- 366 Angeles Basin. A Descriptive Analysis of Organic and Elemental Carbon Concentrations.
- 367 *Atmos. Environ.* 25A: 207–215.
- 368 Watson, J.G., Chow, J.C., Lowenthal, D.H., Pritchett, L.C., Frazier, C.A., Neuroth, G.R. and

369 Robbins, R., 1994. Differences in the Carbon Composition of Source Profiles for Diesel- and
370 Gasoline-powered Vehicles. *Atmos. Environ.* 28: 2493–2505.

371 Wulandari, F., 2002. Reckless Farming in Peatland Forests Major Cause of Fires. The Jakarta
372 Post, August 25. [online] [Accessed 17 July 2013]. Available:
373 <http://www.thejakartapost.com/news/2002/08/25/reckless-farming-peatland-forests-major-caus>
374 [e-fires.html](http://www.thejakartapost.com/news/2002/08/25/reckless-farming-peatland-forests-major-caus).
375

376

Table Captions

377 Table 1. Range of Levoglucosan/Mannosan ratios in PM_{2.5} emitted from several sources and
378 background.

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

	Levoglucosan/Mannosan	Reference
Peatland fire	7.09–14.0	this study
Background	10.7–18.9	this study
Hardwoods burning	13.8–52.3	Engling <i>et al.</i> , 2006
Softwoods burning	2.6–5.0	Engling <i>et al.</i> , 2006
Grasses burning	108–203	Engling <i>et al.</i> , 2006

Table 1.

Figure Captions

399

400 Fig. 1. Map of Bengkalis showing the sampling sites.

401 Fig. 2. Monthly hotspot counts in Riau in 2011 and 2012.

402 Fig. 3. Comparison of OC/EC ratios in PM_{2.5} emitted from peatland fire and other burning

403 sources. Error bar in this study indicates standard deviation.

404 Fig. 4. Abundances (as mass percentage of total carbon) of eight thermally-derived carbon

405 fractions of aerosols sampled at the peatland fire and background sites. Error bars indicate

406 standard deviations.

407 Fig. 5. Carbon fraction composition of PM_{2.5} combustion source profiles. Error bars indicate

408 standard deviations.

409 Fig. 6. GC/MS total ion current (TIC) chromatograph of total extract showing major compounds

410 of peatland fire aerosols.

411 Fig. 7. Comparison of OC/Levoglucosan ratios of PM_{2.5} obtained from peatland fire and other

412 burning sources. Error bar in this study indicates standard deviation.

413

414

415

416

Fig. 1.

Fig. 2.

^a data from Sheesley and Schauer, 2003, ^b data from Lee *et al.*, 2005.

Fig. 3.

Fig. 4.

^a data from Chow *et al.*, 2004.

Fig. 5.

Fig. 6.

^a data from Sheesley and Schauer, 2003, ^b data from Lee *et al.*, 2005.

Fig. 7.