TITLE:

Atmospheric Reactions of PAH Derivatives: Formation and Degradation

AUTHOR(S):

Kameda, Takayuki

CITATION:

Kameda, Takayuki. Atmospheric Reactions of PAH Derivatives: Formation and Degradation. Polycyclic Aromatic Hydrocarbons 2018: 75-91

ISSUE DATE:

2018

URL:

<http://hdl.handle.net/2433/237664>

RIGHT:

This is an author's accepted manuscript (AAM) of a chapter published in 'Polycyclic Aromatic Hydrocarbons'. The final authenticated version is available online at: https://doi.org/10.1007/978-981-10-6775-4_7; The full-text file will be made open to the public on 06 February 2019 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。; This is not the published version. Please cite only the published version.

Polycyclic Aromatic Hydrocarbons: Environmental Behavior and Toxicity in East Asia

Chapter 3.3

Atmospheric Reactions of PAH derivatives: Formation and Degradation

Takayuki Kameda

Graduate School of Energy Science, Kyoto University, Yoshida-Honmachi, Sakyo-ku, Kyoto 606-8501, Japan

Abstract

Polycyclic aromatic compounds are ubiquitous atmospheric pollutants with toxic, mutagenic, and carcinogenic properties. They are produced from chemical reactions of their parent or related compounds in the atmosphere as well as from a wide variety of anthropogenic sources such as fuel combustions. In this chapter, chemical reaction pathways for the atmospheric secondary formation of several polycyclic aromatic hydrocarbon (PAH) derivatives, i.e., gas-phase formation of mutagenic 1- and 2-nitrotriphenylene via OH or NO₃ radical-initiated reactions of the parent triphenylene, formation of carcinogenic 1-nitropyrene from heterogeneous nitration of pyrene on mineral dust aerosols, atmospheric formation of hydroxynitropyrenes from a photochemical reaction of 1-nitropyrene, and photochemical degradation of selected nitrated and oxygenated PAHs on airborne particles under simulated solar UV-irradiation, are addressed.

Keywords: polycyclic aromatic compounds, secondary formation, photoreaction, heterogeneous reaction, gas-phase reaction

3.3.1 Introduction

Polycyclic aromatic hydrocarbons (PAHs) and their derivatives (e.g., nitrated, oxygenated, and hydroxylated derivatives: NPAHs, OPAHs, and OHPAHs) are ubiquitous atmospheric pollutants with toxic, mutagenic, and carcinogenic properties (Schuetzle, 1983; Salmeen et al., 1984). These compounds are produced from a wide variety of anthropogenic sources such as the incomplete combustion of fossil fuels used in industrial plants, through heating, and in diesel powered vehicles (Schuetzle et al., 1982; Bamford et al., 2003; Bezabeh et al., 2003). NPAHs, which generally exhibit higher mutagenicity and carcinogenicity than their parent PAHs, are also generated from atmospheric reactions of PAHs released in the gas-phase with radical species such as OH and NO₃ radicals and nitrogen oxides (Zielinska et al., 1989; Ciccioli et al., 1996; Reisen and Arey, 2005; Atkinson and Arey, 1994; Sasaki et al., 1997). OPAHs and OHPAHs, which may show biological effects beyond mutagenicity and carcinogenicity, are also known to be formed secondarily via oxidation of PAHs and NPAHs in the atmosphere (Cvrčková, et al., 2006; Warner et al., 2004). These PAH derivatives generally have lower volatility than the parent PAHs. As a result, they are more likely to be deposited on airborne particles, especially fine particles which are easily suspended into the atmosphere and can easily be breathed into the human body. Thus, it is necessary to understand the secondary formation processes of PAH derivatives in order to fully understand their effects on humans. In this chapter, chemical reaction pathways for the secondary formation of PAH derivatives are addressed.

3.3.2 Formation of nitrotriphenylenes by radical-initiated reactions

As described above, several types of NPAH are formed via gas-phase reactions of semi-volatile PAHs. For example, 2-nitropyrene (2-NP) is formed by the gas-phase reaction of pyrene with OH radicals in the presence of NO₂, and 2-nitrofluoranthene (2-NFR) is formed by OH or NO₃ radical-initiated reactions of fluoranthene in gas phase (Atkinson and Arey, 1994). Benzanthrone, which has very low vapor pressure, was also found to react with radical species and NO₂ to yield a nitrated compound in the gas phase

(Phousongphouang and Arey, 2003). Nitrotriphenylenes (NTPs), which include the strongly mutagenic isomer 2-nitrotriphenylene (2-NTP), have been found in airborne particles as well (Ishii, et al., 2000; 2001; Kameda et al., 2004; Kawanaka et al., 2005). The possibility of the atmospheric formation of NTPs via reactions of triphenylene was pointed out based on seasonal and diurnal changes in the concentration of ambient NTPs (Ishii, et al., 2001; Kameda et al., 2004). Because the strongly mutagenic 2-NTP has been found in air at concentrations comparable to those of 1-nitropyrene (1-NP) and 2-NFR, which are the most abundant airborne NPAHs (Ishii, et al., 2000; 2001; Kameda et al., 2004; Kawanaka et al., 2005), the contribution of 2-NTP to the mutagenicity of airborne particles may be significant. From the point of view of public hygiene, it is important to acquire more detailed data about the environmental occurrence of NTPs. Thus, the formation of 1-nitrotriphenylene (1-NTP) and 2-NTP by gas-phase OH or NO₃ radical-initiated reaction of triphenylene, was demonstrated using a flow reaction system (Kameda et al., 2006). Nitration of triphenylene with N₂O₅ in CCl₄ was also examined in order to determine the isomer distributions of NTPs formed via NO₃ radical-initiated nitration of triphenylene, and in order to predict the rate constants of the gas-phase OH or NO₃ radical-initiated reactions of triphenylene.

The formation of 1- and 2-NTP was clearly shown by HPLC analysis of the products of the gas-phase reaction of triphenylene initiated by the OH radicals. A 2-NTP/1-NTP ratio of 1.22 was obtained for OH radical-initiated nitration. In contrast, the NO₃ radical-initiated reaction predominantly gave 2-NTP, with traces of 1-NTP. Because the amount of 1-NTP formed in the NO₃ radical-initiated reaction was too small to determine, a precise 2-NTP/1-NTP value could not be calculated and was thus assumed to be a value greater than 1.5. Preferential production of 2-NTP was also observed in the nitration of triphenylene with N₂O₅ in CCl₄; for which the yields of 1- and 2-NTP were 6 (±2) % and 35 (±2) %, respectively. It has been reported that the reactions of several kinds of PAH with N₂O₅ in CCl₄ are similar in terms of nitro-isomer distribution to gas-phase NO₃ radical-initiated nitration (Phousongphouang and Arey, 2003; Zielinska et al., 1986). Thus, the analogous isomer distribution of NTP (the larger yield of 2-NTP over 1-NTP) in the gas-phase reaction, to that in CCl₄ liquid-phase nitration does not contradict previous

findings regarding the nitration of PAHs. The mean 2-NTP/1-NTP ratio in samples of airborne particles was > 1.55 . This value was similar to the ratios from the radical-initiated reactions, and was much higher than that of the diesel exhaust particulate (DEP) samples (2-NTP/1-NTP = 0.37). This finding indicates that the atmospheric radical-initiated reactions significantly contribute to the formation of airborne NTPs, especially 2-NTP. The gas-phase formation of NPAHs via OH or NO₃ radical-initiated reaction involves the addition of an OH or NO₃ radical to the PAH at the carbon atom with the highest electron density, followed by ortho-addition of NO₂. This is followed by a loss of water or nitric acid. For triphenylene, the carbon at the 1-position is the most electron-rich (Barker et al., 1955; Radner, 1983); therefore, preferential formation of 2-NTP over 1-NTP is expected in the gas-phase radical-initiated reaction (Fig. 1).

Figure 1

The rate constants of gas-phase reactions of triphenylene with OH and NO₃ radicals at 298 K were predicted to be $(8.6 \pm 1.2) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ and $(6.6 \pm 1.5) \times 10^{-29} [\text{NO}_2] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, respectively using a relative-rate method in a CCl₄ liquid phase-system (Kameda et al., 2013). Based on the ambient concentrations of 2-NTP and the rate constant obtained for the reaction of triphenylene with the radicals, the atmospheric loss rate of 2-NTP relative to 2-NFR (which is the most abundant NPAH and is also produced from radical reactions) was successfully estimated. This is to say, 2-NTP is less susceptible to decomposition than 2-NFR is under ambient conditions.

3.3.3 Secondary formation of 1-nitropyrene promoted on mineral dust aerosols

One of the most abundant NPAHs is 1-NP, which is formed through the combustion of fossil fuels such as coal and diesel fuel (Schuetzle, 1983; Yang et al., 2010). 1-NP is likely a carcinogen (IARC, 2013) and can also be formed from gas-particle phase heterogeneous reactions (Esteve et al., 2004; Finlayson-Pitts and Pitts, 2000; Inazu et al., 2000; Miet et al., 2009; Nguyen et al., 2009; Ramdahl et al., 1984; Wang et al., 2000). It is

formed by the reaction of pyrene (Py) with gaseous NO₂ on various substrates such as graphite, as a model for soot (Esteve et al., 2004), and a variety of metal oxides, as models for mineral aerosols (Inazu et al., 2000; Miet et al., 2009; Ramdahl et al., 1984; Wang et al., 2000). However, the heterogeneous formation of atmospheric 1-NP has been previously thought to be negligible because the reaction rate and the yield of 1-NP through this process are not sufficient to account for ambient 1-NP concentration (Finlayson-Pitts and Pitts, 2000; Nguyen et al., 2009; Ramdahl et al., 1984; Shiraiwa et al., 2009). Previous studies of heterogeneous NPAH formation used simple inorganic oxides such as SiO₂, Al₂O₃, and TiO₂ as models of mineral dust aerosols (Inazu et al., 2000; Ma et al., 2011; Wang et al., 2000), but these substances lack the complexity of real mineral dust aerosols and thus may not be good models for investigating heterogeneous NPAH formation. Mineral dust is a major component of airborne particulates on a global scale (Cwierny et al., 2008). It is transported by winds from deserts or semiarid regions (Tanaka and Chiba, 2006), which account for 40% of the total world land area (Fernández, 2002). Organic compounds adsorbed on the surface of mineral dust can have important health implications (Falkovich et al., 2004). Thus, the formation of 1-NP from Py and NO₂ on authentic mineral dust was examined (Kameda et al., 2016).

In the NO₂ exposure experiments of particle-bound Py, degradation of Py was measured under 3 ppmv NO₂-air in the dark. On quartz (SiO₂) particles, Py was slowly converted to 1-NP, reaching a yield of ~40% in 12 h (Fig. 2a). On Chinese desert dust (CDD) particles, more than 90% of the initial amount of Py was degraded and the maximum yield of 1-NP was attained after a reaction time of 1 h (Fig. 2b). 1-NP was then gradually converted to dinitropyrenes (DNPs) (Fig. 2b). Other mononitropyrene isomers were not detected. Desert dust is generally composed of various minerals including quartz, corundum (α -Al₂O₃), clay minerals, carbonates, feldspars, and hematite (Fe₂O₃) (Usher et al., 2003). To determine which components contribute to rapid nitration, the percentage of degraded Py (*D*_{Py}) and the yield of 1-NP (*Y*_{1-NP}) were compared during a reaction time of 2 h on various substrates that generally constitute desert dust. The most active components were natural montmorillonites, kaolin, and saponite, as well as Arizona Test Dust (ATD; standard test dust made from Arizona desert sand) and CDD (Table 1). In most of these

cases, the conversion of Py to 1-NP was completed within 2 h (Table 1). DNP formation was observed except on saponite. Kaolin, montmorillonites A and B, and saponite are types of clay minerals. For the other mineral substrates, such as quartz, carbonates (limestone and dolomite), and feldspars; D_{Py} and $Y_{1\text{-NP}}$ were less than 20%, and no DNP was formed during the NO_2 exposure (Table 1). To quantify the rate of degradation of Py on each substrate, the kinetics of the heterogeneous reaction between NO_2 and Py adsorbed on the substrates tested in this study were determined by following the consumption of Py as a function of NO_2 exposure time. The apparent rate constants of the pseudo-first-order reaction, k_{obs} , were (2.9×10^{-4} to 2.5×10^{-3}) s^{-1} on CDD, ATD, and clay minerals and (2.5×10^{-6} to 9.0×10^{-5}) s^{-1} on the other substrates when the concentration of NO_2 was 3 ppmv.

Figure 2

Table 1

The nitration of PAHs is catalysed by acids (Shiri et al., 2010). Thus, the surface acid property of mineral dust may play a role in the heterogeneous nitration of Py. The surface acid properties of solid catalysts, including clay minerals, can be examined using Fourier transform infrared spectroscopy (FT-IR) with pyridine as a probe (Parry, 1963). When pyridine binds to Brønsted acid sites, pyridinium ions are produced, which have an absorption band around 1545 cm^{-1} . In contrast, pyridine molecules coordinated to Lewis acid sites have an absorption band around 1445 cm^{-1} . The band at 1490 cm^{-1} is attributed to both molecules. The spectra of pyridine adsorbed onto some substrates (CDD, ATD, montmorillonites, kaolin, and saponite) have absorption bands at (1445 and 1490) cm^{-1} , while no absorption band is observed around 1545 cm^{-1} , except in the cases of kaolin and montmorillonite K10 (Fig. 3). This suggests that CDD and ATD, as well as clay minerals, have abundant acid sites, particularly Lewis acid sites. On the contrary, the spectra of the other substrates displayed no clear peaks, indicating that they have little to no acid sites on their surfaces. The largest k_{obs} value was obtained for the reaction on montmorillonite K10, an acid-activated clay. These results strongly suggest that substrates showing acidic

surface properties have an accelerating effect on the rate of heterogeneous nitration of PAHs by NO₂.

Figure 3

Lewis acid sites on aluminosilicates are proposed to function as electron acceptors, leading to the formation of aromatic radical cations *via* electron transfer (Laszlo, 1987; Soma and Soma, 1989). The radical cations of several kinds of PAHs, such as Py, perylene, anthracene, and benzo[*a*]pyrene (which form on the surface of aluminosilicates) have been identified by spectroscopic methods, such as electron spin resonance (ESR) (Garcia and Roth, 2002). These cations would couple with surface NO₂ to yield NPAHs (Laszlo, 1987), similar to that with the nitrous acid-catalysed (NAC) nitration mechanism (Ridd, 1991). That is, the rate-determining step would be the subsequent addition of NO₂ to the aromatic radical cation yielding a σ -complex (Wheland intermediate), and the deprotonation of this complex would constitute the final fast step that produces the nitrocompound (Fig. 4). Thus, the finding that the Lewis acid property of the substrates probably plays a role in nitration, suggests that the rapid formation of 1-NP on mineral dust is the result of NO₂ reacting with the radical cations of Py, which form at the surface Lewis acid sites (Fig. 4).

Figure 4

3.3.4 Atmospheric formation of hydroxynitropyrenes from a photochemical reaction of particle-associated 1-nitropyrene

The 1-NP taken up by humans and animals is transformed into various metabolites such as hydroxynitropyrenes (OHNPs), in the presence of cytochrome P450 enzymes (Rosser et al., 1996). Several isomers of OHNP (Fig. 5), such as 1-hydroxy-3-nitropyrene (1-OH-3-NP), 1-hydroxy-6-nitropyrene (1-OH-6-NP), and 1-hydroxy-8-nitropyrene (1-OH-8-NP) have also been observed on airborne particles (Gibson et al., 1986; Kameda et al., 2010) and on diesel exhaust particles (DEP)

(Schuetzle, 1983; Schuetzle et al., 1985; Manabe et al., 1985). Several studies have found that most OHNP isomers have lower mutagenic activity than the parent 1-NP (Rosser et al., 1996; Manabe et al., 1985; Ball et al., 1985). Recently, however, OHNPs such as 1-OH-3-NP, 1-OH-6-NP, and 1-OH-8-NP have been found to act as endocrine disruptors: they act as estrogenic, anti-estrogenic, and anti-androgenic compounds (Kameda et al., 2008; Kameda et al., 2011a) that may cause dysfunction of human and wildlife endocrine systems, abnormal development of reproductive systems, and immunodeficiencies. In view of the influence of OHNPs on human health, we need to learn more about their environmental concentration levels, sources, and behaviors. Therefore, the formation of OHNPs including 1-OH-3-NP, 1-OH-6-NP, and 1-OH-8-NP from photochemical reactions of 1-NP was examined in laboratory experiments to clarify the occurrence of atmospheric OHNPs (Kameda et al., 2011b).

Figure 5

Figure 6 shows a profile of an HPLC analysis with chemiluminescence detection (HPLC/CLD) for the products from photoreactions of 1-NP in methanol. Five chromatographic peaks were observed in the chromatogram (symbolized as A, B, C, D, and E). The retention times of peaks B, C, D, and E were the same as those of authentic 1-OH-6-NP, 1-OH-8-NP, 1-OH-3-NP, and 1-OH-2-NP, respectively. When analyzed by LC/MS/MS, a fraction containing the photo-reaction products also yielded five peaks. By comparing the retention times and the MS/MS spectra of these peaks with those of the authentic standards, four known OHNPs (1-OH-2-NP, 1-OH-3-NP, 1-OH-6-NP, and 1-OH-8-NP) were identified. For these compounds, the molecule-related ion m/z 262 ($[M-H]^-$) together with the characteristic fragment ions m/z 232 ($[M-H-NO]^-$) and 216 ($[M-H-NO_2]^-$) were detected in a full scan analysis. In the study, and for the first time, 1-OH-3-NP, 1-OH-6-NP, and 1-OH-8-NP were found in 1-NP photoreaction products. The unknown compound that was observed in the HPLC/CLD chromatogram was also observed in the LC/MS/MS analysis. This compound also gave a characteristic MS/MS spectrum with a molecule-related ion m/z 262 and fragment ions m/z 232 and m/z 216. The

similarity between the fragmentation patterns of the unknown compound and known OHNPs, indicates that the unknown compound is an isomer of OHNP. The structure of the unknown compound obtained by the preparative scale photoreaction was then determined by analysis of its ^1H -NMR spectrum. On the basis of chemical shifts and coupling patterns, the unknown compound contained in the photo-reaction products was identified as 1-hydroxy-5-nitropyrene (1-OH-5-NP).

Figure 6

All the OHNP isomers, which were found in the 1-NP photo-reaction products, were also identified in ambient airborne particles collected at a typical residential area in Osaka, Japan. In contrast, 1-OH-2-NP and 1-OH-5-NP were not found in Standard Reference Materials (SRM) 1650b and SRM 1975, which are typical DEP samples. The concentrations of the other OHNP isomers in the DEP samples were much lower than the concentration of 1-NP. On the other hand, significantly higher concentration ratios of ΣOHNP ($= 1\text{-OH-3-NP} + 1\text{-OH-6-NP} + 1\text{-OH-8-NP}$) to 1-NP were observed in ambient airborne particles than in the DEP samples. In ambient airborne particles, the mean $\Sigma\text{OHNP}/1\text{-NP}$ concentration ratio of 1.4 was 35 times higher than that in SRM 1650b, and 470 times higher than that in SRM 1975. The diurnal concentration of 1-NP observed at the site in Osaka, increased early in the morning and late in the evening, suggesting that automotive emissions contributed to the occurrence of 1-NP. The OHNP concentrations also rose in the morning and variations of OHNP concentrations similar to those of 1-NP were observed during the daytime. However, the concentrations of OHNPs did not increase in the evening rush hour, and were low at night (i.e., in the absence of sunlight). These results support the idea that atmospheric OHNPs are predominantly formed *via* secondary formation processes; photochemical reactions of 1-NP are expected to have a significant effect on the occurrence of OHNPs in the atmosphere.

3.3.5 Photochemical decomposition of selected nitro- and oxy-polycyclic aromatic hydrocarbons on airborne particles under simulated solar UV-irradiation

As in the case of 1-NP, photo-induced decomposition is a dominant pathway for the degradation of particle-associated PAH derivatives (Finlayson-Pitts and Pitts, 2000; Kamens et al., 1988). Solar radiation in the UV spectral region can modify PAH derivatives to form new compounds that may exhibit different types of biological effects. These might include the disruption of endocrine systems and the production of reactive oxygen species (ROS) in the human body (Chung et al., 2007). Thus, it is necessary to understand the photodecomposition of PAH derivatives in order to understand their effects on humans. However, the rate constants and the quantum yields related to the photolysis of the derivatized PAHs, which are the most significant factors for photodecomposition, have not been studied well. Thus, photodecomposition experiments for selected OPAHs and NPAHs, including 3-nitrobenzanthrone (3-NBA) a nitrated aromatic ketone that is strongly mutagenic (Enya et al., 1997), were conducted on a glass surface. With this as a simple model of airborne particles, the photolysis rate constants and the quantum yields for the PAH derivatives in the system were determined (Kameda et al., 2009). Furthermore, the atmospheric lifetime of the compounds due to photodecomposition was estimated using the actinic flux on the Earth's surface, photolysis rate constants, and quantum yields obtained in the study.

The highest photolysis rate constant was observed for 9-nitroanthracene (9-NA), while 4-nitropyrene (4-NP) and 3-NBA were found to be the most stable of the nitrated compounds under UV-irradiation (Table 2). It is hypothesized that photoreactivity of NPAHs is governed by the orientation of the nitro group; i.e. NPAHs having nitro groups perpendicular to the aromatic ring are more easily photodecomposed than those having parallel ones (Yang et al., 1994; Warner et al., 2004). The fast photodegradations observed for 9-NA, 6-nitrobenzo[*a*]pyrene (6-NBaP), and 7-nitrobenz[*a*]anthracene (7-NBaA), which all have a perpendicular nitro group, were consistent with this hypothesis (Fig. 7). The photoreaction products of NPAHs were reported to include quinoid PAHs (Warner et al., 2004), as well as photoreaction products of PAHs. Benzo[*c*]phenanthrene-5,6-quinone

(BcP-5,6-Q), which has a similar photolysis rate constant to 9-nitrophenanthrene (9-NPh), degraded the fastest of the seven OPAHs tested. Of all the substituted PAHs examined in the study, 1,2-benzanthraquinone (1,2-BAQ) was most resistant to photodecomposition. Although previous studies on the photostability of OPAHs were quite limited, Cvrčková and Ciganek (2005) reported that 9,10-phenanthraquinone (9,10-PQ) was less stable than anthraquinone (AQ) under UV irradiation. This is consistent with the results.

Table 2

Figure 7

The photolysis rate constants of the compounds tested under solar irradiation were estimated based on the actinic flux at the Earth's surface (Demerjian et al., 1980) and on the quantum yields obtained in the study. Atmospheric lifetimes of the compounds due to photodecomposition were calculated to be 0.5–22 h for NPAHs and 4.5–35 h for OPAHs using the obtained photolysis rate constants. OPAHs were found to be more stable against photo-irradiation than were NPAHs. This finding indicates that the risk induced by OPAHs is critical for human health, because we may be continuously exposed to OPAHs due to their longer residence time in the atmosphere. Recently, it has been found that ROS, which can cause severe oxidative stress connected with inflammatory processes, is produced in larger amounts via chain reactions induced by quinoid PAHs in the human body (Chung et al., 2007). Further studies on the biological effects, and on the atmospheric formation and decomposition mechanisms of OPAHs are required.

References

- Atkinson R, Arey J (1994) Atmospheric chemistry of gas-phase polycyclic aromatic hydrocarbons: formation of atmospheric mutagens. *Environ. Health Perspect.* 102: 117-126
- Ball LM, Kohan MJ, Claxton LD, Lewtas J (1984) Mutagenicity of derivatives and metabolites of 1-nitropyrene: activation by rat liver S9 and bacterial enzymes. *Mutat. Res.* 138: 113-125
- Bamford HA, Bezabeh DZ, Schantz MM, Wise SA, Baker JE (2003) Determination and comparison of nitrated-polycyclic hydrocarbons measured in air and diesel particulate reference materials. *Chemosphere* 50: 575-587
- Barker CC, Emmerson RG, Periam JD (1955) The monosubstitution of triphenylene. *J. Chem. Soc.:* 4482-4485
- Bezabeh DZ, Bamford HA, Schantz MM, Wise SA (2003) Determination of nitrated polycyclic aromatic hydrocarbons in diesel particulate-related standard reference materials by using gas chromatography/mass spectrometry with negative ion chemical ionization. *Anal. Bioanal. Chem.* 375: 381-388
- Chung SW, Chung HY, Toriba A, Kameda T, Tang N, Kizu R, Hayakawa K (2007) An environmental quinoid polycyclic aromatic hydrocarbon, acenaphthenequinone, modulates cyclooxygenase-2 expression through reactive oxygen species generation and nuclear factor kappa B activation in A549 cells. *Toxicol. Sci.* 95: 348-355
- Ciccioli P, Cecinato A, Brancaleoni E, Frattoni M, Zacchei P, Miguel AH, de Castro Vasconcellos P (1996) Formation and transport of 2-nitrofluoranthene and 2-nitropyrene of photochemical origin in the troposphere. *J. Geophys. Res.* 101: 19567-19581
- Cvrčková O, Ciganeck M (2005) Photostability of polycyclic aromatic hydrocarbons (PAHs) and nitrated polycyclic aromatic hydrocarbons (NPAHs) in dichloromethane and isooctane solutions. *Polycyclic Aromat. Compd.* 25: 141-156
- Cvrčková O, Ciganeck M, Šimek Z (2006) Anthracene, chrysene, their nitro- and methyl-derivatives photostability in isooctane. *Polycyclic Aromat. Compd.* 26: 331-344

- Cwiertny DM, Young MA, and Grassian VH (2008) Chemistry and photochemistry of mineral dust aerosol. *Annu. Rev. Phys. Chem.* 59: 27-51
- Demerjian KL, Schere KL, Peterson JT (1980) Theoretical estimates of actinic (spherically integrated) flux and photolytic rate constants of atmospheric species in the lower troposphere. *Adv. Environ. Sci. Technol.* 10: 369-459
- Enya T, Suzuki H, Watanabe T, Hirayama T, Hisamatsu Y (1997) 3-nitrobenzanthrone, a powerful bacterial mutagen and suspected human carcinogen found in diesel exhaust and airborne particulates. *Environ. Sci. Technol.* 31: 2772-2776
- Esteve W, Budzinski H, Villenave E (2004) Relative rate constants for the heterogeneous reactions of OH, NO₂ and NO radicals with polycyclic aromatic hydrocarbons adsorbed on carbonaceous particles. Part 1: PAHs adsorbed on 1-2 µm calibrated graphite particles. *Atmos. Environ.* 38: 6063-6072
- Falkovich AH, Schkolnik G, Ganor E, Rudich Y (2004) Adsorption of organic compounds pertinent to urban environments onto mineral dust particles. *J. Geophys. Res.-Atmos.* 109: D02208
- Fernández RJ (2002) Do humans create deserts? *Trends Ecol. Evol.* 17: 6-7
- Finlayson-Pitts BJ, Pitts Jr JN (2000) *Chemistry of the upper and lower atmosphere*, Academic Press, San Diego, CA
- Garcia, H., Roth, H.D. (2002) Generation and reactions of organic radical cations in zeolites. *Chem. Rev.* 102, 3947-4007.
- Gibson TL, Korsog PE, Wolff GT (1986) Evidence for the transformation of polycyclic organic matter in the atmosphere. *Atmos. Environ.* 20: 1575-1578
- Inazu K, Tsutsumi N, Aika KI, Hisamatsu Y (2000) SO₂-enhanced nitration of fluoranthene and pyrene adsorbed on particulate matter in the heterogeneous reaction in the presence of NO₂. *Polycyclic Aromat. Compd.* 20: 191-203

- International Agency for Research on Cancer. (2013) Diesel and Gasoline Engine Exhausts and Some Nitroarenes, IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Vol. 105 (International Agency for Research on Cancer)
- Ishii S, Hisamatsu Y, Inazu K, Aika K (2000) Ambient measurement of nitrotriphenylenes and possibility of nitrotriphenylene formation by atmospheric reaction. *Environ. Sci. Technol.* 34: 1893-1899
- Ishii S, Hisamatsu Y, Inazu K, Aika K (2001) Environmental occurrence of nitrotriphenylene observed in airborne particulate matter. *Chemosphere* 44: 681-690
- Kameda T, Akiyama A, Toriba A, Tachikawa C, Yoshita M, Tang N, Hayakawa K (2008) Evaluation of endocrine disrupting activities of monohydroxylated derivatives of 1-nitropyrene by yeast two-hybrid assay. *J. Health Sci.* 54: 118-122
- Kameda T, Akiyama A, Toriba A, Tachikawa C, Yoshita M, Tang N, Hayakawa K (2011a) Mutagenicities and endocrine-disrupting activities of 1-hydroxy-2-nitropyrene and 1-hydroxy-5-nitropyrene. *J. Health Sci.* 57: 372-377
- Kameda T, Akiyama A, Toriba A, Tang N, Hayakawa K (2010) Determination of particle-associated hydroxynitropyrenes with correction for chemical degradation on a quartz fibre filter during high volume air sampling. *Int. J. Environ. Anal. Chem.* 90: 976-987
- Kameda T, Akiyama A, Toriba A, Tang N, Hayakawa K (2011b) Atmospheric formation of hydroxynitropyrenes from a photochemical reaction of particle-associated 1-nitropyrene. *Environ. Sci. Technol.* 45: 3325-3332
- Kameda T, Inazu K, Asano K, Murota M, Takenaka N, Sadanaga Y, Hisamatsu Y, Bandow H (2013) Prediction of rate constants for the gas phase reactions of triphenylene with OH and NO₃ radicals using a relative rate method in CCl₄ liquid phase-system. *Chemosphere* 90: 766-771
- Kameda T, Inazu K, Hisamatsu Y, Takenaka N, Bandow H (2004) Determination of atmospheric nitro-polycyclic aromatic hydrocarbons and their precursors at a heavy traffic roadside and at a residential area in Osaka, Japan. *Polycyclic Aromat. Compd.* 24: 657-666

- Kameda T, Inazu K, Hisamatsu Y, Takenaka N, Bandow H (2006) Isomer distribution of nitrotriphenylenes in airborne particles, diesel exhaust particles, and the products of gas-phase radical-initiated nitration of triphenylene. *Atmos. Environ.* 40: 7742-7751
- Kameda T, Nakayama Y, Goto T, Koyanagi T, Bandow H, Fujimori K, Toriba A, Tang N, Hayakawa K (2009) Photochemical degradation of selected nitro- and oxy-polycyclic aromatic hydrocarbons on airborne particles under simulated solar UV-irradiation. *Airborne Particulates*, Nova Science Publishers, NY, pp.291-307
- Kameda T, Azumi E, Fukushima A, Tang N, Matsuki A, Kamiya Y, Toriba A, Hayakawa K (2016) Mineral dust aerosols promote the formation of toxic nitropolycyclic aromatic compounds. *Sci. Rep.* 6: 24427
- Kamens RM, Guo Z, Fulcher J, Bell D (1988) Influence of humidity, sunlight, and temperature on the daytime decay of polyaromatic hydrocarbons on atmospheric soot particles. *Environ. Sci. Technol.* 22: 103-108
- Kawanaka Y, Sakamoto K, Wang N, Yun S (2005) Determination of nitroarenes and 3-nitrobenzanthrone in atmospheric particulate matter by gas chromatography/tandem mass spectrometry with negative ion chemical ionization. *Bunseki Kagaku* 54: 685-691
- Laszlo P (1987) Chemical reaction on clays. *Science* 235: 1473-1477
- Ma JZ, Liu YC, He H (2011) Heterogeneous reactions between NO₂ and anthracene adsorbed on SiO₂ and MgO. *Atmos. Environ.* 45: 917-924
- Manabe Y, Kinouchi T, Ohnishi Y (1985) Identification and quantification of highly mutagenic nitroacetoxypyrenes and nitrohydroxypyrenes in diesel-exhaust particles. *Mutat. Res.* 158: 3-18
- Miet K, Le Menach K, Flaud PM, Budzinski H, Villenave E (2009) Heterogeneous reactivity of pyrene and 1-nitropyrene with NO₂: Kinetics, product yields and mechanism. *Atmos. Environ.* 43: 837-843

- Nguyen ML, Bedjanian Y, Guilloteau A (2009) Kinetics of the reactions of soot surface-bound polycyclic aromatic hydrocarbons with NO₂. *J. Atmos. Chem.* 62: 139-150
- Parry EP (1963) An infrared study of pyridine adsorbed on acidic solids characterization of surface acidity. *J. Catal.* 2: 371-379
- Phousongphouang PT, Arey J (2003) Sources of the atmospheric contaminants, 2-nitrobenzanthrone and 3-nitrobenzanthrone. *Atmos. Environ.* 37: 3189-3199
- Radner F (1983) Nitration of polycyclic aromatic hydrocarbons with dinitrogen tetroxide. A simple and selective synthesis of mononitroderivatives. *Acta Chem. Scand. Ser. B* 37: 65-67
- Ramdahl T, Bjorseth A, Lokensgard DM, Pitts JN (1984) Nitration of polycyclic aromatic hydrocarbons adsorbed to different carriers in a fluidized-bed reactor. *Chemosphere* 13: 527-534
- Reisen F, Arey J (2005) Atmospheric reactions influence seasonal PAH and nitro-PAH concentrations in the Los Angeles basin. *Environ. Sci. Technol.* 39: 64-73
- Ridd JH (1991) The range of radical processes in nitration by nitric acid. *Chem. Soc. Rev.* 20: 149-165
- Rosser PF, Ramachandran P, Sangaiah R, Austin RN, Gold A, Ball LM (1996) Role of O-acetyltransferase in activation of oxidised metabolites of the genotoxic environmental pollutant 1-nitropyrene. *Mutat. Res.* 369: 209-220
- Salmeen IT, Pero AM, Zator R, Schuetzle D, Riley TL (1984) Ames assay chromatograms and the identification of mutagens in diesel particle extracts. *Environ. Sci. Technol.* 18: 375-382
- Sasaki J, Aschmann SM, Kwok ESC, Atkinson R, Arey J (1997) Products of the gas-phase OH and NO₃ radical-initiated reactions of naphthalene. *Environ. Sci. Technol.* 31: 3173-3179
- Schuetzle D (1983) Sampling of vehicle emissions for chemical analysis and biological testing. *Environ. Health Perspect.* 47: 65-80

- Schuetzle D, Jensen TE, Ball JC (1985) Polar polynuclear aromatic hydrocarbon derivatives in extracts of particulates: Biological characterization and techniques for chemical analysis. *Environ. Int.* 11: 169-181
- Schuetzle D, Riley TL, Prater TJ, Harvey TM, Hunt DF (1982) Analysis of nitrated polycyclic aromatic-hydrocarbons in diesel particulates. *Anal. Chem.* 54: 265-271
- Shiraiwa M, Garland RM, Pöschl U (2009) Kinetic double-layer model of aerosol surface chemistry and gas-particle interactions (K2-SURF): Degradation of polycyclic aromatic hydrocarbons exposed to O₃, NO₂, H₂O, OH and NO₃. *Atmos. Chem. Phys.* 9: 9571-9586
- Shiri M, Zolfigol MA, Kruger HG, Tanbakouchian Z (2010) Advances in the application of N₂O₄/NO₂ in organic reactions. *Tetrahedron* 66: 9077-9106
- Soma Y, Soma M (1989) Chemical reactions of organic compounds on clay surfaces. *Environ. Health Perspect.* 83: 205-214
- Tanaka TY, Chiba M (2006) A numerical study of the contributions of dust source regions to the global dust budget. *Global Planet. Change* 52: 88-104
- Usher CR, Michel AE, Grassian VH (2003) Reactions on mineral dust. *Chem. Rev.* 103: 4883-4939
- Wang H, Hasegawa K, Kagaya S (2000) The nitration of pyrene adsorbed on silica particles by nitrogen dioxide. *Chemosphere* 41: 1479-1484
- Warner SD, Farant JP, Butler IS (2004) Photochemical degradation of selected nitropolycyclic aromatic hydrocarbons in solution and adsorbed to solid particles. *Chemosphere* 54: 1207-1215
- Yang DTC, Chou A, Chen E, Chiu LH (1994) Photodecomposition of environmental nitro-polycyclic aromatic hydrocarbons. *Polycyclic Aromat. Compd.* 5: 201-208
- Yang XY, Igarashi K, Tang N, Lin J, Wang W, Kameda T, Toriba A, Hayakawa K (2010) Indirect- and direct-acting mutagenicity of diesel, coal and wood burning-derived particulates

and contribution of polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons. *Mutat. Res.-Genet. Toxicol. Environ. Mutag.* 695: 29-34

Zielinska B, Arey J, Atkinson R, Ramdahl T, Winer AM (1989) The nitroarenes of molecular weight 247 in ambient particulate samples collected in southern California. *Atmos. Environ.* 23: 223-229

Zielinska B, Arey J, Atkinson R, Ramdahl T, Winer AM, Pitts Jr JN (1986) Reaction of dinitrogen pentoxide with fluoranthene. *J. Am. Chem. Soc.* 108: 4126-4132

Table 1. Observed pseudo-first order rate constants for the reaction of Py on the substrates examined in this study with 3 ppmv NO₂ (k_{obs}), percentage of degraded Py (D_{Py}), and yields of 1-NP ($Y_{1\text{-NP}}$) (Reproduced from Kameda et al., 2016).

Substrates	$k_{\text{obs}} \times 10^5 \text{ (s}^{-1}\text{)}^*$	$D_{\text{Py}} \text{ (}\%\text{)}^\dagger$	$Y_{1\text{-NP}} \text{ (}\%\text{)}^\dagger$	DNP formation [‡]
Chinese desert dust (CDD)	86 ± 4	96	53	+
Arizona test dust (ATD)	36 ± 1	88	58	+
Kaolin [§]	110 ± 10	98	60	+
Montmorillonite A	53 ± 5	95	89	+
Montmorillonite B	29 ± 4	84	79	+
Saponite	39 ± 3	82	73	-
Potassium feldspar	1.1 ± 0.2	14	10	-
Sodium feldspar	0.30 ± 0.06	12	6	-
Feldspar	0.86 ± 0.14	17	4	-
Limestone	1.4 ± 0.1	18	5	-
Dolomite	0.83 ± 0.15	16	4	-
Calcium sulfate	1.5 ± 0.5	6	0	-
Quartz	1.7 ± 0.1	9	5	-
Aluminum oxide	0.25 ± 0.00	2	1	-
Iron (III) oxide	9.0 ± 3.3	17	0	-
Titanium (IV) oxide	1.4 ± 0.0	14	3	-
Montmorillonite K10	250 ± 20	100	6	+
Graphite [#]	1.9 ± 0.1	9	1	-

* Errors represent one standard error derived from nonlinear least-squares fitting for the Py decay plots.

† Obtained from reactions for 2 h.

‡ Reaction time, 12 h; +, yes; -, no.

§ Note that kaolin consists largely of kaolinite.

|| Acid-activated montmorillonite.

As a control.

Table 2. Quantum yield, photolysis rate constant, and estimated lifetime of the PAH derivatives on the Earth's surface (Reproduced in part from Kameda et al., 2009). See Fig. 7 for compound abbreviations.

Compound	Photolysis rate constant (s^{-1})	Lifetime (h)
BaP	2.22×10^{-5}	74.6
BkF	9.89×10^{-6}	125.3
1-OHPyr	2.38×10^{-4}	9.3
1,3-DNP	8.15×10^{-5}	15.3
1,6-DNP	1.75×10^{-4}	7.6
1,8-DNP	2.23×10^{-4}	5.8
1-NP	7.85×10^{-5}	17.8
1-NPer	1.42×10^{-3}	1.0
2-NA	1.01×10^{-3}	1.6
2-NF	3.26×10^{-4}	6.6
2-NFR	1.05×10^{-4}	17.4
2-NTP	1.02×10^{-4}	21.1
3-NBA	5.23×10^{-5}	22.2
3-NFR	8.56×10^{-5}	17.8
3-NPer	1.65×10^{-4}	9.9
4-NP	7.31×10^{-5}	21.5
6-NBaP	1.38×10^{-3}	1.0
6-NC	5.93×10^{-4}	2.8
7-NBaA	2.12×10^{-3}	0.9
9-NA	3.15×10^{-3}	0.5
9-NPh	4.79×10^{-4}	4.3
1,2-BAQ	4.74×10^{-5}	34.9
1,4-CQ	7.63×10^{-5}	15.8
5,6-CQ	9.86×10^{-5}	11.6
9,10-PQ	2.23×10^{-4}	7.1
AcQ	2.35×10^{-4}	7.0
AQ	8.38×10^{-5}	22.7
B[c]P-5,6-Q	4.33×10^{-4}	4.5

Figure Captions

Figure 1. A schematic for the formation of 2-NTP from OH or NO₃ radical-initiated reactions of triphenylene (Kameda, 2011). [Reproduced with permission from Journal of Health Science Vol. 57 No. 6. Copyright 2011 The Pharmaceutical Society of Japan]

Figure 2: Concentrations of Py and nitropyrenes (1-NP and DNPs) on quartz (a) and CDD (b) (expressed as a percent of the initial Py concentration) after exposure to 3 ppmv NO₂ for the indicated times (Reproduced from Kameda et al., 2016). The data points represent mean values (± 1 SD) of triplicate experiments: circles, Py; squares, 1-NP; diamonds, DNPs (= 1,3-DNP + 1,6-DNP + 1,8-DNP). The curves for Py decay are exponential nonlinear least-squares fits assuming first-order reactions. The curves for nitropyrene formation are for illustrative purposes only.

Figure 3: IR spectra of pyridine adsorbed on the substrates examined in the study (Reproduced from Kameda et al., 2016).

Figure 4: Proposed mechanisms for the nitration of aromatic compounds (ArH) on the acidic surface of mineral dust (Reproduced from Kameda et al., 2016). (a) Schematic illustration of the heterogeneous nitration. The gas-particle interface is divided into a gas-phase with gaseous NO₂ (NO₂(g)), a sorption layer with adsorbed NO₂ (NO₂(s)), a quasi-static surface layer with the aromatic radical cation (ArH⁺·), and a particle bulk. The blue arrow indicates the adsorption and desorption fluxes of NO₂. The red and purple arrows indicate chemical reactions and electron transfer, respectively. This heterogeneous chemistry is based on the Pöschl-Rudich-Ammann (PRA) framework (Shiraiwa et al., 2009). (b) Chemical reaction scheme of the nitroaromatic compound (ArNO₂) formation on Lewis acid sites of the dust surface (DS⁺).

Figure 5. Structure of hydroxynitropyrene.

Figure 6. Chromatograms of standard OHNPs (a) and photoreaction products of 1-NP (b) obtained by the HPLC-chemiluminescence detection system (Kameda, 2011). [Reproduced with permission from Journal of Health Science Vol. 57 No. 6. Copyright 2011 The Pharmaceutical Society of Japan]

Figure 7. Structures of PAH derivatives examined in the photodecomposition experiment.

Figure 1:

Figure 2:

Figure 3:

Figure 4:

b

Figure 5:

Figure 6:

Figure 7:

Figure 7 (continued):

Acenaphthenequinone
(AcQ)

Anthraquinone
(AQ)

9,10-Phenanthrenequinone
(9,10-PQ)

1,2-Benzanthraquinone
(1,2-BAQ)

Benzo[c]phenanthrene-5,6-quinone
(B[c]P-5,6-Q)

1,4-Chrysenequinone
(1,4-CQ)

5,6-Chrysenequinone
(5,6-CQ)