TITLE:

Impact of cumulative steroid dose on infectious diseases after allogeneic hematopoietic stem cell transplantation

AUTHOR(S):

Watanabe, Mizuki; Kanda, Junya; Hishizawa, Masakatsu; Kondo, Tadakazu; Yamashita, Kouhei; Takaori-Kondo, Akifumi

CITATION:

Watanabe, Mizuki ...[et al]. Impact of cumulative steroid dose on infectious diseases after allogeneic hematopoietic stem cell transplantation. *Transplant Infectious Disease* 2019, 21(2): e13049.

ISSUE DATE:

2019-04

URL:

<http://hdl.handle.net/2433/240763>

RIGHT:

This is the peer reviewed version of the following article: Watanabe, M, Kanda, J, Hishizawa, M, Kondo, T, Yamashita, K, Takaori - Kondo, A. Impact of cumulative steroid dose on infectious diseases after allogeneic hematopoietic stem cell transplantation. *Transpl Infect Dis*. 2019; 21:e13049., which has been published in final form at <https://doi.org/10.1111/tid.13049>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.; The full-text file will be made open to the public on 5 April 2020 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。; This is not the published version. Please cite only the published version.

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

Impact of cumulative steroid dose on infectious diseases after allogeneic hematopoietic stem cell transplantation

Mizuki Watanabe¹, Junya Kanda¹, Masakatsu Hishizawa¹, Tadakazu Kondo¹, Kouhei Yamashita¹, Akifumi Takaori-Kondo¹

¹Department of Hematology and Oncology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

Conflict-of-interest disclosure

The authors declare no competing financial interests

Abstract

Systemic steroid is used to treat various transplant-related complications after allogeneic hematopoietic stem cell transplantation (allo-HSCT). However, measures to evaluate its impact on infections are still limited. Hence, we examined the cumulative steroid dose used within 30 days after transplant as a predictor of future risk of infections. This study included 226 patients who underwent their first allo-HSCT at Kyoto University Hospital between 2005 and 2015. Sixty-one patients received transplantation from related donors, 106 received unrelated BMT and 59 received unrelated single-unit CBT. Patients were categorized into 3 groups according to the cumulative steroid dose in terms of prednisolone: no-steroid group (n=174), low-dose group (≤ 7 mg/kg) (n=22) and high-dose group (> 7 mg/kg) (n=30). In a multivariate analysis, high-dose steroid administration was associated with CMV antigenemia (HR 1.91, $P=0.037$) and bacteremia (HR 2.59, $P=0.053$). No impact was found on the occurrence of invasive fungal infection. In conclusion, high-dose cumulative steroid could predict high risks of bacteremia and CMV antigenemia. Additional anti-bacterial agents for fever and regular measurement of CMV antigen are recommended for whom with systemic steroid administration even after neutrophil engraftment.

Text word count; abstract word count, 182; tables, 3; figures, 3; supplemental tables, 0

1 Introduction

Systemic steroid is frequently used as a primary treatment for transplant-related complications such as graft-versus-host disease (GVHD) and non-infectious pulmonary complications after allogeneic hematopoietic stem cell transplantation (HSCT)^{1–3}. The use of systemic steroid along with the occurrence of GVHD has been suggested to be a risk factor for various infectious diseases^{4–8}, which are main causes of transplant-related mortality. Since the number of HSCTs with a higher risk of complications, such as cord blood transplantations (CBT) and HLA mismatch transplantations in older patients, has been increasing^{9–12}, it is important to evaluate the effect of steroid use on clinical outcomes.

The associations between the cumulative dose of steroid and the occurrence of side effects have been discussed in patients with non-hematologic diseases who receive systemic steroid for a prolonged period. The impact of the cumulative dose of steroid on infectious complications has been controversial, although a positive association was noted in patients taking immunosuppressive agents after solid organ transplantations^{13–16}. Similarly, in recipients of HSCT, steroid administration could increase the risk of infectious complications because of the concomitant use of calcineurin inhibitors and the delay of immune reconstitution after HSCT. However, there is little information available regarding the steroid dose. Hence, in the present study, we examined the impact of the cumulative steroid dose on the risk of infectious diseases after HSCT in a single transplant center.

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

Methods

Data collection

A total of 238 patients who underwent their first allogeneic HSCT for hematologic diseases at Kyoto University Hospital from 2005 to 2015 and survived at least 30 days after transplantation were included. Patients who had already started to receive steroid before transplantation were excluded. Patients who had active bacterial, fungal, or viral infection at transplantation or who had had history of invasive fungal infection before transplantation were also excluded. The Institutional Review Board of Kyoto University Hospital, where this study was organized, approved this study.

Treatment Policy and Definition

Definitions

Neutrophil engraftment was diagnosed when an absolute neutrophil count over 500/ μ L was observed for 3 days in a row. Acute GVHD was diagnosed and classified by each physician according to traditional criteria¹⁷.

Invasive fungal infections

β D-glucan was examined once a week, and imaging inspection and blood culture were examined for fever or other suspicious conditions. Diagnoses of invasive fungal infections were categorized into 3 types; possible, probable, and proven, based on the practice guidelines from the Infectious Diseases Society of America (IDSA) and Japanese guidelines^{18–20}.

In our hospital, antifungal prophylaxis was administered in all patients who underwent allo-HSCT. The antifungal agents that were generally used as prophylaxis were oral fluconazole, voriconazole, micafungin and liposomal

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

1 amphotericin B injection, according to each patient's history of fungal infection.

2 All patients were hospitalized in a cleanroom of ISO Class 5 (ISO 14644-1)²¹
3 before and in the early period after day0 and moved to a cleanroom of ISO Class
4 6 (ISO 14644-1)²¹ after they achieved neutrophil engraftment.

5
6 *CMV antigenemia and CMV disease*

7 CMVpp65 antigen examinations were performed using C10/11²² method or
8 C7-HRP²³ method once a week for every patient after transplantation and
9 examined additionally for suspicious symptoms of CMV diseases.

10 In cases with more than 3 positive cells in 2 slides (C10/C11 method) or more
11 than 2 positive cells out of 50000 WBC (C7-HRP method), pre-emptive therapy
12 was given followed by close CMV-antigen monitoring^{22,23}. Diagnosis of CMV
13 end-organ diseases were diagnosed according to published definitions²⁴.

14
15 *Other viremias*

16 Patients were examined by viral PCR detection at the timing of fever of unknown
17 origin or any other symptoms of infection based on the judgment of each
18 physician in charge. Viruses examined in PCR included adenovirus, BK virus,
19 JC virus, varicella zoster virus, human herpes simplex, EB virus and other
20 viruses according to each patient's symptoms.

21
22 *Bacteremia*

23 Two sets of blood culture were examined for each patient with fever or any other
24 symptoms suggesting infectious diseases. As our policy, antibacterial
25 prophylaxis was not applied in every patient, except for those who were at high
26 risk of bacterial infection, such as those with a history of repeated severe

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

1 bacterial infection or a long history of chemotherapeutic treatment.

3 **Endpoints**

4 The endpoint of this study was the incidence of various infectious diseases
5 including invasive fungal infection, cytomegalovirus (CMV) antigenemia, and
6 bacteremia diagnosed from 30 days to 6 months after HSCT. The cumulative
7 steroid dose was calculated as the total amount administered per patient within
8 30 days after transplantation, since the first steroid administration mainly began
9 within this period as a treatment for pre-engraftment or engraftment syndrome
10 and for acute GVHD.

12 **Statistical analysis**

13 Descriptive statistics were used to summarize variables related to patient
14 characteristics. We calculated the cumulative steroid dose within 30 days after
15 HSCT. The landmark day was set at 30 days after transplantation.
16 Prednisolone-equivalent conversion was performed in accordance to the general
17 formula²⁵. Episodes of infectious diseases (invasive fungal infection, CMV
18 antigenemia or disease, and bacteremia) were calculated based on cumulative
19 incidence curves. A competing event was death without infectious disease.
20 Cumulative incidences in the groups were compared using the Gray test. Fine
21 and Gray's proportional hazards model was used to evaluate the effect of
22 cumulative steroid dose on the occurrence of infectious diseases²⁶. The
23 following covariates were considered; recipient's sex, age (<50 or ≥50 years
24 old), disease diagnosis (myeloid malignancies, lymphoid malignancies, or
25 others), year of transplantation (2005-2009 or 2010-2016), disease status
26 (complete remission [CR] or non-CR), donor type (bone marrow transplantation

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

from unrelated donor, peripheral blood stem cell transplantation from related donor, or CBT), conditioning regimen (reduced-intensity or myeloablative), GVHD prophylaxis (tacrolimus or cyclosporine in addition to mycophenolate mofetil or methotrexate), presence or absence of neutrophil engraftment at day 30, and prophylactic administration of levofloxacin. All factors, in addition to the main effect, were selected with a variable retention criterion of $P < 0.05$ in the univariate analysis and analyzed in the multivariate analysis.

Although acute GVHD has been suggested to be a risk factor for infectious diseases after HSCT, we did not include acute GVHD because there was a correlation between acute GVHD and steroid administration (data not shown), and it would be inappropriate to include both in the same model.

All statistical analyses were performed with Stata version 14 (Stata Corp, College Station, TX) and EZR (Saitama Medical Center, Jichi Medical University, Saitama, Japan), which is a graphical user interface for R (The R Foundation for Statistical Computing, version 3.1.1, Vienna, Austria).

Results

Patient characteristics

Sixty-one patients received transplantation from a related donor, 106 received unrelated bone marrow grafts, and 59 received unrelated cord blood units. Their median age was 51 years (range, 17–66). Neutrophil engraftment was achieved in 203 patients (90%) by day 30 and mean neutrophil engraftment day from transplantation in each graft were 21 in bone marrow transplantation, 17 in peripheral blood stem cell transplantation and 25 in cord blood transplantation.

Patients were categorized into 3 groups according to the cumulative steroid dose within 30 days: no steroid group (n = 174), low-dose cumulative steroid group (7mg/kg or less of prednisolone-equivalent dose, n = 22), and high-dose cumulative steroid group (over 7mg/kg of prednisolone-equivalent dose, n =30). The cutoff value of 7mg/kg of prednisolone-equivalent dose approximately stands for initial steroid treatment against acute GVHD in Japan (1mg/kg during 7 days at maximum). The reason for steroid administration was treatment for GVHD in 33 patients, engraftment syndrome in 9, and other reasons including lung complications in 10. Grade II to IV acute GVHD was diagnosed in 96 patients in total. There was no obvious difference in background among the different donor sources. (Table 1).

Invasive fungal infection

We observed 13 cases of invasive fungal infection, including one proven case with candida bloodstream infection and 2 probable and 10 possible cases of pneumonia. The cumulative incidence of invasive fungal infection at 6 months was 5.7%, 4.5%, and 6.7% in the no-administration, low-dose, and high-dose groups, respectively (P = 0.231, Gray test) (Figure 1). Multivariate analysis

showed no association between steroid administration and the occurrence of invasive fungal infection. We found no other significant risk factor.

CMV antigenemia and diseases

Eighty-six HSCT were performed from CMV-antibody (Ab) positive donors to CMV-Ab positive recipients, 10 were from CMV-Ab positive donors to CMV negative recipients and the other 103 were from CMV-Ab negative donors (Table 1).

A total of 105 (46%) patients were diagnosed as CMV antigenemia and 81 (78%) received Ganciclovir as a pre-emptive antiviral therapy. Seven patients were pathologically diagnosed as CMV disease including colitis and hepatitis, all of whom were positive for CMV antigenemia. There were 4 cases of CMV antigenemia with clinically suspected CMV diseases, although they were not definitely diagnosed due to a lack of pathological evidence. No patient died of CMV-related complications. The cumulative incidences of CMV antigenemia at 6 months in the no-administration, low-dose, and high-dose groups were 49.7%, 68.8%, and 69.6%, respectively ($P = 0.038$) (Figure 2). Reason for steroid initiation had little impact on the occurrence of CMV antigenemia (GVHD vs other reasons: HR 2.119, $P=0.089$). Multivariate analysis showed that both a low-dose and high-dose of cumulative steroid administration were associated with CMV reactivation, although the association in the low-dose group was not statistically significant (low-dose vs. no-administration group: HR 1.64, $P=0.100$, high-dose vs. no-administration group: HR 1.91 $P = 0.037$). Other risk factors detected were cord blood unit as a donor source (cord blood unit vs. sibling donor: HR 1.62, $P = 0.018$) and recipient age over 50 years at transplantation (age ≥ 50 vs. <50 : HR 1.62, $P = 0.007$) (Table 2).

1

2 **Viral infections other than CMV**

3 A total of 15 cases were diagnosed as viremia including Adenovirus in 1 patient,
4 BK virus in 2, Epstein Barr virus in 1, Varicella Zoster virus in 3, and human
5 herpes virus 6 in 7. Ten patients were in the no-administration group and there
6 was no association between viremia and the cumulative steroid dose.

8 **Bacteremia**

9 The cumulative incidences of bacteremia at 6 months in the no-administration,
10 low-dose, and high-dose groups were 9.3%,15.8%, and 21.7%, respectively (P =
11 0.224) (Figure 3). Detected microbes at the first onset of bacteremia were
12 gram-negative rods in 15 cases, gram-positive cocci in 7 cases, and
13 gram-positive rods in 1 case. Reason for steroid initiation had little impact on the
14 occurrence of bacteremia (GVHD vs other reasons: HR 4.89, P=0.14).
15 Administration of levofloxacin showed no apparent prophylactic effect on
16 bacteremia (HR 0.73, P=0.574).
17 Multivariate analysis showed that the high-dose group was marginally
18 associated with an increased risk of bacteremia (low-dose vs. no-administration
19 group: HR 2.13, P =0.240, high-dose vs. no-administration group: HR 2.59, P =
20 0.053). Regarding the microbes detected, there was no significant difference
21 among the three groups. The other major risk factor for bacteremia was a
22 recipient age over 50 years at transplantation, which had a HR of 2.69 (age >=50
23 vs. <50: P = 0.021) (Table 3).

25 **Other bacterial infections**

26 The other infectious events proven as bacterial complications were 4 cases

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

- 1 *Clostridium difficile* colitis, 2 cases of pneumonia (1 of *Pseudomonas*
- 2 *aeruginosa*, 1 of *Stenotrophomonas maltophilia*), 1 cellulitis of
- 3 *Coagulase-negative staphylococcus*, and 1 endophthalmitis of
- 4 *Coagulase-negative staphylococcus*.

Discussion

In the present study, we examined the impact of the cumulative dose of steroid on infectious complications after HSCT and found associations between steroid dose and both CMV and bacterial infections following HSCT.

Although acute GVHD and systemic steroid have been reported to be risk factors for invasive fungal infection after HSCT^{27–30}, the cumulative steroid dose was not associated with fungal infection in our study. All patients in our hospital continued prophylactic treatment with antifungal drugs according to the risk of fungal infection, following Japanese and European guidelines³¹. Only 13 of 226 patients had invasive fungal infection over 10 years, although our cohort included a relatively large number of cord blood transplantations. This suggests that fungal infection could be avoided regardless of the occurrence of acute GVHD, steroid use, and donor source by appropriate clinical practice.

With regard to CMV-related complications, steroid use was strongly associated with CMV antigenemia regardless of the cumulative dose, which is similar to previous reports^{4,32,33}. Almost all the patients in our cohort were seropositive before transplant and thus CMV antigen levels must be measured regularly after HSCT. Another risk factor for CMV antigenemia was cord blood unit as a donor source, although the HR was lower than previously reported and there were no CMV-related deaths. Older patients also had a higher risk of CMV antigenemia. Contrary to a previous report⁴, myeloablative conditioning was not found to be a risk factor for CMV antigenemia, which is probably due to the difference in the conditioning regimen or the medication used for GVHD prophylaxis.

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

High-dose, but not low-dose, cumulative steroid administration was a risk factor for bacterial infection. Anti-bacterial prophylaxis and preemptive therapies for fever of undetected origin might be better considered for patients after HSCT receiving a high cumulative dose of steroid, regardless of their neutrophil count. An advanced age at transplant was another risk factor for bacterial infection after HSCT, which was consistent with previous reports³⁴.

The present study has several limitations. First, this is a retrospective study of small population with heterogeneous background in a single transplant center. Second, the loads of viruses other than CMV were not regularly measured and the timing of the examination was determined by each physician in charge. Finally, information on blood sugar levels was not collected, although blood sugar levels were checked regularly and treated by continuous intravenous insulin infusion, which minimized the effect of hyperglycemia on bacterial infections.

In conclusion, our study confirmed that the cumulative steroid dose could be a good prognostic marker for CMV antigenemia and bacterial infection after HSCT. These post-transplant complications must be detected and managed in the early period, particularly in elderly patients who are receiving a high cumulative dose of steroid.

ACKNOWLEDGEMENTS

We are grateful to Emi Furusaka, Tomoko Okuda and Megumi Oka for their expert data-management and secretarial assistance, and to all the members of the transplant teams at Kyoto University Hospital for their dedicated care of the

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

1 patients and donors.

2 This work was supported in part by the Takeda Science Foundation (JK).

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

Reference

1. Jamil MO, Mineishi S. State-of-the-art acute and chronic GVHD treatment. *Int J Hematol.* 2015;101(5):452-466. doi:10.1007/s12185-015-1785-1
2. Sung AD, Chao NJ. Concise Review: Acute Graft-Versus-Host Disease: Immunobiology, Prevention, and Treatment. *Stem Cells Transl Med.* 2013;2(1):25-32. doi:10.5966/sctm.2012-0115
3. Magenau J, Reddy P. Next generation treatment of acute graft-versus-host disease. *Leukemia.* 2014;28(12):2283-2291. doi:10.1038/leu.2014.195
4. Cohen L, Yeshurun M, Shpilberg O, Ram R. Risk factors and prognostic scale for cytomegalovirus (CMV) infection in CMV-seropositive patients after allogeneic hematopoietic cell transplantation. *Transpl Infect Dis.* 2015;17(4):510-517. doi:10.1111/tid.12398
5. Faraci M, Lanino E, Morreale G, et al. Bacteremias and invasive fungal diseases in children receiving etanercept for steroid-resistant acute GVHD. *Bone Marrow Transplant.* 2011. doi:10.1038/bmt.2010.80
6. Greco R, Crucitti L, Noviello M, et al. Human Herpesvirus 6 Infection Following Haploidentical Transplantation: Immune Recovery and Outcome. *Biol Blood Marrow Transplant.* 2016. doi:10.1016/j.bbmt.2016.09.018
7. Mihu CN, Schaub J, Kesh S, et al. Risk Factors for Late Staphylococcus Aureus Bacteremia after Allogeneic Hematopoietic Stem Cell Transplantation: A Single-Institution, Nested Case-Controlled Study. *Biol Blood Marrow Transpl.* 2008;14:1429-1433. doi:10.1016/j.bbmt.2008.09.005
8. Yoon HS, Lee JH, Choi ES, et al. Cytomegalovirus infection in children

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

- 1 who underwent hematopoietic stem cell transplantation at a single center:
- 2 A retrospective study of the risk factors. *Pediatr Transplant*.
- 3 2009;13(7):898-905. doi:10.1111/j.1399-3046.2008.01084.x
- 4 9. Sawada A, Inoue M, Koyama-Sato M, et al. Umbilical cord blood as an
- 5 alternative source of reduced-intensity hematopoietic stem cell
- 6 transplantation for chronic epstein-barr virus-associated T or natural killer
- 7 cell lymphoproliferative diseases. *Biol Blood Marrow Transplant*.
- 8 2014;20(2):214-221. doi:10.1016/j.bbmt.2013.10.026
- 9 10. Anasetti C. Use of alternative donors for allogeneic stem cell
- 10 transplantation. *Hematology Am Soc Hematol Educ Program*.
- 11 2015;2015(1):220-224. doi:10.1182/asheducation-2015.1.220
- 12 11. Lorentino F, Labopin M, Fleischhauer K, et al. The impact of HLA
- 13 matching on outcomes of unmanipulated haploidentical HSCT is
- 14 modulated by GVHD prophylaxis. *Blood Adv*. 2017;1(11):669-680.
- 15 doi:10.1182/bloodadvances.2017006429
- 16 12. Hambach L, Stadler M, Dammann E, Ganzer A, Hertenstein B. Increased
- 17 risk of complicated CMV infection with the use of mycophenolate mofetil in
- 18 allogeneic stem cell transplantation. *Bone Marrow Transpl*.
- 19 2002;29(11):903-906. doi:10.1038/sj.bmt.1703583
- 20 13. Del Rincón I, Battafarano DF, Restrepo JF, Erikson JM, Escalante A.
- 21 Glucocorticoid dose thresholds associated with all-cause and
- 22 cardiovascular mortality in rheumatoid arthritis. *Arthritis Rheumatol*. 2014.
- 23 doi:10.1002/art.38210
- 24 14. Dixon WG, Kezouh A, Bernatsky S, Suissa S. The influence of systemic
- 25 glucocorticoid therapy upon the risk of non-serious infection in older
- 26 patients with rheumatoid arthritis: A nested case-control study. *Ann*

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

- 1
- 2
- 3
- 4
- 5
- 6 1 *Rheum Dis.* 2011;70(6):956-960. doi:10.1136/ard.2010.144741
- 7
- 8 2 15. STANBURY RM, GRAHAM EM. Systemic corticosteroid therapy---side
- 9
- 10 3 effects and their management. *Br J Ophthalmol.* 1998;82(6):704-708.
- 11
- 12 4 doi:10.1136/bjo.82.6.704
- 13
- 14 5 16. Opelz G, Döhler B. Association between steroid dosage and death with a
- 15
- 16 6 functioning graft after kidney transplantation. *Am J Transplant.*
- 17
- 18 7 2013;13(8):2096-2105. doi:10.1111/ajt.12313
- 19
- 20 8 17. Przepiorka D, DJ W, Martin P, et al. 1994 Consensus Conference on
- 21
- 22 9 *Acute GvHD Grading.* Vol 15.; 1995.
- 23
- 24 10 18. Patterson TF, Thompson GR, Denning DW, et al. Practice guidelines for
- 25
- 26 11 the diagnosis and management of aspergillosis: 2016 update by the
- 27
- 28 12 infectious diseases society of America. *Clin Infect Dis.* 2016;63(4):e1-e60.
- 29
- 30 13 doi:10.1093/cid/ciw326
- 31
- 32 14 19. Pappas PG, Kauffman CA, Andes DR, et al. Clinical Practice Guideline for
- 33
- 34 15 the Management of Candidiasis: 2016 Update by the Infectious Diseases
- 35
- 36 16 Society of America. *Clin Infect Dis.* 2015;62(4):e1-e50.
- 37
- 38 17 doi:10.1093/cid/civ933
- 39
- 40 18 20. De Pauw B, Walsh TJ, Donnelly JP, et al. Revised Definitions of Invasive
- 41
- 42 19 Fungal Disease from the European Organization for Research and
- 43
- 44 20 Treatment of Cancer/Invasive Fungal Infections Cooperative Group and
- 45
- 46 21 the National Institute of Allergy and Infectious Diseases Mycoses Study
- 47
- 48 22 Group (EORTC/MSG) C. *Clin Infect Dis.* 2008;46(12):1813-1821.
- 49
- 50 23 doi:10.1086/588660
- 51
- 52 24 21. INTERNATIONAL STANDARD. 2016;2015.
- 53
- 54 25 22. Mori T, Okamoto S, Matsuoka S, et al. Risk-adapted pre-emptive therapy
- 55
- 56 26 for cytomegalovirus disease in patients undergoing allogeneic bone
- 57
- 58
- 59
- 60

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

- marrow transplantation. 2000;(August 1999):765-769.
23. Kanda Y, Mineishi S, Saito T, et al. Pre-emptive therapy against cytomegalovirus (CMV) disease guided by CMV antigenemia assay after allogeneic hematopoietic stem cell transplantation: A single-center experience in Japan. *Bone Marrow Transplant*. 2001;27(4):437-444. doi:10.1038/sj.bmt.1702805
24. Ljungman P, Griffiths P, Paya C. Definitions of Cytomegalovirus Infection and Disease in Transplant Recipients. 2002;34.
25. Casavant MJ. *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. Vol 288. 13th ed. (Laurence L. Brunton, PhD Bruce A. Chabner BCK, ed.); 2002. doi:10.1001/jama.288.16.2052
26. Fine JP, Gray RJ, Fine JP, Gray RJ. A Proportional Hazards Model for the Subdistribution of a Competing Risk Stable URL : <http://www.jstor.org/stable/2670170> All use subject to <http://about.jstor.org/terms> A Proportional Hazards Model for the Subdistribution of a Competing Risk. 1999;94(446):496-509.
27. Liu Y-C, Chien S-H, Fan N-W, et al. Incidence and risk factors of probable and proven invasive fungal infection in adult patients receiving allogeneic hematopoietic stem cell transplantation. *J Microbiol Immunol Infect*. 2015. doi:10.1016/j.jmii.2015.01.002
28. Miyakoshi S, Kusumi E, Matsumura T, et al. Invasive Fungal Infection Following Reduced-Intensity Cord Blood Transplantation for Adult Patients with Hematologic Diseases. *Biol Blood Marrow Transplant*. 2007;13(7):771-777. doi:10.1016/j.bbmt.2007.02.012
29. Mielcarek M, Storer BE, Boeckh M, et al. Initial therapy of acute graft-versus-host disease with low-dose prednisone does not compromise

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

- 1 patient outcomes. *Blood*. 2009. doi:10.1182/blood-2008-07-168401
- 2
- 3
- 4
- 5
- 6 1
- 7
- 8 2 30. Matsumura-Kimoto Y, Inamoto Y, Tajima K, et al. Association of
- 9
- 10 3
- 11 Cumulative Steroid Dose with Risk of Infection after Treatment for Severe
- 12 4
- 13 Acute Graft-versus-Host Disease. *Biol Blood Marrow Transplant*. 2016.
- 14 5
- 15 doi:10.1016/j.bbmt.2016.02.020
- 16 6 31. Maertens J, Marchetti O, Herbrecht R, et al. European guidelines for
- 17 7
- 18 antifungal management in leukemia and hematopoietic stem cell
- 19 8
- 20 transplant recipients: Summary of the ECIL 32009 update. *Bone Marrow*
- 21 9
- 22 *Transplant*. 2011;46(5):709-718. doi:10.1038/bmt.2010.175
- 23
- 24 10 32. George B, Kerridge IH, Gilroy N, et al. A risk score for early
- 25 11
- 26 cytomegalovirus reactivation after allogeneic stem cell transplantation
- 27 12
- 28 identifies low-, intermediate-, and high-risk groups: Reactivation risk is
- 29 13
- 30 increased by graft-versus-host disease only in the intermediate-risk group.
- 31 14
- 32 *Transpl Infect Dis*. 2012;14(2):141-148.
- 33 15
- 34 doi:10.1111/j.1399-3062.2011.00706.x
- 35
- 36 16 33. Osarogiagbon RU, Defor TE, Weisdorf MA, Erice A, Weisdorf DJ. CMV
- 37 17
- 38 antigenemia following bone marrow transplantation: Risk factors and
- 39 18
- 40 outcomes. *Biol Blood Marrow Transplant*. 2000;6(3):280-288.
- 41 19
- 42 doi:10.1016/S1083-8791(00)70010-3
- 43
- 44 20 34. Frère P, Baron F, Bonnet C, et al. Infections after allogeneic hematopoietic
- 45 21
- 46 stem cell transplantation with a nonmyeloablative conditioning regimen.
- 47 22
- 48 *Bone Marrow Transplant*. 2006. doi:10.1038/sj.bmt.1705255
- 49 23
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

Watanabe, et al. Impact of cumulative steroid on infections after allo-SCT

1
2
3
4
5
6 1 **Figure legends**

7 2 **Figure 1 Cumulative incidence of invasive fungal infection**

8 3 **Figure 2 Cumulative incidence of CMV antigenemia**

9 4 **Figure 3 Cumulative incidence of bacteremia**

10
11
12
13
14 5
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table1 Patient characteristics

Group by cumulative steroid dose within 30 days			No administration (n=174)			Low (<=7mg/kg PSL) (n=22)		High (>7mg/kg PSL) (n=30)		Variance
			value			value		value		
			n*1	n	%*2	n	%	n	%	P-Value
Age*3 median(range)				51 (17-66)		47 (21-66)		48 (20-66)		0.651
Gender	Male	126	103	59.2		9	40.9	14	46.7	0.144
	Female	100	71	40.8		13	59.1	16	53.3	
Donor source	Sibling	61	47	27.0		7	31.8	7	23.3	0.930
	Unrelated BM	106	81	46.6		9	40.9	16	53.3	
	Unrelated CB	59	46	26.4		6	27.3	7	23.3	
Disease	AML/MDS	134	113	64.9		11	50.0	10	33.3	0.015
	ALL/other leukemias	50	30	17.2		8	36.4	12	40.0	
	Malignant lymphoma	35	25	14.4		3	13.6	7	23.3	
	Aplastic anemia	7	6	3.4		0	0.0	1	3.3	
Disease status	CR	94	72	41.4		11	50.0	11	36.7	0.652
	non CR	132	102	58.6		11	50.0	19	63.3	
Conditioning intensity	Myeloablative	112	86	49.4		11	50.0	15	50.0	1.000
	Reduced intensity	114	88	50.6		11	50.0	15	50.0	
Neutrophil engraftment at Day30	No	20	18	10.5		1	4.5	1	3.4	0.477
	Yes	203	154	89.5		21	95.5	28	96.6	

Transplant Infectious Disease

1	levofloxacin	NO	181	142	84.0	14	63.6	25	89.3	0.079
2	prophylaxis	Yes	38	27	16.0	8	36.4	3	10.7	
3										
4										
5										
6										
7		CI	19	12	6.9	2	9.1	5	16.7	
8		CI+MMF	35	27	15.5	4	18.2	4	13.3	
9	GVHD prophylaxis	CI+MTX	137	107	61.5	13	59.1	17	56.7	0.755
10		CI+MMF+MTX	35	28	16.1	3	13.6	4	13.3	
11										
12										
13										
14		I	38	30	34.5	6	30.0	2	7.4	
15	GVHD grade at	II	76	47	54.0	11	55.0	18	66.7	
16	onset	III	14	6	6.9	2	10.0	6	22.2	0.092
17		IV	6	4	4.6	1	5.0	1	3.7	
18										
19										
20	CMV resopositivity	Donor+/ Recipient+	86	63	41.2	9	45.0	14	53.8	
21		Donor+/ Recipient-	10	9	5.9	0	0.0	1	3.8	
22		Donor-/ Recipient+	85	64	41.8	10	50.0	11	42.3	0.527
23		Donor-/ Recipient-	18	17	11.1	1	5.0	0	0.0	
24										
25		Acute GVHD	32			13	59.1	19	63.3	
26	Reason for steroid	Engraftment syndrome	9			3	13.6	6	20.0	
27		Others	11			6	27.3	5	16.7	

*1n indicates the number of patients with each characteristics

*2% indicates the percentage of patients in each steroid group

*3Age indicates patients' age at transplantation

Calcinerin inhibitors include Tacrolimus and Cyclosporin

Abbreviation: AML, acute myeloid leukemia; MDS, myelodysplastic syndrome; ALL, acute lymphoblastic leukemia; CR, complete remission; BM, bone marrow; CB, cord blood; GVHD, graft-versus-host disease; CI, Calcinerin inhibitor; MMF, mycophenolate mofetil; MTX, methotrexate; PSL, prednisolone

Table 2 Univariate and multivariate analysis of CMV antigenemia

Variables		Univariate analysis			Multivariate analysis		
		HR	95% CI	P-Value	HR	95% CI	P-Value
Age*1	<50	1.00		reference	1.00		reference
	≥50	1.46	(1.01-2.09)	0.042	1.62	(1.14-2.30)	0.007
Gender	Male	1.00		reference			
	Female	0.88	(0.60-1.28)	0.499			
Year of trasnplant	2005-2009	1.00		reference			
	2010-2015	1.19	(0.81-1.74)	0.373			
Donor source	Sibling	1.00		reference	1.00		reference
	Unrelated BM	1.10	(0.68-1.78)	0.687			
	Unrelated CB	1.64	(1.00-2.68)	0.047	1.62	(1.09-2.40)	0.018
Disease	AML/MDS	1.00		reference			
	ALL/other leukemias	1.43	(0.88-2.31)	0.140			
	Malignant lymphoma	0.93	(0.50-1.75)	0.824			
	Aplastic anemia	1.46	(0.54-3.93)	0.453			
Disease status	CR	1.00		reference			
	non CR	1.18	(0.80-1.75)	0.394			

Conditioning regimen	Myeloablative	1.00		reference		
	Reduced intensity	1.19	(0.81-1.74)	0.369		
GVHD prophylaxis	CI	1.00		reference		
	CI+MMF	1.00	(0.50-1.99)	0.994		
	CI+MTX	0.65	(0.35-1.21)	0.178		
	CI+MMF+MTX	1.10	(0.54-2.22)	0.792		
Neutrophil engraftment at day30	NO	1.00				
	Yes	1.46	(0-86-2.50)	0.164		
steroid group	No administration	1.00		reference	1.00	reference
	Low-dose* ²	1.58	(0.87-2.87)	0.140	1.64 (0.91-2.96)	0.100
	High-dose* ³	1.78	(1.02-3.12)	0.044	1.91 (1.04-3.50)	0.037

*¹Age indicates patients' age at transplantation

*²Low-dose indicates group who undertook low cumulative dose of steroid (≤ 7 mg/kg of prednisolone)

*³High-dose indicates group who undertook high cumulative dose of steroid (> 7 mg/kg of prednisolone)

Calcinerin inhibitors include Tacrolimus and Cyclosporin

Abbreviation: HR, hazard ratio; AML, acute myeloid leukemia; MDS, myelodysplastic syndrome; ALL, acute

lymphoblastic leukemia; CR, complete remission; BM, bone marrow; CB, cord blood; GVHD, graft-versus-host disease;

CI, Calcinerin inhibitor; MMF, mycophenolate mofetil; MTX, methotrexate; PSL, prednisolone

Table 3 Univariate and multivariate analysis of bacteremia

Variables		Univariate analysis			Multivariate analysis		
		HR	95% CI	P-Value	HR	95% CI	P-Value
Age*1	<50	1.00		reference	1.00		reference
	≥50	2.40	(1.07-5.38)	0.034	2.69	(1.16-6.22)	0.021
Gender	Male	1.00		reference			
	Female	1.20	(0.52-2.78)	0.671			
Year of trasnplant	2005-2009	1.00		reference			
	2010-2015	1.11	(0.48-2.53)	0.813			
Donor source	Sibling	1.00		reference			
	Unrelated BM	1.28	(0.46-3.55)	0.633			
	Unrelated CB	1.18	(0.37-3.72)	0.778			
Disease	AML/MDS	1.00		reference			
	ALL/other leukemias	1.06	(0.36-2.77)	0.917			
	Malignant lymphoma	1.52	(0.52-3.96)	0.419			
	Aplastic anemia						
Disease status	CR	1.00		reference			
	non CR	1.63	(0.66-3.98)	0.287			
Conditioning regimen	Myeloablative	1.00		reference			
	Reduced intensity	1.70	(0.72-4.03)	0.226			

		CI	1.00		reference		
		CI+MMF	1.61	(0.31-9.13)	0.568		
		CI+MTX	0.72	(0.15-3.37)	0.674		
		CI+MMF+MTX	1.16	(0.24-6.58)	0.864		
	Neutrophil	NO	1.00				
	engraftment at day30	Yes	3.86	(0.23-64.05)	0.346		
	Levofloxacin	NO	1.00				
	prophylaxis	Yes	0.73	(0.250-2.159)	0.574		
		No administration	1.00		reference	1.00	reference
	steroid group	Low-dose ^{*2}	1.74	(0.50-6.07)	0.390	2.13	(0.60-7.51) 0.240
		High-dose ^{*3}	2.27	(0.87-5.93)	0.097	2.59	(0.99-6.78) 0.053

*1Age indicates patients' age at transplantation

*2Low-dose indicates group who undertook low cumulative dose of steroid (<=7mg/kg of prednisolone)

*3High-dose indicates group who undertook high cumulative dose of steroid (>7mg/kg of prednisolone)

Calcinerin inhibitors include Tacrolimus and Cyclosporin

Abbreviation: HR, hazard ratio; AML, acute myeloid leukemia; MDS, myelodysplastic syndrome; ALL, acute lymphoblastic leukemia; CR, complete remission; BM, bone marrow; CB, cord blood; GVHD, graft-versus-host disease; CI, Calcinerin inhibitor; MMF, mycophenolate mofetil; MTX, methotrexate; PSL, prednisolone

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Figure 1. Cumulative incidence of invasive fungal infection

Figure 1

338x190mm (96 x 96 DPI)

Figure 2. Cumulative incidence of CMV antigenemia

Figure 2

338x190mm (96 x 96 DPI)

Figure 3. Cumulative incidence of bacteremia

Figure 3

338x190mm (96 x 96 DPI)