

# The duality of abnormal extremals on subriemannian Cartan structures

By

YUMIKO KITAGAWA\*

## Abstract

A subriemannian manifold  $(M, D, g)$  is a differential manifold  $M$  equipped with a sub-bundle  $D$  of the tangent bundle  $TM$  of  $M$ .

We emphasize abnormal extremals (abnormal geodesics) in subriemannian geometry and in particular show results of abnormal extremals on Cartan distributions of growth  $(2, 3, 5)$ . The subriemannian geometry is very important in differential geometry and it is closely related to the control theory.

## § 1. Introduction

This paper is a survey paper on subriemannian geometry and author's thesis [10] with related topics, based on the talk by the author at RIMS Symposium “Theory of singularities of smooth mappings and around it” (Nov.25-29, 2013), organized by Takashi Nishimura. We emphasize abnormal extremals (abnormal geodesics) in subriemannian geometry and in particular show results of abnormal extremals on Cartan distributions. The subriemannian geometry is very important in differential geometry and it is closely related to the control theory.

First of all we give a survey on the problem of length-minimizing paths and explain the abnormal extremals (abnormal geodesics) that does not appear in riemannian geometry but does in subriemannian geometry.

A subriemannian structure on a manifold  $M$  is a pair  $(D, g)$  such that  $D$  is a smooth distribution on  $M$  and  $g$  is a riemannian metric on  $D$ . A subriemannian manifold is a triple  $(M, D, g)$  such that  $M$  is a manifold and  $(D, g)$  is a subriemannian structure

---

Received June 30, 2014. Revised September 19, 2014.

2010 Mathematics Subject Classification(s): Primary 58A15; Secondary 58A17, 58A20, 58A30

*Key Words*: subriemannian geometry, control system, cartan theory.

\*National Institute of Technology, Oita College, Oita 870-0152, Japan.

e-mail: [kitagawa@oita-ct.ac.jp](mailto:kitagawa@oita-ct.ac.jp)

on  $M$ . In particular, if  $D = TM$  then  $(M, D, g)$  is nothing but a riemannian manifold  $(M, g)$ .

Riemannian geometry tells us that a minimizer (i.e., a shortest path) between two points of a riemannian manifold  $(M, g)$  is a geodesic, provided that the curve is parametrized by arc-length, and the geodesics are characterized to be the curves satisfying the geodesic equation expressed in local coordinates as:

$$\ddot{x}^i + \sum \Gamma_{jk}^i \dot{x}^j \dot{x}^k = 0,$$

where  $\Gamma_{jk}^i$  denotes the Christoffel symbol. Conversely, every geodesic is locally length minimizing. In the formulation of symplectic geometry, the geodesics  $x(t)$  are the projections to the base manifold  $M$  of the integral curves  $(x(t), p(t))$  of the Hamiltonian vector field  $\vec{E}$  defined on the cotangent bundle  $T^*M$ , where  $E$  is the energy function associated to the metric  $g$ .

Now in subriemannian geometry, it is also of fundamental importance to study minimizers between two points of a subriemannian manifold  $(M, D, g)$ . Since the metric  $g$  is defined only on the subbundle  $D$  of  $TM$  in this subriemannian case, there is no canonical means to define the length of a general curve  $\gamma : [a, b] \rightarrow M$ . But we can well speak of the length of  $\gamma$  if  $\gamma$  is an integral curve of  $D$ , that is, if  $\dot{\gamma}(t) \in D_{\gamma(t)}$  for all  $t$ .

On the other hand Chow's theorem tells that if  $M$  is connected and if  $D$  is non-holonomic (in other word, bracket-generating), then any two points of  $M$  can be joined by a piecewise smooth integral curve of  $D$  (see Theorem 2.4).

Hence, especially for a nonholonomic subriemannian manifold  $(M, D, g)$ , it makes sense and is important to study the minimizers (length minimizing piecewise smooth integral curves) between two points of the subriemannian manifold  $(M, D, g)$ . However, contrary to the riemannian case, this problem is very subtle, mainly because the space  $C_D(p, q)$  of all integral curves of  $D$  joining  $p$  and  $q$  may have singularities, while the space  $C(p, q)$  of all curves joining  $p$  and  $q$  has no singularity and is a smooth infinite dimensional manifold, which makes difficult to apply directly the method of variation to the subriemannian case.

For a subriemannian manifold  $(M, D, g)$  we define a *normal biextremal* to be an integral curve of the Hamiltonian vector field  $\vec{E}$  associated to the Hamiltonian function  $E : T^*M \rightarrow \mathbf{R}$ , where  $E$  is the energy function associated with the subriemannian metric  $g$ . We then define a *normal extremal* to be the projection to  $M$  of a normal biextremal(Definition 5.1). Then, as in riemannian geometry, a normal extremal is locally a minimizer(Theorem 5.2)

However, R. Montgomery ([13], [14]) and I. Kupka [11] discovered that there exists a minimizer which is not a normal extremal, and hence called it abnormal. The appearance of abnormal minimizers is a surprising phenomenon never arising in riemannian geometry but peculiar to subriemannian geometry.

If  $D$  is a distribution on  $M$ , then the annihilator bundle  $D^\perp$ , considered as a submanifold of the symplectic manifold  $T^*M$ , carries a (singular) characteristic distribution  $\text{Ch}(D^\perp)$ . An integral curve of this characteristic system  $\text{Ch}(D^\perp)$  contained in  $D^\perp \setminus \{\text{zero section}\}$  is called an *abnormal biextremal*, of which the projection to  $M$  is called an *abnormal extremal* (Definition 7.1).

A rigorous application of the Pontryagin Maximum Principle of Optimal Control Theory to subriemannian geometry shows that a minimizer of subriemannian manifold  $(M, D, g)$  is either a normal extremal of  $(D, g)$  or an abnormal extremal of  $D$ .

This settled the long discussions that had been made until 1990's by many mathematicians with erroneous statements, and gave a right way to treat the problem of length-minimizing paths in subriemannian geometry. The difference between riemannian geometry and subriemannian geometry lies in particular on the existence of abnormal geodesics. It is known that abnormal geodesics are characterized as singular curves of so-called endpoint mapping in control theory (see Montgomery[14] and §9 of this paper). We will give a survey on the problem of length-minimizing paths mainly following Liu and Sussmann [12]. We then consider this problem in a concrete case of the standard Cartan distribution. Referring to [16], we will carry out detailed computation of extremals, which will well illustrate how normal and abnormal extremals appear in subriemannian geometry.

Let  $D \subset TY$  be a distribution on a five dimensional manifold  $Y$  with growth  $(2, 3, 5)$ . Distributions with growth  $(2, 3, 5)$  were studied by Cartan in the famous paper [6]. In fact they are called systems of *Cartan type* in [5]. In general, the space of singular paths (or abnormal extremals) of a given distribution plays an important role as it is an intrinsic invariant of the distribution. For spaces with  $(2, 3, 5)$ -distributions, the construction of the space of singular paths and the existence of natural double fibration was mentioned for the first time in unpublished lecture notes by Bryant [4].

It is known that for any point  $y$  of  $Y$  and for any direction of  $D_y$ , there exists uniquely a singular  $D$ -path (or an immersed abnormal extremal) through  $y$  with the given direction. Then the immersive singular  $D$ -paths form another five dimensional manifold  $X$ . Moreover we remark that  $X$  is endowed with a natural cone structure  $C \subset TX$  induced from the Cartan prolongation  $E$  of  $D$  on a six dimensional manifold  $Z$  and a projection  $Z \rightarrow X$ .

It seems to be natural to consider what are "abnormal geodesics" for the cone structure and whether the original space  $Y$  can be regarded as the moduli space of the "abnormal geodesics" of  $X$ , after a proper formulation. Because the cone structure  $C \subset TX$  is not a distribution, so it is impossible to apply the non-holonomic geometry or subriemannian geometry directly to our situation. However, we can regard the cone structure as a control system and therefore we can consider the endpoint mapping and

singular curves of cone structure. Thus we establish the duality on abnormal geodesics or singular paths on Cartan distribution.

Then we are naturally led to consider the corresponding objects for the cone structure  $C \subset TX$  to singular paths for  $D \subset TY$  and to obtain the “inverse” construction from the space  $X$  to the original space  $Y$ . To realise this, we regard the cone structure  $C \subset TX$  as a control system  $\mathbb{C} : E \rightarrow TX \rightarrow X$  on  $X$  defined by the differential map  $E \hookrightarrow TZ \rightarrow TX$  of the projection  $Z \rightarrow X$  and consider the space of singular  $\mathbb{C}$ -paths. Then we show that the original space  $Y$  is identified with the space of singular  $\mathbb{C}$ -paths on  $X$ , at least locally, while  $X$  is identified with the space of singular  $D$ -paths (See Theorem 10.1). In fact, a singular  $\mathbb{C}$ -path on  $X$  consists of singular  $D$ -paths on  $Y$  which pass through a fixed point of  $Y$ .

In the author’s thesis [10], we studied homogeneous subriemannian structures, in particular, we classified homogeneous contact subriemannian structures. In our paper [8], we have studied cone structures which arise naturally from Cartan distributions, in terms of control systems. The usage of the notion of control systems enables us clarify the duality of abnormal geodesics on subriemannian Cartan structures. Since control systems are generalizations of distributions, it is very natural to study homogeneous control systems and subriemannian structures on them, which will be treated in forthcoming papers.

## § 2. Nonholonomic distributions

Let  $M$  be a differentiable manifold. A subbundle  $D$  of its tangent bundle  $TM$  of  $M$  of rank  $r$  is alternatively called a distribution on  $M$  of dimension  $r$ , since it gives a law which assigns to every point  $p \in M$  an  $r$ -dimensional subspace  $D_p$  of the tangent space  $T_pM$ . A section of  $D$  on an open set  $U \subset M$  is a local vector field  $X$  defined on  $U$  such that  $X_p \in D_p$  for all  $p \in U$ . A local basis of  $D$  on  $U$  is a system of sections  $X_1, \dots, X_r$  of  $D$  defined on  $U$  such that  $\{(X_1)_p, \dots, (X_r)_p\}$  forms a basis of  $D_p$  for all  $p \in U$ . It is clear that for any point  $p_0 \in M$  there is an local basis of  $D$  defined on a neighborhood of  $p_0$ . If  $\{X_1, \dots, X_r\}$  is a local basis of  $D$  on  $U$ , then any section  $X$  of  $D$  on  $U$  is uniquely written:

$$X = f_1X_1 + \cdots + f_rX_r$$

with some functions  $f_1, \dots, f_r$  on  $U$ , and we say that  $D$  is locally generated, or defined, by  $X_1, \dots, X_r$ .

Let  $D^\perp$  denote the annihilators of  $D$ , that is,  $D^\perp = \bigcup_{p \in M} D_p^\perp$  with

$$D_p^\perp = \{\alpha \in T_p^*M; \langle \alpha, v \rangle = 0 \text{ for all } v \in D_p\}.$$

Clearly  $D^\perp$  is a subbundle of the cotangent bundle  $T^*M$  of rank  $s$ , where  $s = \dim M - r$ . If  $\{\omega^1, \dots, \omega^s\}$  is a local basis of  $D^\perp$ , we say that  $D$  is locally defined by the Pfaff system  $\{\omega^1, \dots, \omega^r\}$  or by the Pfaff equations:

$$\omega^1 = \dots = \omega^s = 0.$$

In this sense, a distribution is also called a differential system or a Pfaff system.

Given an  $r$ -dimensional distribution  $D$  on  $M$ , one of the most important problems that has been studied since the nineteenth century is to study integral manifolds of  $D$ . An immersed submanifold  $f : S \rightarrow M$  is called an integral manifold of  $D$  if

$$f_*T_s S \subset D_{f(s)} \quad \text{for all } s \in S.$$

Evidently the dimension of an integral manifold is  $\leq r$ . However, it is not always the case that there exists an  $r$ -dimensional integral manifold.

**Definition 2.1.** A distribution  $D$  of dimension  $r$  on  $M$  is called completely integrable if about every point  $p_0 \in M$  there is a coordinate system  $(U, (x^1, \dots, x^n))$  such that all the submanifolds of  $U$  given by  $x^{r+1} = \text{const}, x^{r+2} = \text{const}, \dots, x^n = \text{const}$  are integral manifolds of  $D$ .

As is well-known, the Frobenius theorem gives a criterion for  $D$  to be completely integrable:

**Theorem 2.2** (Frobenius). *A distribution  $D$  on  $M$  is completely integrable if and only if  $D$  is involutive, that is,  $D$  satisfies the condition: "For any open set  $U \subset M$ , the Lie bracket  $[X, Y]$  of sections  $X, Y$  of  $D$  on  $U$  is also a section of  $D$ ." Moreover, if  $D$  is completely integrable then the manifold  $M$  is a disjoint union  $\bigcup_\lambda L_\lambda$  of the maximal connected  $r$ -dimensional integral manifolds  $L_\lambda$  of  $D$ , each  $L_\lambda$  being called a leaf of  $D$ .*

The problem of finding integral manifolds of distributions which are not completely integrable are treated by Cartan-Kähler theory.

Now let us proceed to consider integral curves of  $D$ . In order to well analyze the length functional we had better expand the class of curves to consider to that of the absolutely continuous curves: A continuous curve  $\gamma : I \rightarrow M$ ,  $I$  being an interval  $[a, b]$  of  $\mathbf{R}$ , is absolutely continuous if it has a derivative for almost all  $t$ , and if in any coordinate system the components of this derivative are measurable functions. We then define an integrable curve of  $D$  to be an absolutely continuous curve  $\gamma : I \rightarrow M$  such that  $\dot{\gamma} \in D_{\gamma(t)}$  for almost all  $t \in I$ . An integral curve of  $D$  is also called an integral path, a  $D$ -arc, or a horizontal curve.

If  $\{X_1, \dots, X_r\}$  is a local basis of  $D$  defined on an open set  $U \subset M$ , then a curve  $\gamma : I \rightarrow U$  is an integral curve of  $D$  if

$$(*) \quad \dot{\gamma}(t) = c_1(t)(X_1)_{\gamma(t)} + \cdots + c_r(t)(X_r)_{\gamma(t)}$$

for some functions  $c_1(t), \dots, c_r(t)$ . Conversely if the function  $c_1(t), \dots, c_r(t)$  and  $\gamma(t_0)$  at some  $t_0 \in I$  are assigned then the curve  $\gamma(t)$  is determined by the ordinary differential equation (\*). In control theory  $c_1, \dots, c_r$  are interpreted as control parameters and  $D$  (or  $X_1, \dots, X_r$ ) is regarded as a control system.

If two points  $p, q \in M$  can be joined by an integral curve of  $D$ , we say that  $q$  is reachable from  $p$ . If  $D$  is completely integrable then the set of all points reachable from  $p$  is the leaf passing through  $p$ .

Let us now introduce a class of distributions which are in a sense at the opposite end from the completely integrable distributions.

**Definition 2.3.** A distribution  $D$  on  $M$  is called nonholonomic or bracket-generating if for any local basis  $X_1, \dots, X_r$  of  $D$  on  $U$  the collection of all vector fields  $\{X_i, [X_i, X_j], [X_i, [X_j, X_k]], \dots\}$  generated by Lie brackets of the  $X_i$  spans the whole tangent bundle  $TU$ .

This definition can be rephrased as follows: Let  $\underline{D}$  denote the sheaf of germs of section of  $D$ . Define the sheaves  $\{\mathcal{D}^k\}_{k \geq 1}$  inductively by setting first  $\mathcal{D}^1 = \underline{D}$  and then

$$\mathcal{D}^{k+1} = \mathcal{D}^k + [\mathcal{D}^1, \mathcal{D}^k] \quad (k \geq 1).$$

Then  $D$  is completely integrable if  $\mathcal{D}^1 = \mathcal{D}^2$ , and nonholonomic if  $\bigcup \mathcal{D}^k = \underline{TM}$ .

The following theorem of Chow [7] is fundamental.

**Theorem 2.4 (Chow).** *Let  $M$  be a connected manifold and  $D$  a nonholonomic distribution on  $M$ , then there exists for any two points  $p, q \in M$  a piecewise smooth integral curve by which  $p$  and  $q$  can be joined.*

A detailed proof can be also found in [17], or in [15].

### § 3. Subriemannian distance

If  $(M, D, g)$  is a subriemannian manifold, and  $p \in M$ ,  $v \in D_p$ , we define the *length*  $\|v\|_g$  of  $v$  by

$$\|v\|_g = g_p(v, v)^{\frac{1}{2}}$$

If  $\gamma : [a, b] \rightarrow M$  is an integral curve of  $D$ , then we define the *length* of  $\gamma$  by

$$\|\gamma\|_g = \int_a^b \|\dot{\gamma}(t)\|_g dt.$$

If  $\gamma$  is not an integral curve, we agree to define  $\|\gamma\|_g = +\infty$ . We then define a function  $d_g : M \times M \rightarrow \mathbf{R} \cup \{\infty\}$  by

$$d_g(p, q) = \inf\{\|\gamma\|_g; \partial\gamma = (p, q)\},$$

where we denote  $\partial\gamma = (\gamma(a), \gamma(b))$ .

If  $M$  is connected and  $D$  is bracket-generating, then  $d_g : M \times M \rightarrow \mathbf{R}$  is a metric function on  $M$  and the topology on  $M$  that the metric determines coincides with the original manifold topology of  $M$ . The first assertion follows from Chow's theorem and the second assertion follows from the Ball-Box Theorem ([15], See p.29 ). The distance  $d_g : M \times M \rightarrow \mathbf{R}$  is called *subriemannian distance* or Carnot-Caratheodory metric.

If an integral curve  $\gamma : [a, b] \rightarrow M$  of  $D$  satisfies

$$d_g(\gamma(a), \gamma(b)) = \|\gamma\|_g,$$

$\gamma$  is called a *minimizer*. Concerning minimizers, here we cite the following two theorems ([15], p.10 ):

**Theorem 3.1** (Local existence). *If  $D$  is a nonholonomic distribution on a manifold  $M$ , then any point  $p$  of  $M$  is contained in a neighbourhood  $U$  such that every  $q$  in  $U$  can be connected to  $p$  by a minimizer.*

**Theorem 3.2** (Global existence). *Let  $M$  be a connected manifold and  $D$  a non-holonomic smooth distribution on  $M$ , and suppose that  $M$  is complete relative to the subriemannian distance function. Then any two points of  $M$  can be joined by a minimizer.*

#### § 4. Hamiltonian formalism

If  $M$  is a manifold and  $k \in \{0, 1, \dots, \infty\}$ , we use  $C^k(M)$  to denote the set of all real-valued functions on  $M$  that are class  $C^k$ , and  $V^k(M)$  to denote the set of all vector fields of class  $C^k$  on  $M$ .

If  $N$  is a symplectic manifold with symplectic 2-form  $\Omega$ , and  $H \in C^1(N)$ , we use  $\vec{H}$  to denote the *Hamiltonian vector field* associated to  $H$ .  $\vec{H}$  is the vector field  $V$  on  $N$  such that  $\Omega(X, V) = \langle dH, X \rangle$  for every vector field  $X$  on  $N$ . If  $H \in C^k(N)$  and  $k \geq 1$ , then vector field  $\vec{H}$  is of class  $C^{k-1}$ . If  $H, K \in C^1(N)$ , then the *Poisson bracket*  $\{H, K\}$  is the directional derivative of  $K$  in the direction of  $\vec{H}$ , i.e.,

$$\{H, K\} = \langle dK, \vec{H} \rangle = \Omega(\vec{H}, \vec{K}).$$

Then we have the following formulas

$$\{H, KL\} = \{H, K\}L + \{H, L\}K,$$

$$\{H, \{K, L\}\} + \{K, \{L, H\}\} + \{L, \{H, K\}\} = 0,$$

and

$$\overrightarrow{HK} = \overrightarrow{H}K + K\overrightarrow{H}.$$

Note also the fact that the map  $H \rightarrow \overrightarrow{H}$  is a Lie algebra homomorphism from  $(C^\infty(N), \{, \})$  to  $(V^\infty(N), [, ])$ .

The cotangent bundle  $T^*M$  of a manifold  $M$  has a natural symplectic structure determined by the 2-form  $\Omega_M = d\omega_M$ , where  $\omega_M$  is the Liouville form given by

$$\omega_M(x, \lambda)(v) = \langle \lambda, d\pi_M^*(v) \rangle \quad \text{for } v \in T_{(x, \lambda)}(T^*M),$$

$\pi_M^*$  being the projection  $T^*M \rightarrow M$ . Relative to a coordinate chart

$$T^*\kappa = (x^1, \dots, x^n, \lambda_1, \dots, \lambda_n)$$

induced by a chart  $\kappa = (x^1, \dots, x^n)$  on  $M$ , we have the formulas

$$\omega_M = \sum_j \lambda_j dx^j,$$

$$\Omega_M = \sum_j d\lambda_j \wedge dx^j,$$

$$\overrightarrow{H} = \sum_j \left( \frac{\partial H}{\partial \lambda_j} \frac{\partial}{\partial x^j} - \frac{\partial H}{\partial x^j} \frac{\partial}{\partial \lambda_j} \right),$$

$$\{H, K\} = \sum_j \left( \frac{\partial H}{\partial \lambda_j} \frac{\partial K}{\partial x^j} - \frac{\partial H}{\partial x^j} \frac{\partial K}{\partial \lambda_j} \right).$$

To each vector field  $X$  on  $M$  we associated the function  $H_X : T^*M \rightarrow \mathbf{R}$  given by

$$H_X(q, \lambda) = \langle \lambda, X(q) \rangle \quad \text{for } \lambda \in T_q^*M.$$

Then  $H_X$  is of class  $C^k$  if and only if  $X$  is. Moreover,

$$d\pi_M^*(\overrightarrow{H}_X(x, \lambda)) = X(x) \quad \text{for all } (x, \lambda) \in T^*M$$

The identity

$$\{H_X, H_Y\} = H_{[X, Y]}$$

holds for  $X, Y \in V^1(M)$ , and therefore the map  $X \rightarrow H_X$  is a Lie algebra homomorphism from  $(V^\infty(M), [, ])$  to  $(C^\infty(T^*M), \{, \})$ .

If  $X \in V^1(M)$  then the vector field  $\overrightarrow{H}_X$  is called the *Hamiltonian lift* of  $X$ .

### § 5. Normal extremals

Let  $(M, D, g)$  be a subriemannian manifold. If  $(p, \lambda) \in T^*M$ , then the restriction  $\lambda|_{D_p}$  of  $\lambda$  to the subspace  $D_p$  of  $T_pM$  has well-defined norm, since  $D_p$  is an inner product space. We will use  $\|\lambda\|_g$  to denote this norm. The function  $E : T^*M \rightarrow \mathbf{R}$  given by

$$E(x, \lambda) = -\frac{1}{2}\|\lambda\|_g^2$$

is the energy function of the subriemannian structure  $(D, g)$ .

**Definition 5.1.** A normal biextremal of a subriemannian structure  $(D, g)$  is a curve  $\Gamma : I \rightarrow T^*M$  such that

- (i)  $\Gamma$  is an integral curve of the Hamiltonian vector field  $\overrightarrow{E}$ , namely

$$\dot{\Gamma}(t) = \overrightarrow{E}_{\Gamma(t)}$$

- (ii)  $E$  does not vanish along  $\Gamma$ .

A normal extremal is a curve in  $M$  which is a projection of a normal biextremal.

**Theorem 5.2.** *Let  $(M, D, g)$  be a subriemannian manifold. Then every normal extremal is locally length minimizing.*

This theorem is non-trivial, but the proof is similar to that of riemannian case. However, contrary to the riemannian case, the converse of the theorem does not hold. There appeared several papers asserting that every minimizer of a subriemannian manifold is a normal extremal. But Kupka [11] and Montgomery [14] proved that there exists a subriemannian manifold and a minimizer of the subriemannian manifold which is not a normal extremal. Such a minimizer is called an abnormal minimizer. In the following sections we will give a characterization of the abnormal minimizers.

### § 6. Characteristic system

Let  $(N, \Omega)$  be a symplectic manifold. For a submanifold  $S$  of  $N$  we define the characteristic system (bundle)  $\text{Ch}(S)$  of by

$$\text{Ch}(S) = TS \cap (TS)^\perp,$$

that is, the fibre  $\text{Ch}(S)_s$  on  $s \in S$  is given by

$$\text{Ch}(S)_s = T_s S \cap (T_s S)^\perp,$$

where

$$(T_s S)^\perp = \{v \in T_s N; \Omega(v, u) = 0 \text{ for all } u \in T_s S\}.$$

Let  $F_1, \dots, F_r$  be local defining equations of  $S$ , say, defined on a neighbourhood  $U$  of  $s_0 \in S$  such that  $(dF_1)_s, \dots, (dF_r)_s$  are linearly independent for  $s \in U$  and

$$U \cap S = \{F_1 = \dots = F_r = 0\}.$$

From the very definition of Hamiltonian vector field we see immediately that

$$\{(\vec{F}_1)_s, \dots, (\vec{F}_r)_s\}$$

forms a basis of  $(T_s S)^\perp$  for  $s \in U$ . Hence we have

$$\text{Ch}(S)_s = T_s S \cap \langle (\vec{F}_1)_s, \dots, (\vec{F}_r)_s \rangle.$$

Let  $\Omega_S = \iota_S^* \Omega$ , where  $\iota_S : S \rightarrow N$  is the canonical inclusion, and let:

$$\text{Null}_s(\Omega_S) = \{v \in T_s S; \Omega_S(v, u) = 0 \text{ for all } u \in T_s S\}.$$

Then it is clear that

$$\text{Ch}(S)_s = \text{Null}_s(\Omega_S).$$

We then have:

**Proposition 6.1.** *For a submanifold  $S$  of a symplectic manifold  $(N, \Omega)$ , the characteristic system  $\text{Ch}(S) = \bigcap_{s \in S} \text{Ch}(S)_s \subset TS$  is given by:*

$$\begin{aligned} \text{Ch}(S)_s &= T_s S \cap (T_s S)^\perp \\ &= (T_s S) \cap \langle (\vec{F}_1)_s, \dots, (\vec{F}_r)_s \rangle \\ &= \text{Null}_s(\Omega_S) \end{aligned}$$

*If  $\dim \text{Ch}(S)_s$  is constant, then  $\text{Ch}(S)$  is a completely integrable subbundle of  $TS$ .*

The last assertion of the proposition follows from the exactness of the symplectic form.

## § 7. Abnormal extremals

Let  $(M, D, g)$  be a subriemannian manifold. We denote by  $D^\perp$  the annihilator bundle of  $D$  and by  $\text{Ch}(D^\perp)$  its characteristic system.

**Definition 7.1.** An abnormal biextremal of  $(M, D, g)$  is an curve  $\Gamma : I \rightarrow D^\perp \setminus \{O\}$  ( $O$  denoting the zero section) such that  $\dot{\Gamma}(t) \in \text{Ch}(D^\perp)_{\Gamma(t)}$  for almost all  $t \in I$ . An abnormal extremal of  $(M, D, g)$  is a curve in  $M$  which is a projection of an abnormal biextremal.

It should be remarked that the above definition does not depend on the metric  $g$  but depends only on  $(M, D)$ .

If  $\{X_1, \dots, X_r\}$  is a local basis of  $D$  defined on  $U \subset M$ , then  $H_{X_1}, \dots, H_{X_r}$  give defining equations of  $D^\perp$  on  $(\pi_M^*)^{-1}U$ . Hence by Proposition 6.1, we have

$$\text{Ch}(D^\perp)_z = T_z D^\perp \cap \langle (\overrightarrow{H_{X_1}})_z, \dots, (\overrightarrow{H_{X_r}})_z \rangle.$$

Therefore a curve  $\Gamma : I \rightarrow (\pi_M^*)^{-1}U \setminus \{O\}$  is an abnormal biextremal of  $(M, D)$  if and only if

$$\begin{cases} \text{(i) } H_{X_i}(\Gamma(t)) = 0 & \text{for all } t \in I \text{ and } i = 1, \dots, r \\ \text{(ii) } \dot{\Gamma}(t) \in \langle (\overrightarrow{H_{X_1}})_{\Gamma(t)}, \dots, (\overrightarrow{H_{X_r}})_{\Gamma(t)} \rangle & \text{for almost all } t \in I \end{cases}$$

By using the Pontryagin Maximam Principle on Control system, it is shown that the following theorem holds (see [12], p.81, Appendix B).

**Theorem 7.2.** *Let  $(M, D, g)$  be a subriemannian manifold, and let  $\gamma : [a, b] \rightarrow M$  be a length-minimizer parametrized by arc-length. Then  $\gamma$  is a normal extremal or an abnormal extremal.*

### § 8. Extremals on the standard Cartan distribution

As was shown by Cartan [6], a generic Pfaff system defined by three Pfaff equations in the space of five variables, that is, a tangent distribution  $D$  of rank 2 on  $\mathbf{R}^5$  enjoys interesting properties: Its automorphism group makes a Lie group of dimension not greater than 14, and if the maximal dimension is attained, then the automorphism group is locally isomorphic to the exceptional simple Lie group  $G_2$  and the tangent distribution  $D$  is locally isomorphic to the standard Cartan distribution defined as follows: Let  $(x^1, x^2, x^3, x^4, x^5)$  be the standard coordinates of  $\mathbf{R}^5$  and let the vector fields  $X_1, \dots, X_5$  be given by:

$$\begin{aligned} X_1 &= \frac{\partial}{\partial x^1} - \frac{1}{2}x^2 \frac{\partial}{\partial x^3} - (x^3 - \frac{1}{2}x^1x^2) \frac{\partial}{\partial x^4} \\ X_2 &= \frac{\partial}{\partial x^2} + \frac{1}{2}x^1 \frac{\partial}{\partial x^3} - (x^3 + \frac{1}{2}x^1x^2) \frac{\partial}{\partial x^5} \\ X_3 &= \frac{\partial}{\partial x^3}, \quad X_4 = \frac{\partial}{\partial x^4}, \quad X_5 = \frac{\partial}{\partial x^5}. \end{aligned}$$

These vector fields satisfy the following bracket relations:

$$\begin{cases} [X_1, X_2] = X_3 \\ [X_1, X_3] = X_4 \\ [X_2, X_3] = X_5 \\ \text{The others are trivial} \end{cases}$$

The dual basis  $\omega^1, \dots, \omega^5$  of  $X_1, \dots, X_5$  is given by:

$$\begin{cases} \omega^1 = dx^1 \\ \omega^2 = dx^2 \\ \omega^3 = dx^3 - \frac{1}{2}(x^1 dx^2 - x^2 dx^1) \\ \omega^4 = dx^4 + (x^3 - \frac{1}{2}x^1 x^2) dx^1 \\ \omega^5 = dx^5 + (x^3 + \frac{1}{2}x^1 x^2) dx^2. \end{cases}$$

Then we have the following the structure equations:

$$\begin{cases} d\omega^1 = 0 \\ d\omega^2 = 0 \\ d\omega^3 + \omega^1 \wedge \omega^2 = 0 \\ d\omega^4 + \omega^1 \wedge \omega^3 = 0 \\ d\omega^5 + \omega^2 \wedge \omega^3 = 0. \end{cases}$$

Let us take  $D$  to be the tangent distribution spanned by  $X_1$  and  $X_2$ , that is,

$$\Gamma(D) = \langle X_1, X_2 \rangle = \{\omega^3 = \omega^4 = \omega^5 = 0\}.$$

Then, choosing a subriemannian metric  $g$  on  $D$  so that  $\{X_1(p), X_2(p)\}$  forms an orthonormal basis of  $D_p$ , we consider the subriemannian manifold  $(\mathbf{R}^5, D, g)$ .

Let us determine the normal extremals and the abnormal extremals of this subriemannian manifold.

If  $(x^1, x^2, x^3, x^4, x^5, p_1, p_2, p_3, p_4, p_5)$  are the local coordinates in  $T^*\mathbf{R}^5$ , the energy function  $E$  of  $(D, g)$  is given by

$$\begin{aligned} E = & -\frac{1}{2}[\{p_1 - \frac{1}{2}x^2 p_3 - (x^3 - \frac{1}{2}x^1 x^2)p_4\}^2 \\ & + \{p_2 + \frac{1}{2}x^1 p_3 - (x^3 + \frac{1}{2}x^1 x^2)p_5\}^2]. \end{aligned}$$

Then the Hamiltonian vector field  $\vec{E}$  is given by

$$\begin{aligned} \vec{E} = & -A \frac{\partial}{\partial x^1} - B \frac{\partial}{\partial x^2} + (\frac{1}{2}x^2 A - \frac{1}{2}x^1 B) \frac{\partial}{\partial x^3} + (x^3 - \frac{1}{2}x^1 x^2) A \frac{\partial}{\partial x^4} \\ & + (x^3 + \frac{1}{2}x^1 x^2) B \frac{\partial}{\partial x^5} + \{\frac{1}{2}x^2 p_4 A + (\frac{1}{2}p_3 - \frac{1}{2}x^2 p_5) B\} \frac{\partial}{\partial p_1} \\ & + \{(\frac{1}{2}x^1 p_4 - \frac{1}{2}p_3) A - \frac{1}{2}x^1 p_5 B\} \frac{\partial}{\partial p_2} + (-p_4 A - p_5 B) \frac{\partial}{\partial p_3}, \end{aligned}$$

where

$$(8.1) \quad A = p_1 - \frac{1}{2}x^2p_3 - (x^3 - \frac{1}{2}x^1x^2)p_4$$

$$(8.2) \quad B = p_2 + \frac{1}{2}x^1p_3 - (x^3 + \frac{1}{2}x^1x^2)p_5.$$

Then we see that a normal biextremal of  $(D, g)$  satisfies

$$(8.3) \quad \dot{x}^1 = -A$$

$$(8.4) \quad \dot{x}^2 = -B$$

$$(8.5) \quad \dot{x}^3 = \frac{1}{2}x^2A - \frac{1}{2}x^1B$$

$$(8.6) \quad \dot{x}^4 = (x^3 - \frac{1}{2}x^1x^2)A$$

$$(8.7) \quad \dot{x}^5 = (x^3 + \frac{1}{2}x^1x^2)B$$

$$(8.8) \quad \dot{p}_1 = \frac{1}{2}x^2p_4A + (\frac{1}{2}p_3 - \frac{1}{2}x^2p_5)B$$

$$(8.9) \quad \dot{p}_2 = (-\frac{1}{2}p_3 + \frac{1}{2}x^1p_4)A - \frac{1}{2}x^1p_5B$$

$$(8.10) \quad \dot{p}_3 = -p_4A - p_5B$$

$$(8.11) \quad \dot{p}_4 = 0$$

$$(8.12) \quad \dot{p}_5 = 0$$

Differentiating the equation (8.1), and substituting (8.3),(8.4),(8.5),(8.8),(8.10),(8.11) into it, we have

$$(8.13) \quad \ddot{x}^1 = p_3\dot{x}^2.$$

Similarly differentiating (8.2), and substituting (8.3),(8.4),(8.5),(8.9),(8.10),(8.12) into it, we have

$$(8.14) \quad \ddot{x}^2 = -p_3\dot{x}^1.$$

On the other hand,  $p_4, p_5$  are constant by (8.11), (8.12). Then integrating (8.10), we have:

$$(8.15) \quad p_3 = p_4x^1 + p_5x^2 + C,$$

where  $C$  is a constant. Therefore the second order differential equations with respect to  $x^1$  and  $x^2$  are given in the formulae (8.13), (8.14) and (8.15). These equations for  $(x^1, x^2)$  can be written in the following form:

$$\begin{pmatrix} \ddot{x}^1 \\ \ddot{x}^2 \end{pmatrix} = p_3 \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} \dot{x}^1 \\ \dot{x}^2 \end{pmatrix},$$

where  $p_3$  is a linear function given by (8.15). Since the acceleration vector  $\begin{pmatrix} \ddot{x}^1 \\ \ddot{x}^2 \end{pmatrix}$  is

obtained by the rotation of  $\frac{\pi}{2}$  of the velocity vector  $\begin{pmatrix} \dot{x}^1 \\ \dot{x}^2 \end{pmatrix}$  with the scalar multiplication of  $p_3$ , this equation represents the equation of motion of an electron moving in a plane under a magnetic field whose direction is perpendicular to the plane and whose magnitude is given by the linear function  $p_3 = p_4x^1 + p_5x^2 + C$ . By a change of local coordinates:

$$\begin{cases} x = p_4x^1 + p_5x^2 + C \\ y = p_5x^1 - p_4x^2, \end{cases}$$

we have

$$\begin{cases} \ddot{x} = -x\dot{y} \\ \ddot{y} = x\dot{x}. \end{cases}$$

Then we also have

$$\dot{y} = \frac{1}{2}x^2 + k,$$

where  $k$  is a constant. By substituting this equation into  $\ddot{x} = -x\dot{y}$ , we have

$$\ddot{x} = -\frac{1}{2}x^3 - kx.$$

So we have

$$\frac{1}{2}\{\dot{x}\}^2 = -\frac{1}{8}x^4 - \frac{1}{2}kx^2,$$

and

$$\dot{x} = \pm\sqrt{-\frac{1}{4}x^4 - kx^2}.$$

Since

$$-\frac{1}{4}x^2(x^2 + 4k) \geq 0$$

we see  $k \leq 0$ . If  $k = 0$ , we have  $x = \dot{x} = 0$ . Therefore  $x^1$  and  $x^2$  run along the line

$$p_4x^1 + p_5x^2 + C = 0.$$

If  $k < 0$ ,

$$p_4x^1 + p_5x^2 + C$$

moves periodically between  $-2\sqrt{-k}$  and  $2\sqrt{-k}$ .

Now we will give the differential equations that an abnormal extremal  $\Gamma : I(=[\alpha, \beta]) \rightarrow T^*\mathbf{R}^5 \setminus \{O\}$  of  $D$  must satisfy. If we choose the local coordinates

$(x^1, x^2, x^3, x^4, x^5, p_1, p_2, p_3, p_4, p_5)$  in  $T^*\mathbf{R}^5$ , the Hamiltonian function  $H_{X_1}$  and  $H_{X_2}$  can be expressed as

$$H_{X_1} = p_1 - \frac{1}{2}x^2p_3 - (x^3 - \frac{1}{2}x^1x^2)p_4,$$

$$H_{X_2} = p_2 + \frac{1}{2}x^1p_3 - (x^3 + \frac{1}{2}x^1x^2)p_5.$$

By the definition of an abnormal extremal of  $(M, D)$ ,  $H_{X_1}$  and  $H_{X_2}$  should vanish along the curve  $\Gamma$ . Hence we have:

$$p_1 - \frac{1}{2}x^2p_3 - (x^3 - \frac{1}{2}x^1x^2)p_4 = 0,$$

$$p_2 + \frac{1}{2}x^1p_3 - (x^3 + \frac{1}{2}x^1x^2)p_5 = 0.$$

Now the Hamiltonian lift of  $\vec{H}_{X_1}$  of  $X_1$  and  $\vec{H}_{X_2}$  of  $X_2$  can be expressed as:

$$\begin{aligned} \vec{H}_{X_1} &= \frac{\partial}{\partial x^1} - \frac{1}{2}x^2 \frac{\partial}{\partial x^3} - (x^3 - \frac{1}{2}x^1x^2) \frac{\partial}{\partial x^4} \\ &\quad - \frac{1}{2}x^2p_4 \frac{\partial}{\partial p_1} - (\frac{1}{2}x^2p_4 - \frac{1}{2}p_3) \frac{\partial}{\partial p_2} + p_4 \frac{\partial}{\partial p_3}, \end{aligned}$$

$$\begin{aligned} \vec{H}_{X_2} &= \frac{\partial}{\partial x^2} + \frac{1}{2}x^1 \frac{\partial}{\partial x^3} - (x^3 + \frac{1}{2}x^1x^2) \frac{\partial}{\partial x^5} \\ &\quad - (\frac{1}{2}p_3 - \frac{1}{2}x^2p_5) \frac{\partial}{\partial p_1} + \frac{1}{2}x^1p_5 \frac{\partial}{\partial p_2} + p_5 \frac{\partial}{\partial p_3}. \end{aligned}$$

Then the following conditions must be satisfied:

$$\dot{\Gamma}(t) = a^1(t)(\vec{H}_{X_1})_{\Gamma(t)} + a^2(t)(\vec{H}_{X_2})_{\Gamma(t)},$$

where  $a^1(t)$  and  $a^2(t)$  are some functions on  $I$ .

Therefore if  $\Gamma(t) = (x(t), p(t))$  is an abnormal extremal of  $(M, D)$  then  $(x(t), p(t))$

satisfies the following equations,

$$(8.16) \quad \dot{x}^1 = a^1$$

$$(8.17) \quad \dot{x}^2 = a^2$$

$$(8.18) \quad \dot{x}^3 = -\frac{1}{2}a^1x^2 + \frac{1}{2}a^2x^1$$

$$(8.19) \quad \dot{x}^4 = -a^1(x^3 - \frac{1}{2}x^1x^2)$$

$$(8.20) \quad \dot{x}^5 = -a^2(x^3 + \frac{1}{2}x^1x^2)$$

$$(8.21) \quad \dot{p}_1 = -\frac{1}{2}a^1x^2p_4 - a^2(\frac{1}{2}p_3 - \frac{1}{2}x^2p_5)$$

$$(8.22) \quad \dot{p}_2 = a^1(-\frac{1}{2}x^2p_4 + \frac{1}{2}p_3) + \frac{1}{2}a^2x^1p_5$$

$$(8.23) \quad \dot{p}_3 = a^1p_4 + a^2p_5$$

$$(8.24) \quad \dot{p}_4 = 0$$

$$(8.25) \quad \dot{p}_5 = 0$$

$$(8.26) \quad p_1 - \frac{1}{2}x^2p_3 - (x^3 - \frac{1}{2}x^1x^2)p_4 = 0$$

$$(8.27) \quad p_2 + \frac{1}{2}x^1p_3 - (x^3 + \frac{1}{2}x^1x^2)p_5 = 0$$

Differentiating the equation (8.27), and substituting (8.16), (8.17), (8.18), (8.22), (8.23), (8.25) into it, we have

$$p_3\dot{x}^1 = 0.$$

Similarly differentiating (7.26), and substituting (8.16), (8.17), (8.18), (8.21), (8.23), (8.24) into it, we have

$$p_3\dot{x}^2 = 0.$$

From these equations on account of (8.16), (8.17), it follows that if  $(a^1(t_0), a^2(t_0)) \neq 0$  at  $t_0$ , then  $p_3 = 0$  around  $t_0$ . Therefore we may assume  $p_3 \equiv 0$ , and we have

$$\dot{p}_3 = (a^1(t)p_4 + a^2(t)p_5) = 0.$$

Hence we have

$$\begin{pmatrix} \dot{x}^1 \\ \dot{x}^2 \end{pmatrix} = \begin{pmatrix} a^1 \\ a^2 \end{pmatrix} = \varphi \begin{pmatrix} p_5 \\ -p_4 \end{pmatrix},$$

where  $\varphi$  is a function along the abnormal biextremal. If we set

$$\psi = \int_{\alpha}^t \varphi(s) ds,$$

we have

$$\begin{pmatrix} x^1(t) \\ x^2(t) \end{pmatrix} = \psi(t) \begin{pmatrix} p_5 \\ -p_4 \end{pmatrix} + \begin{pmatrix} q^1 \\ q^2 \end{pmatrix},$$

where  $q^1 = x^1(\alpha)$ ,  $q^2 = x^2(\alpha)$ . Then  $x^3, x^4, x^5$  are obtained by integrating (8.18), (8.19), (8.20). Thus the lines in  $(x^1, x^2)$ -space give rise to the abnormal extremals.

### § 9. Control system

We need the following generalized notion of control systems [1]. Let  $M$  be a manifold. A *control system* on  $M$  is given by a locally trivial fibration  $\pi_{\mathcal{U}} : \mathcal{U} \rightarrow M$  over  $M$  and a fibered map  $F : \mathcal{U} \rightarrow TM$  over the identity of  $M$ . In this paper, we assume that the fibre  $U$  of  $\pi_{\mathcal{U}} : \mathcal{U} \rightarrow M$  is an open subset of  $\mathbf{R}^r$ . Note that we mainly treat a local case where  $\pi_{\mathcal{U}}$  is trivial. However our results do not depend on any choice of trivialisation of  $\pi_{\mathcal{U}}$ .

For a given control system, an  $L^\infty$  (measurable, essentially bounded) map  $c : [a, b] \rightarrow \mathcal{U}$  on an interval is called an *admissible control* if the curve  $\gamma := \pi_{\mathcal{U}} \circ c : [a, b] \rightarrow M$  satisfies  $\dot{\gamma}(t) = F(c(t))$ , for almost every  $t \in [a, b]$ . Then the Lipschitz curve  $\gamma$  is called a *trajectory*. In what follows we use the term “path” for a smooth ( $C^\infty$ ) immersive trajectory regarded up to parametrisation.

The *initial* (resp. *end*) *point* of  $\gamma$  is given by  $\gamma(a) = \pi_{\mathcal{U}}(c(a))$  (resp.  $\gamma(b) = \pi_{\mathcal{U}}(c(b))$ ). Note that, in our terminology, the term “control” contains the data on both the “control” in usual sense and (the initial point of) the trajectory. Therefore trajectory is determined uniquely by the control, while the control is not unique for the trajectory in general.

If we write locally  $x(t) = \gamma(t)$  and  $c(t) = (x(t), u(t))$ , under a local triviality  $\mathcal{U}|_V \cong V \times U$  over an open set  $V \subset M$ , then the control system is expressed by a family of vector fields,  $\{f_u\}_{u \in U}$ ,  $f_u(x) = F(x, u)$  over  $V \subset M$  and by the equation  $\dot{x}(t) = f_{u(t)}(x(t))$ . In this local situation or in the case where the fibration  $\pi_{\mathcal{U}}$  is globally trivial, for a given initial point,  $u(t)$  is called a control defining the trajectory  $x(t)$  as usual. Note that, for a given initial point  $q_0$  and a given  $L^\infty$  control  $u(t)$ , the Cauchy problem  $f_u(x) = F(x, u), x(0) = q_0$ , has a unique Lipschitz solution (trajectory)  $x(t)$  depending smoothly on  $q_0$  by the classical Carathéodory theorem.

Now, given a control system and given a point  $q_0 \in M$ , we denote by  $\mathcal{C}_{ad}$  the set of all admissible controls  $c : [a, b] \rightarrow \mathcal{U}$  with the initial point  $\pi_{\mathcal{U}}(c(a)) = q_0$ . Then it is known that  $\mathcal{C}_{ad}$  is a Banach manifold ([2]). The *endpoint mapping*  $\mathcal{E} : \mathcal{C}_{ad} \rightarrow M$  is defined by  $\mathcal{E}(c) := \pi_{\mathcal{U}}(c(b))$ . The control  $c$  with the initial point  $\pi_{\mathcal{U}}(c(a)) = q_0$  is called *singular* or *abnormal*, if it is a singular point of  $\mathcal{E}$ , namely if the differential  $\mathcal{E}_* : T_c \mathcal{C}_{ad} \rightarrow T_{\mathcal{E}(c)} M$  is not surjective. If  $c$  is a singular control, then the trajectory  $\gamma = \pi_{\mathcal{U}} \circ c$  is called a *singular trajectory* or an *abnormal extremal* ([2][3]).

Let  $D \subset TZ$  be a Cartan distribution. Then, it is known that for any point  $y$  of  $Y$  and for any direction  $\ell \in D_y$ , there exists uniquely a singular  $D$ -path (an immersed abnormal extremal) through  $y$  with the given direction  $\ell$ . Thus the singular  $D$ -paths form another five dimensional manifold  $X$ .

And let  $Z = PD = (D - 0)/\mathbf{R}^\times$  be the space of tangential lines in  $D$ ,  $\dim(Z) = 6$ . Then  $Z$  is naturally foliated by the liftings of singular  $D$ -paths, and we have locally double fibrations:

$$Y \xleftarrow{\pi_Y} Z \xrightarrow{\pi_X} X.$$

Let  $E \subset TZ$  be the Cartan prolongation of  $D \subset TY$ : For each  $(y, \ell) \in Z$ ,  $\ell \in T_y Y$ ,  $E_{(y, \ell)} := \pi_{Y*}^{-1}(\ell)$ . Then  $E$  is a distribution with growth  $(2, 3, 4, 5, 6)$ .

If we put  $L = \text{Ker}(\pi_{Y*})$ ,  $K = \text{Ker}(\pi_{X*})$ , then we have a decomposition  $E = L \oplus K$  by integrable subbundles.

A cone field  $C \subset TX$  is defined by setting, for each  $x \in X$ ,

$$C_x := \bigcup_{z \in \pi_X^{-1}(x)} \pi_{X*}(L_z) \subset T_x X$$

Thus we have distributions  $D, E, L, K$  and the cone field  $C$  from the double fibrations:

$$\begin{array}{ccccc} TY & \xleftarrow{\pi_{Y*}} & TZ & \xrightarrow{\pi_{X*}} & TX \\ \cup & & \cup & & \cup \\ D & \xleftarrow{\pi_{Y*}|_E} & E = L \oplus K & \xrightarrow{\pi_{X*}|_E} & C \\ \downarrow & & \downarrow & & \downarrow \\ Y & \xleftarrow{\pi_Y} & Z & \xrightarrow{\pi_X} & X \end{array}$$

We regard the cone field as a control system over  $X$ :

$$\mathbb{C} : L \xrightarrow{\pi_{X*}|_L} TX \rightarrow X.$$

Then we have the following main theorem([8]):

### § 10. Duality and asymmetry

**Theorem 10.1.** *Singular paths of the control system*

$$\mathbb{C} : L \xrightarrow{\pi_{X*}|_L} TX \rightarrow X$$

are given by  $\pi_X$ -images of  $\pi_Y$ -fibres.

*Proof.* Let  $\gamma : I \rightarrow X$  be a singular  $\mathbb{C}$ -trajectory. Then, by the next Lemma 10.2 ([8]),  $\gamma$  lifts to a Lipschitz abnormal bi-extremal  $\beta : I \rightarrow L \times_X T^*X$  for  $\mathbb{C}$ . Then, by Lemma 10.2 ([8]),  $\gamma$  lifts to a singular  $\mathbb{E}$ -trajectory  $\tilde{\gamma}$  for the projection  $\pi_X : Z \rightarrow X$ ,  $\pi_X \circ \tilde{\gamma} = \gamma$ . Since  $\gamma$  is immersive,  $\tilde{\gamma}$  is also immersive. By Proposition 10.3,  $\tilde{\gamma}$  is either a  $\pi_Y$ -fibre or the lift of a singular  $D$ -trajectory. In this case,  $\tilde{\gamma}$  must be a  $\pi_Y$ -fibre, and we have that  $\gamma$  is the  $\pi_X$ -image of a  $\pi_Y$ -fibre up to parametrisation. □

Therefore, for any  $x \in X$  and for any direction  $\ell \in C_x$ , there exists uniquely a singular  $\mathbb{C}$ -paths passing through  $x$  with the direction  $\ell$  at  $x$ .

The original space  $Y$  is identified with the space of singular paths on the space  $X$ , the space of singular paths on  $Y$ .

**Lemma 10.2.** ([8]) *A path on  $X$  is a singular  $\mathbb{C}$ -path if and only if it is a singular  $\mathbb{E}/\pi_X$ -path. Any singular  $\mathbb{E}/\pi_X$ -path on  $X$  lifts to a  $\mathbb{E}$ -path on  $Z$  via  $\pi_X : Z \rightarrow X$ .*

**Proposition 10.3.** ([8]) *Let  $D$  be a distribution with growth  $(2, 3, 5)$  on a five dimensional manifold  $Y$ , and  $(Z, E)$  the Cartan prolongation of  $D$ . Then any singular  $E$ -path (un-parametrised immersed  $E$ -trajectory) is either a fibre of  $\pi_Y : Z \rightarrow Y$  or the lift of a singular  $D$ -path. Any  $\pi_Y$ -fibre is a regular singular path of  $E$ . The lift of any singular  $D$ -path is a totally irregular singular path of  $E$ .*

A singular path  $x(t)$  for  $E \subset TZ$  is called regular singular if it is associated with an abnormal bi-extremal  $(x(t), p(t), u(t))$  such that  $p(t) \in E^{(2)\perp} \setminus E^{(3)\perp} \subset T^*Z$ .

A singular path  $x(t)$  for  $E \subset TZ$  is called totally irregular singular if any associated abnormal bi-extremals satisfies that  $p(t) \in E^{(3)\perp} \subset T^*Z$ .

Recently the double fibration  $(Y, D) \xleftarrow{\pi_Y} (Z, E) \xrightarrow{\pi_X} (X, C)$  with  $G_2$ -symmetry have been explicitly constructed in [9]. Our Theorems 10.1, Proposition 10.3 ([8]) show, from the viewpoint of geometric control theory the duality and asymmetry hold, not only for  $G_2$ -case, but for general Cartan distributions.

### § 11. Local characterization of abnormal extremal

The main tool to show Lemma 10.2 and Proposition 10.3 is the following: We see that if a control is singular then its restriction to any subinterval is singular. In fact, as a part of *Pontryagin maximal principle*, the general characterization of singular controls is known. To state the characterization, consider the fibre product of  $\pi_{\mathcal{U}} : \mathcal{U} \rightarrow M$  and  $\pi_M^* : T^*M \rightarrow M$ :

$$\mathcal{U} \times_M T^*M := \{(w, \varphi) \in \mathcal{U} \times T^*M; \pi_{\mathcal{U}}(w) = \pi_M^*(\varphi)\},$$

(extended cotangent bundle), endowed with the natural projections  $\Pi_{\mathcal{U}} : \mathcal{U} \times_M T^*M \rightarrow \mathcal{U}$  and  $\Pi_M^* : \mathcal{U} \times_M T^*M \rightarrow T^*M$ . Regarding the local triviality  $\mathcal{U}|_V \cong V \times U$  over a local coordinate neighborhood  $V \subset M$ ,  $\mathcal{U} \times_M T^*M$  is identified with  $(\mathcal{U}|_V) \times_V (T^*M|_V)$  and with  $T^*M|_V \times U$ .

We define the Hamiltonian function  $H : \mathcal{U} \times_M T^*M \rightarrow \mathbf{R}$  of the control system  $F : \mathcal{U} \rightarrow TM$  by

$$H(x, p, u) := \langle p, F(x, u) \rangle, \quad ((x, u), (x, p)) \in \mathcal{U} \times_M T^*M.$$

A singular control  $(x(t), u(t))$  is characterized by the liftability to an abnormal biextremal  $(x(t), p(t), u(t))$  satisfying the constrained Hamiltonian equation

$$\begin{cases} \dot{x}_i(t) = \frac{\partial H}{\partial p_i}(x(t), p(t), u(t)), & (1 \leq i \leq m) \\ \dot{p}_i(t) = -\frac{\partial H}{\partial x_i}(x(t), p(t), u(t)), & (1 \leq i \leq m) \\ \frac{\partial H}{\partial u_j}(x(t), p(t), u(t)) = 0, & (1 \leq j \leq r), \quad p(t) \neq 0, \end{cases}$$

where  $m$  is the dimension of manifold  $M$ .

## References

- [1] A. A. Agrachev, *Geometry of Optimal Control Problems and Hamiltonian Systems*, Non-linear and Optimal Control Theory, Springer-Verlag (2004), pp. 1–59.
- [2] A.A. Agrachev, Y.L. Sachcov, *Control Theory from Geometric Viewpoint*, Encyclopaedia of Mathematical Sciences **87**, Springer-Verlag, (2004).
- [3] B. Bonnard, M. Chyba, *Singular Trajectories and the Role in Control Theory*, Springer-Verlag (2003).
- [4] R. L. Bryant, *Élie Cartan and geometric duality*, A lecture given at the Institut d'Élie Cartan on 19 June 1998.
- [5] R.L. Bryant, L. Hsu, *Rigidity of integral curves of rank 2 distributions*, Inventiones Mathematicae, **114** (1993), 435–461.
- [6] E. Cartan, *Les systèmes de Pfaff à cinq variables et les équations aux dérivées partielles du second ordre*, Ann. Sci. Ecole Norm. Sup. (3), **27** (1910), 109–192.
- [7] W. L. Chow, *Über System Von Linearen Partiellen Differentialgleichungen Erster Ordnung*, Math. Ann. **117** (1940–1941), 98–105.
- [8] G. Ishikawa, Y. Kitagawa, W. Yukuno, Duality of singular paths for (2,3,5)-distributions, to appear in Journal of Dynamical and Control Systems.
- [9] G. Ishikawa, Y. Machida, M. Takahashi, *Singularities of tangent surfaces in Cartan's split  $G_2$ -geometry*, Hokkaido University Preprint Series in Mathematics #1020, (2012).
- [10] Y. Kitagawa, *The Infinitesimal automorphisms of a homogeneous subriemannian contact manifold*, Thesis, Nara Women's University (2005).

- [11] I. Kupka, *Abnormal extremals*, preprint, (1992).
- [12] W. Liu, H.J. Sussman, *Shortest paths for sub-Riemannian metrics on rank-two distributions*, *Memoirs of Amer. Math. Soc.*, **118–564**, Amer. Math. Soc., (1995).
- [13] R. Montgomery, *Abnormal Minimizers*, *SIAM J. Control and Optimization*, **32-6**, (1994), 1605–1620.
- [14] R. Montgomery, *A Survey of Singular Curves in Subriemannian Geometry*, *Journal of Dynamical and Control Systems*, **1** (1995), 49–90.
- [15] R. Montgomery, *A Tour of Subriemannian Geometries, Their Geodesics and Applications*, *Mathematical Surveys and Monographs*, **91**, Amer. Math. Soc. (2002).
- [16] M. Okada, *Minimizers on Sub-Riemannian manifolds, (in Japanese)*, Mc Thesis Kyoto Univ. of Edu., (1996).
- [17] N. Tanaka, *A differential geometric study on strongly pseudo-convex manifolds, Lectures in Mathematics*, Department of Mathematics, Kyoto Univ., **9**, (1975).