

1 **Lymphocyte-Area Under the Curve as a predictive factor for viral infection**
2 **after allogenic hematopoietic stem cell transplantation**

3 Mizuki Watanabe¹, Junya Kanda^{1†}, Masakatsu Hishizawa¹, Tadakazu Kondo¹, Kouhei
4 Yamashita¹, Akifumi Takaori-Kondo¹

5

6 ¹Department of Hematology and Oncology, Graduate School of Medicine, Kyoto University,
7 Kyoto, Japan

8

9 †Correspondence: Junya Kanda, M.D,

10 Department of Hematology and Oncology, Graduate School of Medicine, Kyoto University,
11 Kyoto, Japan, 54 shogoin-kawaramachi, Kyoto, Kyoto, Japan, 606-8507.

12 Tel: +81-75-751-3152; Fax: +81-75-751-3153; e-mail: jkanda16@kuhp.kyoto-u.ac.jp

13

14 **Conflict-of-interest disclosure**

15 The authors declare no competing financial interests.

16

17 This work was supported in part by the Takeda Science Foundation (JK).

18

19 **Running head:** Lymphocyte-AUC as a predictive factor for viral infection after allo-HSCT

20

21

22

23 Text word count, 2864; abstract word count, 287; 4 tables; 2 figures; 0 supplemental tables;

24 33 references

25

26

1 **Abstract**

2 *Background*

3 Viral infection is a serious complication that can greatly affect patient mortality
4 and morbidity after allogeneic hematopoietic stem cell transplantation (allo-HSCT).
5 For the early identification of patients who are at high risk for viral infection, we
6 evaluated the impact of lymphocyte-Area Under the Curve (AUC) as a new
7 predictive factor for early immune reconstitution after allo-HSCT against viral
8 infection.

9 *Method*

10 This study included 286 patients who underwent their first allo-HSCT at Kyoto
11 University Hospital between 2005 and 2017. Lymphocyte-AUC from day 0 to day
12 15 was calculated in the analysis of HHV-6, and that from day 0 to day 30 was
13 calculated in the analysis of other viruses (cytomegalovirus (CMV), adenovirus,
14 BK virus, JC virus, and varicella zoster virus). The risk factors for each viral
15 reactivation/infection were assessed using multivariate analysis.

16 *Results*

17 The median age at transplantation was 51 (range, 17–68) years. The median
18 lymphocyte-AUC was 63 (range, 0–5620)/ μ L at day 15 and 3880 (range, 0–
19 118260)/ μ L at day 30. An increase in lymphocyte-AUC was significantly
20 associated with a high frequency of HHV6 reactivation ($P=0.033$) and a low
21 frequency of CMV antigenemia ($P=0.014$). No apparent association was found
22 between lymphocyte-AUC and reactivation/infection of other viruses. Aplastic
23 anemia as a primary disease (HR, 5.34; $P<0.001$) and cord blood as a donor
24 source (HR, 3.05; $P=0.006$) were other risk factors for HHV-6 reactivation. The
25 occurrence of acute graft-versus-host disease (HR 2.21, $P<0.001$) and recipient
26 age (HR 1.55, $P=0.017$) were also risk factors for CMV antigenemia. Higher

1 lymphocyte-AUC at day 30 was significantly associated with low treatment-
2 related mortality (HR 0.47, P=0.045).

3 *Conclusion*

4 Lymphocyte-AUC may be a good predictive factor for immune reconstitution
5 against CMV reactivation. It also provides valuable information for predicting
6 HHV-6 reactivation and treatment-related mortality.

7 **Key words**

8 *lymphocyte-AUC, HHV-6, CMV antigenemia, viral reactivation, immune*
9 *reconstitution*

10

1 **Highlights**

2

3 ● High lymphocyte-AUC was associated with a lower frequency of CMV
4 antigenemia.

5 ● High lymphocyte-AUC was associated with a lower risk of treatment-related
6 mortality.

7 ● Lymphocyte-AUC could be a prognostic factor of immune-reconstitution after
8 HSCT.

9 ● High lymphocyte-AUC was associated with a high frequency of HHV-6
10 reactivation

11 ● Rapid recovery of lymphocyte after HSCT might be associated with HHV-6
12 expansion.

13

14

15

1 **Introduction**

2 Viral infections continue to be serious complications that negatively impact patient
3 survival after allogenic hematopoietic stem cell transplantation (allo-HSCT). After
4 allo-HSCT, patients often develop reactivation of and infection by various latent
5 viruses, including cytomegalovirus (CMV), varicella zoster virus (VZV), human
6 herpes virus-6 (HHV-6), adenovirus (ADV), BK virus (BKV), and JC virus (JCV),
7 due to their prolonged and strongly immunosuppressed background¹.

8

9 Since both the number of transplantations from various stem cell sources such
10 as cord blood unit and the number of transplantations performed for high-risk
11 patients are increasing, the management of viral infection is becoming even more
12 important to improve clinical outcomes of HSCT. However, preventive measures
13 and effective treatments against these viruses are still limited and remain largely
14 dependent on immune reconstitution in the recipients themselves. However, as
15 seen in the prophylactic administration of acyclovir/valacyclovir against VZV^{1,2}
16 and pre-emptive therapies against CMV infections diagnosed via serum antigen
17 or real-time polymerase chain reaction (PCR)^{3,4}, early intervention leads to
18 favorable outcomes. It is important to identify high-risk patients for viral infection
19 in the early stage after HSCT. Hence, in this study, we assessed a new biomarker,
20 lymphocyte-AUC, as a new predictive factor for immune reconstitution after allo-
21 HSCT by evaluating its impact on viral reactivation/infection.

22

1 **Methods**

2 **Data collection**

3 A total of 286 patients who underwent their first allogeneic HSCT for
4 hematological diseases at a single center of Kyoto University Hospital between
5 2005 and 2017 were reviewed. Lymphocyte-AUC is defined as the sum of serial
6 absolute lymphocyte counts under the lymphocyte count-time curve⁵. In the
7 analysis of HHV-6 reactivation, lymphocyte-AUC values from day 0 to day 15
8 were calculated in patients who survived over 15 days after the transplant, as
9 most cases of HHV-6 virus reactivation occurred from day 15 to day 30.
10 Regarding the analysis of other viruses (CMV, ADV, BKV, JCV, and VZV),
11 lymphocyte-AUC values from day 0 to day 30 were calculated in patients who
12 survived over 30 days after transplant, since infection by these viruses was
13 mostly seen from 30 days after HSCT.

14 This study was approved by the Institutional Review Board of Kyoto University
15 Hospital and written informed consent was obtained from every patient.

16

17 **Viral detection and treatment**

18 *CMV antigenemia and CMV virus infection*

19 CMVpp65 antigen was examined once weekly in every patient after an increase
20 in the neutrophil count was ascertained and was examined additionally in patients
21 with suspicious signs and symptoms of CMV diseases. Most of the patients were
22 examined via the C10/C11 method, while some patients were assessed via the
23 C7-HRP method. The results of the C7-HRP method are known to be highly
24 correlated with those of the C10/11 method⁶. Both methods were performed as
25 previously reported⁷⁻¹⁰. In cases where more than 2 positive cells within 2 slides
26 (within 50000 WBC in C10/C11) were detected, pre-emptive therapy was given
27 followed by close monitoring of CMV antigen^{6,8}.

1

2 *HHV-6 preventive measures, reactivation, and infection*

3 The HHV-6 viral load was determined quantitatively after transplantation by
4 multiplex PCR designed for multiple viral detection¹¹ whenever a patient
5 developed symptoms suspicious for HHV-6 reactivation. In patients who received
6 CBT within the past seven years, PCR was examined consistently (every one or
7 two weeks until 2 months after transplantation).

8 For patients who received CBT within the past three years, foscarnet infusion was
9 started at a maintenance dose (90mg/kg/day, adjusted by the patient's kidney
10 function) to prevent severe HHV-6 reactivation when patients were administered
11 systemic steroid for immune reactions such as engraftment syndrome or acute
12 graft-versus-host disease (aGVHD). Foscarnet at a curative dose (180mg/kg/day,
13 adjusted by the patient's kidney function) was injected when HHV-6 infection,
14 including HHV-6 encephalitis, was diagnosed¹². For patients with only HHV-6
15 reactivation, who were diagnosed as serum HHV-6 positive without any
16 symptoms, treatment was initiated based on the physician's discretion,
17 considering the detected viral dose (approximately 10^3 copies/ml) and the
18 patient's background.

19

20 *Adenovirus, BK virus, and JC virus viral infection*

21 When symptoms indicative of urinary tract infection such as hematuria emerged,
22 serum and urinary levels of ADV, BKV, and JCV were examined by multiplex
23 PCR¹¹. For ADV, patients were also subjected to additional examinations when
24 they developed hepatitis, fever, or other symptoms of infection of undetectable
25 origin. For patients in whom ADV and BKV were detected in serum, systemic
26 cidofovir injection was initiated at 1 mg/kg, three times a week. Meanwhile, for
27 those in whom BKV and ADV were detected only in the urine, bladder instillation

1 of cidofovir was preferred at 5 mg/kg for two days in a row¹³⁻¹⁵.

2

3 **Endpoints**

4 The primary endpoint of this study was the occurrence of reactivation and
5 infection with various viruses (CMV, VZV, HHV-6, ADV, BKV, and JCV) diagnosed
6 within 180 days after HSCT.

7

8 **Statistical analysis**

9 Descriptive statistics were used to summarize variables related to the patient
10 characteristics. Viral reactivation/infection, treatment-related mortality and
11 disease relapse that occurred by day180 were calculated based on cumulative
12 incidence curves^{16,17}. Overall survival was evaluated by the Kaplan-Meier method.
13 Competing events were defined as deaths without a diagnosis of viral
14 reactivation/infection. Lymphocyte-AUC was estimated by collecting the area
15 under the curve of lymphocyte counts in each patient from day 1 until either day
16 15 (for HHV-6) or day 30 (for other viruses). These landmark days (day 15, day
17 30) were determined based on a preceding analysis in which over 75% of onset
18 cases were detected between day 15 and day 30 in HHV6 reactivation and after
19 day 30 in CMV antigenemia. Fine and Gray's proportional hazards model¹⁸ was
20 used to evaluate the impact of lymphocyte-AUC on viral reactivation/infection in
21 each patient. The following possible covariates were considered; recipient's sex,
22 age at transplant (<50 years old or ≥50 years old), disease diagnosis (myeloid
23 malignancies, lymphoid malignancies, and others), disease status (complete
24 remission [CR] or non-CR), donor type (bone marrow transplantation from
25 unrelated donor, peripheral blood stem cell transplantation from related donor,
26 cord blood transplant), conditioning regimen (reduced-intensity or myeloablative),
27 GVHD prophylaxis (tacrolimus or cyclosporine in addition to mycophenolate

1 mofetil or methotrexate), and the occurrence of acute GVHD by day 30 (only for
2 CMV antigenemia). All covariate factors with a variable retention criterion of
3 $P < 0.05$ in the univariate analysis were selected and analyzed together with
4 lymphocyte-AUC in the multivariate analysis. All statistical analyses were
5 performed with EZR (Saitama Medical Center, Jichi Medical University, Saitama,
6 Japan), which is a graphical user interface for R (The R Foundation for Statistical
7 Computing, version 3.1.1, Vienna, Austria)¹⁹.

8

1 **Result**

2 **Patient characteristics**

3 Two-hundred-eighty-six patients were reviewed in the analysis of HHV-6
4 reactivation and 283 were examined for other viral reactivation/infection (3
5 patients died between day 15 and day 30). Seventy-eight patients received
6 transplantation from a related donor, 129 from unrelated bone marrow grafts,
7 and 79 from unrelated cord blood units. Their median age at transplantation
8 was 51 (range, 17–68) years. The median lymphocyte-AUC was 63 (range, 0–
9 5620)/ μL at day 15 and 3880 (range, 0–118260)/ μL at day 30. No apparent
10 difference in lymphocyte-AUC was seen between different donor sources.
11 We categorized the patients into 3 groups according to their lymphocyte-AUC
12 count by day 15 and day 30. However, in the analysis of HHV-6 reactivation, the
13 first tertile was 0/ μL , since 129 patients showed no lymphocyte recovery by day
14 15. Hence, we used the second tertile of 230/ μL as a threshold to categorize
15 patients into two groups in the analysis of lymphocyte-AUC by day 15:
16 lymphocyte-AUC \leq 230/ μL (n=189) and lymphocyte-AUC $>$ 230/ μL (n=97)
17 (Table 1). In the analysis of CMV antigenemia and infection, patients were
18 categorized into three groups according to the first (2710/ μL) and the second
19 (5250/ μL) tertile: low lymphocyte-AUC (n=93), middle lymphocyte- AUC (n=93),
20 and high lymphocyte-AUC (n=97) (Table 2).

21

22 **HHV-6 reactivation/infection**

23 HHV-6 reactivation was detected in 48 of the 286 patients (cumulative
24 incidence: 17.5% on day 180), of whom 8 patients developed virologically
25 diagnosed HHV-6 encephalitis with typical neurological symptoms and viral
26 detection in spinal fluid with or without positive findings in magnetic resonance
27 imaging. Nine patients received foscarnet injection as prophylaxis from the 1st to

1 4th weeks from transplantation after CBT, of whom 5 were diagnosed as HHV-6
2 viremia after cessation of foscarnet.

3 Multivariate analysis showed that a high lymphocyte-AUC was significantly
4 associated with HHV-6 reactivation (high AUC group vs. low-plus-middle AUC
5 group: HR, 1.83; P=0.048) (Figure 1). Other risk factors detected were aplastic
6 anemia as a primary disease (HR, 5.34; P<0.001) and cord blood as a donor
7 source (HR, 3.05; P=0.006) (Table 3). In the sub-analysis of patients with a
8 history of HHV-6 viremia, there was no significant difference in lymphocyte-AUC
9 between the HHV-6 encephalitis group and no-encephalitis group (median
10 lymphocyte-AUC value, encephalitis group, 530/ μ L; no-encephalitis group
11 249/ μ L; P=0.248). Foscarnet treatment had no prophylactic effect on HHV-6
12 viremia (incidence: patients with foscarnet prophylaxis vs. without; 55.6% vs.
13 37.1%).

14 Since HHV-6 reactivation has been epidemiologically suggested to be
15 associated with immune reactions before engraftment including pre-engraftment
16 immune reaction in CBT²⁰, we performed an additional analysis to examine the
17 association between lymphocyte-AUC and the occurrence of immune-related
18 reactions by day 15. High lymphocyte-AUC was associated with the occurrence
19 of immune-related reactions (Odds ratio 2.02, P=0.015). However, in a
20 stratification analysis, high-lymphocyte AUC was significantly associated with
21 HHV-6 reactivation in patients both with and without an immune reaction by
22 day15 (high AUC group vs. low-plus-middle AUC group, patients with immune
23 reaction; HR, 2.41, P=0.047, patients without immune reaction; HR, 2.51,
24 P=0.018). Meanwhile, in another stratification analysis, immune-related
25 reactions showed no apparent association with HHV-6 reactivation in groups of
26 patients with both a high lymphocyte-AUC and low-plus-middle lymphocyte-
27 AUC (patients with an immune reaction vs. patients without an immune

1 reaction, high AUC group; HR, 1.73, P=0.160, low-plus-middle AUC group; HR,
2 1.83, P=0.169)

4 **CMV antigenemia**

5 CMV antigenemia was detected in 146 of the 284 patients (cumulative
6 incidence: 54.7% by day 180). Nine cases of CMV end organ infection
7 occurred, 6 of which were diagnosed as CMV-related colitis/gastritis, and one
8 each as retinitis, hepatitis, and pneumonia. Foscarnet was used for 9 patients
9 as a prophylaxis for HHV-6. This was discontinued after day 30. No other
10 agents were used for HHV-6 or CMV prophylaxis for the remaining 277 patients.
11 In a multivariate analysis, the high AUC-lymphocyte group with AUC 5250/ μ L or
12 above had a lower risk for CMV antigenemia than the low lymphocyte-AUC
13 group (HR, 0.61; P=0.052). Meanwhile, the risk for CMV antigenemia was not
14 significantly different between the middle lymphocyte-AUC group under 5250/ μ L
15 and the low lymphocyte-AUC group (HR, 1.13; P=0.560) (Figure 2). Other risk
16 factors detected in the multivariate analysis were age 50 years or older (vs.
17 <50, HR, 1.55; P=0.017) and the occurrence of acute GVHD by day 30 (vs. no
18 occurrence of acute GVHD; HR, 2.21; P<0.001) (Table 4).

19 There was no association between preceding HHV-6 reactivation and the
20 occurrence of CMV antigenemia (cumulative incidence of CMV reactivation
21 after day 30: patients with history of HHV-6 reactivation by day 30 vs. those
22 without, HR 1.07; P=0.746).

24 **Reactivation of other viruses**

25 A total of 27 cases in 20 patients were diagnosed as various viral reactivations,
26 including ADV viremia (n=7), BKV viremia (n=13), JCV viremia (n=5), VZV
27 viremia (n=1), and EBV viremia (n=1). Nine cases represented multiple viral

1 coinfections (ADV/BKV n=4; BKV/JCV n=4; and ADV/BKV /JCV n=1). No
2 apparent association was noted between these viral infections and the
3 lymphocyte-AUC.
4 Regarding the frequencies of sequential infections of these viruses, 6 of 45
5 patients with a history of HHV-6 viremia by day 30 had a subsequent infection
6 with ADV, BKV or JCV, and 3 of 238 patients without history of HHV-6 viremia
7 had these infections. The cumulative incidence of ADV, BKV or JCV reactivation
8 after day 30 was significantly higher in patients with a history of HHV-6
9 reactivation by day 30 than in those without (HR, 11.1; P=0.001).

10

11 **Overall survival, relapse and treatment-related mortality**

12 No apparent association was observed between lymphocyte-AUC at day 15 and
13 overall survival (high-AUC group vs. low-plus-middle-AUC group, HR, 0.81
14 P=0.386), relapse (high-AUC group vs. low-plus-middle-AUC group, HR, 1.01;
15 P=0.974) or treatment-related mortality (high-AUC group vs. low-plus-middle-
16 AUC group, HR, 0.77; P=0.477) were found.

17 Also, neither overall survival (high-AUC group vs. low-AUC group, HR, 0.66;
18 P=0.110, middle-AUC group vs. low-AUC group, HR, 0.63; P=0.095) nor
19 relapse (high-AUC group vs. low-AUC group, HR, 0.821; P=0.581, middle-AUC
20 group vs. low-AUC group, HR, 1.25; P=0.512) was significantly associated with
21 lymphocyte-AUC at day 30. However, treatment-related mortality was
22 associated with lymphocyte-AUC at day 30 (high-AUC group vs. low-AUC
23 group, HR, 0.47; P=0.045, middle-AUC group vs. low-AUC group, HR, 0.33;
24 P=0.013).

25

1 **Discussion**

2 In this study, we evaluated lymphocyte-AUC at day 15 and day 30 as a predictive
3 factor for reactivation of and infection by several viruses. HHV-6 and CMV are the
4 two major viruses that cause various complications during the management of
5 HSCT, negatively affecting patient mortality and morbidity. We found that
6 lymphocyte-AUC can be used to identify patients at high risk for reactivation of
7 these viruses.

8

9 In the analysis of HHV-6 reactivation, high lymphocyte-AUC was strongly
10 associated with viral reactivation. Since early intervention with antiviral agents is
11 necessary to reduce HHV-6 reactivation and subsequent virus-related
12 complications²¹⁻²⁴, regular examination of the plasma level of HHV-6 viral load is
13 strongly recommended for all patients, especially in those who show rapid growth
14 of lymphocytes by day 15. In previous studies, HHV-6 reactivation was associated
15 with a myeloablative conditioning regimen, cord blood transplantation, and
16 immune reactions^{21,25}. Contrary to our expectation, an early immune reaction
17 before engraftment had less of an impact on HHV-6 reactivation than lymphocyte-
18 AUC despite the temporary administration of systemic steroid to treat it. This
19 finding that HHV-6 reactivation occurred with the rapid growth of lymphocytes
20 regardless of an immune reaction and the preceding use of systemic steroid by
21 day 15 might provide insights into the mechanism of HHV-6 growth after
22 transplantation. Although it is not known whether the preceding HHV-6 growth
23 increased the lymphocyte counts or the rapid growth of lymphocytes stimulated
24 HHV-6 growth, HHV-6 expansion was accompanied by lymphocyte growth. This
25 is consistent with previous reports which suggested that an inflammatory
26 background caused by various sources of pathogenesis and the upregulation of
27 several chemokines were associated with HHV-6 reactivation²⁶⁻²⁸. Viral latency

1 of HHV-6 and its interaction with lymphocytes and chemokines in growth
2 mechanisms remain to be disclosed. Our limited data (N=49) on lymphocyte
3 subsets examined from day 15 to day 21 after transplantation failed to clarify
4 which constituent of lymphocytes contributed to the growth of HHV-6 (data not
5 shown). However, our data suggested that rapid and early growth of lymphocytes
6 is a predictor of HHV-6 reactivation after HSCT.

7
8 Regarding CMV antigenemia, only the high lymphocyte-AUC group with AUC of
9 5250/ μ L or higher showed a low predicted risk of virus reactivation, indicating that
10 sufficient recovery of lymphocytes is required for immunity against CMV
11 reactivation. CMV antigen must be screened regularly if the lymphocyte-AUC is
12 still low, regardless if a single-point blood count at day 30 shows that the patient's
13 immunity appears to have recovered. Our findings also showed that the
14 occurrence of acute GVHD was associated with CMV reactivation, which is
15 consistent with previous reports^{29,30}.

16
17 In the analysis of viral infections other than HHV-6 and CMV, HHV-6 reactivation
18 influenced the subsequent occurrence of ADV, BKV and/or JCV, which is
19 compatible with the findings in a previous study³¹. This suggests that HHV-6
20 infection may directly influence subsequent ADV/BKV/JCV infection or may
21 simply reflect the severity of the immunocompromised status. Further prospective
22 analysis is required to tackle this clinically important topic of coinfection and
23 sequential viral infection in patients after HSCT.

24
25 As for overall survival and treatment-related mortality, only a low lymphocyte-AUC
26 under 2710/ μ L was suggested to be associated with an elevated risk for
27 treatment-related mortality. The two major causes of treatment-related mortality

1 after HSCT are the occurrence of GVHD and complications caused by various
2 pathogens including bacteria, viruses and fungi. Considering that lymphocyte-
3 AUC at day 30 was not associated with the occurrence of acute GVHD or chronic
4 GVHD (data not shown), the high risk of treatment-related mortality for low
5 lymphocyte-AUC seems to reflect the immature immune reconstitution. Our study
6 suggests that lymphocyte-AUC at day 30 may be a good predictor of general
7 immune reconstitution, including antiviral immunity against CMV antigenemia.

8
9 However, our study had several limitations. First, data on lymphocyte subsets
10 were limited. Since various lineages of lymphocyte reconstitution have been
11 suggested to be associated with HHV-6 reactivation^{32,33}, they should be
12 evaluated more precisely to further clarify the interaction between HHV-6 and
13 lymphocytes. Second, since the number of cases with HHV-6 infection such as
14 encephalitis in our hospital was limited, the impact of lymphocyte-AUC on HHV-
15 6 infection was not examined. Studies with a larger cohort are required to
16 examine the impact of lymphocyte-AUC on symptomatic HHV-6 reactivation.

17
18 In conclusion, increases in lymphocyte-AUC at day 15 and day 30 may help to
19 identify patients who are at high risk for HHV-6 reactivation, low risk for CMV
20 reactivation and treatment-related mortality, respectively. A prospective clinical
21 study of pre-emptive therapy with antiviral agents against HHV-6 for patients with
22 high lymphocyte-AUC at day 15 is expected in the future.

23 24 **ACKNOWLEDGEMENTS**

25 We are grateful to Emi Furusaka, Tomoko Okuda, and Megumi Oka for their
26 expert data management and secretarial assistance, and to all the members of
27 the transplant teams at Kyoto University Hospital for their dedicated care of the

1 patients and donors.

2 This work was supported in part by the Takeda Science Foundation (JK).

3

4

5

6

7

1 References

- 2 1. Tomblyn M, Chiller T, Einsele H, et al. Guidelines for Preventing
3 Infectious Complications among Hematopoietic Cell Transplantation
4 Recipients: A Global Perspective. *Biol Blood Marrow Transplant.*
5 2009;15(10):1143-1238. doi:10.1016/j.bbmt.2009.06.019
- 6 2. Styczynski J, Reusser P, Einsele H, et al. Management of HSV, VZV and
7 EBV infections in patients with hematological malignancies and after SCT:
8 Guidelines from the Second European Conference on Infections in
9 Leukemia. *Bone Marrow Transplant.* 2009;43(10):757-770.
10 doi:10.1038/bmt.2008.386
- 11 3. Ljungman P. CMV infections after hematopoietic stem cell transplantation.
12 *Bone Marrow Transplant.* 2008;42:70-72. doi:10.1038/bmt.2008.120
- 13 4. Chaer F El, Shah DP, Chemaly RF. How I Treat How I treat resistant
14 cytomegalovirus infection in hematopoietic cell transplantation recipients.
15 *Blood.* 2017;128(23):2624-2637. doi:10.1182/blood-2016-06-
16 688432.Before
- 17 5. Kimura SI, Wada H, Sakamoto K, et al. L-index as a novel index to
18 evaluate both the intensity and duration of lymphopenia after allogeneic
19 hematopoietic stem cell transplantation. *Transpl Infect Dis.*
20 2012;14(4):364-373. doi:10.1111/j.1399-3062.2012.00727.x
- 21 6. https://www.jshct.com/guideline/pdf/guideline_CMV_2.pdf. Monogr Vol/31.
22 2011;31.
- 23 7. Takenaka K, Gondo H, Tanimoto K, et al. Increased incidence of
24 cytomegalovirus (CMV) infection and CMV- associated disease after
25 allogeneic bone marrow transplantation from unrelated donors. 1997:241-
26 248.
- 27 8. Kanda Y, Mineishi S, Saito T, et al. Pre-emptive therapy against

- 1 cytomegalovirus (CMV) disease guided by CMV antigenemia assay after
2 allogeneic hematopoietic stem cell transplantation: A single-center
3 experience in Japan. *Bone Marrow Transplant.* 2001;27(4):437-444.
4 doi:10.1038/sj.bmt.1702805
- 5 9. Boeckh BM, Bowden RA, Goodrich JM, Pettinger M, Meyerst JD.
6 Cytomegalovirus Antigen Detection in Peripheral Blood Leukocytes After
7 Allogeneic Marrow Transplantation. 2018;80(5):1358-1364.
- 8 10. Mori T, Okamoto S, Matsuoka S, et al. Risk-adapted pre-emptive therapy
9 for cytomegalovirus disease in patients undergoing allogeneic bone
10 marrow transplantation. 2000;(August 1999):765-769.
- 11 11. Inazawa N, Hori T, Hatakeyama N, et al. Large-scale multiplex
12 polymerase chain reaction assay for diagnosis of viral reactivations after
13 allogeneic hematopoietic stem cell transplantation. *J Med Virol.* 2015.
14 doi:10.1002/jmv.24161
- 15 12. Ljungman P, de la Camara R, Cordonnier C, et al. Management of CMV,
16 HHV-6, HHV-7 and Kaposi-sarcoma herpesvirus (HHV-8) infections in
17 patients with hematological malignancies and after SCT. *Bone Marrow
18 Transplant.* 2008;42(4):227-240. doi:10.1038/bmt.2008.162
- 19 13. Cesaro S, Hirsch HH, Faraci M, et al. Cidofovir for BK Virus–Associated
20 Hemorrhagic Cystitis: A Retrospective Study. *Clin Infect Dis.*
21 2009;49(2):233-240. doi:10.1086/599829
- 22 14. Sakurada M, Kondo T, Umeda M, Kawabata H, Yamashita K, Takaori-
23 Kondo A. Successful treatment with intravesical cidofovir for virus-
24 associated hemorrhagic cystitis after allogeneic hematopoietic stem cell
25 transplantation: A case report and a review of the literature. *J Infect
26 Chemother.* 2016;22(7):495-500. doi:10.1016/j.jiac.2016.01.013
- 27 15. Nagafuji K, Aoki K, Henzan H, et al. Cidofovir for treating adenoviral

- 1 hemorrhagic cystitis in hematopoietic stem cell transplant recipients. *Bone*
2 *Marrow Transplant*. 2004;34(10):909-914. doi:10.1038/sj.bmt.1704682
- 3 16. Storer BE. ESTIMATION OF FAILURE PROBABILITIES IN THE
4 PRESENCE OF COMPETING RISKS : NEW REPRESENTATIONS OF
5 OLD ESTIMATORS. 1999;706(May 1998):695-706.
- 6 17. Gray RJ, Annals T, Sep N. Competing Risk. 1988;16(3):1141-1154.
- 7 18. Fine JP, Gray RJ, Fine JP, Gray RJ. A Proportional Hazards Model for
8 the Subdistribution of a Competing Risk Stable URL :
9 <http://www.jstor.org/stable/2670170> All use subject to
10 <http://about.jstor.org/terms> A Proportional Hazards Model for the
11 Subdistribution of a Competing Risk. 1999;94(446):496-509.
- 12 19. Kanda Y. Investigation of the freely available easy-to-use software “EZR”
13 for medical statistics. *Bone Marrow Transplant*. 2013;48(3):452-458.
14 doi:10.1038/bmt.2012.244
- 15 20. Miyashita N, Endo T, Onozawa M, et al. Risk factors of human
16 herpesvirus 6 encephalitis/myelitis after allogeneic hematopoietic stem
17 cell transplantation. *Transpl Infect Dis*. 2017;19(3):1-10.
18 doi:10.1111/tid.12682
- 19 21. Ogata M, Satou T, Kadota JI, et al. Human herpesvirus 6 (HHV-6)
20 reactivation and HHV-6 encephalitis after allogeneic hematopoietic cell
21 transplantation: A multicenter, prospective study. *Clin Infect Dis*.
22 2013;57(5):671-681. doi:10.1093/cid/cit358
- 23 22. Dulery R, Salleron J, Dewilde A, et al. Early Human Herpesvirus Type 6
24 Reactivation after Allogeneic Stem Cell Transplantation: A Large-Scale
25 Clinical Study. *Biol Blood Marrow Transplant*. 2012;18(7):1080-1089.
26 doi:10.1016/j.bbmt.2011.12.579
- 27 23. Aoki J, Numata A, Yamamoto E, Fujii E, Tanaka M, Kanamori H. Impact

- 1 of Human Herpesvirus-6 Reactivation on Outcomes of Allogeneic
2 Hematopoietic Stem Cell Transplantation. *Biol Blood Marrow Transplant.*
3 2015;21(11):2017-2022. doi:10.1016/j.bbmt.2015.07.022
- 4 24. Pichereau C, Desseaux K, Janin A, et al. The Complex Relationship
5 between Human Herpesvirus 6 and Acute Graft-versus-Host Disease.
6 *Biol Blood Marrow Transplant.* 2012;18(1):141-144.
7 doi:10.1016/j.bbmt.2011.07.018
- 8 25. Jeulin H, Agrinier N, Guery M, et al. Human herpesvirus 6 infection after
9 allogeneic stem cell transplantation: Incidence, outcome, and factors
10 associated with HHV-6 reactivation. *Transplantation.* 2013;95(10):1292-
11 1298. doi:10.1097/TP.0b013e318289958b
- 12 26. Sashihara J, Tanaka-Taya K, Tanaka S, et al. High incidence of human
13 herpesvirus 6 infection with a high viral load in cord blood stem cell
14 transplant recipients. *Blood.* 2002;100(6):2005-2011.
- 15 27. Shimazu Y, Kondo T, Ishikawa T, Yamashita K, Takaori-Kondo A. Human
16 herpesvirus-6 encephalitis during hematopoietic stem cell transplantation
17 leads to poor prognosis. *Transpl Infect Dis.* 2013;15(2):195-201.
18 doi:10.1111/tid.12049
- 19 28. Razonable RR. Infections due to human herpesvirus 6 in solid organ
20 transplant recipients. *Curr Opin Organ Transplant.* 2010;15(6).
21 [https://journals.lww.com/co-](https://journals.lww.com/co-transplantation/Fulltext/2010/12000/Infections_due_to_human_herpesvirus_6_in_solid.2.aspx)
22 [transplantation/Fulltext/2010/12000/Infections_due_to_human_herpesviru](https://journals.lww.com/co-transplantation/Fulltext/2010/12000/Infections_due_to_human_herpesvirus_6_in_solid.2.aspx)
23 [s_6_in_solid.2.aspx.](https://journals.lww.com/co-transplantation/Fulltext/2010/12000/Infections_due_to_human_herpesvirus_6_in_solid.2.aspx)
- 24 29. Osarogiagbon RU, Defor TE, Weisdorf MA, Erice A, Weisdorf DJ. CMV
25 antigenemia following bone marrow transplantation: Risk factors and
26 outcomes. *Biol Blood Marrow Transplant.* 2000;6(3):280-288.
27 doi:10.1016/S1083-8791(00)70010-3

- 1 30. George B, Kerridge IH, Gilroy N, et al. A risk score for early
2 cytomegalovirus reactivation after allogeneic stem cell transplantation
3 identifies low-, intermediate-, and high-risk groups: Reactivation risk is
4 increased by graft-versus-host disease only in the intermediate-risk
5 group. *Transpl Infect Dis.* 2012;14(2):141-148. doi:10.1111/j.1399-
6 3062.2011.00706.x
- 7 31. Quintela A, Escuret V, Roux S, et al. HHV-6 infection after allogeneic
8 hematopoietic stem cell transplantation: From chromosomal integration to
9 viral co-infections and T-cell reconstitution patterns. *J Infect.*
10 2016;72(2):214-222. doi:10.1016/j.jinf.2015.09.039
- 11 32. de Koning C, Admiraal R, Nierkens S, Boelens JJ. Human herpesvirus 6
12 viremia affects T-cell reconstitution after allogeneic hematopoietic stem
13 cell transplantation. *Blood Adv.* 2018;2(4):428-432.
14 doi:10.1182/bloodadvances.2017012724
- 15 33. Eliassen E, Di Luca D, Rizzo R, Barao I. The interplay between natural
16 killer cells and human herpesvirus-6. *Viruses.* 2017;9(12):14-16.
17 doi:10.3390/v9120367

18
19

1 **Figure legends**

2 **Figure 1 Cumulative incidence of HHV-6 reactivation**

3 **Figure 2 Cumulative incidence of CMV antigenemia**

Table1

Table1 Patient characteristics (Day15)

		Group by Lymphocyte-AUC by day15 (n=286)	Low, Middle AUC(<230/μl) (n=189)		High AUC (≥230/μl) (n=97)		Variance
		Total n ^{*1}	value		value		P-Value
			n	% ^{*2}	n	%	
Age ^{*3} median(range)		51 (17-68)	52 (18-68)		50 (17-68)		0.581
Gender	Male	168	108	57.1	60	61.9	0.526
	Female	118	81	42.9	37	38.1	
Donor source	Sibling	78	51	27.0	27	27.8	<0.05
	Unrelated BM	129	99	52.4	30	30.9	
	Unrelated CB	79	39	20.6	40	41.2	
Disease	AML/MDS	172	115	60.8	57	58.8	0.838
	ALL/other leukemias	61	41	21.7	20	20.6	
	Malignant lymphoma	45	25	13.2	20	20.6	
	Aplastic anemia	8	8	4.2	0	0.0	
Disease status	CR	130	79	41.8	51	52.6	0.068
	non CR	156	110	58.2	46	47.4	
Conditioning intensity	Myeloablative	149	101	53.4	48	49.5	0.535
	Reduced intensity	137	88	46.6	49	50.5	
GVHD prophylaxis	CI	23	7	3.7	16	16.5	<0.05
	CI+MMF	56	28	14.8	28	28.9	
	CI+MTX	161	119	63.0	42	43.3	
	CI+MMF+MTX	44	34	18.0	10	10.3	
	regimens containing ATG	2	1	0.5	1	1.0	

*1n indicates the number of patients with each characteristics

*2% indicates the percentage of patients in each group

*3Age indicates patients' age at transplantation

Calcinerin inhibitors include Tacrolimus and Cyclosporin

Abbreviation: AML, acute myeloid leukemia; MDS, myelodysplastic syndrome; ALL, acute lymphoblastic leukemia; CR, complete remission; BM, bone marrow; CB, cord blood; GVHD, graft-versus-host disease; CI, Calcinerin inhibitor; MMF, mycophenolate mofetil; MTX, methotrexate; ATG, antithymocyte globulin

Table2

Table2 Patient characteristics (Day30)

Group by Lymphocyte-AUC by day30 (n=283)		Low AUC(<2710/μl) (n=93)			Middle AUC (≥2710/μl, <5250/μl) (n=93)		High AUC (≥5250/μl) (n=97)		Variance
		Total	value		value		value		
		n ^{*1}	n	% ^{*2}	n	%	n	%	P-Value
Age ^{*3} median(range)		51 (17-68)	52 (20-68)		51 (18-68)		49 (17-68)		0.581
Gender	Male	117	29	31.2	44	47.3	44	45.4	0.050
	Female	166	64	68.8	49	52.7	53	54.6	
Donor source	Sibling	77	19	20.4	22	23.7	36	37.1	<0.05
	Unrelated BM	128	30	32.3	44	47.3	54	55.7	
	Unrelated CB	78	44	47.3	27	29.0	7	7.2	
Disease	AML/MDS	169	59	63.4	56	60.2	54	55.7	0.208
	ALL/other leukemias	61	15	16.1	22	23.7	24	24.7	
	Malignant lymphoma	45	13	14.0	14	15.1	18	18.6	
	Aplastic anemia	8	6	6.5	1	1.1	1	1.0	
Disease status	CR	130	33	35.5	52	55.9	45	46.4	<0.05
	non CR	153	60	64.5	41	44.1	52	53.6	
Conditioning intensity	Myeloablative	146	47	50.5	48	51.6	51	52.6	0.908
	Reduced intensity	137	46	49.5	45	48.4	46	47.4	
GVHD prophylaxis	CI	22	10	10.8	10	10.8	2	2.1	<0.05
	CI+MMF	55	27	29.0	20	21.5	8	8.2	
	CI+MTX	161	45	48.4	44	47.3	72	74.2	
	CI+MMF+MTX	43	11	11.8	19	20.4	13	13.4	
	regimens containing ATG	2	0	0.0	0	0.0	2	2.1	
GVHD (by day30) grade at onset	I	18	4	4.3	6	6.5	8	8.2	0.577
	II	51	13	14.0	22	23.7	16	16.5	
	III	10	3	3.2	2	2.2	5	5.2	
	IV	2	0	0.0	2	2.2	0	0.0	

*1n indicates the number of patients with each characteristics

*2% indicates the percentage of patients in each group

*3Age indicates patients' age at transplantation

Calcineurin inhibitors include Tacrolimus and Cyclosporin

Abbreviation: AML, acute myeloid leukemia; MDS, myelodysplastic syndrome; ALL, acute lymphoblastic leukemia; CR, complete remission; BM, bone marrow; CB, cord blood; GVHD, graft-versus-host disease; CI, Calcineurin inhibitor; MMF, mycophenolate mofetil; MTX, methotrexate; ATG, antithymocyte globulin

Table 3 Univariate and multivariate analysis of HHV-6 reactivation

Variables		Univariate analysis			Multivariate analysis		
		HR	95% CI	P-Value	HR	95% CI	P-Value
Age ^{*1}	<50	1.00		reference			
	≥50	0.63	(0.32-1.28)	0.201			
Gender	Male	1.00		reference			
	Female	1.29	(0.95-1.75)	0.102			
Donor source	Sibling	1.00		reference	1.00		reference
	Unrelated BM	0.54	(0.21-1.40)	0.204			
	Unrelated CB	4.53	(2.17-9.45)	<0.001	3.05	(1.38-6.72)	0.006
Disease	AML/MDS	1.00		reference	1.00		reference
	ALL/other leukemias	0.78	(0.36-1.70)	0.527			
	Malignant lymphoma	1.12	(0.52-2.42)	0.779			
	Aplastic anemia	3.24	(1.29-8.16)	0.012	5.34	(2.38-12.00)	<0.001
Disease status	CR	1.00		reference			
	non CR	0.65	(0.39-1.08)	0.096			
Conditioning regimen	Myeloablative	1.00		reference			
	Reduced intensity	0.91	(0.52-1.60)	0.749			
GVHD prophylaxis	CI	1.00		reference	1.00		reference
	CI+MMF	2.37	(0.91-6.16)	0.077			
	CI+MTX	0.26	(0.09-0.75)	0.013	0.35	(0.15-0.84)	0.019
	CI+MMF+MTX	0.67	(0.21-2.12)	0.493			
	regimens containing ATG	2.07	(0.35-12.33)	0.421			
Lymphocyte-AUC group	Low,Middle-AUC ^{*2}	1.00		reference	1.00		reference
	High-AUC ^{*3}	2.44	(1.40-4.23)	0.002	1.83	(1.01-3.34)	0.048

^{*1}Age indicates patients' age at transplantation

^{*2}Low,Middle-AUC indicates group of patients with lymphocyte-AUC under 230/μl

^{*3}High-AUC indicates group of patients with lymphocyte-AUC of 230/μl or over

Calcinerin inhibitors include Tacrolimus and Cyclosporin

Abbreviation: AML, acute myeloid leukemia; MDS, myelodysplastic syndrome; ALL, acute lymphoblastic leukemia; CR, complete remission; BM, bone marrow; CB, cord blood; GVHD, graft-versus-host disease; CI, Calcinerin inhibitor; MMF, mycophenolate mofetil; MTX, methotrexate; ATG, antithymocyte globulin

Table 4 Univariate and multivariate analysis of CMV antigenemia

Variables		Univariate analysis			Multivariate analysis		
		HR	95% CI	P-Value	HR	95% CI	P-Value
Age* ¹	<50	1.00		reference	1.00		reference
	≥50	1.46	(1.01-2.09)	0.042	1.55	(1.08-2.21)	0.017
Gender	Male	1.00		reference			
	Female	0.90	(0.77-1.08)	0.273			
Donor source	Sibling	1.00		reference			
	Unrelated BM	1.08	(0.71-1.63)	0.731			
	Unrelated CB	1.47	(0.22-1.78)	0.075			
Disease	AML/MDS	1.00		reference			
	ALL/other leukemias	1.30	(0.84-2.00)	0.237			
	Malignant lymphoma	0.93	(0.54-1.60)	0.800			
	Aplastic anemia	1.13	(0.41-3.07)	0.817			
Disease status	CR	1.00		reference			
	non CR	1.01	(0.75-1.35)	0.960			
Conditioning regimen	Myeloablative	1.00		reference			
	Reduced intensity	0.98	(0.70-1.36)	0.882			
GVHD prophylaxis	CI	1.00		reference			
	CI+MMF	0.83	(0.44-1.54)	0.549			
	CI+MTX	0.66	(0.37-1.17)	0.154			
	CI+MMF+MTX	1.06	(0.56-2.00)	0.847			
	regimens containing ATG	1.14	(0.67-1.93)	0.618			
aGVHD by	no	1.00		reference	1.00		reference
	occurrence	1.94	(1.37-2.75)	<0.001	2.21	(1.49-3.29)	<0.001
Lymphocyte-AUC group	Low-AUC* ²	1.00		reference	1.00		reference
	Middle-AUC* ³	1.27	(0.87-1.84)	0.212	1.13	(0.74-1.73)	0.560
	High-AUC* ⁴	0.63	(0.40-0.98)	0.041	0.61	(0.37-1.01)	0.052

*¹Age indicates patients' age at transplantation

*²Low-AUC indicates group of patients with lymphocyte-AUC under 2710/μl

*³Middle-AUC indicates group of patients with lymphocyte-AUC of 2710/μl or over and under 5250/μl

*⁴High-AUC indicates group of patients with lymphocyte-AUC of 5250/μl or over

Calcinerin inhibitors include Tacrolimus and Cyclosporin

Abbreviation: AML, acute myeloid leukemia; MDS, myelodysplastic syndrome; ALL, acute lymphoblastic leukemia; CR, complete remission; BM, bone marrow; CB, cord blood; GVHD, graft-versus-host disease; CI, Calcinerin inhibitor; MMF, mycophenolate mofetil; MTX, methotrexate; ATG, antithymocyte globulin




Figure 1. Cumulative incidence of HHV-6 reactivation




Figure 2. Cumulative incidence of CMV antigenemia