TITLE:

Black spruce assimilates nitrate in boreal winter

AUTHOR(S):

Koyama, Lina A; Kielland, Knut

CITATION:

Koyama, Lina A ...[et al]. Black spruce assimilates nitrate in boreal winter. *Tree Physiology* 2019, 39(4): 536-543

ISSUE DATE:

2019-04

URL:

<http://hdl.handle.net/2433/241700>

RIGHT:

This is a pre-copyedited, author-produced PDF of an article accepted for publication in 'Tree Physiology' following peer review. The version of record 'Lina A Koyama, Knut Kielland; Black spruce assimilates nitrate in boreal winter, *Tree Physiology*, 39(4) 536-543' is available online at: <https://doi.org/10.1093/treephys/tpy109>; The full-text file will be made open to the public on 21 November 2019 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

1 TITLE

2 Black spruce assimilates nitrate in boreal winter

3 AUTHORS

4 Lina A. Koyama¹5 Knut Kielland²

6 AFFILIATIONS AND ADDRESSES OF THE AUTHORS:

7 1. Graduate School of Informatics, Kyoto University

8 2. Institute of Arctic Biology, University of Alaska Fairbanks,

9 AUTHOR FOR CORRESPONDENCE:

10 Name: Lina A. Koyama

11 Address: Laboratory of Biosphere Informatics,

12 Department of Social Informatics,

13 Graduate School of Informatics, Kyoto University,

14 Kyoto 606-8501 Japan

15 E-mail: linak@bre.soc.i.kyoto-u.ac.jp

16 Phone number: +81-75-753-3297

17 Fax number: +81-75-753-3133

18 RUNNING HEAD

19 Black spruce assimilates nitrate in boreal winter

20

21 ABSTRACT

22 Winter has long been considered a dormant season in boreal forests regarding plant
23 physiological activity such as nutrient acquisition. However, biogeochemical data clearly show that
24 soil can remain unfrozen with substantial rates of nutrient transformation for several weeks
25 following autumn snowfall. Here we examined nitrate (NO_3^- -N) assimilation by black spruce (*Picea*
26 *mariana*) during summer and winter in Interior Alaska to test our hypothesis that this boreal species
27 is able to assimilate NO_3^- -N, even at the very low temperatures typical of early winter. Nitrate
28 reductase activity (NRA) was measured in current year needles and fine roots of black spruce as an
29 indicator of NO_3^- -N assimilation in the summer and winter at two boreal forest sites. NO_3^- -N
30 concentration in the needles and roots were also measured to determine whether NO_3^- -N was
31 available in plant tissue for the enzyme. NRA and NO_3^- -N were detected in needles and roots in the
32 winter as well as the summer. The results of a generalized linear mixed model (GLMM) showed that
33 season had minimal effects on NRA and NO_3^- -N concentration in this species. Additionally, the
34 effect of incubation temperature for the NRA assays was tested at 30 °C and -3 °C for samples
35 collected in the winter. Substantial enzyme activity was detected in winter-collected samples, even
36 in incubations conducted at -3 °C. These results indicate that this dominant tree species in the boreal
37 forests of Interior Alaska, black spruce, has the capacity to assimilate NO_3^- -N below freezing
38 temperatures, suggesting that the physiological activity required for N resource acquisition may
39 extend beyond the typical growing season. Our findings coupled to biogeochemical evidence for
40 high microbial activity under the snow also indicate that winter N acquisition should be taken into
41 account when estimating the annual N budgets of boreal forest ecosystems.

43 KEYWORDS

44 nitrate (NO_3^- -N), nitrate reductase activity (NRA), non-growing seasons, Taiga

INTRODUCTION

Nitrogen is among the most important limiting factors of plant productivity in the boreal forests of Interior Alaska (Yarie and Van Cleve 2006). The annual plant N requirement is only partly supplied by the major N source for plants, soil inorganic N (Valentine et al. 2006, Lisuzzo et al. 2008). In other words, there are marked discrepancies between the current estimates of inputs of inorganic N available to plants (via N-mineralization, N-fixation, and dry/wet deposition) and their annual N uptake rates or requirements (Kielland 2001, Valentine et al. 2006, Lisuzzo et al. 2008). The direct uptake of N in the form of amino acids further narrows the growing season gap between supply and demand (Persson and Näsholm 2001, Kielland et al. 2006a). Moreover, some of the discrepancies may be explained by uptake during the shoulder seasons, i.e., the period between growing season and mid winter, which has been largely ignored in the estimation of N flux including above-mentioned works. Kielland et al. (2006b) used over-winter incubations to demonstrate that boreal forest soils have a substantial capacity for N mineralization during the cold season and concluded that conventional measures have greatly underestimated the annual flux of inorganic N because they have been restricted to the growing season (May–September). In this study, we focused on N use by boreal plants during the winter to discuss the possible contribution of winter for N acquisition by plants.

Two inorganic forms of N (nitrate [NO_3^- -N] and ammonium [NH_4^+ -N]) in soils are available to most plant species. NO_3^- -N assimilation has been investigated in a variety of plant species using nitrate reductase activity (NRA) as an index (e.g., Smirnov et al. 1984, Gebauer et al. 1988). Nitrate reductase (NR) catalyzes the reduction of NO_3^- -N to NO_2^- -N, which is the first and rate-limiting step of plant NO_3^- -N assimilation. Measurements of NRA can be used to estimate plant NO_3^- -N use without disturbing the soil, which is not the case for experimental manipulations, such as the application of ^{15}N tracers. Numerous studies have shown that both external environmental changes and internal physiological shifts in plants can cause temporal changes in NO_3^- -N assimilation (cf. Högberg et al. 1986, Gebauer et al. 1987, Schmidt et al. 1991, Högberg et al. 1992, Stadler and Gebauer 1992, Ohlson and Högbom 1993, Pearson and Ji 1994, Troelstra et al. 1995, Koyama et al. 2008). However, these studies measured NRA during the growing seasons of the

species examined. NRA has rarely been examined in the winter, although Koyama et al. (2008) investigated the nitrate assimilation of a temperate, evergreen *Quercus* species during the winter, and the NRA of several evergreen coniferous species growing in temperate forests actually appears to be higher in the winter than in the summer (M. Ueda pers. comm.).

In boreal forests, where winter air temperatures can fall to below -40 °C, the seasonal patterns of enzyme activity may differ from warmer regions. On the other hand, Kielland et al. (2006b) demonstrated that soils from black spruce stands exhibited significant nitrification in late winter to spring. Hence, any potential to take up and/or assimilate NO_3^- -N at very low temperatures in boreal tree species will influence the current estimates of N flux in these cold, high latitude forests. Furthermore, recent global changes in climate could make uptake/assimilation activity during the winter of even greater relevance in nutrient budget calculations.

In this study, we compared winter and summer NRA and NO_3^- -N concentrations in the needles and fine roots of black spruce (*Picea mariana* (Mill.) Britton, Sterns and Poggenb.) in a boreal forest located in Interior Alaska, USA. We also simulated ambient soil temperatures during wintertime enzyme incubations to determine the extent to which enzyme activity is maintained under near-natural conditions in an attempt to explore the idea that black spruce can maintain physiological activities at low temperatures.

MATERIALS AND METHODS

Study Site

The study was conducted in late-successional black spruce forests in Interior Alaska, USA (64°52'N, 147°50'W). During the study years, the temperature at the nearby weather station ranged from -39.13–33.02 °C, averaging 0.37 °C (Fig. 1; Van Cleve et al. 2016). The average annual precipitation at the site was 437 mm, of which 35 % fell as snow. The ground was covered with snow from mid-October through late April. The mean annual NO_3^- and total inorganic N deposition in this site from 2009 to 2016 were 0.54 ± 0.10 and 0.99 ± 0.34 kg-N ha⁻¹, respectively, which are lower by an order of magnitude than the US average (National Atmospheric Deposition Program (NRSP-3) 2017; <http://nadp.sws.uiuc.edu>).

Sample Collection and Laboratory Analysis

Experiment 1: effects of season, site, and tissue on NRA and NO_3^- -N concentration

The first experiment to compare the effects of seasons, sites, and tissues was conducted in two late-successional black spruce forests near the campus of the University of Alaska, Fairbanks, and two sites were located approximately 2 km away from each other (site 1: 64°51'36"N, 147°53'12"W and site 2: 64°51'49"N, 147°50'43"W). Plant sample collection was conducted at site 1 in July (summer) and December (winter) in 2009 and at site 2 in July (summer) 2015 and November (winter) 2016. At the time of summer sampling, the air temperatures in 2009 and 2015 were approximately 17 °C and 15 °C and the surface soil temperatures were approximately 13 °C and 14 °C, respectively. During winter sampling, the air temperatures were approximately -20 °C and -10 °C in 2009 and 2016, and the surface soil temperatures were about -2 °C and -1 °C, respectively. The snow depth reached approximately 14 cm and 8 cm in 2009 and 2016, respectively.

Current year needles and fine roots (diameter: < 2 mm) of black spruce were collected in the summer (July 2009 and 2015) and winter (December 2009 and November 2016) from five mature trees. These needles and roots were used in NRA and NO_3^- -N concentration assays.

Experiment 2: effects of incubation temperature on NRA

In the second experiment to test the effects of incubation temperatures on NRAs, both needle and root samples from site 1 were incubated at two temperatures: 30 °C and -3 °C. The incubation at 30 °C provided optimal conditions for enzymatic catalysis. Incubation at -3 °C simulated the soil temperature on the day of sampling. We did not run tests at air temperature on the day of sampling (-20 °C) because this was well below the freezing temperature of the incubation buffers.

The assay of $\text{NRA}(+\text{NO}_3)$, $\text{NRA}(-\text{NO}_3)$ and NO_3^- -N concentration

We measured two types of NRA as indices of plant NO_3^- -N use: $\text{NRA}(+\text{NO}_3)$ and

NRA($-\text{NO}_3$). NRA($+\text{NO}_3$) is a measure of the nitrate reduction capacity with a non-limiting nitrate supply; NRA($-\text{NO}_3$) is the nitrate reduction rate of nitrate absorbed by plants, which is considered to be the closest approximation of the *in situ* NO_3^- -N assimilation rate (Thomas and Hilker, 2000). Both NRA assays were conducted with modified versions of the Jaworski procedure (Jaworski, 1971, Thomas and Hilker, 2000, Koyama and Kielland, 2011). NRA($+\text{NO}_3$) was measured as the rate of nitrite (NO_2^- -N) production in an incubation buffer containing a non-limiting concentration of NO_3^- -N. NRA($-\text{NO}_3$) was determined in parallel measurements using an incubation buffer without additional NO_3^- -N, which allowed us to examine the relative magnitude of *in situ* NO_3^- -N assimilation.

Current year needles and fine roots (diameter: < 2 mm) were sampled from five mature black spruce trees on each sampling occasion. Needle samples were collected from the surface of the crown at various heights, and the sampled needles were mostly exposed to adequate light due to low canopy density (Fujino pers. comm.). Root samples were washed in tap water and then in deionized water to remove the soil. Fine root samples were randomly collected from root tips, thus possibly ectomycorrhizal (ECM) fungal tissue were mixed with spruce root tissue. Approximately 100 mg (fresh weight) of needles and roots were cut into small fragments (each *ca.* 2 mm long) and transferred to test tubes. The incubation buffer (5 mL) was added to the needles and roots, and the tube contents were vacuum infiltrated. The composition of the incubation buffer for NRA($+\text{NO}_3$) was as follows: $0.1 \text{ mol L}^{-1} \text{ KNO}_3$, $0.1 \text{ mol L}^{-1} \text{ KH}_2\text{PO}_4$, 1.5 % 1-propanol; the pH was adjusted to *ca.* 7.5 using an NaOH solution. The concentration of NO_3^- -N was determined by a preliminary optimization process in which different concentrations of NO_3^- -N were added to the incubation buffer. A supply of varying NO_3^- -N concentration ranging from 0.00 mM to 0.25 mM in incubation buffer yielded a peak of NRA at 0.10 mM of NO_3^- -N supply (Appendix 1). The incubation buffer for NRA($-\text{NO}_3$) contained all of the reagents other than KNO_3 . The samples were incubated for 1 h in darkness, and NO_2^- -N concentration in the incubation buffer was measured at the end point. Before the measurement, enzyme activity was terminated by placing the sample vials in hot water (>80 °C). The concentration of NO_2^- -N in the incubation buffer was measured colorimetrically following diazotization (Keeney and Nelson, 1982). The confounding effects of plant pigments were

accounted for by subtracting the absorbance of controls to which N-naphthylethylene diamine dihydrochloride was not added (Gebauer et al. 1998). A fraction of each leaf sample was oven-dried at 105 °C and then weighed to calculate the activity per unit dry weight.

For tissue NO_3^- -N concentration measurements, the aliquots of needle and root samples were dried and ground. Approximately 100 mg of ground sample was extracted with 10 mL deionized water for 1 h at 45 °C. The extract was filtered, and the concentration of NO_3^- -N in the extract was colorimetrically analyzed in an AutoAnalyzerIII (BLTec, Osaka, Japan). Plant pigments in extracts might cause an overestimation of NO_3^- -N concentration, and other unknown compounds in the extracts might inhibit the reduction of NO_3^- -N to NO_2^- -N, which is colorimetrically measured in the AutoAnalyzerIII (data not shown). Again, the confounding effects of plant pigments were taken into account by subtracting the absorbance of controls to which N-naphthylethylene diamine dihydrochloride was not added. In addition, a standard addition method was applied to compensate for the effects of pigments and other compounds in the extract as necessary when the sample composition was unknown or complex and might affect the analytical signal (Harris 2007). In this method, standard solutions of known concentrations were added to each extract, and from the increases in signal (i.e., absorbance), the concentration in the original extract was calculated.

Statistical Analysis

For experiment 1, we fitted a generalized linear mixed model (GLMM) with a gamma distribution to evaluate the effects of Season (summer or winter), Site (site 1 or site 2), and Tissue (needle or root) on $\text{NRA}(\text{+NO}_3)$, $\text{NRA}(\text{-NO}_3)$, or NO_3^- concentrations, following a Shapiro-Wilk test to test the normality of data. Five individual trees were included as random effects. Two of $\text{NRA}(\text{+NO}_3)$ and seven of NO_3^- concentrations data were below the detection limit; to fit the model with a gamma distribution, 1×10^{-10} and 1×10^{-6} were substituted for zero for these samples that presented values below the detection limit, respectively. All possible subsets of the explanatory variables and their interactions were compared with Akaike Information Criterion (AIC) for each of the response variables, $\text{NRA}(\text{+NO}_3)$, $\text{NRA}(\text{-NO}_3)$, or NO_3^- -N concentrations.

For experiment 2, we fitted a generalized linear mixed model (GLMM) with a gamma

distribution to evaluate effects of the variables Incubation temperature (-3°C or 30°C) and Tissue (needle or root) on $\text{NRA}(+\text{NO}_3)$ or $\text{NRA}(-\text{NO}_3)$. Five individual trees were included as random effects. All possible subsets of the explanatory variables and their interactions were compared with Akaike Information Criterion (AIC) for each of response variable, $\text{NRA}(+\text{NO}_3)$ or $\text{NRA}(-\text{NO}_3)$.

It should be noted that in both experiments, the link function ‘inverse’ was applied for GLMM with gamma distribution, and consequently a positive value of the coefficient implies a negative effect of the explanatory variable on the response variable. It is worth noting that all of the regression coefficients can be compared to each other, although they were not standardized, as all of the variables are categorical variables with an equal number of categories: two in each experiment. All statistical analyses were conducted using the statistical platform R (ver. 3.3.3; <http://www.R-project.org>), and the lme4 package (version 1.1-13) was used for fitting GLMM.

RESULTS

Experiment 1: effects of season, site, and tissue on NRA and NO_3^- -N concentration

Both $\text{NRA}(+\text{NO}_3)$ and $\text{NRA}(-\text{NO}_3)$ were detected in the needles and fine roots of black spruce in experiment 1 (Fig. 2a, b). The best performing model fitted for $\text{NRA}(+\text{NO}_3)$ had Season, Site, and their interaction as explanatory variables. However, only Site had the coefficient with a P-value lower than 0.05, indicating that zero was not included within the 95 % Wald confidence interval (CI) of estimated coefficient (Table 1). The best performing model fitted for $\text{NRA}(-\text{NO}_3)$ had all of the explanatory variables and their interactions except the interaction of Season $\times$ Site $\times$ Tissue. However, the coefficient for the Season and the interaction Season $\times$ Tissue exhibited P-values higher than 0.05, indicating that zero was included within the 95 % Wald CI of estimated coefficients.

NO_3^- -N was also detected in most needle and fine root samples (Fig. 2c). The best performing model fitted for NO_3^- -N concentration had Season, Tissue, and their interaction as explanatory variables. However, Season and Tissue had a P-value higher than 0.05, indicating that zero was included within the 95 % Wald CI of estimated coefficients.

Experiment 2: effects of incubation temperature on NRA

Both NRA(+NO₃) and NRA(-NO₃) were detected in current year needles and fine roots, even at the low incubation temperature (-3 °C; Fig. 3). Both the Incubation temperature and Tissue were selected for the best performing model fitted for NRA(+NO₃), but their interaction was not (Table 2). Moreover, both the coefficient for the Incubation temperature and Tissue showed P-values lower than 0.05, indicating that zero was not included within the 95 % Wald CI of estimated coefficient. On the other hand, the best performing model for NRA(-NO₃) had only Incubation temperature as a coefficient, and the P-value for that was higher than 0.05, indicating that zero was included within the 95 % Wald CI of estimated coefficient.

DISCUSSION

Nitrate Assimilation of Black Spruce in Winter and Summer

Winter is generally considered to be a season of dormancy in boreal forests due to the extremely low temperatures, reduced light intensity, and short photoperiods. However, we have demonstrated that black spruce trees in Interior Alaska are able to assimilate NO₃⁻-N in the winter as well as in the summer (Fig. 2), indicating that, in the winter, (i) black spruce induced NR and (ii) NO₃⁻-N was available for NR.

The disparity between our findings and the accepted winter dormancy concept could be attributed to a *de novo* induction of the enzyme during the storage time in our experiments. However, the *in vivo* NRA was measured under the premise that NR had not been induced *de novo* during the storage period after sample collection. This premise was based on the known light requirement for NR induction (Lillo et al. 2004); our samples were stored in complete darkness. Furthermore, Högberg et al. (1986) found that the shoot NRA of *Deschampsia flexuosa* declined for the first 30 min of storage and remained stable thereafter. Thus, we suggest that NR was not newly induced in our detached samples during storage, and that the NRA detected by our measurements was not likely to be the result of artificially inflated rates of enzyme induction following sample collection.

The results of GLMM fitting and model selection showed that Season had no significant effect on NRA(+NO₃), NRA(-NO₃), or NO₃⁻-N concentration (Table 1). Site and the interaction

between Season $\times$ Site were selected as effective variables in the best models for both of NRA(+NO₃) and NRA(-NO₃) (Table 1). In addition, Tissue and the interaction between Site $\times$ Tissue were also selected in the best model for NRA(-NO₃). On the other hand, only the interaction between Season $\times$ Tissue was selected as a variable influencing NO₃⁻-N concentration. Taken together, these results indicated that season was not a significant factor affecting NO₃⁻-N use by black spruce. Because newly expanding leaves contain higher concentrations of N than fully expanded leaves, winter buds of temperate evergreen species most likely receive N transported from old tissues (Silla and Escudero 2003, Koyama et al. 2008). Consequently, we surmise that the wintertime acquisition of N by the needles of black spruce may play a role in the preparation of additional N sources for the new needles that flush in late spring in these sites.

Lambers et al. (2008) demonstrated that the NRA shoot/root ratio generally increases with NO₃⁻-N availability in temperate and subtropical species. Our results showed that black spruce assimilate nitrate in their current year needles. Some previous studies revealed prior assimilation of nitrate in the roots of coniferous species (Peuke and Tischner 1991, Gebauer and Schulze 1997, Yao et al. 2011), and our results were contrary to these observations. Assuming soil NO₃⁻-N availability was lower at site 2 based on the site differences of NRAs, the results were consistent with the relationship between the allocation of NRA and soil NO₃⁻-N availability in temperate and subtropical species (Lambers et al. 2008), because both NRA(+NO₃) and NRA(-NO₃) were higher in roots than in needles at site 2.

Effects of Incubation Temperature on Winter NRA

NRA(+NO₃) was detected even at the low incubation temperature, -3 °C, both in needles and roots (Fig. 3), although the low incubation temperature significantly reduced the activity in comparison with the samples incubated at the high temperature (30 °C; Table 2). Roots showed lower NRA(+NO₃) than needles, and this allocation pattern did not differ by the incubation temperature. On the other hand, neither incubation temperature nor tissue influenced NRA(-NO₃), implying the low incubation temperature did not inhibit NRA(-NO₃). Thus, the enzyme in winter needles is clearly capable of catalyzing NO₃⁻-N reduction at very low temperatures.

Early studies attempted to optimize the incubation conditions for *in vivo* NRA assays (Nicholas et al. 1976, Al Ghabi and Hipkin 1984, Gebauer et al. 1984). Optimal temperatures were found to be in the range 28–33 °C (Sym, 1984, Högberg et al. 1992), and even higher optimal temperatures (40–50 °C) were reported for some crop species (Chopra, 1983). Högberg et al. (1992) showed that NRA increased with temperature, reaching an optimum at approximately 25 °C. At the coldest incubation temperature they applied (approximately 0 °C), the NRA was very low (Högberg et al. 1992). However, these results were obtained from temperate species and/or herbaceous taxa, such as barley (*Hordeum vulgare*) and *D. flexuosa*. No trials were conducted on boreal evergreen tree species. Furthermore, earlier experiments were conducted during the growing season, but never in winter. The distinct NR responses to temperature in boreal species may well represent an adaptation to cold climates.

We tested only two temperatures in our study, which did not allow us to examine the functional responses of the NR enzyme to temperature. We were also unable to measure enzyme activity in the needle samples at ambient winter air temperatures (-20 °C), which would have frozen the incubation buffer. Accordingly, we cannot rule out the possibility that we overestimated the activity of the enzyme in winter needle samples. Nevertheless, our study has reduced the level of NRA overestimation attributable to the conventional incubation temperature (30 °C).

Ecological Implications

Based on the results showing the capacity of black spruce to use NO_3^- -N, we conclude that this species is able to assimilate NO_3^- -N in the winter. In earlier studies that showed significant species difference in the capacity to assimilate nitrate, coniferous species or gymnosperms were considered to have low capacities for nitrate assimilation (Smirnoff et al. 1984, Gebauer et al. 1998, Hayashi-Tang et al. 2012), and the our results were consistent with these earlier studies. However, considering that black spruce maintained the capacity to assimilate nitrate in winter, this observation indicates that they are capable of using nitrate for a longer period than deciduous species.

We were not able to estimate the magnitude of NO_3^- -N uptake in winter from our results of NRA and NO_3^- -N concentration, because NO_3^- -N, unlike NH_4^+ -N can be stored in plant tissues. It is

possible that the plants had absorbed and stored NO_3^- -N during previous seasons and then subsequently assimilated the stored ions during the following winters. Our results showed lower root NO_3^- -N concentration in the winter than in the summer (Fig. 2c), and this may be a consequence of plant usage of internally stored NO_3^- in winter. The quantitative evaluation of winter N acquisition to the whole N budget requires further investigation.

Kielland et al. (2006b) suggested that the restriction of soil process measurements to the growing season greatly underestimated the annual flux of soil N in Interior Alaska. Our data corroborate this viewpoint; the N assimilation that we measured during the winter strongly indicated that the annual N budgets of boreal ecosystems should be reexamined.

The assimilation of NO_3^- -N is an energy consuming process (Bloom et al. 1992). When excess light energy is available beyond that required for carbon assimilation, it may be used for NO_3^- -N assimilation, thereby reducing the damaging effects of photoinhibition. The putative winter NO_3^- -N assimilation of boreal black spruce may function as a sink for the surplus light energy absorbed by photosynthetic pigments. This proposal is certainly worthy of further exploration, especially considering climate change may alter the relationship between temperature and light condition.

New information on plant physiological performance during winter, such as photosynthesis (Miyazawa and Kikuzawa 2004, Saarinen et al. 2016) and N use (Koyama et al. 2008, Onipchenko et al. 2009, Ueda et al. 2010) is relevant to current considerations of recent climate change (Makoto et al. 2014, Sanders-Demott and Templer 2017). The effects have generally been considered in terms of the direct influence of higher temperatures, changes in habitat availability, and extensions of plant growth periods (Walther et al. 2002, Cleland et al. 2007, Bokhorst et al. 2008, Miller-Rushing and Primack 2008, Polgar and Primack 2011). With the current lack of information on the responses of plant NRA to the range of temperatures during the boreal winter, we are not in a position to estimate the influence of shorter and warmer winters on plant N acquisition. However, our data clearly show that the influence of a changing winter climate on ecosystem N budgets should be taken into account in considerations of the effects of climate warming.

325 FUNDING

326 This research was partly supported by a Grant-in-Aid for Young Scientists (B) from the
327 Japan Society for the Promotion of Science (No. 21780149) and a Grant-in-Aid for Young Scientists
328 (A) from the Japan Society for the Promotion of Science (No. 25712017) awarded to LAK.

329

330 ACKNOWLEDGEMENTS

331 We would like to thank Drs. N. Tokuchi and M. Ueda for inspiring us to conduct this study.
332 We would also like to thank Mr. K. Olson and Ms. Audrey Mutschlecner for their assistance with
333 our field and laboratory work.

334 REFERENCES

- 335 Al Gharbi A, Hipkin CR (1984) Studies on nitrate reductase in British angiosperms I. A comparison
336 of nitrate reductase activity in ruderal, woodland-edge and woody species. *New Phytol*
337 97:629-639.
- 338 Bloom AJ, Sukrapanna SS, Warner RL (1992) Root respiration associated with ammonium and
339 nitrate absorption and assimilation by barley. *Plant Physiol.* 99:1294-1301.
- 340 Bokhorst S, Bjerke J, Bowles F, Melillo J, Callaghan T, Phoenix G (2008) Impacts of extreme
341 winter warming in the sub-Arctic: growing season responses of dwarf shrub heathland.
342 *Global Change Biol.* 14:2603-2612.
- 343 Chopra RK (1983) Effects of Temperature on the In vivo Assay of Nitrate Reductase in some C₃ and
344 C₄ Species. *Ann Bot.* 51:617-620.
- 345 Cleland EE, Chuine I, Menzel A, Mooney HA, Schwartz MD (2007) Shifting plant phenology in
346 response to global change. *Trends Ecol Evol.* 22:357-365.
- 347 Gebauer G, Hahn G, Rodenkirchen H, Zuleger M (1998) Effects of acid irrigation and liming on
348 nitrate reduction and nitrate content of *Picea abies* (L.) Karst. and *Oxalis acetosella* L. *Plant*
349 *Soil.* 199:59-70.
- 350 Gebauer G, Melzer A, Rehder H (1984) Nitrate content and nitrate reductase activity in *Rumex*
351 *obtusifolius*: 1. Differences in organs and diurnal changes. *Oecologia.* 63:136-142.
- 352 Gebauer G, Rehder H, Wollenweber B (1988) Nitrate, nitrate reduction and organic nitrogen in
353 plants from different ecological and taxonomic groups of Central Europe. *Oecologia.*
354 75:371-385.
- 355 Gebauer G, Schuhmacher MI, Krstic B, Rehder H, Ziegler H (1987) Biomass production and nitrate
356 metabolism of *Atriplex hortensis* L. (C₃ plant) and *Amaranthus retroflexus* L. (C₄ plant) in
357 cultures at different levels of nitrogen supply. *Oecologia.* 72:303-314.
- 358 Gebauer G, Schulze E -D, (1997) Nitrate nutrition of Central European forest trees. In:
359 Rennenberg H, Eschrich W, Ziegler H (eds) *Trees- Contributions to Modern Tree*
360 *Physiology.* Backhuys Publishers, Leiden, Netherlands, pp 273-291.

- 361 Harris D (2007) Quality assurance and calibration methods: Standard addition. In: Harris D (eds)
362 Quantitative Chemical Analysis. W. H. Freeman and Company, New York, pp 87-90.
- 363 Hayashi-Tang, M., S. Porder and G.M. Lovett. (2012) Species differences in nitrate reductase
364 activity are unaffected by nitrogen enrichment in northeastern US forests. *Forest Ecol Manag.*
365 275:52-59.
- 366 Högberg P, Granström A, Johansson T, Lundmark-Thelin A, Näsholm T (1986) Plant nitrate
367 reductase activity as an indicator of availability of nitrate in forest soils. *Can J Forest Res.*
368 16:1165-1169.
- 369 Högberg P, Högbom L, Näsholm T (1992) Shoot nitrate reductase activities of field-layer species in
370 different forest types. II. Seasonal variation and effects of temperature. *Scand J Forest Res.*
371 7:1-14.
- 372 Jaworski EG (1971) Nitrate reductase assay in intact plant tissues. *Biochem Bioph Res Co.*
373 43:1274-1279.
- 374 Keeney DR, Nelson DW (1982) Nitrogen - Inorganic forms. In: Page AL, Miller RH, Keeney DR
375 (eds) *Methods of Soil Analysis Part 2*. ASA and SSSA, Madison, WI, pp 643-698.
- 376 Kielland K (2001) Short-circuiting the nitrogen cycle: Strategies of nitrogen uptake in plants from
377 marginal ecosystems. In: Ae N, Arihara J, Okada K and Srinivasan A (eds) *Plant Nutrient*
378 *Acquisition: New Perspectives*. Springer-Verlag, Berlin. pp 376-398.
- 379 Kielland K, McFarland J, Olson K (2006a) Amino acid uptake in deciduous and coniferous taiga
380 ecosystems. *Plant Soil*. 288:297-307.
- 381 Kielland K, Olson K, Ruess RW, Boone RD (2006b) Contribution of winter processes to soil
382 nitrogen flux in taiga forest ecosystems. *Biogeochemistry*. 81:349-360.
- 383 Koyama L, Kielland K (2011) Plant physiological responses to hydrologically mediated changes in
384 nitrogen supply on a boreal forest floodplain: a mechanism explaining the discrepancy in
385 nitrogen demand and supply. *Plant Soil*. 342:129-139.
- 386 Koyama L, Tokuchi N, Fukushima K, Terai M, Yamamoto Y (2008) Seasonal changes in nitrate use
387 by three woody species: the importance of the leaf-expansion period. *Trees*. 22:851-859.

- 388 Lambers H, Chapin FS, Pons TL (2008) Mineral Nutrition. In: Lambers H, Chapin FS, Pons TL
389 (eds) Plant Physiological Ecology. Springer Science+Business Media, New York, pp
390 255-320.
- 391 Lillo C, Meyer C, Lea U, Provan F, Oltedal S (2004) Mechanism and importance of
392 post-translational regulation of nitrate reductase. *J Exp Bot.* 55:1275-1282.
- 393 Lisuzzo NJ, Kielland K, Jones JB (2008) Hydrologic controls on nitrogen availability in a
394 high-latitude, semi-arid floodplain. *Ecoscience.* 15:366-376.
- 395 Makoto K, Kajimoto T, Koyama L, Kudo G, Shibata H, Yanai Y, Cornelissen JHC (2014) Winter
396 climate change in plant–soil systems: summary of recent findings and future perspectives.
397 *Ecol Res.* 29:593-606.
- 398 Miller-Rushing AJ, Primack RB (2008) Global warming and flowering times in Thoreau's Concord:
399 a community perspective. *Ecology.* 89:332-341.
- 400 Miyazawa Y, Kikuzawa K (2004) Winter photosynthesis by saplings of evergreen broad-leaved
401 trees in a deciduous temperate forest. *New Phytol.* 165:857-866.
- 402 Nicholas JC, Harper JE, Hageman RH (1976) Nitrate Reductase Activity in Soybeans (*Glycine max*
403 [L.] Merr.): I. Effects of Light and Temperature. *Plant Physiol.* 58:731-735.
- 404 Ohlson M, Högbom L (1993) Species-specific dynamics in nitrate reductase activity in coexisting
405 swamp forest plants. *J Ecol.* 81:739-744.
- 406 Onipchenko V, Makarov M, Logtestijn R, Ivanov V, Akhmetzhanova A, Tekeev D, Ermak A,
407 Salpagarova F, Kozhevnikova A, Cornelissen J (2009) New nitrogen uptake strategy:
408 specialized snow roots. *Ecology Letters.* 12:758-764.
- 409 Pearson J, Ji YM (1994) Seasonal variation of leaf glutamine synthetase isoforms in temperate
410 deciduous trees strongly suggests different functions for the enzymes. *Plant Cell Environ.*
411 17:1331-1337.
- 412 Persson J, Nasholm T (2001) Amino acid uptake: a widespread ability among boreal forest plants.
413 *Ecol Lett.* 4:434-438.
- 414 Peuke AD, Tischner R (1991) Nitrate uptake and reduction of aseptically cultivated spruce seedlings,
415 *Picea abies* (L.) karst. *J Exp Bot.* 42:723-728.

- 416 Polgar CA, Primack RB (2011) Leaf-out phenology of temperate woody plants: from trees to
417 ecosystems. *New Phytol.* 191:926-941.
- 418 Saarinen T, Rasmus S, Lundell R, Kauppinen O-K, Hänninen H (2016) Photosynthetic and
419 phenological responses of dwarf shrubs to the depth and properties of snow. *OIKOS*.
420 125:364-373.
- 421 Sanders-DeMott R, Templer PH (2017) What about winter? Integrating the missing season into
422 climate change experiments in seasonally snow covered ecosystems. *Methods in Ecology*
423 *and Evolution.* 8:1183-1191.
- 424 Schmidt B, Strack D, Weidner M (1991) Nitrate reductase in needles, roots and trunk wood of
425 spruce trees (*Picea abies* (L.) Karst.). *Trees.* 5:215-226.
- 426 Silla F, Escudero A (2003) Uptake, demand and internal cycling of nitrogen in saplings of
427 Mediterranean *Quercus* species. *Oecologia.* 136:28-36.
- 428 Smirnoff N, Todd P, Stewart G (1984) The occurrence of nitrate reduction in the leaves of woody
429 plants. *Ann Bot.* 54:363-374.
- 430 Stadler J, Gebauer G (1992) Nitrate reduction and nitrate content in ash trees (*Fraxinus excelsior*
431 L.): distribution between compartments, site comparison and seasonal variation. *Trees.*
432 6:236-240.
- 433 Sym GJ (1984) Optimisation of the in-vivo Assay Conditions for Nitrate Reductase in Barley
434 (*Hordeum vulgare* L. cv. Igri). *J Sci Food Agr.* 35:725-730.
- 435 Thomas FM, Hilker C (2000) Nitrate reduction in leaves and roots of young pedunculate oaks
436 (*Quercus robur*) growing on different nitrate concentrations. *Environ Exp Bot.* 43:19-32.
- 437 Troelstra SR, Wagenaar R, Smant W, De-Boer W (1995) Soil nitrogen transformations and nitrate
438 utilization by *Deschampsia flexuosa* (L.) Trin. at two contrasting heathland sites. *Plant Soil.*
439 176:81-93.
- 440 Ueda M, Mizumachi E, Tokuchi N (2010) Winter nitrate uptake by the temperate deciduous tree
441 *Quercus serrata*. *Journal of Forest Research.* 15:411-414.
- 442 Valentine DW, Kielland K, Chapin III FS, McGuire AD, Van Cleve K (2006) Patterns of
443 Biogeochemistry in Alaskan Boreal Forests. In: Chapin III FS, Oswald MW, van Cleve K,

- 444 Viereck LA, Verbyla DL (eds) Alaska's changing boreal forest. Oxford University Press, Inc,
445 New York, pp 241-266.
- 446 Van Cleve, Keith; Chapin, F. Stuart; Ruess, Roger W. 2017. Bonanza Creek LTER: Hourly Air
447 Temperature Measurements (sample, min, max) at 50 cm and 150 cm from 1988 to Present
448 in the Bonanza Creek Experimental Forest near Fairbanks, Alaska, Bonanza Creek LTER -
449 University of Alaska Fairbanks. BNZ:1, <http://www.lter.uaf.edu/data/data-detail/id/1>.
450 doi:10.6073/pasta/006bae44c88f7d8b6fabe8cfebee86ff, the last accessed date: 2017-03-08
- 451 Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee T, Fromentin J-M,
452 Hoegh-Guldberg O, Bairlein F (2002) Ecological responses to recent climate change. *Nature*.
453 416:389-395.
- 454 Yao B, Cao J, Zhao C, Rengel Z (2011) Influence of ammonium and nitrate supply on growth,
455 nitrate reductase activity and N-use efficiency in a natural hybrid pine and its parents. *J Plant*
456 *Ecol*. 4:275-282.
- 457 Yarie J, Van Cleve K (2006) Controls over Forest Production in Interior Alaska. In: Chapin III FS,
458 Oswald MW, van Cleve K, Viereck LA, Verbyla DL (eds) Alaska's changing boreal forest.
459 Oxford University Press, Inc, New York, pp 171-188.
- 460

FIGURE CAPTIONS

Figure 1: Temperature changes in 2009, 2015, and 2016. The temperatures recorded at an adjacent long-term ecological research (LTER) site (64°44'30"N, 148°18'50"W) are presented as daily means (closed circles: 2009, open circles: 2015, open squares: 2016) and ranges (continuous bars: 2009, dashed bars: 2015, dotted bars: 2016). Arrows indicate the sampling days (closed arrows: 2009, open arrows: summer 2015 and winter 2016). Data were obtained from the Bonanza Creek LTER Database.

Figure 2: Seasonal differences in nitrate reductase activities (NRA) and NO_3^- -N concentration in current year needles and fine roots of black spruce (*P. mariana*) in two sites. (a) $\text{NRA}(+\text{NO}_3)$ assayed with incubation buffer containing NO_3^- -N, (b) $\text{NRA}(-\text{NO}_3)$ assayed with incubation buffer containing no NO_3^- -N, and (c) NO_3^- -N concentration. Both NRA measurements were made at 30 °C. Samples were collected from five individual trees in each site.

Figure 3: The effects of incubation temperature on NRA in current year needles and fine roots of black spruce (*P. mariana*) collected in December 2009 and (a) supplied with NO_3^- -N [$\text{NRA}(+\text{NO}_3)$], or (b) not supplied with NO_3^- -N [$\text{NRA}(-\text{NO}_3)$]. Samples were collected from five individual trees in site 1.

Fig. 1

Fig. 2

(a)

(b)

Fig. 3

Table 1 Explanatory variables and the coefficients of variables for the generalized linear mixed model (GLMM) to describe the effects of season, site, and tissue on NRA(+NO₃), NRA(-NO₃) and NO₃⁻-N concentration in black spruce. The coefficients for the best performing models are shown, and the models were selected to have the lowest Akaike Information Criterion (AIC) by comparing AIC for each of possible subset of explanatory variables (See Appendices 2–4 for details). P values indicate the probability of including zero value of coefficients within 95 % Wald confidence interval.

	Variable type	NRA(+NO ₃)			NRA(-NO ₃)			NO ₃ ⁻ -N concentration		
		Coefficient	Std Error	P value	Coefficient	Std Error	P value	Coefficient	Std Error	P value
(Intercept)		12.94	5.24	0.014	23.17	4.51	< 0.001	285.20	93.70	0.002
Season [†]	categorical	-5.01	6.15	0.415	-0.29	6.09	0.962	-38.84	127.45	0.761
Site ^{††}	categorical	78.78	37.17	0.034	53.52	7.14	< 0.001	-	-	-
Tissue ^{†††}	categorical	-	††††	-	-6.55	2.35	0.005	-7.16	134.99	0.958
Season × Site	interaction	-68.30	38.04	0.073	-23.85	8.20	0.004	-	-	-
Site × Tissue	interaction	-	-	-	-29.38	4.82	< 0.001	-	-	-
Season × Tissue	interaction	-	-	-	4.95	3.33	0.137	1139.10	572.22	0.047
Season × Site × Tissue	interaction	-	-	-	-	-	-	-	-	-

[†]: The regression parameter estimates for these categorical variables were measured as departures from summer to winter.

^{††}: The regression parameter estimates for these categorical variables were measured as departures from site 1 to site 2.

^{†††}: The regression parameter estimates for these categorical variables were measured as departures from current year needles to fine roots.

^{††††}: Variables with blank (-) in coefficients were not used in the selected best performing model based on AIC.

Table 2 Explanatory variables and the coefficients of variables for the generalized linear mixed model (GLMM) to describe the effects of incubation temperature and tissue on NRA(+NO₃) and NRA(-NO₃) in black spruce. The coefficients for the best performing models are shown, and the models were selected to have the lowest Akaike Information Criterion (AIC) by comparing AIC for each of possible subset of explanatory variables (See Appendices 5–6 for details). P values indicate the probability of including zero value of coefficients within 95 % Wald confidence interval.

	Variable type	NRA(+NO ₃)			NRA(-NO ₃)		
		Coefficient	Std Error	P value	Coefficient	Std Error	P value
(Intercept)		10.98	1.28	< 0.001	21.80	2.50	< 0.001
Incubation temperature [†]	categorical	-5.04	1.53	0.001	-0.26	1.76	0.883
Tissue ^{††}	categorical	7.45	1.36	< 0.001	-	-	-
Incubation temperature × Tissue	interaction	- ^{†††}	- ^{†††}	-	-	-	-

[†]: The regression parameter estimates for these categorical variables were measured as departures from the incubation temperature -3 °C and 30 °C.

^{††}: The regression parameter estimates for these categorical variables were measured as departures from current year needles to fine roots.

^{†††}: Variables with blank (-) in coefficients were not used in the selected best performing model based on AIC.

Appendix 1: Effects of NO_3^- -N concentration in incubation buffer on needle NRA of black spruce (*P. mariana*). The incubation buffers had same composition other than NO_3^- -N concentration. Different symbols indicate different individual trees (n = 3).

Appendix 2 Comparison of candidate generalized linear mixed models (GLMM) to test the effects of season, site, tissue, and their interactions on NRA(+NO₃) of black spruce. The variables (fixed effects) and their interactions applied in each model are indicated by “+”, and the models are rank ordered from most to least supported based on AIC.

Season	Site	Tissue	Season × Site	Site × Tissue	Tissue × Season	Season × Site × Tissue	df	AIC	ΔAIC
+	+		+				34	-155.33	
+	+	+	+				33	-154.15	1.18
+	+	+	+	+			32	-152.80	1.35
+	+						35	-152.17	0.63
+	+	+	+		+		32	-152.15	0.02
	+						36	-151.94	0.20
+	+	+					34	-150.89	1.06
+	+	+	+	+	+		31	-150.81	0.08
	+	+					35	-150.74	0.07
+	+	+		+			33	-149.48	1.26
	+	+		+			34	-149.37	0.11
+	+	+	+	+	+	+	30	-149.33	0.04
+	+	+			+		33	-148.94	0.38
+							36	-148.14	0.80
+	+	+		+	+		32	-147.59	0.55
+		+					35	-146.99	0.61
		+					36	-146.55	0.44
+		+			+		34	-145.02	1.52

Appendix 3 Comparison of candidate generalized linear mixed models (GLMM) to test the effects of season, site, tissue, and their interactions on NRA(-NO₃) of black spruce. The variables (fixed effects) and their interactions applied in each model are indicated by “+”, and the models are rank ordered from most to least supported based on AIC.

Season	Site	Tissue	Season × Site	Site × Tissue	Tissue × Season	Season × Site × Tissue	df	AIC	ΔAIC
+	+	+	+	+	+		31	-239.95	
+	+	+	+	+			32	-239.82	0.13
+	+	+	+	+	+	+	30	-237.96	1.86
+	+	+		+	+		32	-236.32	1.65
+	+	+		+			33	-236.31	0.01
	+	+		+			34	-235.32	0.99
+	+	+	+				33	-217.90	17.42
+	+	+	+		+		32	-215.90	2.00
+	+	+					34	-212.09	3.81
	+	+					35	-211.70	0.39
+	+	+			+		33	-210.10	1.59
+	+		+				34	-208.12	1.98
		+					36	-206.09	2.03
+		+					35	-205.50	0.59
+		+			+		34	-203.51	1.99
	+						36	-202.83	0.69
+	+						35	-202.27	0.56
+							36	-195.16	7.11

Appendix 4 Comparison of candidate generalized linear mixed models (GLMM) to test the effects of season, site, tissue, and their interactions on NO_3^- -N concentration of black spruce. The variables (fixed effects) and their interactions applied in each model are indicated by “+”, and the models are rank ordered from the most to least supported based on AIC.

Season	Site	Tissue	Season × Site	Site × Tissue	Tissue × Season	Season × Site × Tissue	df	AIC	ΔAIC
+		+			+		33	-388.70	
		+					35	-387.25	1.45
+	+	+			+		32	-387.24	0.01
	+						35	-386.70	0.54
+							35	-386.58	0.12
+	+	+		+	+		31	-386.28	0.31
	+	+					34	-385.88	0.39
+		+					34	-385.86	0.03
+	+	+	+		+		31	-385.25	0.61
+	+						34	-385.24	0.01
	+	+		+			33	-384.67	0.57
+	+	+	+	+	+		30	-384.58	0.09
+	+	+					33	-384.46	0.12
+	+	+	+	+	+	+	29	-383.60	0.86
+	+		+				33	-383.35	0.25
+	+	+		+			32	-383.25	0.10
+	+	+	+				32	-382.56	0.70
+	+	+	+	+			31	-381.37	1.18

Appendix 5 Summary of generalized linear mixed models (GLMM) comparisons to test the effects of incubation temperature, tissue, and their interactions on NRA(+NO₃) of black spruce. Models are rank ordered from the most to least supported based on AIC.

Incubation temperature	Tissue	Incubation temperature × Tissue	df	AIC	ΔAIC
+	+		15	-80.40	
+	+	+	14	-79.21	0.23
	+		16	-76.17	0.05
+			16	-62.37	1.45

Appendix 6 Summary of generalized linear mixed models (GLMM) comparisons to test the effects of incubation temperature, tissue, and their interactions on NRA(-NO₃) of black spruce. Models are rank ordered from the most to least supported based on AIC.

Incubation temperature	Tissue	Incubation temperature × Tissue	df	AIC	ΔAIC
	+		16	-109.62	
+			16	-109.61	0.01
+	+		15	-107.63	1.98
+	+	+	14	-106.25	1.38