TITLE:

Propagation of Boundary-Induced Discontinuity in Stationary Radiative Transfer

AUTHOR(S):

Kawagoe, Daisuke; Chen, I-Kun

CITATION:

Kawagoe, Daisuke ...[et al]. Propagation of Boundary-Induced Discontinuity in Stationary Radiative Transfer. Journal of Statistical Physics 2018, 170(1): 127-140

ISSUE DATE:

2018-01

URL:

<http://hdl.handle.net/2433/242789>

RIGHT:

This is a post-peer-review, pre-copyedit version of an article published in Journal of Statistical Physics. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s10955-017-1922-8>; この論文は出版社版でありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

J. Stat. Phys. manuscript No.
(will be inserted by the editor)

Propagation of boundary-induced discontinuity in stationary radiative transfer

Daisuke Kawagoe · I-Kun Chen

Received: date / Accepted: date

Abstract We consider the boundary value problem of the stationary transport equation in the slab domain of general dimensions. In this paper, we discuss the relation between discontinuity of the incoming boundary data and that of the solution to the stationary transport equation. We introduce two conditions posed on the boundary data so that discontinuity of the boundary data propagates along positive characteristic lines as that of the solution to the stationary transport equation. Our analysis does not depend on the celebrated velocity averaging lemma, which is different from previous works. We also introduce an example in two dimensional case which shows that piecewise continuity of the boundary data is not a sufficient condition for the main result.

Keywords Stationary transport equation · Boundary value problem · Discontinuity · Characteristic lines

Mathematics Subject Classification (2000) 70B05

1 Introduction

We consider the stationary transport equation:

$$\xi \cdot \nabla_x f(x, \xi) + \mu_t(x) f(x, \xi) = \mu_s(x) \int_{S^{d-1}} p(x, \xi, \xi') f(x, \xi') d\sigma_{\xi'},$$
$$(x, \xi) \in \Omega \times S^{d-1}, \quad (1)$$

The second author is supported in part by JSPS KAKENHI grant number 15K17572.

Daisuke Kawagoe
Yoshida-Honmachi, Sakyo, Kyoto 6068501, Japan
Tel.: +81-075-7533385
Fax: +81-075-7533380
E-mail: d.kawagoe@acs.i.kyoto-u.ac.jp

I-Kun Chen
No. 1, Sec. 4, Roosevelt Rd., Taipei 10617, Taiwan
E-mail: ikun.chen@gmail.com

where Ω is the slab domain $\mathbb{R}^{d-1} \times (0, 1)$ of the dimension $d \geq 2$ and S^{d-1} is the unit sphere in $\mathbb{R}^d$. The stationary transport equation describes propagation of particles interacting only with the media, for example neutron (Case and Zweifel (1967)) or photon (Chandrasekhar (1960)). The function $f(x, \xi)$ describes the density of particles at the point $x \in \Omega$ with direction $\xi \in S^{d-1}$. The coefficient μ_t and the product $\mu_s p$ characterize the effect of the media; they are called the attenuation coefficient and the scattering indicatrix, respectively.

We pose the incoming boundary condition as follows. Let

$$\Gamma_- := \{(x, \xi) \in \partial\Omega \times S^{d-1} | n(x) \cdot \xi < 0\},$$

where $n(x)$ is the outer normal vector at $x \in \partial\Omega$. Then, for a given function f_0 on Γ_- , a solution f to the stationary transport equation (1) must satisfy

$$f(x, \xi) = f_0(x, \xi), \quad (x, \xi) \in \Gamma_-. \quad (2)$$

In this paper, we discuss the relation between discontinuity of the boundary data and that of the solution to the stationary transport equation. Aoki et al. (2001) emphasize importance and significance of this analysis.

We assume that two coefficients μ_t and μ_s are nonnegative bounded continuous functions on Ω satisfying

$$\inf_{x \in \Omega} (\mu_t(x) - \mu_s(x)) > 0. \quad (3)$$

We note that, from the assumption above, we have

$$\underline{\mu}_t := \inf_{x \in \Omega} \mu_t(x) > 0$$

and

$$M := \sup_{x \in \Omega} \left(\frac{\mu_s(x)}{\mu_t(x)} \right) < 1,$$

which are crucial and facilitate our analysis dramatically. Although the assumption (3) does not hold in some relevant mathematical models, the linearized Boltzmann equation and the linearized BGK equation, in gas dynamics, it is reasonable in the case of radiative transfer.

We also assume that the integral kernel p is a nonnegative bounded continuous function on $\Omega \times S^{d-1} \times S^{d-1}$ which satisfies

$$\int_{S^{d-1}} p(x, \xi, \xi') d\sigma_{\xi'} = 1$$

for all $(x, \xi) \in \Omega \times S^{d-1}$.

We regard the directional derivative $\xi \cdot \nabla_x f(x, \xi)$ as

$$\xi \cdot \nabla_x f(x, \xi) := \frac{d}{dt} f(x + t\xi, \xi)|_{t=0}$$

and the measure $d\sigma_{\xi'}$ is the Lebesgue measure on the sphere S^{d-1} .

We introduce some notations. Let

$$X := (\Omega \times S^{d-1}) \cup \Gamma_-.$$

We introduce two functions $\tau_{\pm}$ on X defined by

$$\tau_{\pm}(x, \xi) := \inf\{t > 0 | x \pm t\xi \notin \Omega\}.$$

Let

$$S_{\pm}^{d-1} := \{\xi = (\xi_1, \xi_2, \dots, \xi_d) \in S^{d-1} | \xi_d \gtrless 0\}$$

and let $\Gamma_{-, \xi}$ and $\Gamma_{-, x}$ be projections of Γ_- on $\partial\Omega$ and $S_{\pm}^{d-1}$ respectively, that is,

$$\Gamma_{-, \xi} := \{x \in \partial\Omega | n(x) \cdot \xi < 0\}, \quad \xi \in S_{\pm}^{d-1}$$

and

$$\Gamma_{-, x} := \{\xi \in S^{d-1} | n(x) \cdot \xi < 0\}, \quad x \in \partial\Omega.$$

At last, let $\text{disc}(f)$ and $\text{disc}(f_0)$ be the set of discontinuous points for a function f on X and for a function f_0 on Γ_- , respectively.

The main result in this paper is as follows:

Theorem 1. *Suppose that a boundary data f_0 is bounded and that it satisfies at least one of the following two conditions:*

1. $f_0(\cdot, \xi)$ is continuous on $\Gamma_{-, \xi}$ for almost all $\xi \in S_{\pm}^{d-1}$,
2. $f_0(x, \cdot)$ is continuous on $\Gamma_{-, x}$ for almost all $x \in \partial\Omega$.

Then, there exists a unique solution f to the boundary value problem (1)-(2) and the following relation holds:

$$\text{disc}(f) = \{(x_* + t\xi_*, \xi_*) | (x_*, \xi_*) \in \text{disc}(f_0), 0 \leq t < \tau_+(x_*, \xi_*)\}.$$

Here, we call a bounded function f on X a solution to the boundary value problem (1)-(2) if it satisfies the stationary transport equation (1) for all $(x, \xi) \in \Omega \times S^{d-1}$ and the boundary condition (2) for all $(x, \xi) \in \Gamma_-$. This theorem means that discontinuity of the boundary data propagates along positive characteristic lines as that of the solution to the stationary transport equation.

Remark 1. If the boundary data f_0 is bounded continuous on Γ_- , then there exists a unique solution f . Moreover, since $\text{disc}(f_0)$ is the empty set by assumption, $\text{disc}(f)$ is also the empty set, which implies that the solution f is also bounded continuous on X .

Anikonov et al. (1993) have shown this property assuming condition 1 in a three dimensional bounded convex domain with piecewise continuous coefficients. They also assumed so-called general convexity to these pieces. They made use of this property to solve the inverse problem to determine the coefficient μ_t from the knowledge of the boundary measurements $f|_{\Gamma_+}$, where

$$\Gamma_+ := \{(x, \xi) \in \partial\Omega \times S^{d-1} | n(x) \cdot \xi > 0\}$$

and $n(x)$ is again the outer normal vector at $x \in \partial\Omega$.

On the other hand, Aoki et al. (2001) have shown the same property assuming condition 2 in the two dimensional half plane domain with $\mu_t = \mu_s = 1$ and $p = 1/2\pi$. We note that this assumption violates ours because $\mu_t - \mu_s = 0$ for all $x \in \Omega$. They also assume that the boundary data is periodic with respect to the spacial variable and is independent of the directional variable. We have succeeded in modifying their assumptions. Their analysis is based on the celebrated velocity averaging lemma, which works in L^2 -framework only when p is constant. On the other hand, p is not constant in our setting. So, we cannot apply their approach directly. However, we overcome this point by L^∞ -based discussion.

We mention some relevant works in gas dynamics. In the time dependent case, Kim (2011) has investigated formation and propagation of discontinuity of solutions to the Boltzmann equation in non-convex domains with three kinds of boundary conditions, which

are the incoming boundary condition, the diffuse boundary condition and the bounce-back boundary condition. Also, Guo and Yang (2016) have discussed existence and BV-regularity for transport equation with a combination of the incoming boundary condition and the diffuse boundary condition in non-convex domains. On the other hand, in the stationary case, Aoki et al. (2001) have numerically analyzed discontinuity of solutions to the BGK equation in a two dimensional square domain with the diffuse boundary condition with discontinuous temperature. Esposito et al. (2013) also have discussed discontinuity of solutions to the Boltzmann equation with non-isothermal diffuse boundary condition in non-convex domains.

The ingredient of the rest part in this paper is as follows. In Section 2, we derive integral equations from the boundary value problem, and we show existence and uniqueness of solutions to derived integral equations. In Section 3, we discuss regularity of the solution to integral equations. Especially, we decompose the solution into two parts, the discontinuous part and the continuous part. In Section 4, we check the equivalence between the boundary value problem and integral equations. In other words, we prove that the solution to integral equations indeed satisfies the stationary transport equation under the assumption in Theorem 1. In Section 5, we introduce an example in two dimensional case which shows that piecewise continuity of the boundary data is not a sufficient condition for the main result.

2 Existence and uniqueness of solutions to the stationary transport equation

In this section, we derive integral equations from the boundary value problem, and we show existence and uniqueness of solutions to derived integral equations.

For $(x, \xi) \in X$, by integrating the stationary transport equation (1) from x along the negative characteristic line $\{x - t\xi \mid t > 0\}$ until the line touches the boundary $\partial\Omega$ and by taking the boundary condition (2) into consideration, we obtain the following integral equations: when $\xi_d \neq 0$,

$$\begin{aligned} f(x, \xi) = & \exp\left(-M_t(x, \xi; \tau_-(x, \xi))\right) f_0(x - \tau_-(x, \xi)\xi, \xi) \\ & + \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp\left(-M_t(x, \xi; s)\right) \\ & \times \int_{S^{d-1}} p(x - s\xi, \xi, \xi') f(x - s\xi, \xi') d\sigma_{\xi'} ds, \end{aligned} \quad (4)$$

and when $\xi_d = 0$,

$$f(x, \xi) = \int_0^\infty \mu_s(x - s\xi) \exp\left(-M_t(x, \xi; s)\right) \int_{S^{d-1}} p(x - s\xi, \xi, \xi') f(x - s\xi, \xi') d\sigma_{\xi'} ds, \quad (5)$$

where

$$M_t(x, \xi; s) := \int_0^s \mu_t(x - r\xi) dr.$$

We call a bounded function f on X a solution to integral equations (4)-(5) if it satisfies integral equations (4)-(5) for all $(x, \xi) \in X$.

We note that, although solutions to the boundary value problem (1)-(2) satisfy integral equations (4)-(5), the converse does not hold in general. However, as we will see later in Section 4, under the assumption in Theorem 1, the solution to integral equations (4)-(5) is also the solution to the boundary value problem (1)-(2). Therefore, our current task is to find a solution to integral equations (4)-(5).

Proposition 1. *The solution to integral equations (4)-(5) is unique, if it exists.*

Proof. Let f_1 and f_2 be two solutions to integral equations (4)-(5). Then the difference $\tilde{f} := f_1 - f_2$ is also bounded on X and satisfies the following integral equation:

$$\tilde{f}(x, \xi) = \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp(-M_t(x, \xi; s)) \int_{S^{d-1}} p(x, \xi, \xi') \tilde{f}(x - s\xi, \xi') d\sigma_{\xi'} ds$$

for all $(x, \xi) \in X$. Then, we have

$$\begin{aligned} |\tilde{f}(x, \xi)| &\leq \left(\sup_{(x, \xi) \in X} |\tilde{f}(x, \xi)| \right) \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp(-M_t(x, \xi; s)) ds \\ &\leq \left(\sup_{(x, \xi) \in X} |\tilde{f}(x, \xi)| \right) \int_0^\infty \frac{\mu_s(x - s\xi)}{\mu_t(x - s\xi)} \left| \frac{d}{ds} \exp(-M_t(x, \xi; s)) \right| ds \\ &\leq M \left(\sup_{(x, \xi) \in X} |\tilde{f}(x, \xi)| \right) \end{aligned}$$

for all $(x, \xi) \in X$. We emphasize that the supremum in this paper is not the essential supremum, which enables us to justify the pointwise discussion. Therefore,

$$\sup_{(x, \xi) \in X} |\tilde{f}(x, \xi)| \leq M \left(\sup_{(x, \xi) \in X} |\tilde{f}(x, \xi)| \right). \quad (6)$$

Since $M < 1$, the inequality (6) implies $\sup_{(x, \xi) \in X} |\tilde{f}(x, \xi)| = 0$, that is, $f_1 = f_2$ for all $(x, \xi) \in X$. $\square$

At last, we prove existence of a solution by iteration. This strategy is standard in the field of radiative transfer. For example, see Anikonov et al. (1993). Let us define a family of functions $\{f^{(n)}\}_{n \geq 0}$ on X as follows:

$$f^{(0)}(x, \xi) := \begin{cases} \exp(-M_t(x, \xi; \tau_-(x, \xi))) f_0(x - \tau_-(x, \xi)\xi, \xi), & \xi_d \neq 0, \\ 0, & \xi_d = 0, \end{cases} \quad (7)$$

and

$$\begin{aligned} f^{(n+1)}(x, \xi) &:= \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp(-M_t(x, \xi; s)) \\ &\quad \times \int_{S^{d-1}} p(x - s\xi, \xi, \xi') f^{(n)}(x - s\xi, \xi') d\sigma_{\xi'} ds. \end{aligned} \quad (8)$$

Now we prove that the sum $f := \sum_{n=0}^\infty f^{(n)}$ is a solution to integral equations (4)-(5). To begin with, we prove that $f := \sum_{n=0}^\infty f^{(n)}$ is indeed defined on X . Especially, we show the following two propositions.

Proposition 2. *Suppose that the boundary data f_0 is bounded on Γ_- . Then, $f^{(n)}$ is also bounded on X for all $n \geq 0$.*

Proof. We use the induction on n .

For $(x, \xi) \in \Omega \times S^{d-1}$ and $\xi_d = 0$, we have

$$|f^{(0)}(x, \xi)| = 0 \leq \sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)|,$$

and for otherwise, we have

$$\begin{aligned} |f^{(0)}(x, \xi)| &\leq \exp\left(-M_t(x, \xi; \tau_-(x, \xi))\right) |f_0(x - \tau_-(x, \xi)\xi, \xi)| \\ &\leq \sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)|. \end{aligned}$$

These estimates imply that $f^{(0)}$ is bounded on X .

Now, we assume that $f^{(n)}$ is bounded on X for some $n \in \mathbb{N}$. Then,

$$\begin{aligned} |f^{(n+1)}(x, \xi)| &\leq \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp\left(-M_t(x, \xi; s)\right) \\ &\quad \times \int_{S^{d-1}} p(x - s\xi, \xi, \xi') |f^{(n)}(x - s\xi, \xi')| d\sigma_{\xi'} ds \\ &\leq \left(\sup_{(x, \xi) \in X} |f^{(n)}(x, \xi)| \right) \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp\left(-M_t(x, \xi; s)\right) ds \\ &\leq M \left(\sup_{(x, \xi) \in X} |f^{(n)}(x, \xi)| \right) \end{aligned} \quad (9)$$

for all $(x, \xi) \in X$. This inequality implies that $f^{(n+1)}$ is defined and bounded on X . This completes the proof. $\square$

Proposition 3. Suppose that the boundary data f_0 is bounded on Γ_- . Then, the sum $\sum_{n=0}^{\infty} f^{(n)}(x, \xi)$ converges absolutely and uniformly on X .

Proof. From the inequality (9), we have

$$\begin{aligned} \sup_{(x, \xi) \in X} |f^{(n)}(x, \xi)| &\leq M \left(\sup_{(x, \xi) \in X} |f^{(n-1)}(x, \xi)| \right) \\ &\leq M^n \left(\sup_{(x, \xi) \in X} |f^{(0)}(x, \xi)| \right) \leq M^n \left(\sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)| \right) \end{aligned}$$

for all $n \geq 0$. Thus,

$$\begin{aligned} \sum_{n=0}^{\infty} |f^{(n)}(x, \xi)| &\leq \sum_{n=0}^{\infty} \sup_{(x, \xi) \in X} |f^{(n)}(x, \xi)| \\ &\leq \sum_{n=0}^{\infty} M^n \left(\sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)| \right) \\ &= \frac{1}{1-M} \left(\sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)| \right) < \infty, \end{aligned}$$

which implies absolute and uniform convergence of the sum $\sum_{n=0}^{\infty} f^{(n)}(x, \xi)$ on X . $\square$

From Proposition 2 and Proposition 3, the sum $f(x, \xi) = \sum_{n=0}^{\infty} f^{(n)}(x, \xi)$ converges absolutely and uniformly on X and satisfies

$$\begin{aligned} f(x, \xi) &= f^{(0)}(x, \xi) + \sum_{n=0}^{\infty} f^{(n+1)}(x, \xi) \\ &= f^{(0)}(x, \xi) + \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp(-M_t(x, \xi; s)) \\ &\quad \times \int_{S^{d-1}} p(x - s\xi, \xi, \xi') \sum_{n=0}^{\infty} f^{(n)}(x - s\xi, \xi') d\sigma_{\xi'} ds \\ &= f^{(0)}(x, \xi) + \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp(-M_t(x, \xi; s)) \\ &\quad \times \int_{S^{d-1}} p(x - s\xi, \xi, \xi') f(x - s\xi, \xi') d\sigma_{\xi'} ds \end{aligned}$$

for all $(x, \xi) \in X$, which is the pair of integral equations (4)-(5). Thus, the sum $f(x, \xi) = \sum_{n=0}^{\infty} f^{(n)}(x, \xi)$ is the solution to integral equations (4)-(5).

3 Regularity of the solution

In this section, we discuss regularity of the solution to integral equations (4)-(5). We decompose the solution f into two parts as below:

$$f(x, \xi) = F_0(x, \xi) + F_1(x, \xi),$$

where

$$\begin{aligned} F_0(x, \xi) &:= f^{(0)}(x, \xi), \\ F_1(x, \xi) &:= \sum_{n=1}^{\infty} f^{(n)}(x, \xi). \end{aligned}$$

From now, we observe discontinuity of F_0 and prove continuity of F_1 . This decomposition is the main idea in our analysis.

3.1 Discontinuity of F_0

First, we prove the following proposition.

Proposition 4.

$$\text{disc}(F_0) = \{(x_* + t\xi_*, \xi_*) | (x_*, \xi_*) \in \text{disc}(f_0), 0 \leq t < \tau_+(x_*, \xi_*)\}.$$

Proof. Let us recall the explicit formula of F_0 (7): when $\xi_d \neq 0$,

$$F_0(x, \xi) = \exp\left(-\int_0^{\tau_-(x, \xi)} \mu_t(x - r\xi) dr\right) f_0(x - \tau_-(x, \xi)\xi, \xi).$$

τ_- is continuous on X with $\xi_d \neq 0$ because of its explicit formula:

$$\tau_-(x, \xi) = \begin{cases} x_d/\xi_d, & \xi_d > 0, \\ (x_d - 1)/\xi_d, & \xi_d < 0. \end{cases}$$

Thus, we have when $\xi_d \neq 0$,

$$(x, \xi) \in \text{disc}(F_0) \Leftrightarrow (x - \tau_-(x, \xi)\xi, \xi) \in \text{disc}(f_0),$$

which implies that the statement holds when $\xi_d \neq 0$.

Thus, only we have to check is continuity of F_0 at $(x, \xi) \in \Omega \times S^{d-1}$ with $\xi_d = 0$. In this setting, continuity of F_0 with respect to x is obvious from the explicit formula of it (7). So, we focus on continuity of F_0 with respect to ξ . Since

$$\lim_{\xi_d \rightarrow 0} \tau_-(x, \xi) = \infty$$

for all $x \in \Omega$ and since f_0 is bounded on Γ_- , we have

$$\begin{aligned} \lim_{\xi_d \rightarrow 0} |F_0(x, \xi)| &\leq \left(\sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)| \right) \lim_{\xi_d \rightarrow 0} \exp \left(- \int_0^{\tau_-(x, \xi)} \mu_t(x - r\xi) dr \right) \\ &\leq \left(\sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)| \right) \lim_{\xi_d \rightarrow 0} \exp \left(- \underline{\mu}_t \tau_-(x, \xi) \right) = 0, \end{aligned}$$

which means $\lim_{\xi_d \rightarrow 0} F_0(x, \xi) = 0$ for all $x \in \Omega$. So F_0 is continuous at $(x, \xi) \in \Omega \times S^{d-1}$ with $\xi_d = 0$. This completes the proof. $\square$

3.2 Continuity of F_1

Secondly, we prove continuity of F_1 . To do so, we prove by induction that functions $f^{(n)}$, defined above, are bounded continuous on X for all $n \geq 1$. After that, we know from Proposition 3 that the sum $\sum_{n=1}^{\infty} f^{(n)}$ converges uniformly on X , which implies that it is also bounded continuous on X .

Lemma 1. *Under the assumption in Theorem 1, $f^{(1)}$ is bounded continuous on X .*

Proof. Boundedness of $f^{(1)}$ was proved in Section 2, so here we prove continuity of it. By substituting the explicit formula of $f^{(0)}$ (7) for one appeared in the recursive formula (8) with $n = 0$, we have

$$\begin{aligned} f^{(1)}(x, \xi) &= \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp(-M_t(x, \xi; s)) \\ &\quad \times \int_{S^{d-1}} p(x - s\xi, \xi, \xi') f^{(0)}(x - s\xi, \xi') d\sigma_{\xi'} ds \\ &= \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp(-M_t(x, \xi; s)) G(x - s\xi, \xi) ds, \end{aligned}$$

where

$$G(x, \xi) = G_+(x, \xi) + G_-(x, \xi),$$

and

$$G_{\pm}(x, \xi) := \int_{S_{\pm}^{d-1}} p(x, \xi, \xi') \exp(-M_t(x, \xi'; \tau_-(x, \xi))) f_0(x - \tau_-(x, \xi')\xi', \xi') d\sigma_{\xi'}. \quad (10)$$

Then, we introduce the following lemma, whose proof will be appeared later.

Lemma 2. *Under the assumption in Theorem 1, G is bounded continuous on $\Omega \times S^{d-1}$.*

Remark 2. Lemma 2 is the very key idea in this paper. Aoki et al. (2001) proved the same lemma when p is constant, which led them to success.

Admitting Lemma 2, we continue to prove Lemma 1. Let $\tilde{G}$ be the zero extension of G to $\mathbb{R}^d \times S^{d-1}$, that is,

$$\tilde{G}(x, \xi) := \begin{cases} G(x, \xi), & (x, \xi) \in \Omega \times S^{d-1}, \\ 0, & \text{otherwise.} \end{cases}$$

Also let $\tilde{\mu}_t$ be the zero extension of μ_t to $\mathbb{R}^d$ and let $\tilde{M}_t$ be the corresponding M_t . Then, $f^{(1)}$ can be written as the following.

$$f^{(1)}(x, \xi) = \int_0^\infty \tilde{\mu}_s(x - s\xi) \exp\left(-\tilde{M}_t(x, \xi; s)\right) \tilde{G}(x - s\xi, \xi) ds,$$

Since the integrand is dominated by

$$\left(\sup_{x \in \Omega} \mu_s(x)\right) \left(\sup_{(x, \xi) \in \Omega \times S^{d-1}} |G(x, \xi)|\right) \exp(-\mu_t s),$$

which is integrable with respect to s on the half line $[0, \infty)$, and since the integrand is continuous at each point $(x, \xi) \in \Omega \times S^{d-1}$ for almost all $s \in [0, \infty)$, we apply the dominated convergence theorem to prove continuity of $f^{(1)}$ on X . $\square$

Proof of Lemma 2. Since boundedness of G_+ and G_- is obvious from their formulae (10), we focus on discussing continuity of them. At first, we fix a point $(\bar{x}, \bar{\xi}) \in \Omega \times S^{d-1}$ and prove continuity of G_+ at the point $(\bar{x}, \bar{\xi})$.

We first suppose that the boundary data f_0 satisfies condition 1. Since τ_- is continuous on X , the integrand

$$p(x, \xi, \xi') \exp\left(-M_t(x, \xi'; \tau_-(x, \xi'))\right) f_0(x - \tau_-(x, \xi')\xi', \xi')$$

is continuous at $(\bar{x}, \bar{\xi}) \in \Omega \times S^{d-1}$ for almost all $\xi' \in S_+^{d-1}$. Furthermore, the integrand is bounded by

$$\left(\sup_{(x, \xi, \xi')} p(x, \xi, \xi')\right) \left(\sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)|\right).$$

Therefore, we can apply the dominated convergence theorem to conclude that G_+ is bounded continuous on $\Omega \times S^{d-1}$.

Next, we suppose that f_0 satisfies condition 2. By changing variables of integration $y_0 = (y_1, y_2, \dots, y_{d-1}, 0) = x - \tau_-(x, \xi')\xi'$, we have

$$\begin{aligned} G_+(x, \xi) &= \int_{\mathbb{R}^{d-1}} p\left(x, \xi, \frac{x - y_0}{|x - y_0|}\right) \exp\left(-M_t\left(x, \frac{x - y_0}{|x - y_0|}; |x - y_0|\right)\right) \\ &\quad \times f_0\left(y_0, \frac{x - y_0}{|x - y_0|}\right) \frac{x_d}{|x - y_0|^d} dy_1 dy_2 \cdots dy_{d-1}, \end{aligned}$$

where $x_d/|x - y_0|^d$ is the Jacobian of this change.

By condition 2, for almost all $(y_1, y_2, \dots, y_{d-1}) \in \mathbb{R}^{d-1}$, the integrand

$$p\left(x, \xi, \frac{x-y_0}{|x-y_0|}\right) \exp\left(-M_t\left(x, \frac{x-y_0}{|x-y_0|}; |x-y_0|\right)\right) f_0\left(y_0, \frac{x-y_0}{|x-y_0|}\right) \frac{x_d}{|x-y_0|^d}$$

is continuous at $(\bar{x}, \bar{\xi}) \in \Omega \times S^{d-1}$. Furthermore, the integrand is uniformly bounded by

$$\left(\left(\sup_{(x, \xi, \xi') \in \Gamma_-} p(x, \xi, \xi')\right) \left(\sup_{(x, \xi) \in \Gamma_-} |f_0(x, \xi)|\right)\right) / (|\bar{x} - y_0| - \varepsilon/2)^d$$

on the neighborhood $B_{\varepsilon/2}(\bar{x}) \times S^{d-1}$, where $B_{\varepsilon/2}(\bar{x})$ is the closed ball with centre $\bar{x}$ and radius $\varepsilon/2$, and $\bar{x}_d > \varepsilon > 0$. Since the dominant is integrable with respect to $y_1, y_2, \dots$, and y_{d-1} , we can apply the dominated convergence theorem to conclude that G_+ is bounded continuous at $(\bar{x}, \bar{\xi}) \in \Omega \times S^{d-1}$.

Thus, it follows from the discussion above that G_+ is bounded continuous on $\Omega \times S^{d-1}$ if the boundary data f_0 satisfies the assumption in Theorem 1. In the same way, we can show that G_- is also bounded continuous on $\Omega \times S^{d-1}$. After all, G itself is also bounded continuous on $\Omega \times S^{d-1}$. $\square$

Remark 3. Our technique, changing variables of integration, seems available in the case of a general convex domain Ω with C^1 boundary $\partial\Omega$. However, the authors have not found the Jacobian of this change in the general case.

Lemma 3. Suppose that the function $f^{(n)}$, defined by recursive formulae (7)-(8), is bounded continuous on X for some $n \in \mathbb{N}$. Then, the successive function $f^{(n+1)}$ is also bounded continuous on X .

Proof. As the proof of Lemma 1, let $\tilde{f}^{(n)}$ be the zero extention of $f^{(n)}$ to $\mathbb{R}^d \times S^{d-1}$. Also let $\tilde{\mu}_s$ and $\tilde{p}$ be the zero extensions of μ_s and p to $\mathbb{R}^d$ and $\mathbb{R}^d \times S^{d-1} \times S^{d-1}$, respectively. Then, we have

$$f^{(n+1)}(x, \xi) = \int_0^\infty \tilde{\mu}_s(x - s\xi) \exp\left(-\tilde{M}_t(x, \xi; s)\right) \times \int_{S^{d-1}} \tilde{p}(x - s\xi, \xi, \xi') \tilde{f}^{(n)}(x - s\xi, \xi') d\sigma_{\xi'} ds$$

for all $(x, \xi) \in X$. Since $f^{(n)}$ is continuous on X , the integrand

$$\tilde{\mu}_s(x - s\xi) \exp\left(-\tilde{M}_t(x, \xi; s)\right) \int_{S^{d-1}} \tilde{p}(x - s\xi, \xi, \xi') \tilde{f}^{(n)}(x - s\xi, \xi') d\sigma_{\xi'}$$

is also continuous at each point $(x, \xi) \in X$ for almost all $s \in [0, \infty)$. In addition, this integrand is dominated by

$$\left(\sup_{x \in \Omega} \mu_s(x)\right) \left(\sup_{(x, \xi) \in X} |f^{(n)}(x, \xi)|\right) \exp\left(-\underline{\mu}_t s\right),$$

which is integrable with respect to s . Thus, we can apply Lebesgue's convergence theorem to prove continuity of $f^{(n+1)}$ on X . $\square$

By Lemma 1 and Lemma 3, $f^{(n)}$ is bounded continuous on X for all $n \in \mathbb{N}$, and by Proposition 3, the sum $\sum_{n=1}^\infty f^{(n)}(x, \xi)$ converges uniformly on X , which implies that the function F_1 is also bounded continuous on X .

Thus, we succeed to separate the solution into two parts, the discontinuous part F_0 and the continuity part F_1 .

4 Equivalence between the boundary value problem and derived integral equations

In this section, we check the equivalence between the boundary value problem (1)-(2) and integral equations (4)-(5). As we noted in Section 2, although solutions to the boundary value problem (1)-(2) satisfy integral equations (4)-(5), the converse does not hold in general. So, we see that, under the assumption in Theorem 1, the solution to integral equations (4)-(5) is indeed the solution to the boundary value problem (1)-(2). In other words, we prove the following two propositions.

Proposition 5. *Let f be the solution to integral equations (4)-(5). Suppose that the boundary data f_0 satisfies the assumption in Theorem 1. Then, the directional derivative $\xi \cdot \nabla_x f(x, \xi)$ is defined for all $(x, \xi) \in \Omega \times S^{d-1}$ and f satisfies the stationary transport equation (1).*

Proposition 6. *The solution f to integral equations (4)-(5) satisfies the boundary condition (2).*

Proof of Proposition 5. Making use of the following equality

$$\tau_-(x + t\xi, \xi) = \tau_-(x, \xi) + t$$

for $(x, \xi) \in \Omega \times S^{d-1}$ and $t \in \mathbb{R}$ such that $x + t\xi \in \Omega$, we have

$$f^{(0)}(x + t\xi, \xi) = \begin{cases} \exp\left(-M_t(x + t\xi, \xi; \tau_-(x, \xi) + t)\right) f_0(x - \tau_-(x, \xi)\xi, \xi), & \xi_d \neq 0, \\ 0, & \xi_d = 0. \end{cases}$$

When $\xi_d \neq 0$, since

$$M_t(x + t\xi, \xi; \tau_-(x, \xi) + t) = M_t(x, \xi; \tau_-(x, \xi)) - M_t(x, \xi; -t),$$

we have

$$\begin{aligned} \xi \cdot \nabla_x f^{(0)}(x, \xi) &= \frac{dM_t}{dt}(x, \xi; -t)|_{t=0} \exp\left(-M_t(x, \xi; \tau_-(x, \xi))\right) f_0(x - \tau_-(x, \xi)\xi, \xi) \\ &= -\mu_t(x) \exp\left(-M_t(x, \xi; \tau_-(x, \xi))\right) f_0(x - \tau_-(x, \xi)\xi, \xi) \\ &= -\mu_t(x) f^{(0)}(x, \xi). \end{aligned}$$

When $\xi_d = 0$, it is obvious that $\xi \cdot \nabla_x f^{(0)}(x, \xi) = 0 = -\mu_t(x) f^{(0)}(x, \xi)$.

Thus, in both cases, we have

$$\xi \cdot \nabla_x f^{(0)}(x, \xi) = -\mu_t(x) f^{(0)}(x, \xi) \quad (11)$$

for all $(x, \xi) \in \Omega \times S^{d-1}$.

Since not only $f^{(1)}$ but also G , appeared in the proof of Lemma 1, is bounded continuous on $\Omega \times S^{d-1}$, the function

$$\begin{aligned} f^{(1)}(x + t\xi, \xi) &= \int_0^{\tau_-(x + t\xi, \xi)} \mu_s(x + t\xi - s\xi) \exp\left(-M_t(x + t\xi, \xi; s)\right) G(x + t\xi - s\xi, \xi) ds \\ &= \int_{-t}^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp\left(M_t(x, \xi; -t) - M_t(x, \xi; s)\right) G(x - s\xi, \xi) ds \end{aligned}$$

is differentiable with respect to t at $t = 0$ for all $(x, \xi) \in \Omega \times S^{d-1}$ and

$$\begin{aligned} \xi \cdot \nabla_x f^{(1)}(x, \xi) &= \mu_s(x, \xi) G(x, \xi) \\ &\quad - \mu_t(x) \int_0^{\tau_-(x, \xi)} \mu_s(x - s\xi) \exp\left(-M_t(x, \xi; s)\right) G(x - s\xi, \xi) ds \\ &= \mu_s(x) \int_{S^{d-1}} p(x, \xi, \xi') f^{(0)}(x, \xi') d\sigma_{\xi'} - \mu_t(x) f^{(1)}(x, \xi) \end{aligned} \quad (12)$$

for all $(x, \xi) \in \Omega \times S^{d-1}$.

Since the functions $f^{(n)}$ are bounded continuous on X for all $n \in \mathbb{N}$ by Lemma 3, the following relation holds from the direct calculation:

$$\xi \cdot \nabla_x f^{(n+1)}(x, \xi) = \mu_s(x) \int_{S^{d-1}} p(x, \xi, \xi') f^{(n)}(x, \xi') d\sigma_{\xi'} - \mu_t(x) f^{(n+1)}(x, \xi) \quad (13)$$

for all $(x, \xi) \in \Omega \times S^{d-1}$ and for all $n \in \mathbb{N}$.

By Proposition 3, we can sum up right hand sides of (11), (12), and (13) to obtain

$$\begin{aligned} \xi \cdot \nabla_x f(x, \xi) &= \xi \cdot \nabla_x \sum_{n=0}^{\infty} f^{(n)}(x, \xi) = \sum_{n=0}^{\infty} \xi \cdot \nabla_x f^{(n)}(x, \xi) \\ &= \sum_{n=0}^{\infty} \mu_s(x) \int_{S^{d-1}} p(x, \xi, \xi') f^{(n)}(x, \xi') d\sigma_{\xi'} - \sum_{n=0}^{\infty} \mu_t(x) f^{(n)}(x, \xi) \\ &= \mu_s(x) \int_{S^{d-1}} p(x, \xi, \xi') \sum_{n=0}^{\infty} f^{(n)}(x, \xi') d\sigma_{\xi'} - \mu_t(x) \sum_{n=0}^{\infty} f^{(n)}(x, \xi) \\ &= \mu_s(x) \int_{S^{d-1}} p(x, \xi, \xi') f(x, \xi') d\sigma_{\xi'} - \mu_t(x) f(x, \xi). \end{aligned}$$

for all $(x, \xi) \in \Omega \times S^{d-1}$, which is the stationary transport equation (1) itself. Thus, the directional derivative $\xi \cdot \nabla_x f(x, \xi)$ is defined for all $(x, \xi) \in \Omega \times S^{d-1}$ by termwise differentiation and the original function f satisfies the stationary transport equation (1). $\square$

Proof of Proposition 6. For all $(x, \xi) \in \Gamma_-$,

$$f^{(n)}(x, \xi) = \begin{cases} f_0(x, \xi), & n = 0, \\ 0, & n \geq 1. \end{cases}$$

Therefore,

$$f(x, \xi) = \sum_{n=0}^{\infty} f^{(n)}(x, \xi) = f_0(x, \xi)$$

for all $(x, \xi) \in \Gamma_-$. $\square$

From proposition 5 and Proposition 6, it follows that the boundary value problem (1)-(2) and integral equations (4)-(5) are equivalent in this setting.

5 Example for nonequivalence between the stationary transport equation and derived integral equations

In this section, we introduce an example in two dimensional case which shows that piecewise continuity of the boundary data is not a sufficient condition for the main result. Let $d = 2$ and fix $\bar{x} = (\bar{x}_1, \bar{x}_2) \in \Omega$. We introduce the polar coordinate to S^1 :

$$\xi(\theta) = (\cos \theta, \sin \theta), \quad \theta \in [0, 2\pi).$$

We note that, by this coordinate, S_+^1 and S_-^1 are identified with intervals $(0, \pi)$ and $(\pi, 2\pi)$, respectively. We introduce pieces of Γ_- by

$$\begin{aligned} \Gamma_{-,1} &:= \{(x, \xi(\theta)) \in \Gamma_- | x_2 = 0, x_1 \geq \bar{x}_1 - \bar{x}_2 \cot \theta, \theta \in (0, \pi)\}, \\ \Gamma_{-,2} &:= \{(x, \xi(\theta)) \in \Gamma_- | x_2 = 0, x_1 < \bar{x}_1 - \bar{x}_2 \cot \theta, \theta \in (0, \pi)\}, \\ \Gamma_{-,3} &:= \{(x, \xi(\theta)) \in \Gamma_- | x_2 = 1, \theta \in (\pi, 2\pi)\}. \end{aligned}$$

We note that $\Gamma_- = \Gamma_{-,1} \cup \Gamma_{-,2} \cup \Gamma_{-,3}$ and $\Gamma_{-,1} \cap \Gamma_{-,2} = \Gamma_{-,2} \cap \Gamma_{-,3} = \Gamma_{-,3} \cap \Gamma_{-,1} = \emptyset$.

We take the boundary data f_0 as follows:

$$f_0(x, \xi) := \begin{cases} 1, & (x, \xi) \in \Gamma_{-,1}, \\ 0, & (x, \xi) \in \Gamma_{-,2} \cup \Gamma_{-,3}. \end{cases}$$

The boundary data f_0 is obviously bounded and constant on each $\Gamma_{-,i}$, $i = 1, 2, 3$, which implies that the boundary data f_0 is indeed piecewise continuous.

With this boundary data f_0 , we define a family of functions $\{f^{(n)}\}_{n \geq 0}$ on X by recursive formulae (7)-(8). Through the same discussion in Section 2, we see that the sum $f = \sum_{n=0}^{\infty} f^{(n)}$ is still the unique solution to the integral equations (4)-(5). However, the directional derivative $\xi \cdot \nabla_x f(x, \xi)$ of this sum f is not defined at $(\bar{x}, \xi)$ for all $\xi \in S^1$.

In this setting, the function G_- , defined by the formula (10) in Section 3, is identically zero, which implies that G_- is continuous in $\Omega \times S^1$, while G_+ , also defined by the formula (10) in Section 3, is discontinuous with respect to x at $(\bar{x}, \xi)$ for all $\xi \in S^1$. Thus, $G = G_+ + G_-$ is also discontinuous with respect to x at $(\bar{x}, \xi)$ for all $\xi \in S^1$. Although G is discontinuous with respect to x at one point, $f^{(1)}$ is continuous on X . This implies that the directional derivative $\xi \cdot \nabla_x f^{(n)}(x, \xi)$ is defined for all $(x, \xi) \in \Omega \times S^1$ and for all $n \geq 2$, whereas $\xi \cdot \nabla_x f^{(1)}(x, \xi)$ is not defined at $(\bar{x}, \xi) \in \Omega \times S^1$. Since $\xi \cdot \nabla_x f^{(0)}(x, \xi)$ is defined for all $(x, \xi) \in \Omega \times S^1$, we have to conclude that $\xi \cdot \nabla_x \sum_{n=0}^{\infty} f^{(n)}(x, \xi)$ is not defined at $(\bar{x}, \xi) \in \Omega \times S^1$, which means that the sum $f(x, \xi) = \sum_{n=0}^{\infty} f^{(n)}(x, \xi)$ is not a solution to the boundary value problem (1)-(2). This conclusion implies that this boundary value problem (1)-(2) has no solution.

Acknowledgements The authors would like to thank Emeritus Professor Kazuo Aoki for suggesting this problem.

References

- Anikonov, D. S., Prokhorov, I. V., Kovtanyuk, A. E., 1993. Investigation of scattering and absorbing media by the methods of X-ray tomography, *J. Inv. Ill-Posed Problems*, 1, no. 4, pp. 259–281.
Aoki, K., Bardos, C., Dogbe, C., Golse, F., 2001. A note on the propagation of boundary induced discontinuities in kinetic theory, *Math. Models Methods Appl. Sci.*, 11, no. 9, pp. 1581–1595.

- Aoki, K., Takata, S., Aikawa, H, Golse F., 2001. A rarefied gas flow caused by a discontinuous wall temperature *Phys. Fluids*, 13, no. 9, pp. 2645–2661.
- Case, K. M., Zweifel, P. F., 1967. *Linear Transport Theory*, Reading, Mass.: Addison-Wesley Pub. Co.
- Chandrasekhar, S. 1960. (1950) *Radiative Transfer*, Reprint, New York: Dover Publications, Inc.
- Esposito, R., Guo, Y., Kim, C., Marra, R., 2013. Non-isothermal boundary in the Boltzmann theory and Fourier law, *Commun. Math. Phys.*, 323, pp. 177–239.
- Kim, C., 2011. Formation and propagation of discontinuity for Boltzmann equation in non-convex domains, *Commun. Math. Phys.*, 308, pp. 641–701.
- Guo, Y., Yang, X., 2016, Existence and BV-regularity for neutron transport equation in nonconvex domain, *SIAM J. Math. Anal.*, 48, no. 5, pp. 3467–3495.