TITLE:

Ambient fine and coarse particles in Japan affect nasal and bronchial epithelial cells differently and elicit varying immune response

AUTHOR(S):

Onishi, Toshinori; Honda, Akiko; Tanaka, Michitaka; Chowdhury, Pratiti H.; Okano, Hitoshi; Okuda, Tomoaki; Shishido, Daiki; ... Yasuda, Makoto; Hirano, Shigeru; Takano, Hirohisa

CITATION:

Onishi, Toshinori ...[et al]. Ambient fine and coarse particles in Japan affect nasal and bronchial epithelial cells differently and elicit varying immune response. Environmental Pollution 2018, 242(Part B): 1693-1701

ISSUE DATE:

2018-11

URL:

<http://hdl.handle.net/2433/243828>

RIGHT:

© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>; The full-text file will be made open to the public on 1 November 2020 in accordance with publisher's 'Terms and Conditions for Self-Archiving'. ; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。 ; This is not the published version. Please cite only the published version.

Ambient fine and coarse particles in Japan affect nasal and bronchial epithelial cells differently and elicit varying immune response

Toshinori Onishi^{1, 2}, Akiko Honda^{1*}, Michitaka Tanaka¹, Pratiti H Chowdhury¹, Hitoshi Okano¹, Tomoaki Okuda³, Daiki Shishido³, Yoshihiro Terui³, Shuichi Hasegawa⁴, Takayuki Kameda⁵, Susumu Tohno⁵, Masahiko Hayashi⁶, Chiharu Nishita-Hara⁶, Keiichiro Hara⁶, Kozo Inoue⁷, Yasuda Makoto², Shigeru Hirano², Hirohisa Takano¹

¹ Environmental Health Division, Department of Environmental Engineering, Graduate School of Engineering, Kyoto University, Kyoto, Japan

² Department of Otolaryngology-Head and Neck Surgery, Kyoto Prefectural University of Medicine, Kyoto, Japan

³ Faculty of Science and Technology, Keio University, Kanagawa, Japan

⁴ Center for Environmental Science in Saitama, Japan

⁵ Graduate School of Energy Science, Kyoto University, Japan

⁶ Fukuoka Institute of Atmospheric Environment and Health, Fukuoka University, Japan

⁷ Tokyo Dylec Corp, Japan

***Corresponding author:** Dr. Akiko Honda, Environmental Health Division, Department of Environmental Engineering, Graduate School of Engineering, Kyoto University. C Cluster, Katsura Campus, Nishikyo-ku, Kyoto 615-8540, Japan.

Tel.: +81-75-383-3345; Fax: +81-75-383-3344

E-mail: akko@health.env.kyoto-u.ac.jp

23

24 **Abstract**

25 Ambient particulate matter (PM) epidemiologically exacerbates respiratory and immune
26 health, including allergic rhinitis (AR) and bronchial asthma (BA). Although fine and coarse
27 particles can affect respiratory tract, the differences in their effects on the upper and lower
28 respiratory tract and immune system, their underlying mechanism, and the components
29 responsible for the adverse health effects have not been yet completely elucidated. In this
30 study, ambient fine and coarse particles were collected at three different locations in Japan
31 by cyclone technique. Both particles collected at all locations decreased the viability of nasal
32 epithelial cells and antigen presenting cells (APCs), increased the production of IL-6, IL-8,
33 and IL-1 β from bronchial epithelial cells and APCs, and induced expression of dendritic and
34 epithelial cell (DEC) 205 on APCs. Differences in inflammatory responses, but not in
35 cytotoxicity, were shown between both particles, and among three locations. Some
36 components such as Ti, Co, Zn, Pb, As, OC (organic carbon) and EC (elemental carbon)
37 showed significant correlations to inflammatory responses or cytotoxicity. These results
38 suggest that ambient fine and coarse particles differently affect nasal and bronchial epithelial
39 cells and immune response, which may depend on particles size diameter, chemical
40 composition and source related particles types.

41

42 **Keywords:** ambient particulate matter, cyclone technique, respiratory cells, immune cells,
43 inflammatory responses

44 **Capsule:** We showed for the first time in the world that ambient fine and coarse particles

collected from Japan by the new technique using cyclone have different effects on the epithelium cells of the upper and lower respiratory tract and elicit varying immune response, which may depend on particles size diameter, chemical composition and source related particles types.

1. Introduction

The health effect of ambient particulate matter (PM) is still a problem worldwide. PM is a complex mixture of particles having different chemical components such as solid and liquid materials that contain elemental carbon (EC), organic carbon (OC), inorganic salts, and metals and biological components such as endotoxin and β -glucan and has a compound effect on biological reactions (Schins et al., 2004; Cachon et al., 2014; Honda et al., 2017). Generally, the fine fraction of PM (aerodynamic diameter $< 2.5 \mu\text{m}$) in urban atmosphere is a complex mixture of primary particles emitted from combustion sources and secondary particles that form in the atmosphere from gaseous components (Marcazzan et al., 2001; Sharma et al., 2007; Sevastyanova et al., 2008; Zerbi et al., 2008). The coarse fraction of PM (aerodynamic diameter $> 2.5 \mu\text{m}$) generally includes mineral particles of crustal material, sea salt particles, fly ash, and adsorbed species such as endotoxin (Schins et al., 2004; Perez et al., 2007). These components can differ depending on the sources, geographical areas, and seasons. In addition, PM composition depends on factors such as atmospheric photochemical reaction and physical redistribution (Vecchi et al., 2004; Samoli et al., 2008). PM epidemiologically exacerbates respiratory and immune health such as allergic rhinitis (AR) and bronchial asthma (BA) (Tecer et al., 2008) in addition to cardiovascular diseases

and cancer (Kappos et al., 2004). Clinically, AR and BA have a close relationship: about 80% of patients with BA have complications of AR (Bachert et al., 2002). In general, coarse particles and limited fine particles can affect upper respiratory tract, whereas fine particles and limited coarse particles can affect lower respiratory tract (Heyder J., 1986). However, the difference in the effects of fine and coarse particles on the upper or lower respiratory tract and immune responses related to them, as well as their underlying mechanisms have not yet been clarified. Moreover, the components of PM responsible for the adverse health effects have not yet been elucidated owing to their complexity (Lindbom et al., 2006; Hong et al., 2016).

A large amount of fine and coarse particles is needed to evaluate the adverse health effects by *in vivo* and/or *in vitro* studies. However, it is difficult to collect a sufficient amount of PM by conventional filter collection method with extraction. Because of different extraction efficiency and loss of PM constituents, the exposure experiment using PM extracts has a possibility that would not reflect the actual biological response. Our previous study disclosed extracts efficiency of PM_{2.5} and discussed the problem (Chowdhury et al., 2018). On the other hand, the cyclone technique enables collection of a sufficient amount of PM (fine and coarse particles themselves) for *in vivo* and/or *in vitro* assays enabling the analysis of the effects of ambient particles on respiratory health without the use of a filter or extraction process (Okuda et al., 2015, 2018).

In this study, we investigated the effects of ambient fine and coarse particles collected at three Japanese locations by cyclone technique on nasal epithelial cells (RPMI-2650), bronchial epithelial cells (BEAS-2B), and bone marrow derived antigen presenting cells

(APCs) from NC/Nga mice. Our aim was to estimate the different effects of ambient fine and coarse particles on respiratory and immune cells, their underlying mechanism, and the components which can be responsible for the respiratory and immune health such as AR and BA.

2. Materials and Methods

2.1. Sampling of PM

Samples of fine and coarse particles were collected at an urban area in Fukuoka City, at a suburban of the metropolitan area in Kazo City, Saitama Prefecture (Saitama), and a capital area in Yokohama City in Japan (Suppl. Figure S1) during February to March 2017. The particles as references were obtained by National Institute for Environmental Studies in Japan. One reference (CRM#8) is ethanol-treated vehicle exhaust particulates (Okamoto., 1987) and another (CRM#28) is irradiated atmospheric dust collected by a ventilation filter of the building in Beijing (Mori et al., 2008). Okuda (2013) has indicated CRM#8 consists mainly of fine (or ultrafine) particles, while CRM#28 consist mainly of coarse particles.

The collection was conducted with a high-volume PM sampler using the virtual impactor and cyclone technique with no filter or extraction process (Okuda et al., 2018). The air flow volume per given time for the inlet (virtual impactor) is 1,300 L/min. The total volume of air sampled was determined from the measured volumetric flow rate and the sampling time. The mass concentration of particles in the ambient air was computed as the total mass of collected particles divided by the total volume of air sampled. After sampling, the particles in the amber bottles were collected using a stainless spatula. We previously confirmed size distribution

and morphology of ambient particles collected by cyclone (Suppl. Figure S2). Particles were dissolved in sterile phosphate-buffered saline (PBS) and ultrasonicated at the concentration of 10 mg/mL. Finally, we adjusted at concentrations of 0, 7.5, and 75 µg/mL using medium, PBS (1%) and Dimethyl sulfoxide (DMSO) (0.1%) for the cell exposure experiment in this study. Medium for BEAS-2B cells is serum-free. Similarly, we did not add serum in medium for RPMI-2650 cells to evaluate under the same condition of exposure.

2.2. Chemical, mineralogical and biochemical investigation

The collected particles was characterized by ion chromatography for Anion species (Cl^- , NO_3^- , and SO_4^{2-}) and cation species (Na^+ , NH_4^+ , K^+ , Mg^{2+} , and Ca^{2+}), thermal-optical method (IMPROVE protocol) for OC1-4 and EC1-3, high performance liquid chromatography (HPLC) for polycyclic aromatic hydrocarbons (PAHs) (Chrysene, Benz[a]anthracene, Benzo[b]fluoranthene, Benzo[k]fluoranthene, and Benzo[a]pyrene), and inductively coupled plasma mass spectrometry (ICP-MS) for metals (Al, Si, Ti, V, Cr, Mn, Fe, Ni, Cu, Zn and Pb). The procedure of chemical characterization mentioned above were generally described in several previous papers (Okuda., 2013; Okuda et al., 2013, 2014). Endotoxin and β-glucan have induced inflammatory responses from respiratory cells and immune cells (Veranth et al., 2004; Carmona et al., 2010; Neveu et al., 2011). In this study, we investigated the effect of endotoxin and β-glucan as substances derived from biological components in PM. We performed an endotoxin test and a β-glucan test (both from Associates of Cape Cod, Falmouth, MA, USA) following the manufacturer's instructions.

2.3. Cell Cultures and PM exposure

2.3.1. Upper and Lower Respiratory cells

The RPMI-2650, derived from squamous cell carcinoma of nasal septum was used as model of human nasal epithelial cells which are cells of the upper respiratory tract. These cells display consistent growth and high stability throughout continued culturing *in vitro* with no alteration to the normal diploid karyotype (Moorhead, 1965). The cell line was purchased from the European Collection of Cell Cultures (Salisbury, Wiltshire, United Kingdom) and maintained in Eagle's minimal essential medium (DS Pharma Biomedicals, Osaka, Japan) supplemented with 10% heat-inactivated fetal bovine serum (MP Biomedicals, Eschwege, Germany), 2 mM L-glutamine, 100 U/mL penicillin and 100 µg/mL streptomycin (Sigma, St Louis, Missouri). As representative of the cells of the lower respiratory tract, the BEAS-2B, derived from human bronchial epithelial cells, was purchased from the European Collection of Cell Cultures and maintained in LHC-9 medium (Thermo Scientific, Waltham, Massachusetts) which is serum-free medium containing Gentamicin. RPMI-2650 cells and BEAS-2B cells were maintained by subculture in 37°C at 5% CO₂ in medium.

2.3.2. Immune cells

Ten-week-old male SPF NC/NgaTndCrlj mice were purchased from Charles River (Osaka, Japan). NC/Nga mice are atopy-prone mice. APCs were obtained after sacrificing mice by cervical dislocation and exsanguination. The procedures used in all animal studies were approved by the Animal Research Committee at Kyoto University. APCs were differentiated using a modification of the protocol provided by Lutz et al (1999). We confirmed APCs by

the expression of about 80% of CD11c which is a molecule specifically expressed in dendritic cells. Bone marrow cells ($4 \times 10^5/\text{mL}$) were cultured in R10 which is RPMI 1640 (Thermo Scientific) supplemented with 10% heat-inactivated fetal bovine serum (MP Biomedicals), 100 U/mL penicillin, 100 $\mu\text{g}/\text{mL}$ streptomycin (Sigma), and 50 mM 2-mercaptoethanol (Thermo Scientific) containing Granulocyte-Macrophage Colony-Stimulating Factor (GM-CSF).

2.4. Experimental Protocol

The RPMI-2650 cells, BEAS-2B cells, and APCs were exposed to ambient fine and coarse particles at concentrations of 0, 7.5, or 75 $\mu\text{g}/\text{mL}$ and reference particles at 75 $\mu\text{g}/\text{mL}$ only for 24 h. We evaluated the cell viability, the cytokine release, and dendritic and epithelial cell (DEC) 205 on the cell surface. All control cells were treated with each medium.

2.4.1. Cell Viability

We measured the viability of the RPMI-2650 cells, BEAS-2B cells, and APCs by WST-1 assay using the Premix WST-1 Cell Proliferation Assay System (TaKaRa Bio, Shiga, Japan) as previously described (Honda et al., 2017). The results are expressed as the percentage of exposed group to control cells (0 $\mu\text{g}/\text{mL}$).

2.4.2. Quantification of Pro-Inflammatory Cytokines in the Culture Supernatants

The amounts of IL-6 and IL-8 release in the supernatants from the RPMI-2650 cells and BEAS-2B cells and those of IL-6 and IL-1 β release in the supernatants from APCs were

measured by ELISA (Thermo Scientific), according to the manufacturer's protocol as previously described (Honda et al., 2017). The detection limits of IL-6 and IL-8 from RPMI-2650 cells and BEAS-2B cells, and IL-6 and IL-1 β from APCs were <2.2 pg/mL, <9.8 pg/mL, <1.9 pg/mL, <1.6 pg/mL, <16 pg/mL and 10 pg/mL, respectively.

2.4.3. Expression of DEC205 on APCs cell surface

We measured the expression of DEC205 on the APCs' surface by the FACS analysis, the following monoclonal antibodies were used: Mouse BD Fc Block purified anti-mouse CD16/CD32 (Becton Dickinson), DEC205 (NLDC-145, PE-conjugated; Bio-Legend, San Diego, California), Rat IgG2a, k Isotype Control (RTK2758, PE-conjugated; BioLegend). The fluorescence was measured by a FACSCalibur (Becton Dickinson) as previously described (Honda et al., 2017).

2.5. Statistical Analysis

The data are presented as the mean $\pm$ standard error of the mean (SEM) for each experimental group (n =3 or 4). The significance of variation among different groups was determined by one-way analysis of variance. Differences among groups were analyzed using Tukey's multiple comparison test. A P-value < 0.05 was considered to indicate a significant difference. Relationships between components in PM and cell viability or cytokine release were tested using Pearson's correlation, with a two-tailed significance study using SPSS software. A P < 0.01 and R > 0.9 was shown as a high degree of correlation.

3. Results

3.1. The characterization of collected ambient particles

This cyclone system achieved 50% collection efficiency with components having the following aerodynamic cut-off diameters: virtual impactor, 2.4 μm ; fine-particle cyclone, 0.18-0.30 μm ; and coarse-particle cyclone, 0.7 μm . Particles smaller than 2.4 μm flowed to the fine side at the virtual impactor part, and thus, fine particles were 0.30-2.4 μm and coarse particles were 2.4 μm or more in size. The mean concentrations of fine particles in Fukuoka, Saitama, and Yokohama were 3.0, 5.9 and 9.9 $\mu\text{g}/\text{m}^3$, respectively while those of coarse particles were 1.5, 4.3 and 14.2 $\mu\text{g}/\text{m}^3$, respectively. Note that these mass concentrations of fine and coarse particles were expressed as the weights of particles collected by the cyclones per sampled air volume. The concentrations of components of both particles in the prepared solution are shown (Suppl. Table S1-S3). Both particles had different proportions of metal components between Fukuoka and the other locations. In Fukuoka, instead of a small amount of metal components, amounts of Na^+ , Cl^- and SO_4^{2-} were high (Suppl. Figure S3).

The content of endotoxin in fine and coarse particles was 0.080 and 0.060 (Fukuoka), 2.50 and 2.97 (Saitama), 4.29 and 6.59 (Yokohama) EU/mL, respectively. Endotoxin in both particles at Fukuoka was lower than those collected at Saitama and Yokohama. The highest content of endotoxin was found in coarse particles at Yokohama. The contents of β -glucan in fine and coarse particles were 317.3 and 1106 (Fukuoka), 294.4 and 666.0 (Saitama), 774.6 and 1060 (Yokohama) pg/mL, respectively. β -glucan in fine particles collected at all locations was lower than those in coarse particles at each location. The highest content of β -glucan was present in coarse particles at Fukuoka.

221

222 **3.2. Biological effects of ambient particles on the Nasal Epithelial Cells**

223 We examined the effects of exposure (24 h) to ambient fine and coarse particles on the
224 viability of RPMI-2650 cells. A significant decrease ($p < 0.01$ vs. control) in the viability of
225 cells was seen upon exposure to both particles collected at all locations in a concentration-
226 dependent manner as compared to the control (unexposed cells). No difference in cytotoxicity
227 caused by ambient fine and coarse particles from each location was detected except for
228 sample collected at Yokohama at a concentration of $7.5 \mu\text{g/mL}$ (Figure 1).

229 We investigated the effects of 24 h exposure of ambient fine and coarse particles on the pro-
230 inflammatory responses via release of IL-6 and IL-8 from RPMI-2650 cells. None of the
231 particles, at any location were able to evoke detectable cytokine release from RPMI-2650
232 cells.

233

234 **3.3. Biological effects of ambient particles on the Bronchial Epithelial Cells**

235 We examined the effects of exposure to ambient fine and coarse particles for 24 h on the
236 viability of BEAS-2B cells. BEAS-2B cells showed no decrease in viability upon exposure
237 to both particles collected at all locations at any concentration, when compared to control
238 (unexposed cells) (Figure 2A).

239 We investigated the effects of ambient fine and coarse particles on the pro-inflammatory
240 responses via release of IL-6 and IL-8 from BEAS-2B cells after exposure to each particle
241 for 24 h. Both particles collected at all locations increased IL-6 release, in a dose-dependent
242 manner, especially at the concentration of $75 \mu\text{g/mL}$ (Figure 2B; $p < 0.01$ vs. control).

Comparison between fine and coarse particles at the same location indicated that coarse particles at Fukuoka induced higher production of IL-6 than fine particles (Figure 2B; $p < 0.01$ vs. fine particles at $75 \mu\text{g/mL}$). Comparison among the three locations indicated that fine particles collected at Saitama or Yokohama had greater effect than those at Fukuoka (Figure 2B; $p < 0.01$ vs. fine particles at Fukuoka at $75 \mu\text{g/mL}$). Coarse particles collected at Fukuoka had significantly larger effect than those collected at Saitama (Figure 2B; $p < 0.05$ vs. coarse particles at Fukuoka at $75 \mu\text{g/mL}$). The production of IL-8 showed a pattern similar to that of IL-6. The results of IL-8 different from IL-6 are shown. Coarse particles collected at Fukuoka, fine particles collected at Saitama and both particles collected at Yokohama significantly increased the levels of IL-8 at a concentration of $7.5 \mu\text{g/mL}$ (Figure 2B; $p < 0.05$ or 0.01 vs. control). The fine particles collected at Yokohama showed significantly marked induction of IL-8 than coarse particles (Figure 2B; $p < 0.05$ vs. fine particles at $75 \mu\text{g/mL}$). Fine particles collected at Yokohama had greater effect than those collected at Saitama (Figure 2B; $p < 0.01$ vs. fine particles at Saitama at $75 \mu\text{g/mL}$).

3.4. Biological effects of ambient particles on the APCs

We examined the effects of ambient fine and coarse particles on the viability of APCs after exposure to PM for 24 h. Exposure to both particles resulted in decrease in viability of APCs in a dose-dependent manner as compared to control (unexposed cells) (Figure 3A). A significant decrease in the viability of cells was seen upon exposure to both particles collected at all locations at a concentration of $75 \mu\text{g/mL}$ and for those at Saitama and Yokohama at the concentration of $7.5 \mu\text{g/mL}$ (Figure 3A; $p < 0.05$ or 0.01 vs. control). Coarse particles

collected at Fukuoka slightly decreased the viability of cells at a concentration of 7.5 $\mu\text{g/mL}$ (Figure 3A; $p < 0.05$ vs. control). No difference in cytotoxicity was detected between fine and coarse particles collected at each location.

We investigated the effects of ambient fine and coarse particles on the pro-inflammatory responses analyzed as IL-6 and IL-1 β released from APCs after exposure to each particle for 24 h. Both particles collected at all locations increased IL-6 release in a dose-dependent manner especially at a concentration of 75 $\mu\text{g/mL}$ (Figure 3B; $p < 0.01$ vs. control). Both particles collected at Saitama and Yokohama also significantly increased the levels of IL-6 at the concentration of 7.5 $\mu\text{g/mL}$ (Figure 3B; $p < 0.01$ vs. control). Coarse particles collected at Fukuoka induced higher production of IL-6 than the fine particles (Figure 3B; $p < 0.01$ vs. fine particles at 75 $\mu\text{g/mL}$). Both particles collected at Saitama and Yokohama induced higher production of IL-6 than that collected at Fukuoka (Figure 3B; $p < 0.05$ or 0.01 vs. fine particles at Fukuoka at 7.5 and 75 $\mu\text{g/mL}$). Both particles at all locations increased IL-1 β release, in a dose –dependent manner, especially at the concentration of 75 $\mu\text{g/mL}$ similar to what was observed in case of IL-6 (Figure 3B; $p < 0.01$ vs. control). Coarse particles collected at Saitama and both particles at Yokohama at a concentration of 7.5 $\mu\text{g/mL}$ increased the levels of IL-1 β (Fig 3B; $p < 0.05$ or 0.01 vs. control). Coarse particles collected at Fukuoka induced lower production of IL-1 β than the fine particles (Figure 3B; $p < 0.05$ vs. fine particles at 75 $\mu\text{g/mL}$). Both particles at Fukuoka resulted in greater induction of IL-1 β than those at Saitama and Yokohama (Figure 3B; $p < 0.05$ or 0.01 vs. both particles at Fukuoka at 75 $\mu\text{g/mL}$, respectively).

The expression patterns of DEC205 in APCs were examined in order to evaluate the effects

of exposure to ambient fine and coarse particles for 24 h on the maturation and activation of APCs. DEC205 is a member of the macrophage mannose receptor family. This molecule is known to mediate the capture and internalization of ligands for subsequent processing and presentation by APCs (Jiang et al., 1995). Both particles collected at all locations increased the ratio of DEC205-positive cells at concentrations of 7.5 and 75 $\mu\text{g/mL}$ (Figure 3C; $p < 0.01$ vs. control). Although, there were no significant differences in the ratio of DEC205-positive cells exposed to the both particles collected at the same location, fine particles collected at Yokohama significantly increased the ratio of DEC205-positive cells when compared with those at Fukuoka (Figure 3C; $p < 0.05$ vs. fine particles at Fukuoka at 7.5 $\mu\text{g/mL}$).

3.5. Correlation between ambient particles components and biological responses

Determination of the PM components responsible for impacts on examined cell lines by assessing cytotoxicity, pro-inflammatory cytokine release is very important. We evaluated Pearson's correlation coefficients between both particles compounds and the cell viability of RPMI-2650 cells and APCs, and both particles compounds and cytokine release from BEAS-2B cells and APCs. Our analysis showed negative correlations between the cell viability of RPMI-2650 cells and Ca^{2+} , Zn, and OC (Suppl. Figure S4). Positive correlations were observed between IL-6 and IL-8 release from BEAS-2B cells and IL-6 release from APCs and multiple components, including EC, OC, and metals such as Ti and Co (Suppl. Figure S5, S6A). Whereas, positive correlations between IL-1 β from APCs and multiple components, including ions, EC, OC, and metals such as As and Pb were shown (Suppl.

309 Figure S6B).

310

311 4. Discussion

312 In the present study, we conducted experiments using ambient fine and coarse particles
313 collected by cyclone technique. Ambient both particles decreased the cell viability of RPMI-
314 2650 cells and APCs, and induced production of pro-inflammatory cytokines from BEAS-
315 2B cells and APCs. There was a significant difference in the inflammatory response elicited
316 by fine and coarse particles at Fukuoka. Inflammatory responses induced by particles
317 collected at Saitama and Yokohama were similar to each other but different from those
318 observed for particles collected at Fukuoka. Ambient both particles also induced expression
319 of DEC205 on APCs. Viability of RPMI-2650 cells correlated negatively with Ca^{2+} , Zn, and
320 OC3 and OC4. IL-6 and IL-8 release correlated positively with Ti, Fe, Co, Cr, Mn, V, and Zn,
321 and OC3, OC4, and EC2, whereas IL-1 β release correlated positively with As, Pb, and OC2
322 and EC1.

323 Ambient both particles at all locations decreased viability of RPMI-2650 cells but did not
324 induce pro-inflammatory cytokines. Nasal epithelial cells are the first epithelial barrier in the
325 nasal cavity imparting protection from inhaled xenobiotics. Previous studies have indicated
326 that diesel exhaust particles (DEP) or PM10 do not show cytotoxicity nor induce production
327 of pro-inflammatory cytokines from RPMI-2650 cells (Lindbom et al., 2006) and that DEP
328 does not reduce cell viability but decrease the barrier function by reducing zonula occludens
329 -1 (ZO-1) expression in RPMI-2650 cells (Fukuoka et al., 2015). If the barrier function
330 decreases even without cytotoxicity, there is a possibility that PM further weaken the barrier

331 function due to cytotoxicity. It is possible that particles collected in this study weakened the
332 barrier function, thereby allowing easy invasion by allergens.

333 It has been reported that primary nasal epithelial cells have lesser Toll-like receptor (TLR)
334 expression compared to alveolar epithelial cells, and more distribution of Toll-interacting
335 protein (TOLLIP; an inhibitor of TLR signaling) which may be one of the reasons for non-
336 induction of pro-inflammatory cytokines by these cells (Moncayo et al., 2014). The low
337 inflammatory responses due to the low expression of TLR and the high expression of
338 TOLLIP on nasal epithelial cells may lead to colonization and coexistence of many resident
339 bacterial groups in the nasal cavity environment. The epithelial cells in the nasal cavity are
340 always exposed to external environment and various bacteria or chemicals therein. Hence,
341 tolerance to these bacteria without induction of inflammation is required to maintain
342 homeostasis.

343 In case of BEAS-2B cells, though ambient both particles at all locations elevated the levels
344 of IL-6 and IL-8 release, none of them decreased the cell viability. Ambient both particles
345 strongly induced pro-inflammatory cytokine production compared to both reference particles.

346 Bronchial epithelial cells also act as a physical barrier and generate biological and
347 immunological responses against inhaled xenobiotics. IL-6 and IL-8 are the major pro-
348 inflammatory cytokines induced by response to environmental insults and are important key
349 molecules causing acute inflammation in the respiratory system by stimulating lymphocytes
350 (Thacker, 2006). Previous studies have indicated that various chemicals and allergens
351 stimulate IL-6 or IL-8 production from BEAS-2B cells (Honda et al., 2014; Totlandsdal et
352 al., 2012; Park et al., 2009). Ambient both particles collected by cyclone technique can

353 exacerbate allergic inflammation by inducing inflammatory cytokines.

354 Extracts from PM on filter have been used for experiments dealing with health effects of PM.

355 To our knowledge, this is the first experimental demonstration of the effects of allergic

356 inflammation by ambient fine and coarse particles collected by cyclone technique.

357 Fuentes-Mattei et al. (2010) have showed that PM_{2.5} organic extracts from urban area in

358 Puerto Rico at 50 µg/mL decrease about 20% of cell viability and produce about 650 pg/mL

359 of IL-6 and 235 pg/mL of IL-8 from BEAS-2B cells, but those from rural area do not decrease

360 cell viability and induce the lower production of IL-6 compared to those from urban area.

361 Gualtieri et al. (2010) have showed that PM_{2.5} and PM₁₀ aqueous extracts from urban area

362 in Milan at 10 µg/cm² (in this study, 75 µg/mL=13.7 µg/cm²) do not decrease cell viability

363 and produce about 80 pg/mL and 400 pg/mL of IL-8 from BEAS-2B, respectively. We have

364 previously indicated that exposure to PM_{2.5} collected in winter and subjected to organic

365 extraction rather than aqueous extraction causes an inflammatory response via IL-6

366 production from bronchial epithelial cells and PM_{2.5} extracts at Fukuoka subjected to

367 organic extraction, produced about 110 pg/mL of IL-6 at a concentration of 75 µg/mL which

368 was about 2 times greater than the level of control (Honda et al., 2017). These same extracts

369 collected in another season produced about 15 pg/mL of IL-6 and about 200 pg/mL of IL-8

370 from bronchial epithelial cells, and the levels of these cytokines were less than control

371 (Chowdhury et al., 2018). In the present experiment, fine particles collected by cyclone

372 technique at Fukuoka at 75 µg/mL produced about 250 pg/mL of IL-6 and about 670 pg/mL

373 of IL-8 in BEAS-2B cells. The levels of IL-6 and IL-8 produced by fine particles collected

374 using cyclone technique was about 6 and 2.5 times more than the level of control, respectively.

375 Although we could not strictly compare these results because it is not exactly the same
376 particle, it is suggested that ambient particles collected by cyclone technique could induce
377 larger inflammatory response than that induced by PM collected by the conventional filter
378 technique with extraction. Cyclone technique is an efficient method for collecting particles
379 that are subjected to exposure studies. Currently, there exist only a few assessments of health
380 effect conducted using ambient particles collected by cyclone technique and hence, further
381 research in this area should be encouraged (Ogino et al., 2017).

382 In the present study, ambient both particles collected at all locations induced expression of
383 DEC205 on APCs. The current study is the first report that ambient particles collected by
384 cyclone technique activates APCs via DEC205 expression. Several studies have shown that
385 carbon black nanoparticles and Asian dust particles can promote the maturation/activation
386 and function of DEC205 on APCs and may be related to their enhancing effects on allergic
387 diseases or responses (Honda et al., 2014; Koike et al., 2008). Ambient both particles induced
388 pro-inflammatory cytokines such as IL-6 and IL-1 β from APCs as well as BEAS-2B cells.
389 Ambient both particles strongly induced pro-inflammatory cytokine production compared to
390 both reference particles. IL-1 β has been described as a potent pro-inflammatory cytokine and
391 a mediator of a wide range of systemic human diseases (Dinarello, 2005; Koh et al., 2006;
392 Allantaz et al., 2007; James et al., 2011). The difference in reaction of IL-6 and IL-1 β
393 observed in this study may be due to the difference in the transcription factors of cytokines
394 or in the sensitivity to exposure components. APCs play important roles in allergens-related
395 airway inflammation (Lambrecht et al., 2012). APCs are activated upon invasion of the upper
396 or lower respiratory tract by ambient particles collected by cyclone technique. As a result,

397 ambient particles can affect allergic inflammation not only by induction DEC205 expression,
398 but also through induction of pro-inflammatory cytokines.

399 Some experimental and epidemiological studies have indicated that PM10 may exhibit a
400 similar or higher pro-inflammatory potential than PM2.5 and lead to adverse pulmonary
401 responses, which may require hospitalization (Monn et al., 1999; Becker et al., 2005;
402 Brunekreef et al., 2005; Gerlofs et al., 2007; Camatini et al., 2008). One of the reasons for
403 this may be attributed to the greater presence of microbial factors such as endotoxins or β -
404 glucan in PM10 as compared to PM2.5. Previous studies have shown endotoxin and β -glucan
405 to be associated with the inflammatory effects of PM both in *in vitro* and *in vivo* (Douwes et
406 al., 2003; Becker et al., 2005; Jalava et al., 2008). The U.S. Environmental Protection Agency
407 (1995) has noted that PM10 deposits in the upper airways of the lungs and may be more
408 relevant for asthmatic responses and irritation. On the other hand, PM2.5 is more often the
409 cause of lower respiratory symptoms such as cough and sputum compared to PM10
410 (Schwartz et al., 2000). In this way, although both PM can cause harmful health effects, the
411 one that causes greater harm has not been elucidated. In the present study, almost no
412 difference of cytotoxicity on RPMI-2650 cells and APCs was observed between the fine and
413 coarse particles. However, coarse particles collected at Fukuoka induced higher production
414 of IL-6 and IL-8 from BEAS-2B cells and IL-6 from APCs than fine particles, but lesser
415 production of IL-1 β from APCs. Fine particles collected at Yokohama induced higher
416 production of IL-8 from BEAS-2B cells than coarse particles. Our analysis did not show
417 strong correlations between pro-inflammatory cytokine release and microbial factors.
418 Previous studies have suggested that endotoxin at 2000 EU/mL and β glucan at the

concentration of “ $\mu\text{g/mL}$ ” induces pro-inflammatory responses in airway epithelial cells or APCs (Carmona et al. 2010, Veranth et al. 2004). As the level of endotoxin and βglucan in our study was very low compared to those of the previous studies, we suspect that the level of endotoxin and βglucan in our study failed to noticeable correlation with inflammation in the cells. Apart from microbial factors, previous studies have suggested that co-exposure of SO_4^{2-} or constituent of cedar pollen enhances inflammatory responses (Hiyoshi et al., 2005; Ichinose et al., 2005; Yamada et al., 2012). Not only biological factors but also other compounds contained in PM or allergen substance can contribute to the production of pro-inflammatory cytokines.

Yokohama is the most populated city and located in one of the three major industrial areas in Japan. Saitama is close to Yokohama, and relatively the secondary particle tends to be the main component of the formation of fine particles, so secondary air pollution is considered. (Takegawa et al., 2006; Miyakawa et al., 2008). Yokohama and Saitama are located in urban/suburban and industrial areas, respectively. Fe, Cr, Mn, Zn and Co are emitted by the steel industry and V is mainly emitted by oil combustion (Lin et al., 2005; Querol et al., 2006; Japan Ministry of the Environment, 2014). Titanium originates from the chemical industry dealing with nanomaterial particles such as titanium dioxide (Chao et al., 2011). As these metals in particles were correlated with IL-6 and IL-8 production in Yokohama and Saitama, anthropogenic metals might contribute to respiratory disorders. On the other hand, Fukuoka is closest to Mainland China, therefore there are a possibility that the influence of transboundary pollution from China. (kaneyasu et al., 2014; Takami et al., 2016). It has been reported that As and Pb are abundantly contained in coal which is the main energy source in

China (Hioki et al., 2009). Generally, atmospheric concentration of As in Japan is very low, and there are few specific sources of As (Taniguchi et al. 2016). At least, the present study demonstrated that As contained much in the particles in Fukuoka showed a high correlation with IL-1 β release. This indicated that Fukuoka may be influenced by transboundary pollution from China. However, the other biological responses induced by the particles collected in Fukuoka was lower than the particles in other locations. Hence, particles caused by transboundary contamination may have lower biological activity than particles having a source in our country.

Previous studies have reported that metals and their compounds mentioned above induce IL-6, IL-8 or IL-1 β *in vivo* and/or *in vitro* studies. For example, Ti or CoCl₂ induce the production of IL-6 or IL-8 from bronchial epithelial cells or human lung microvascular endothelial cells, and IL-1 β are induced from alveolar macrophages by intratracheal administration of As in mice (Huaux et al., 1995; Carter et al., 1997; Schmalz et al., 1998; Ming et al., 2000; Pascal et al., 2004; Kyoung et al., 2006; Eun-Jung et al., 2008, 2010). It is possible that metals such as Ti, Fe, Cr, Mn, Co, V, Zn, Pb and As can induce IL-6, IL-8 or IL-1 β from BEAS-2B cells and APCs, but further investigation is needed to identify the responsible components. The components in OC and EC have not yet been well elucidated (Grabowsky et al., 2011; Ikemori et al., 2009). However, low- and less- volatile organic carbons (OC3 and OC4) may play an important role in the inflammatory reaction in each cell. Previous studies have reported metals or ions such as Fe, Zn, Cr, Mn, V and Cu induce toxicity of airway epithelial cells (Riley et al., 2003; Honda A., 2015). The concentrations of Fe, Cr, V and Cu in our study was low compared to those of the previous studies, but Mn and

Zn was equivalent concentrations. In this study, we showed only high correlation between Zn and toxicity, however it is a possible that the other metals such as Fe, Cr, Mn, V and Cu also induce toxicity of airway epithelial cells.

5. Conclusion

Exposure to ambient particles collected by cyclone technique reduced cellular viability in RPMI-2650 cells and APCs, induced pro-inflammatory responses in BEAS-2B cells and APCs, and induced the maturation/activation of APCs. There was correlation between of some chemical components and biological responses. These chemical components affected differently between nasal and bronchial epithelial cells and elicited varying immune response. In addition, these effects can differ depending on the diameter of the particles and/or collection locations.

Figure 1. Effects of ambient particles and reference particles on the viability of RPMI-2650 cells. Data are presented as the percentage of the viability of the control. Data are mean $\pm$ standard error of the mean (SEM) of 4 individual cultures. *P<0.05, **P<0.01 vs. 0 μ g/mL, #P<0.05 vs. Fine particles at 7.5 μ g/mL.

481

482

Figure 2. (A): Effects of ambient particles and reference particles on the viability of BEAS-2B cells. Data are presented as the percentage of the viability of the control. (B): IL-6 and IL-8 production from BEAS-2B cells in response to ambient particles and reference particles. Data are mean $\pm$ standard error of the mean (SEM) of 4 individual cultures. * $P < 0.05$, ** $P < 0.01$ vs. 0 $\mu\text{g/mL}$, $^{\$}P < 0.05$ vs. Fine particles at Fukuoka at 7.5 $\mu\text{g/mL}$, $^{\dagger\dagger}P < 0.01$ vs. Fine particles at Fukuoka at 75 $\mu\text{g/mL}$, $^{\#}P < 0.05$, $^{##}P < 0.01$ vs. Coarse particles at Fukuoka at 75 $\mu\text{g/mL}$, $^{\Phi\Phi}P < 0.01$ vs. Fine particles at Saitama at 75 $\mu\text{g/mL}$, $^{\Psi}P < 0.05$ vs. Fine particles at Yokohama at 75 $\mu\text{g/mL}$.

481

Figure 3. (A): Effects of ambient particles and reference particles on the viability of APCs. Data are presented as the percentage of the viability of the control. (B): IL-6 and IL-1β production from APCs in response to ambient particles and reference particles. (C): Effects of ambient particles and reference particles on the DEC205 expression of APCs. Data are presented as positive cells expressed % events. Data are mean $\pm$ standard error of the mean (SEM) of 4 individual cultures. * $P < 0.05$, ** $P < 0.01$ vs. 0 $\mu\text{g/mL}$, \$ $P < 0.05$, \$\$ $P < 0.01$ vs. Fine particles at Fukuoka at 7.5 $\mu\text{g/mL}$, † $P < 0.05$, †† $P < 0.01$ vs. Fine particles at Fukuoka at 75

500 $\mu\text{g/mL}$, ⁰P<0.05, ⁰⁰P<0.01 vs. Coarse particles at Fukuoka at 7.5 $\mu\text{g/mL}$, [#]P<0.05, ^{##}P<0.01
501 vs. Coarse particles at Fukuoka at 75 $\mu\text{g/mL}$..

502

503 **Funding sources**

504 This work was supported by the Environmental Research and Technology Development Fund
505 (5-1651) of the Ministry of the Environment, Japan, the Matching Planner Program from
506 Japan Science and Technology Agency, JST (MP28116789653), JKA foundation (27-127),
507 Steel Foundation for Environmental Protection Technology, and the Keio Leading-edge
508 Laboratory of Science and Technology (KLL) Specified Research Projects.

509

510 **Acknowledgement:** This study was supported by the Environmental Research and
511 Technology Development Fund (5-1651) of the Ministry of the Environment, Japan. The
512 sample collection, preparation, and measurement were performed under the CYCLEX
513 (CYCLone collection of PM2.5 followed by EXposure experiments) project supported by
514 Ministry of the Environment, Japan.

515 We thank Kiyoe Itoi and Yufuko Kobayashi for their technical assistance.

516

517 **Supplementary data:** The following are the supplementary data related to this article:

518

519 **Disclosure:** The authors declare no competing financial interest.

520

521 **Reference**

- 522 Allantaz F, Chaussabel D, Banchereau J, Pascual V. 2007. Microarray-based identification
523 of novel biomarkers in IL-1 mediated disease. *Curr Opin Immunol.*19, 623-632.
- 524 Bachert C, van Cauwenberge P, Khaltaev N. 2002. Allergic rhinitis and its impact on asthma.
525 In collaboration with the World Health Organization. Executive summary of the workshop
526 report. 7-10 December 1999, Geneva, Switzerland. *Allergy.* 57 (9), 841-855.
- 527 Becker S, Mundandhara S, et al. 2005. Regulation of cytokine production in human alveolar
528 macrophages and airway epithelial cells in response to ambient air pollution particles: further
529 mechanistic studies. *Toxicol Appl Pharmacol.* 207 (2), 269-275.
- 530 Brunekreef B, Forsberg B. 2005. Epidemiological evidence of effects of coarse airborne
531 particles on health. *Eur Respir J.* 26, 309-318.
- 532 Cachon BF, Firmin S, Verdin A, Ayi-Fanou L, Billet S, Cazier F, et al. 2014.
533 Proinflammatory effects and oxidative stress within human bronchial epithelial cells exposed
534 to atmospheric particulate matter (PM_{2.5} and PM_{>2.5}) collected from Cotonou, Benin.
535 *Environmental Pollution.* 185, 340-351.
- 536 Camatini M, Gualtieri M, et al. 2008. Seasonal and dimensional differences in pro-
537 inflammatory potency of the atmospheric particulate matter of Milan on lung cell lines.
538 *Toxicol Lett.* 180S,S97.
- 539 Carter JD, Ghio AJ, Samet JM, Devlin RB. 1997. Cytokine production by human airway
540 epithelial cells after exposure to an air pollution particle is metal-dependent. *Toxicol Appl*
541 *Pharmacol.* 146 (2), 180-188.
- 542 Chao Min Teh, Abdul Rahman Mohamed. 2011. Roles of titanium dioxide and ion-doped
543 titanium dioxide on photocatalytic degradation of organic pollutants (phenolic compounds

544 and dyes) in aqueous solutions. *Journal of Alloys and Compounds*. 509 (5), 1648-1660.

545 Chowdhury H Pratiti, Okano H, Honda A, Kudou H, Kitamura G, Ito S, Ueda K, Takano H.

546 2018. Aqueous and organic extract of PM2.5 collected in different seasons and cities of Japan

547 differently affect respiratory and immune systems. *Environmental Pollution*. 235, 223-234.

548 Dinarello CA. 2005. Blocking IL-1 in systemic inflammation. *J Exp Med*. 201, 1355-1359.

549 Douwes J, Thorne P, Pearce N, Heederik D. 2003. Bioaerosol health effects and exposure

550 assessment: progress and prospects. *Ann Occup Hyg*. 47, 187-200.

551 Eun-Jung Park, Jongheop Yi, Kyu-Hyuck Chung, Doug-Young Ryu, Jinhee Choi, Kwangsik

552 Park. 2008. Oxidative stress and apoptosis induced by titanium dioxide nanoparticles in

553 cultured BEAS-2B cells. *Toxicol Lett*. 180, 222-229.

554 Eun-JungPark, Hero Kim, Younghun Kim, Jongheop Yi, Kyunghee Choi, Kwangsik Park.

555 2010. Inflammatory responses may be induced by a single intratracheal instillation of iron

556 nanoparticles in mice. *Toxicology*. 275, 65-71.

557 Fuentes-Mattei E, Rivera E, Gioda A, et al. 2010. Use of human bronchial epithelial cells

558 (BEAS-2B) to study immunological markers resulting from exposure to PM2.5 organic

559 extract from Puerto Rico. *Toxicol Appl Pharmacol*. 243, 381-389.

560 Fukuoka A, Matsushita K, Morikawa T, Takano H, Yoshimoto T. 2015. Diesel exhaust

561 particles exacerbate allergic rhinitis in mice by disrupting the nasal epithelial barrier. *Clinical*

562 *& experimental allergy*. 46, 142-152.

563 Gerlofs-Nijland M, Dormans J.A, et al. 2007. Toxicity of coarse and fine particulate matter

564 from sites with contrasting traffic profiles. *Inhalation Toxicology*. 19, 1055-1069.

565 Grabowsky J, Streibel T, Sklorz M, Chow JD, Watson JG, Mamakos A, Zimmermann R.

566 2011. Hyphenation of a carbon analyzer to photo-ionization mass spectrometry to unravel
567 the organic composition of particulate matter on a molecular level. *Anal Bioanal Chem.* 401,
568 3153–3164.

569 Gualtieri M, Ovrevik J, Holme J, et al. 2010. Differences in cytotoxicity versus pro-
570 inflammatory potency of different PM fractions in human epithelial lung cells. *Toxicology*
571 *in vitro.* 24, 29-39.

572 Heyder J, Gebhart J, Rudolf G, Schiller CF, Stahlhofen W. 1986. Deposition of particles in
573 the human respiratory tract in the size range 0.005-15 μm . *J Aerosol Sci.* 17 (5), 811-825.

574 Hioki T, Kimoto T, Hasegawa S, et al. 2009. Analysis of Long-range Transported and Air
575 Pollution with Trace Metal Concentration Ratio in Aerosols collected at Matsuyama, Osaka
576 and Tsukuba, Japan. *J Jpn Soc Atmos Environ.* 44 (2), 91-100.

577 Hiyoshi K, Ichinose T, Sadakane K, Takano H, Nishikawa M, Mori I, Yanagisawa R,
578 Yoshida S, Kumagai Y, Tomura S, Shibamoto T. 2005. Asian sand dust enhances ovalbumin-
579 induced eosinophil recruitment in the alveoli and airway of mice. *Environ Res.* 99, 361-368.

580 Honda A, Fukushima W, Oishi M, et al. 2017. Effects of components of PM_{2.5} collected in
581 Japan on the respiratory and immune systems. *International Journal of Toxicology.* 36 (2),
582 153-164.

583 Honda A, Matsuda Y, Murayama R, et al. 2014. Effects of Asian sand dust particles on the
584 respiratory and immune system. *J Appl Toxicol.* 3 (3), 250-257

585 Honda A, Tsuji K, Matsuda Y, Hayashi T, Fukushima W, Sawahara T, Kudo H, Murayama
586 R, Takano H. 2015. Effects of air pollution-related heavy metals on the viability and
587 inflammatory responses of human airway epithelial cells. *International Journal of Toxicology.*

- 588 34 (2), 195-203.
- 589 Hong Z, Guo Z, Zhang R, Xu J, Dong W, Zhuang G, Deng C. 2016. Airborne fine particulate
590 matter induces oxidative stress and inflammation in human nasal epithelial cells. *Tohoku J*
591 *exp med.* 239, 117-125.
- 592 Huaux F, Lasfargues G, Lauwerys R, Lison D. 1995. Lung Toxicity of Hard Metal Particles
593 and Production of Interleukin-1, Tumor Necrosis Factor- α , Fibronectin, and Cystatin-c by
594 Lung Phagocytes. *Toxicol Appl Pharmacol.* 132 (1), 53-62.
- 595 Ichinose T, Hiyoshi K, Yoshida S, Takano H, Inoue K, Nishikawa M, Mori I, Kawazato H,
596 Yasuda A, Shibamoto T. 2009. Asian sand dust aggravates allergic rhinitis in guinea pigs
597 induced by Japanese cedar pollen. *Inhal. Toxicology.* 21, 985-993.
- 598 Ichinose T, Nishikawa M, Takano H, Sera N, Sadakane K, Mori I, Yanagisawa R, Oda T,
599 Tamura H, Hiyoshi K, Quan H, Tomura S, Shibamoto T. 2005. Pulmonary toxicity induced
600 by intratracheal instillation of Asian yellow dust (Kosa) in mice. *Environ Toxicol.* 20, 48-56.
- 601 Ikemori F, Yamagami M. 2009. Thermograms of organic element of airborne particles and
602 PM2.5 using IMPROVE method. Nagoya city Environmental Chem Res Lab Report. 39, 55-
603 57.
- 604 Jalava PI, Salonen RO, Pennanen AS, et al. 2008. Effects of solubility of urban air fine and
605 coarse particles on cytotoxic and inflammatory responses in RAW264.7 macrophage cell
606 lines. *Toxicol Appl Pharmacol.* 229, 146-160.
- 607 James H, Michelle H, Caitriona R, Shijuan G, et al. 2011. Autophagy controls IL-1 β
608 secretion by targeting pro-IL-1 β for degradation. *Journal of Biological Chemistry.* 286 (11),
609 9587-9597.

- 610 Japan Ministry of the Environment. 2014. [https://www.env.go.jp/chemi/report/h24-](https://www.env.go.jp/chemi/report/h24-02/index.html)
- 611 [02/index.html](https://www.env.go.jp/chemi/report/h24-02/index.html). Updated March 2014.
- 612 Jiang W, Swiggard WJ, Heufler C, et al. 1995. The receptor DEC205 expressed by dendritic
- 613 cells and thymic epithelial cells is involved in antigen processing. *Nature*. 375 (6527), 151-
- 614 155.
- 615 Kaneyasu N, Yamamoto S, Sato K, Takami A, Hayashi M, Hara K, Kawamoto K, Okuda T,
- 616 Hatakeyama S. 2014. Impact of long-range transport of aerosols on the PM2.5 composition
- 617 at a major metropolitan area in the northern Kyushu area of Japan. *Atmospheric Environment*.
- 618 97, 416-425.
- 619 Koh KK, Han SH, Quon MJ. 2005. Inflammatory markers and the metabolic syndrome:
- 620 insights from therapeutic interventions. *J Am College Caardiol*. 46, 1978-1985.
- 621 Koike E, Takano H, Inoue K, Yanagisawa R, Kobayashi T. 2008. Carbon black nanoparticles
- 622 promote the maturation and function of mouse bone marrow-derived dendritic cells.
- 623 *Chemosphere*. 73 (3), 371-376.
- 624 Kyoung S. Kim, Rajagopal V, Gonsalves C, Johnson C, Vijay K. Kalra. 2006. A Novel Role
- 625 of Hypoxia-Inducible Factor in Cobalt Chloride- and Hypoxia-Mediated Expression of IL-8
- 626 Chemokine in Human Endothelial Cells. *J Immunol*. 177, 7211-7224.
- 627 Lacasana M, Esplugues A, and Ballester F. 2005. Exposure to ambient air pollution and
- 628 prenatal and early childhood health effects. *Eur. J. Epidemiol*. 20 (2), 183–199.
- 629 Lambrecht BN, Hammad H. The airway epithelium in asthma. *Nat. Med*. 2012; 18:684-692.
- 630 Lin C-C, Chen S-J, Huang K-L. et al. 2005. Characteristics of Metals in
- 631 Nano/Ultrafine/Fine/Coarse Particles Collected Beside a Heavily Trafficked Road.

- 632 Environmental science & technology. 39 (21), 8113-8122.
- 633 Lindbom J, Mats G, Goran B, Andreas D, et al. 2006. Exposure to wear particles generated
634 from studded tires and pavement induces inflammatory cytokine release from human
635 macrophages. *Chem Res Toxicol*. 19, 521-530.
- 636 Lutz MB, Kukutsch N, Ogilvie AL, et al. 1999. An advanced culture method for generating
637 large quantities of highly pure dendritic cells from mouse bone marrow. *J Immunol Methods*.
638 223 (1), 77-92.
- 639 Marcazzan G, Vaccaro S, Valli G, Vecchi R. 2001. Characterization of PM10 and PM2.5
640 particulate matter in the ambient air of Milan (Italy). *Atmospheric Environment*. 35 (27),
641 4639-4650.
- 642 Ming-YieLiu, Wen-ChuanHsieh, Bei-Chang. 2000. In vitro aberrant gene expression as the
643 indicator of lead-induced neurotoxicity in U-373MG cells. *Toxicology*. 147 (1), 65-71.
- 644 Miyakawa T, Takegawa T, Kondo Y. 2008. Photochemical evolution of submicron aerosol
645 chemical composition in the Tokyo megacity region in summer. *Journal of geophysical*
646 *research*. 113, D14304. Doi:10.1029/2007JD009493.
- 647 Moncayo-Nieto OL, Wilkinson TS, Brittan M, et al. 2014. Differential response to the
648 bacteria, and TOLLIP expression, in the human respiratory tract. *BMJ Open Resp Res*.
649 1:e000046.
- 650 Monn C, Becker S. 1999. Cytotoxicity and induction of proinflammatory cytokines from
651 human monocytes exposed to fine (PM2.5) and coarse particles (PM10-2.5) in outdoor and
652 indoor air. *Toxicol Appl Pharmacol*. 155, 245-252
- 653 Moorhead PS. 1965. Human tumor cell line with a quasi-diploid karyotype (RPMI2650). *Exp*

- 654 Cell Res. 39 (1), 190-196.
- 655 Mori I, Sun Z, Ukachi M, Nagano K, McLead CW, Cox AG, Nishikawa M. 2008.
- 656 Development and certification of the new NIES CRM 28: urban aerosols for the
- 657 determination of multielements. Anal Bioanal Chem. 391, 1997-2003.
- 658 Ogino K, Nagaoka K, Okuda T. et al. 2017. PM2.5 - induced airway inflammation and
- 659 hyperresponsiveness in NC/Nga mice. Environ Toxicol. 32 (3), 1047-1054.
- 660 Okamoto K. 1987. A new certified reference material, vehicle exhaust particulates.
- 661 Analytical sciences. 3, 191-192.
- 662 Okuda T, Isobe R. 2017. Improvement of a High-volume Aerosol Particle Sampler for
- 663 Collecting Submicron Particles through the Combined Use of a Cyclone with a Smoothened
- 664 Inner Wall and a Circular Cone Attachment. Asian Journal of Atmospheric Environment. 11
- 665 (2), 131-137.
- 666 Okuda T, Isobe R, Nagai Y, Okahisa S, Funato K, Inoue K. 2015. Development of a high-
- 667 Volume PM2.5 Particle Sampler Using Impactor and Cyclone Techniques. Aerosol and Air
- 668 Quality Research. 15, 759-767.
- 669 Okuda T, Schauer J, Shafer M. 2014. Improved methods for elemental analysis of
- 670 atmospheric aerosols for evaluating human health impacts of aerosols in East Asia.
- 671 Atmospheric Environment. 97, 552-555
- 672 Okuda T, Shishido D, Terui Y, Fujioka K, Isobe R, Iwaki Y, Funato K, Inoue K. 2018.
- 673 Development of a high-volume simultaneous sampler for fine and coarse particles using
- 674 virtual impactor and cyclone techniques. Asian Journal of Atmospheric Environment. 12 (1),
- 675 78-86.

- 676 Okuda T, Takada H, Kumata H, Nakajima F, et al. 2013. Inorganic Chemical Characterization
677 of Aerosols in Four Asian Mega-Cities. *Aerosol and Air Quality Research*. 13 (2), 436-449.
- 678 Okuda T. 2013. Measurement of the specific surface area and particle size distribution of
679 atmospheric aerosol reference materials. *Atmospheric Environment*. 75, 1-5.
- 680 Park EJ, Park K. 2009. Induction of pro-inflammatory signals by 1-nitropyrene in cultured
681 BEAS-2B cells. *Toxicol Lett*. 184, 126–133.
- 682 Pascal LE, Tessier DM. 2004. Cytotoxicity of chromium and manganese to lung epithelial
683 cells in vitro. *Toxicol Lett*. 147 (2), 143-151.
- 684 Perez IR, Serrano J, Alfaro-Moreno E, Baumgardner D, Garcia-Cuellar C, Martin del Campo
685 JM, Raga GB, Castillejos M, Colin RD, Osornio-Vargas AR. 2007. Relations between PM10
686 composition and cell toxicity: a multivariate and graphical approach. *Chemosphere*. 67,
687 1218–1228.
- 688 Querol X, Zhuang X, Alastuey A, Viana M, Lv W, Wang Y, Lopez A, Zhu Z, Wei H, Xu S.
689 2006. Speciation and sources of atmospheric aerosols in a highly industrialised emerging
690 mega-city in Central China. *J Environ Monit*. 8, 1049-1059.
- 691 Riley MR, Boesewetter DE, Kim AM, Sirvent FP. 2003. Effects of metals Cu, Fe, Ni, V, and
692 Zn on rat lung epithelial cells. *Toxicology*. 190 (3), 171-184.
- 693 Samoli E, Peng R, Ramsay T, Pipikou M, Touloumi G, Dominici F, Burnett R, Cohen A,
694 Krewski D, Samet J, Katsouyanni K. 2008. Acute effects of ambient particulate matter on
695 mortality in Europe and North America: results from the APHENA study. *Environmental*
696 *Health Perspective*. 116 (11), 1480–1486.
- 697 Schins RPF, Lightbody JH, Borm PJA, Shi T, Donaldson K, Stone V. 2004. Inflammatory

698 effects of coarse and fine particulate matter in relation to chemical and biological constituents.
699 Toxicology and Applied Pharmacology. 195, 1-11.
700 Schmalz G, Schuster U, Schweikl H. 1998. Influence of metals on IL-6 release in vitro.
701 Biomaterials. 19, 1689-1694.
702 Schwartz J, Lucas M. 2000. Fine particles are more strongly associated than coarse particles
703 with acute respiratory health effects in schoolchildren. Epidemiology. 11 (1), 6-10.
704 Sevastyanova O, Novakova Z, Hanzalova K, Binkova B, Sram R, Topinka J. 2008. Temporal
705 variation in the genotoxic potential of urban air particulate matter. Mutation Research. 649,
706 179–186.
707 Sharma H, Jain VK, Khan ZH. 2007. Characterization and source identification of polycyclic
708 aromatic hydrocarbons (PAHs) in the urban environment of Delhi. Chemosphere. 66, 302–
709 310.
710 Takegawa N, Miyakawa T, Kondo Y, Blame DR, Kanaya Y, Koike M, Fukuda M, Komazaki
711 Y, Miyazaki Y, Shimono A, Takeuchi T. 2006. Evolution of submicron organic aerosol in
712 polluted air exported from Tokyo. Geophysical research letters. 33. L15814.
713 Doi:10.1029/2006GL025815.
714 Takami A, Miyoshi T, Irei S, Yoshino A, Sato K, Shimizu A, Hayashi M, Hara K, Kaneyasu
715 N, Hatakeyama S. 2016. Analysis of Organic Aerosol in Fukuoka, Japan Using a PMF
716 Method. Aerosol and Air Quality Research. 16, 314–322.
717 Taniguchi Y, Shimada K, Takami A, Lin NH, Chan CK, Kim YP, Hatakeyama S. 2016.
718 Transboundary and Local Air Pollutants in Western Japan Distinguished on the Basis of
719 Ratios of Metallic Elements in Size-Segregated Aerosols. Aerosol and Air Quality Research.

- 720 17 (12), 3141-3150.
- 721 Tecer LK, Alagha O, Karaca F, Tuncel G, Eldes N. 2008. Particulate Matter (PM_{2.5}, PM_{10-2.5},
722 and PM₁₀) and Children's Hospital Admissions for Asthma and Respiratory Diseases. J
723 Toxicol and Env Health. 71 (8), 512-520
- 724 Thacker EL. 2006. Lung inflammatory responses. Vet Res. 37 (3), 469-486.
- 725 Totlandsdal AI, Herseth JJ, Bølling AK, Kubátová A, Braun A, Cochran RE, Refsnes M,
726 Ovrevik J, Låg M. 2012. Differential effects of the particle core and organic extract of diesel
727 exhaust particles. Toxicol Lett. 208, 262–268.
- 728 U.S. Environmental Protection Agency. 1995. Air Quality Criteria for Particulate Matter.
729 EPA/600/AP-95/001b. Research Triangle Park, NC: U.S. Environmental Protection Agency.
- 730 Vecchi R, Marcazzan G, Ceriani M, Antoniazzi C. 2004. The role of atmospheric dispersion
731 in the seasonal variation of PM₁ and PM_{2.5} concentration and composition in the urban area
732 of Milan (Italy). Atmospheric Environment. 38 (27), 4437-4446.
- 733 Yamada P, Hatta T, Du M, Wakimizu K, Han J, Maki T, Isoda H. 2012. Inflammatory and
734 degranulation effect of yellow sand on RBL-2H3 cells in relation to chemical and biological
735 constituents. Ecotoxicol Environ Saf. 84, 9–17.
- 736 Zerbi G, Ferruggiari A, Fustella G, Tommasini M, Mantecca P, Gualtieri M, Cetta F,
737 Camatini M. 2008. Preliminary observation on the interactions between fine atmospheric
738 particulate matter (PM_{2.5}) and human alveolar epithelial cells (A549). Chemical Engineering
739 Transaction. 16, 387–394.