

Relativistic theory of electron-nucleus-radiation coupled dynamics in molecules: Wavepacket approach

Cite as: J. Chem. Phys. **151**, 084102 (2019); <https://doi.org/10.1063/1.5109272>

Submitted: 07 May 2019 . Accepted: 29 July 2019 . Published Online: 22 August 2019

Kota Hanasaki , and Kazuo Takatsuka 



View Online



Export Citation



CrossMark

ARTICLES YOU MAY BE INTERESTED IN

[Compressing physics with an autoencoder: Creating an atomic species representation to improve machine learning models in the chemical sciences](#)

The Journal of Chemical Physics **151**, 084103 (2019); <https://doi.org/10.1063/1.5108803>

[Scattering off molecules far from equilibrium](#)

The Journal of Chemical Physics **151**, 084301 (2019); <https://doi.org/10.1063/1.5111979>

[Similarity transformation of the electronic Schrödinger equation via Jastrow factorization](#)

The Journal of Chemical Physics **151**, 061101 (2019); <https://doi.org/10.1063/1.5116024>

The Journal
of Chemical Physics

Submit Today

The Emerging Investigators Special Collection and Awards
Recognizing the excellent work of early career researchers!



Relativistic theory of electron-nucleus-radiation coupled dynamics in molecules: Wavepacket approach

Cite as: J. Chem. Phys. 151, 084102 (2019); doi: 10.1063/1.5109272

Submitted: 7 May 2019 • Accepted: 29 July 2019 •

Published Online: 22 August 2019



View Online



Export Citation



CrossMark

Kota Hanasaki^{a)}  and Kazuo Takatsuka^{b)} 

AFFILIATIONS

Fukui Institute for Fundamental Chemistry, Kyoto University, Kyoto 606-8103, Japan

^{a)}Electronic mail: hanasaki@fukui.kyoto-u.ac.jp

^{b)}Electronic mail: kaztak@fukui.kyoto-u.ac.jp

ABSTRACT

We propose a general theoretical scheme of relativistic electron-nucleus coupled dynamics of molecules in radiation fields, which is derived from quantum electrodynamical formalism. Aiming at applications to field-induced dynamics in ultrastrong laser pulses to the magnitude of 10^{16} W/cm² or even larger, we derive a nonperturbative formulation of relativistic dynamics using the Tamm-Dancoff expansion scheme, which results in, within the lowest order expansion, a time-dependent Schrödinger equation with the Coulombic and retarded transversal photon-exchange interactions. We also discuss a wavepacket type nuclear dynamics adapted for such dynamics.

© 2019 Author(s). All article content, except where otherwise noted, is licensed under a Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>). <https://doi.org/10.1063/1.5109272>

I. INTRODUCTION

Electrons accelerated close to the speed of light exhibit relativistic effects.^{1–4} In molecular science, relativistic electrons appear in orbitals of heavy atoms where it has been established that not only the inner-shell electrons^{5–7} but also valence electrons are subject to relativistic effects such as the orbital contraction/expansion effects.^{8,9} Another important type of relativistic electrons in molecular science is those accelerated in ultraintense optical fields.^{10,11} The former aspect has been discussed separately in our recent work¹² within the path integral formalism. We herein develop a relativistic “wavepacket” theory for coupled electronic and nuclear dynamics of molecules, which is applicable to dynamics in external fields of nonperturbative intensity regime.

Rapid progress in experimental techniques^{10,11} has realized ultraintense laser pulses of petawatt intensity and femtosecond width,¹³ and a number of advanced light sources are being developed worldwide.¹⁴ The strongest laser field ever reported¹⁵ exceeds 10^{22} W/cm². Nonrelativistic quantum mechanics breaks down in such ultrastrong fields as the vector field amplitude in the Hamiltonian compares $m_e c^2/|e|$, with m_e and $|e|$ being the electronic mass and charge magnitude, respectively. Introducing nondimensional

relativistic scale for a given vector field \mathbf{A} by $a_0 \equiv |e||\mathbf{A}|/(m_e c^2)$, a typical infrared (IR) laser pulse of wavelength $\lambda = 800$ nm ($\hbar\omega = 1.55$ eV) of intensity 2.14×10^{18} W/cm² corresponds to $a_0 = 1$. Indeed, with pulse intensity of order $\sim 10^{16}$ W/cm² or more, experimental studies have demonstrated relativistic behaviors of electrons including relativistic trajectories of photoelectrons,¹⁶ radiation reactions,^{17,18} and multiphoton Thomson scatterings.¹⁹ Molecular dissociations in laser fields of relativistic intensity have also been reported in Refs. 20 and 21. In Ref. 20, CH₄ molecules irradiated by IR laser fields of intensity exceeding 10^{16} W/cm² were observed to undergo multiple tunnel ionizations, followed by rapid fragmentation.²⁰ Such dynamics is particularly challenging in that strong electron-radiation coupling does not allow perturbative approach which is the most standard tool in the quantum electrodynamics (QED). Another interest is in the interplay of such relativistic electron-radiation coupled dynamics and the nuclear dynamics. Taking into account the typical nucleus-to-electron mass ratio of order 10^3 or more, we can assume that the nuclear dynamics is nonrelativistic even in ultrastrong fields with $a_0 \lesssim 100$. Relativistic electronic dynamics, however, should induce unconventional nuclear dynamics such as Coulomb explosion of heavily ionized molecules.²⁰

Relativistic theory of stationary electronic many-body states within the Born-Oppenheimer (BO) approximation has been intensively developed in the so-called quantum chemistry.^{22–24} Numerical studies have revealed that relativistic electrons, which exist in the inner-core levels of heavy element atoms, play several important roles in the chemical properties of molecules.²² Static properties of those molecules containing heavy elements, including their bonding properties^{25–27} and spectroscopic observables,²⁸ have been successfully reproduced by numerical *ab initio* calculations. Formulations of numerical calculations include the original four-component formulation and more tractable two-component formulations based on the Foldy-Wouthuysen transformation^{29–31} or even more sophisticated theories.^{32–35} Such two-component reduction techniques are sometimes indispensable for the study of large systems, whereas its extension to dynamical problems is not straightforward. We therefore have the present formulation remain in the original four-component theory unless specified explicitly. A remarkable merit in the quantum-chemical approach is its large variety of methodologies of constructing (correlated) many-body wavefunctions; techniques for constructing four-component wavefunctions include the variational theory³⁶ and the exact solution technique.³⁷ Multiconfiguration theories to include correlation effects to the self-consistent field orbitals have also been developed.^{38,39} Along with the methodological sophistication, there are a large number of “tools” available for achieving efficient calculations; studies have established a number of efficient basis sets⁴⁰ and calculation techniques.^{41,42} A number of computational packages for general purposes have also been developed.^{43,44}

However, in the BO framework, the involved nuclei are made fixed in space and no time variable is considered in the electronic wavefunctions, let alone in the radiation fields. Despite its great success in the calculations of static molecular properties, the main assumptions presupposed in the BO approximation (and thereby the conventional quantum chemistry) are not valid when the kinematic coupling between electrons and nuclei are strong. Indeed, the physical and chemical significance of the so-called nonadiabatic reactions has long been recognized since the early stage of quantum mechanics^{45–47} to date.^{48–50} Overcoming such breakdown has become even more important in the regime of laser chemistry,⁵¹ which makes it possible to observe⁵² and control (see Ref. 53) the kinematic coupling between electrons and nuclei. In particular, it has been found that the notion of ultrafast nonadiabatic electron wavepacket dynamics is particularly crucial.^{54–59} Therefore, one of the main objectives in this paper is to extend such non-BO wavepacket dynamics to the relativistic regime. Another shortcoming of the conventional type of relativistic quantum chemistry is that it often shows lack of theoretical rigor; predominantly many of those studies have been made based on the Dirac-Coulomb or the Dirac-Breit theory, which are the low-energy static reductions of the exact QED. Many features in the exact QED treatment, including dynamical coupling to the radiation-field and/or energy-dependence of the electron-electron interactions, are missing in those types of quantum chemical approaches, whereas it can easily be expected that those missing components are essential in high-energy dynamics of our interest. We therefore start from the exact QED formulation to construct a framework of relativistic nonadiabatic electron-nucleus coupled dynamics. The most

standard QED formulation would then take a perturbative approach to include electron-radiation coupling effects. We indeed have taken such an approach in our recent study.¹² It is, however, obvious that such an approach is not applicable to dynamics with non-perturbative couplings to the (external) radiation fields, which is our main target in this paper. In order to develop a nonperturbative approach, we consider the Tamm-Dancoff (TD)^{60,61} expansion of the electron-radiation coupled states. We also note that there have been the development of advanced formulations of quantum chemistry that resolves such discrepancy from the exact QED treatment. For example, the effective quantum electrodynamics (eQED) approach^{23,24} constructs an effective Hamiltonian that includes static QED effects such as the vacuum polarization effect and the one-loop electronic self-energy arising from virtual interactions with the radiation field. It is of theoretical interest to make comparison between these static QED theories and our formulation.

The original idea of the Tamm-Dancoff approximation (TDA) was independently developed by Tamm⁶⁰ and Dancoff⁶¹ earlier than the establishment of the covariant perturbation theory.^{62–66} The original idea of Dancoff⁶¹ was to expand the Heisenberg state vector, or the time-independent energy eigenstate, by a well-defined set of basis vectors in various “sectors” to derive a set of Schrödinger-like equations. A problem in this original approach, pointed out by Dyson,⁶⁷ is that the number of interacting particles adds up in a less-controlled manner than what is in the covariant perturbation theory, making the theory nonrenormalizable. Several modifications to overcome such a problem were proposed including Dyson’s formulation.^{67–69} Among them, Cini⁶⁸ and Visscher⁶⁹ developed an alternative formulation closer in idea to the covariant perturbation theory, which shows that divergences arising from lower order expansions (Cini⁶⁸ considered up to what corresponds to the two-loop expansion in diagrammatic language) can be removed using the renormalization technique. Even with those modifications, TDA has several fundamental problems⁷⁰ including apparent violation of covariance by its naive truncation scheme. Wilson and co-workers formulated TDA in the light-cone frame⁷⁰ to overcome these inconveniences. Their light-front TDA approach⁷¹ has then been applied to quantum color dynamics (QCD). To the best of our knowledge, however, it is not clear that such a drastic change in the frame is applicable or suitable for molecular dynamics. We therefore do not use the light-cone frame but stick to more familiar types of space-time frame (the laboratory or the center-of-mass frame) which may sacrifice the formal rigor. Nevertheless, it appears that the Tamm-Dancoff approach provides a method to formulate electron-radiation coupled dynamics of relativistic energy scale based on “equal-time” amplitudes, which describes a many-body system on a spacelike surface with a single timelike variable. We consider that, in our future study, the equal-time representation would work favorably in importing many-body techniques developed in quantum chemistry in which the same type of space-time frame is (implicitly) assumed.

This paper is organized as follows. In Sec. II, we first prepare the fundamental quantities relevant in the present relativistic dynamics. In Sec. III, we formulate a theory based on the Tamm-Dancoff^{60,61} type expansion of the electron-radiation coupled states. This paper concludes in Sec. IV.

II. DEFINITIONS OF FUNDAMENTAL QUANTITIES

We first define fundamental quantities that appear in our discussion. Although this paper is self-contained, a more comprehensive discussion on some of those quantities is shown in our recent work.¹² Throughout this paper, we use the sign convention $(1, -1, -1, -1)$, and the metric tensor $\eta_{\mu\nu}$ is a diagonal tensor with $\eta_{00} = 1$, $\eta_{11} = \eta_{22} = \eta_{33} = -1$. Symbol q_e represents the electronic charge, which takes a negative value; $q_e = -|e|$. We use the Gauss unit for electromagnetic field, and the fine-structure constant is $e^2/\hbar c \approx 1/137$. We use the Hermitian Dirac matrices, α and β . Four-dimensional notation α^μ stands for $\alpha^\mu \doteq (1, \alpha)$, i.e., α^0 is the four-dimensional unit matrix. Other notations, which follow the standard convention in QED,² are described explicitly in the main text and also summarized at the end of the main text.

We consider an electron-nucleus-radiation coupled system in the Coulomb gauge, whose full Hamiltonian reads

$$\begin{aligned} \mathcal{H}^{\text{tot}} = & \sum_a \frac{1}{2M_a} \left(\mathbf{P}_a + \left(\frac{Z_a|e|}{c} \mathbf{A}^{\text{tr}}(\mathbf{R}_a) \right)^2 \right) \\ & + \int d^3\mathbf{x} \frac{1}{2} \left(4\pi c^2 \mathbf{\Pi}^{\text{tr}2} + \frac{1}{4\pi} (\nabla \times \mathbf{A}^{\text{tr}})^2 \right) \\ & + \int d^3\mathbf{x} \psi^\dagger \left(c\alpha \cdot \frac{\hbar}{i} \nabla + \beta m c^2 \right) \psi \\ & + \frac{1}{2} \int d^3\mathbf{x} \int d^3\mathbf{y} \rho_{\text{mat}}(\mathbf{x}, t) \frac{1}{|\mathbf{x} - \mathbf{y}|} \rho_{\text{mat}}(\mathbf{y}, t) \\ & - \frac{1}{c} \int d^3\mathbf{x} \mathbf{J}_{\text{mat}} \cdot \mathbf{A}^{\text{tr}}, \end{aligned} \quad (1)$$

where α and β are the Dirac matrices, \mathbf{P}_a and M_a represent the three-dimensional momentum and mass of the a th nucleus, whereas ψ and \mathbf{A}^{tr} are the field operators of the electronic and transversal radiation fields, respectively. The canonical conjugate of \mathbf{A}^{tr} is represented by $\mathbf{\Pi}^{\text{tr}} = \dot{\mathbf{A}}^{\text{tr}}/(4\pi c^2)$, which is negative of the transversal part of the electric field. Symbols ρ_{mat} and \mathbf{J}_{mat} represent the matter field charge density and current, respectively, as

$$\rho_{\text{mat}}(\mathbf{x}) = q_e \psi^\dagger(\mathbf{x}) \psi(\mathbf{x}) + \sum_a Z_a |e| \hat{f}_a(\mathbf{x}) \quad (2a)$$

and

$$\mathbf{J}_{\text{mat}}(\mathbf{x}) = q_e c \psi^\dagger(\mathbf{x}) \alpha \psi(\mathbf{x}) + \sum_a \frac{1}{2} Z_a |e| (\hat{f}_a(\mathbf{x}) \dot{\mathbf{R}}_a + \dot{\mathbf{R}}_a \hat{f}_a), \quad (2b)$$

with q_e representing the electronic charge ($q_e = -|e|$), \mathbf{R}_a being the a th nuclear (spatial) coordinate, and $Z_a |e| \hat{f}_a(\mathbf{R})$ being the charge distribution of the a th nucleus, which could simply be a delta function $Z_a |e| \delta^3(\mathbf{R} - \mathbf{R}_a)$ or one of the existing model functions.⁷² Possible nontrivial nuclear charge distributions are taken into account only for calculation of the electronic wavefunction, whereas we approximate them by the pointlike model for description of the field-nucleus and the nucleus-nucleus Coulombic interactions. Such an approximation should be appropriate in the energy range where the typical wavelength of the radiation field is much larger than the length scale of the nuclear charge distribution. We then introduce the nuclear potential field

$$U_{\text{nuc}}(\mathbf{x}) = q_e \sum_a \int d^3\mathbf{R} Z_a |e| \frac{1}{|\mathbf{x} - \mathbf{R}|} \hat{f}_a(\mathbf{R}) \quad (3)$$

and a set of electronic “mean-fields” $\bar{\rho}(\mathbf{x}, t)$ and $W_{\text{HF}}(\mathbf{x}, t)$ and introduce a Fermionic eigenvalue equation

$$\begin{aligned} \left[c\alpha \cdot \frac{\hbar}{i} \nabla + \beta m c^2 + U_{\text{nuc}}(\mathbf{x}) \right] \varphi_\ell(\mathbf{x}, t) \\ + \int d^3\mathbf{y} W_{\text{HF}}(\mathbf{x}, \mathbf{y}, t) \varphi_\ell(\mathbf{y}, t) = \epsilon^0_\ell \varphi_\ell(\mathbf{x}, t), \end{aligned} \quad (4)$$

with ϵ^0_ℓ representing the ℓ th orbital energy. The eigenfunctions of Eq. (4) are hereafter referred to as molecular orbitals (MOs). We assume that we are interested in the state with a given configuration \mathcal{I}_{occ} of N_e positive energy MOs (if we are interested in the ground state, for example, the lowest N_e are occupied and occupied orbitals are $\mathcal{I}_{\text{occ}} = \{1, 2, \dots, N_e\}$),

$$\bar{\rho}(\mathbf{x}, t) = \sum_{\ell \in \mathcal{I}_{\text{occ}}} \varphi_\ell^\dagger(\mathbf{x}, t) \varphi_\ell(\mathbf{x}, t). \quad (5)$$

The most natural choice of the mean-field potential W_{HF} must be the Hartree-Fock potential, which is

$$\begin{aligned} W_{\text{HF}}(\mathbf{x}, \mathbf{y})_{ij} = & \delta^3(\mathbf{x} - \mathbf{y}) \delta_{ij} \int d^3\mathbf{x}' \frac{q_e^2}{|\mathbf{x} - \mathbf{x}'|} \bar{\rho}(\mathbf{x}', t) \\ & - \sum_{\ell \in \mathcal{I}_{\text{occ}}} \psi_{\ell j}^\dagger(\mathbf{y}, t) \frac{q_e^2}{|\mathbf{x} - \mathbf{y}|} \psi_{\ell i}(\mathbf{x}, t). \end{aligned} \quad (6)$$

However, we find it more convenient to use its local approximation and hereafter consider the localized Hartree-Fock field, $W_{\text{HF}}^{\text{loc}}$, which is to be obtained by one of existing techniques.^{73–76}

For later convenience, we also introduce MO creation/annihilation operators as

$$\hat{c}_{\ell, m, n, \dots} = \begin{cases} \hat{c}_{r, s, t, \dots} & \text{positive energy states} \\ \hat{b}_{a, b, c, \dots} & \text{negative energy states,} \end{cases} \quad (7)$$

and the whole set of positive (negative) MO indices are denoted by \mathcal{I}^+ (\mathcal{I}).

Our Hamiltonian now reads $\mathcal{H}^{\text{tot}} = H_{\text{nuc}} + H_{\text{mf}}^{\text{el}} + H_{\text{int}}^{\text{el}} + H_{\text{rad}}$ with

$$H_{\text{nuc}} = \sum_a \frac{1}{2M_a} \left(\frac{\hbar}{i} \nabla_a - \frac{Z_a |e|}{c} \mathbf{A}^{\text{tr}}(\mathbf{R}_a) \right)^2 + \frac{1}{2} \sum_{a \neq b} \frac{Z_a Z_b |e|^2}{|\mathbf{R}_a - \mathbf{R}_b|}, \quad (8a)$$

$$\begin{aligned} H_{\text{mf}}^{\text{el}} = & \int d^3\mathbf{x} \psi(\mathbf{x}, t)^\dagger \left[c\alpha \cdot \frac{\hbar}{i} \nabla + \beta m_e c^2 + U_{\text{nuc}}(\mathbf{x}, t) \right] \psi(\mathbf{x}, t) \\ & + \int d^3\mathbf{x} \psi(\mathbf{x}, t)^\dagger W_{\text{HF}}^{\text{loc}}(\mathbf{x}, t) \psi(\mathbf{x}, t), \end{aligned} \quad (8b)$$

$$\begin{aligned} H_{\text{int}}^{\text{el}} = & \frac{1}{2} \int d^3\mathbf{x} \int d^3\mathbf{y} \psi(\mathbf{x}, t)^\dagger \psi(\mathbf{y}, t)^\dagger \frac{q_e^2}{|\mathbf{x} - \mathbf{y}|} \psi(\mathbf{y}, t) \psi(\mathbf{x}, t) \\ & - \int d^3\mathbf{x} \psi(\mathbf{x}, t)^\dagger W_{\text{HF}}^{\text{loc}}(\mathbf{x}, t) \psi(\mathbf{x}, t) \\ & - q_e \int d^3\mathbf{x} \psi(\mathbf{x}, t)^\dagger \alpha \psi(\mathbf{x}, t) \cdot \mathbf{A}^{\text{tr}}, \end{aligned} \quad (8c)$$

$$H_{\text{rad}} = \frac{1}{2} \int d^3\mathbf{x} \left(4\pi c^2 \mathbf{\Pi}^{\text{tr}2} + \frac{1}{4\pi} (\nabla \times \mathbf{A}^{\text{tr}})^2 \right). \quad (8d)$$

Although the Hamiltonian expression Eq. (8a) is common in the literature² and it is indeed sufficient for calculation of perturbation

expansions, we have to rewrite each term in a symmetric form⁷⁷ when we calculate its vacuum expectation value. The current operator $J^\mu(\mathbf{x}, t) = c\psi^\dagger(\mathbf{x}, t)\alpha^\mu\psi(\mathbf{x}, t)$ is therefore to be interpreted as

$$J^{\mu\text{sym.}}(\mathbf{x}, t) = \frac{1}{2} \sum_{ij} \left(\psi_i^\dagger(\mathbf{x}, t)\psi_j(\mathbf{x}, t) - \psi_j(\mathbf{x}, t)\psi_i^\dagger(\mathbf{x}, t) \right) \alpha_{ij}^\mu, \quad (9)$$

when we calculate its vacuum expectation value.

III. THEORY OF ELECTRON-NUCLEUS-RADIATION COUPLED DYNAMICS

Having formulated fundamental quantities, we now develop the Tamm-Dancoff expansion of the electron-nucleus-radiation coupled dynamics.

A. The lowest order expansion in electron-radiation coupled dynamics

We first discuss electron-radiation coupled dynamics for a given nuclear configuration. Couplings to dynamical nuclei are restored in Subsection III B. Following the original idea of Tamm⁶⁰ and Dancoff,⁶¹ we expand the electron-radiation coupled state vector as a superposition of zero, one, two, . . . photon states. We formally introduce notation (N_e, \bar{N}_e, N_{ph}) , where N_e , \bar{N}_e , and N_{ph} represent the number of electrons, positrons, and photons. Each triplet represents a “sector.” We assume that the physical state of our interest is dominated by a small number of “active” sectors around $(N_e, 0, 0)$ and truncate other sectors. We first consider the electron-radiation dynamics for a given nuclear coordinate. We also assume that the mean-field equation [Eq. (4)] is already solved and the relevant MOs are known. We then set the zeroth order Hamiltonian as

$$H_0 = H_{\text{mf}}^{\text{el}} + H_{\text{rad}}, \quad (10)$$

and the residual interaction for the correction is given as Eq. (8c).

We first consider the smallest nontrivial set of “sectors,” which consists of $(N_e, 0, 0)$, $(N_e, 0, 1)$, and $(N_e + 1, 1, 1)$. For simplicity of notation, we also use more intuitive labels for those sectors: 000, 001, and 111, respectively.

A brief note is made here on the $(N_e + 1, 1, 1)$ or 111 sector; it consists of states with one electron-positron pair plus a photon, wherein we are not necessarily assuming existence of “real” electron-positron pair, which would require energy of order ~ 1 MeV. In the lower energy dynamics, those states are regarded as “virtual” states, which are, nevertheless, not simply negligible for its possible roles in four-component dynamics (see Ref. 78, for an illustrative example) and for achieving mathematically consistent description of dynamics.

1. Coordinate representation

Following Ref. 68, we define an “amplitude” [hereafter referred to as the Tamm-Dancoff (TD) amplitude] of the electronic state vector in $(N_e, 0, 0)$ sector as

$$\mathcal{F}_t^{000}(\{\mathbf{x}\}) \equiv \langle 0 | \tilde{\psi}(\mathbf{x}_1, t) \dots \tilde{\psi}(\mathbf{x}_{N_e}, t) | \Psi_t^{\text{int}} \rangle, \quad (11)$$

where the electronic annihilation operator $\tilde{\psi}$, the vacuum state $|0\rangle$, and the time-dependent electronic state vector Ψ_t^{int} are all in the

interaction representation. The time evolution equation reads

$$i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{\mathbf{x}\}) = \langle 0 | [\tilde{\psi}(\mathbf{x}_1, t) \dots \tilde{\psi}(\mathbf{x}_{N_e}, t), H_0] | \Psi_t^{\text{int}} \rangle + \langle 0 | \tilde{\psi}(\mathbf{x}_1, t) \dots \tilde{\psi}(\mathbf{x}_{N_e}, t) H_{\text{int}}^{\text{el}}(t) | \Psi_t^{\text{int}} \rangle. \quad (12)$$

We do not have to take account of all terms in $H_{\text{int}}^{\text{el}}$ of Eq. (8c) to obtain the correct equation of \mathcal{F}_t^{000} . Since there is a strict one-to-one correspondence in the transversal photon exchange and the Coulombic interaction (plus its associated mean-field subtraction terms), we can concentrate on the transversal interaction part

$$H_{\text{int}}^{\text{el tr}} = -q_e \int d^4x \psi^\dagger(x) \boldsymbol{\alpha} \psi(x) \cdot \mathbf{A}^{\text{tr}}(x) \quad (13)$$

to expand the perturbation series. Contributions arising from the remaining part, denoted by $H_{\text{int}}^{\text{el } \prime} \equiv H_{\text{int}}^{\text{el}} - H_{\text{int}}^{\text{el tr}}$, are restored when we evaluate the obtained diagrams by re-interpreting the photon lines as a summation of the Coulombic and transversal interactions. Except for those representing the self-interactions (i.e., the photon lines emitted and absorbed by the same electron), we also attach the mean-field subtraction terms to those photon lines.

For later convenience, we introduce the mean-field electronic propagator S^F by

$$S^F(x, y) \equiv \frac{1}{i\hbar} \langle 0 | \mathcal{T}(\tilde{\psi}(x) \tilde{\psi}^\dagger(y)) | 0 \rangle, \quad (14)$$

with \mathcal{T} being the time-ordering operator.

We now expand the last term in Eq. (12) using $H_{\text{int}}^{\text{el tr}}$ given as Eq. (13). Details of the expansion procedure is given in Appendix A, but in short, we move all creation operators appearing in $H_{\text{int}}^{\text{el tr}}$ to left by application of the equal-time (anti-)commutation rule. We then obtain

$$i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{\mathbf{x}\}) = H_0 \mathcal{F}_t^{000}(\{\mathbf{x}\}) + \sum_{\mathbf{k}\lambda} \left(\sum_j (\Lambda^+ \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{x}_j))_{(j)} + \mathcal{J}_{-\mathbf{k}}^\mu \boldsymbol{\varepsilon}_{\mathbf{k}\lambda\mu} \right) \times \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda) + \sum_{\mathbf{k}\lambda} \sum_{i_h j_p} \int d^3y (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{y}))_{i_h j_p} \times \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}; i_h, j_p; \mathbf{k}\lambda) + (\text{terms arising from } H_{\text{int}}^{\text{el } \prime}), \quad (15)$$

where Λ^+ (Λ^-) is the projection operator to positive (negative) energy states. The index $\mathbf{k}\lambda$ represents the photon wavevector \mathbf{k} and the polarization index λ , whose associated frequency being $\omega_{\mathbf{k}} = c|\mathbf{k}|$. For notational convenience, we combine the polarization vector $\boldsymbol{\varepsilon}_{\mathbf{k}\lambda\mu}$ and all numerical factors associated with the photon annihilation operator \mathbf{A}^{tr} and further contract with the electronic Dirac spinors α^μ to define

$$\boldsymbol{\varepsilon}_{\mathbf{k}\lambda\mu} \equiv q_e \sqrt{\frac{4\pi c^2 \hbar}{2\omega_{\mathbf{k}} \text{Vol.}}} \mathbf{e}_{\mathbf{k}\lambda\mu}, \quad (16a)$$

$$\boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{x}\ell) \equiv \boldsymbol{\varepsilon}_{\mathbf{k}\lambda\mu}(\alpha^\mu)_{(\ell)} e^{i\mathbf{k}\mathbf{x}\ell}, \quad (16b)$$

with Vol. representing the system volume, which is to be taken the limit $\text{Vol.} \rightarrow \infty$ in the end of the calculation while keeping \mathbf{k} summation finite, i.e., replacing summation by integration as $\frac{1}{\text{Vol.}} \sum_{\mathbf{k}} \rightarrow \int d^3\mathbf{k} / (2\pi)^3$. Subscripts (ℓ) attached to spinor matrices indicate

that the corresponding spinor matrix operates on the ℓ th component of the rank- N_e spinor \mathcal{F} as

$$(\alpha)_{(\ell)}\mathcal{F} \equiv \sum_{i'_\ell} (\alpha)_{i_\ell i'_\ell} \mathcal{F}_{i_1, \dots, i'_\ell, \dots, i_{N_e}}. \quad (17)$$

$\mathcal{F}_{\mathbf{k}}^\mu$ symbolizes the Fourier transformation of the vacuum expectation value of the current

$$\mathcal{F}_{\mathbf{k}}^\mu \equiv \int d^3\mathbf{x} \mathcal{J}^\mu(\mathbf{x}, t) e^{-i\mathbf{k}\cdot\mathbf{x}}, \quad (18)$$

where $\mathcal{J}^\mu(\mathbf{x}, t)$ represents the vacuum expectation value of the current operator

$$\begin{aligned} \mathcal{J}^\mu(\mathbf{x}, t) &= \frac{1}{2} \sum_{ij} \langle 0 | \psi_i^\dagger(\mathbf{x}, t) \psi_j(\mathbf{x}, t) - \psi_j(\mathbf{x}, t) \psi_i^\dagger(\mathbf{x}, t) | 0 \rangle \alpha_{ij}^\mu \\ &= -i\hbar \frac{1}{2} \text{Tr} (S^F(\mathbf{x}, t - 0; \mathbf{x}, t) \alpha^\mu + S^F(\mathbf{x}, t; \mathbf{x}, t - 0) \alpha^\mu), \end{aligned} \quad (19)$$

with $t - 0$ in the last side indicating an infinitesimally small time before t . While the corresponding quantity vanishes in the free-particle system,⁶⁴ it generally has a nonzero value in our

model because of the existence of the mean-field potential. It can be evaluated either by direct numerical calculation using a large set of MOs or by perturbation expansion of the mean-field propagator S^F by the free-particle propagator S_0^F as $\mathcal{J}^\mu(\mathbf{x}) \approx -i\hbar \int d\tau d^3\xi \text{Tr}(\alpha^\mu S_0^F(\mathbf{x}, 0; \xi, \tau) S_0^F(\xi, \tau; \mathbf{x}, 0)) W_{\text{HF}}^{\text{loc}}(\xi, \tau) + \dots$. We also note that the term “vacuum” comes from the vacuum state in the interaction picture $|0\rangle$, which is actually affected by the mean-field Hamiltonian. All related ideas in our following discussion, including the vacuum polarization and vacuum energy shift, reflect the mean-field potential as their physical origin.

New amplitudes belonging to the $(N_e, 0, 1)$ and $(N_e + 1, 1, 1)$ sectors appearing in Eq. (15) are defined, respectively, as

$$\mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda) \equiv \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) a_{\mathbf{k}\lambda}(t) | \Psi_t^{\text{int}} \rangle \quad (20)$$

and

$$\begin{aligned} \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{k}\lambda) &\equiv \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) \left(\check{\psi}_<^\dagger(\mathbf{y}_h, t) \right)_{i_h} \\ &\quad \times \left(\check{\psi}_>(\mathbf{y}_p, t) \right)_{j_p} a_{\mathbf{k}\lambda}(t) | \Psi_t^{\text{int}} \rangle. \end{aligned} \quad (21)$$

The derivation of Eq. (15) is given in detail in Appendix A. The time evolution equation for \mathcal{F}^{001} reads

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda) &= H_0 \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda) + \left(\sum_{\ell} (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{x}_\ell))_{(\ell)} + \mathcal{J}_{\mathbf{k}}^\mu \mathfrak{e}_{\mathbf{k}\lambda}^* \right) \mathcal{F}_t^{000} + \sum_{\mathbf{k}'\lambda'} \sum_{\ell} (\Lambda^+ \mathfrak{s}_{\mathbf{k}'\lambda'}(\mathbf{x}_\ell))_{(\ell)} \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) a_{\mathbf{k}\lambda} a_{\mathbf{k}'\lambda'} | \Psi_t^{\text{int}} \rangle \\ &\quad + \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) \int d^3\mathbf{y} \check{\psi}_<^\dagger(\mathbf{y}, t) \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{y}) \check{\psi}_>(\mathbf{y}, t) | \Psi_t^{\text{int}} \rangle \\ &\quad + \sum_{\mathbf{k}'\lambda'} \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) \int d^3\mathbf{y} \check{\psi}_<^\dagger(\mathbf{y}, t) \mathfrak{s}_{\mathbf{k}'\lambda'}(\mathbf{y}) \check{\psi}_>(\mathbf{y}, t) a_{\mathbf{k}\lambda} a_{\mathbf{k}'\lambda'} | \Psi_t^{\text{int}} \rangle + (\text{terms arising from } H_{\text{int}}^{\text{el}'}), \end{aligned} \quad (22)$$

and that for \mathcal{F}^{111} becomes

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{k}\lambda) &= H_0 \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{k}\lambda) + \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) | \Psi_t^{\text{int}} \rangle \delta^3(\mathbf{y}_h - \mathbf{y}_p) (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{y}_h) \Lambda^-)_{j_p i_h} \\ &\quad - \sum_{\ell} (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{x}_\ell) \Lambda^-)_{i_\ell j_h} \delta^3(\mathbf{x}_\ell - \mathbf{y}_h) \mathcal{F}_t^{000}(\{\mathbf{x}\})_{|i_\ell \rightarrow j_p} \\ &\quad + (\text{terms that do not belong to the } (N_e, 0, 0) \text{ or } (N_e, 1, 1) \text{ sector}) + (\text{terms arising from } H_{\text{int}}^{\text{el}'}), \end{aligned} \quad (23)$$

where the expression $\mathcal{F}_{|i_\ell \rightarrow j_p}$ indicates the spinor \mathcal{F} whose ℓ th index i_ℓ is replaced by j_p .

Concentrating on the terms proportional to the $(N_e, 0, 0)$ sector amplitudes only, a formal solution of Eq. (22) can be symbolically written as

$$\mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda) = \left[i\hbar \frac{\partial}{\partial t} - H_0 \right]^{-1} \left(\sum_{\ell} (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{x}_\ell))_{(\ell)} + \mathcal{J}_{\mathbf{k}}^\mu \mathfrak{e}_{\mathbf{k}\lambda}^* \right) \mathcal{F}_t^{000}(\{\mathbf{x}\}), \quad (24)$$

whereas Eq. (23) is solved as

$$\mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{k}\lambda) = \left[i\hbar \frac{\partial}{\partial t} - H_0 \right]^{-1} \left[(\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{y}_h) \Lambda^-)_{j_p i_h} \delta^3(\mathbf{y}_h - \mathbf{y}_p) \mathcal{F}_t^{000}(\{\mathbf{x}\}) - (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{x}_\ell) \Lambda^-)_{i_\ell j_h} \delta^3(\mathbf{x}_\ell - \mathbf{y}_h) \mathcal{F}_t^{000}(\{\mathbf{x}\})_{|i_\ell \rightarrow j_p} \right]. \quad (25)$$

Substituting these expressions into Eq. (15) and restoring the Coulombic and mean-field subtraction terms, we obtain

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{\mathbf{x}\}) &= \left[\sum_j (\hat{h}_j^{\text{mf}} - W_{\text{HF}j}^{\text{loc}}) + V_C + V_{\text{SI}} \right] \mathcal{F}_t^{000}(\{\mathbf{x}\}) \\
 &+ \sum_{\mathbf{k}\lambda} \left(\sum_j (\Lambda^+ \varepsilon_{\mathbf{k}\lambda}(\mathbf{x}_j))_{(j)} + \mathcal{J}_{-\mathbf{k}}^{\mu} \varepsilon_{\mathbf{k}\lambda\mu} \right) \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i\omega_{\mathbf{k}}(t-t')} \\
 &\times \int \prod_k d^3 \mathbf{x}'_k \prod_k C^>(\mathbf{x}_k, t; \mathbf{x}_k, t') \left(\sum_{\ell} (\Lambda^+ \varepsilon_{\mathbf{k}\lambda}^*(\mathbf{x}'_{\ell}))_{(\ell)} + \mathcal{J}_{\mathbf{k}}^{\nu} \varepsilon_{\mathbf{k}\lambda\nu}^* \right) \mathcal{F}_{t'}^{000}(\{\mathbf{x}'\}) \\
 &+ \sum_{\mathbf{k}\lambda} \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i\omega_{\mathbf{k}}(t-t')} \sum_{\ell} \int d^3 \mathbf{y}'_h d^3 \mathbf{y} \prod_k d^3 \mathbf{x}'_k \prod_{k \neq \ell} C^>(\mathbf{x}_k, t; \mathbf{x}'_k, t') \\
 &\times (C^>(\mathbf{x}_{\ell}, t; \mathbf{y}'_h, t') \varepsilon_{\mathbf{k}\lambda}^*(\mathbf{y}'_h) C^<(\mathbf{y}'_h, t'; \mathbf{y}, t) \varepsilon_{\mathbf{k}\lambda}(\mathbf{y}) C^>(\mathbf{y}, t; \mathbf{x}'_{\ell}, t'))_{(\ell)} \mathcal{F}_{t'}^{000}(\{\mathbf{x}'\}) \\
 &+ \sum_{\mathbf{k}\lambda} \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i\omega_{\mathbf{k}}(t-t')} \int d^3 \mathbf{y} d^3 \mathbf{y}' \prod_k d^3 \mathbf{x}'_k \prod_k C^>(\mathbf{x}_k, t; \mathbf{x}'_k, t') \\
 &\times \text{Tr}(C^>(\mathbf{y}, t; \mathbf{y}', t') \varepsilon_{\mathbf{k}\lambda}^*(\mathbf{y}') C^<(\mathbf{y}', t'; \mathbf{y}, t) \varepsilon_{\mathbf{k}\lambda}(\mathbf{y})) \mathcal{F}_{t'}^{000}(\{\mathbf{x}'\}), \tag{26}
 \end{aligned}$$

where \hat{h}_j^{mf} and $W_{\text{HF}j}^{\text{loc}}$ represent the mean-field Hamiltonian in the operator representation and the mean-field potential acting on the j th electronic coordinate in \mathcal{F}_t^{000} , whereas V_C and V_{SI} represent the Coulombic (inter-electronic) interaction and the Coulombic self-interaction,⁷⁹ respectively. Symbols $C^>$ and $C^<$ represent the zeroth order propagators defined as

$$C^>(x, y) \equiv \langle 0 | \hat{\psi}_>(x) \hat{\psi}_>^{\dagger}(y) | 0 \rangle \theta(x^0 - y^0), \tag{27a}$$

$$C^<_i(x, y) \equiv \langle 0 | -(\hat{\psi}_<^{\dagger}(y))_i (\hat{\psi}_<(x))_i | 0 \rangle \theta(y^0 - x^0), \tag{27b}$$

with $\psi_>$ ($\psi_<$) representing the positive (negative) energy part of the electronic field operator. Using those propagators, the formal expression $[i\hbar \frac{\partial}{\partial t} - H_{\text{mf}}]^{-1}$ appearing in Eqs. (24) and (25) is expanded as

$$\begin{aligned}
 &\left[i\hbar \frac{\partial}{\partial t} - H_{\text{mf}} \right]^{-1} \phi_t(\{\mathbf{x}\}, \{\bar{\mathbf{x}}\}) \\
 &= \frac{1}{i\hbar} \int_{-\infty}^t dt' \int \prod_i d^3 \mathbf{x}'_i \int \prod_a d^3 \bar{\mathbf{x}}'_a \prod_i C^>(\mathbf{x}_i, t; \mathbf{x}'_i, t') \\
 &\times \prod_a (-C^<(\bar{\mathbf{x}}'_a, t'; \bar{\mathbf{x}}_a, t)) \phi_{t'}(\{\mathbf{x}'\}, \{\bar{\mathbf{x}}'\}), \tag{28}
 \end{aligned}$$

with ϕ_t representing an arbitrary physical quantity and $\{\mathbf{x}\}$ ($\{\bar{\mathbf{x}}\}$) representing electronic (positronic) coordinates.

Equation (26) describes the time evolution of the electronic system that interacts through the Coulombic interaction and the transversal photon exchange, and yet Eq. (26) also includes apparently divergent self-interaction terms whose renormalization is better formulated in the MO-representation as we discuss below.

2. Molecular orbital representation

We reformulate the above procedure in the MO-representation. The MO-representation of Eq. (16) is defined as

$$(\varepsilon_{\mathbf{k}\lambda})_{\ell m} \equiv \int d^3 \mathbf{y}' \varphi_{\ell}^{\dagger}(\mathbf{y}') \varepsilon_{\mathbf{k}\lambda}(\mathbf{y}') \varphi_m(\mathbf{y}'). \tag{29}$$

We also use a collective notation \mathbf{I} , which represents a set of MO indices of the N_e (positive energy) electronic orbitals as $\mathbf{I} \equiv \{I_1, I_2, \dots, I_{N_e}\}$, and \mathbf{I}'_j is given by replacing the j th component I_j by the r th MO as $\mathbf{I}'_j \equiv \{I_1, I_2, \dots, I_{j-1}, r, I_{j+1}, \dots, I_{N_e}\}$.

We then define amplitudes in 000, 001, and 111 sectors by

$$f_t^{000}(\mathbf{I}) \equiv \langle 0 | \hat{c}_{I_1}(t) \hat{c}_{I_2}(t) \dots \hat{c}_{I_{N_e}}(t) | \Psi_t^{\text{int}} \rangle, \tag{30a}$$

$$f_t^{001}(\mathbf{I}; \mathbf{k}\lambda) \equiv \langle 0 | \hat{c}_{I_1}(t) \hat{c}_{I_2}(t) \dots \hat{c}_{I_{N_e}}(t) a_{\mathbf{k}\lambda}(t) | \Psi_t^{\text{int}} \rangle, \tag{30b}$$

$$f_t^{111}(\mathbf{I}; a, r, \mathbf{k}\lambda) \equiv \langle 0 | \hat{c}_{I_1}(t) \hat{c}_{I_2}(t) \dots \hat{c}_{I_{N_e}}(t) \hat{b}_a(t) \hat{c}_r(t) a_{\mathbf{k}\lambda}(t) | \Psi_t^{\text{int}} \rangle. \tag{30c}$$

The time evolution equations for them [Eqs. (15), (22), and (23)] are rewritten as

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} f_t^{000}(\mathbf{I}) &= \sum_k \varepsilon^0_{\mathbf{I}_k} f_t^{000}(\mathbf{I}) + \sum_{\mathbf{k}\lambda} \left[\sum_{j,r} (\varepsilon_{\mathbf{k}\lambda})_{I_j r} f_t^{001}(\mathbf{I}'_j; \mathbf{k}\lambda) \right. \\
 &\left. + \mathcal{J}_{-\mathbf{k}}^{\mu} \varepsilon_{\mathbf{k}\lambda\mu} f_t^{001}(\mathbf{I}; \mathbf{k}\lambda) \right] + \sum_{r,a} \sum_{\mathbf{k}\lambda} (\varepsilon_{\mathbf{k}\lambda})_{ar} f_t^{111}(\mathbf{I}; a, r; \mathbf{k}\lambda), \tag{31a}
 \end{aligned}$$

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} f_t^{001}(\mathbf{I}; \mathbf{k}\lambda) &= \left(\sum_k \varepsilon^0_{\mathbf{I}_k} + \hbar\omega_{\mathbf{k}} \right) f_t^{001}(\mathbf{I}; \mathbf{k}\lambda) \\
 &+ \sum_{j,r} (\varepsilon_{\mathbf{k}\lambda}^*)_{I_j r} f_t^{000}(\mathbf{I}'_j) + \mathcal{J}_{\mathbf{k}}^{\mu} \varepsilon_{\mathbf{k}\lambda\mu}^* f_t^{000}(\mathbf{I}) + \dots, \tag{31b}
 \end{aligned}$$

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} f_t^{111}(\mathbf{I}; a, r; \mathbf{k}\lambda) &= \left(\sum_k \varepsilon^0_{\mathbf{I}_k} + |\varepsilon^0_a| + \varepsilon^0_r + \hbar\omega_{\mathbf{k}} \right) f_t^{111}(\mathbf{I}; a, r; \mathbf{k}\lambda) \\
 &- \sum_j (\varepsilon_{\mathbf{k}\lambda}^*)_{I_j, a} f_t^{000}(\mathbf{I}'_j) + \mathcal{J}_{-\mathbf{k}}^{\mu} \varepsilon_{\mathbf{k}\lambda\mu}^* f_t^{000}(\mathbf{I}) + \dots. \tag{31c}
 \end{aligned}$$

Substitution of solutions of Eqs. (31b) and (31c) into Eq. (31a) yields the following equation of f_t^{000} :

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} f_t^{000}(\mathbf{I}) &= \sum_k \epsilon^0 \mathbf{I}_k f_t^{000}(\mathbf{I}) - \sum_j \left(W_{\text{HF}j}^{\text{loc}} \right)_{I_j, r} f_t^{000}(\mathbf{I}_j^r) + \frac{1}{2} \sum_{j\ell} \langle I_j I_\ell \| r s \rangle f_t^{000}(\mathbf{I}_{j\ell}^{rs}) + \frac{1}{2} \sum_j \iint d^3 \mathbf{r} d^3 \mathbf{r}' |\varphi_j(\mathbf{r})|^2 \frac{q_e^2}{|\mathbf{r} - \mathbf{r}'|} \delta(\mathbf{r} - \mathbf{r}') \\
 &+ \hat{\Delta}^{\text{vp}} f_t^{000}(\mathbf{I}) + \sum_{j \neq \ell} \sum_{\mathbf{k}\lambda} \sum_{r,s} (\mathfrak{S}_{\mathbf{k}\lambda})_{I_j r} \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i \left(\sum_{k \neq j} \epsilon^0 I_k + \epsilon^0_r + \hbar\omega_{\mathbf{k}} \right) (t-t')/\hbar} (\mathfrak{S}_{\mathbf{k}\lambda}^*)_{I_\ell s} f_{t'}^{000}(\mathbf{I}_{j\ell}^{rs}) \\
 &+ \sum_j \sum_{\mathbf{k}\lambda} \sum_{r,s} (\mathfrak{S}_{\mathbf{k}\lambda})_{I_j r} \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i \left(\sum_{k \neq j} \epsilon^0 I_k + \epsilon^0_r + \hbar\omega_{\mathbf{k}} \right) (t-t')/\hbar} (\mathfrak{S}_{\mathbf{k}\lambda}^*)_{rs} f_{t'}^{000}(\mathbf{I}_j^s) \\
 &- \sum_j \sum_{\mathbf{k}\lambda} \sum_{a,r} (\mathfrak{S}_{\mathbf{k}\lambda}^*)_{I_j a} \int_{-\infty}^t dt' e^{-i \left(\sum_k \epsilon^0 I_k + \epsilon^0_r + \hbar\omega_{\mathbf{k}} \right) (t-t')/\hbar} e^{-i \epsilon^0_a (t'-t)/\hbar} (\mathfrak{S}_{\mathbf{k}\lambda}^*)_{a,r} f_{t'}^{000}(\mathbf{I}_j^r) \\
 &+ \sum_{\mathbf{k}\lambda} \sum_{j,r} (\mathcal{J}_{-\mathbf{k}}^{\mu} \mathfrak{E}_{\mathbf{k}\lambda\mu}) \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i \left(\sum_k \epsilon^0 I_k + \hbar\omega_{\mathbf{k}} \right) (t-t')/\hbar} (\mathfrak{S}_{-\mathbf{k}\lambda}^*)_{I_j, r} f_{t'}^{000}(\mathbf{I}_j^r) \\
 &+ \sum_{\mathbf{k}\lambda} \sum_{j,r} (\mathfrak{S}_{\mathbf{k}\lambda})_{I_j, r} \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i \left(\sum_{k \neq j} \epsilon^0 I_k + \hbar\omega_{\mathbf{k}} + \epsilon^0_r \right) (t-t')/\hbar} (\mathcal{J}_{\mathbf{k}}^{\mu} \mathfrak{E}_{\mathbf{k}\lambda\mu}) f_{t'}^{000}(\mathbf{I}_j^r), \quad (32)
 \end{aligned}$$

with $\langle I_j I_\ell \| r s \rangle$ representing the instantaneous Coulomb-exchange interaction among four MOs I_j , I_ℓ , r , and s (in the physicists' notation). By the symbol $\hat{\Delta}^{\text{vp}} f_t^{000}(\mathbf{I})$, we separated out contributions that purely arise from vacuum polarization and does not have direct effects on the dynamics of molecular electrons. It reads

$$\begin{aligned}
 \hat{\Delta}^{\text{vp}} f_t^{000}(\mathbf{I}) &= \frac{1}{2} \iint d^3 \mathbf{x} d^3 \mathbf{y} \mathcal{J}^0(\mathbf{x}) \frac{1}{|\mathbf{x} - \mathbf{y}|} \mathcal{J}^0(\mathbf{y}) f_t^{000}(\mathbf{I}) - \frac{1}{2} (i\hbar)^2 \iint d^3 \mathbf{x} d^3 \mathbf{y} S^F(\mathbf{x}, t; \mathbf{y}, t-0) S^F(\mathbf{y}, t-0; \mathbf{x}, t) \frac{1}{|\mathbf{x} - \mathbf{y}|} f_t^{000}(\mathbf{I}) \\
 &+ \sum_{\mathbf{k}\lambda} (\mathcal{J}_{-\mathbf{k}}^{\nu} \mathfrak{E}_{\mathbf{k}\lambda\nu}) \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i \left(\sum_k \epsilon^0 I_k + \hbar\omega_{\mathbf{k}} \right) (t-t')/\hbar} (\mathcal{J}_{\mathbf{k}}^{\mu} \mathfrak{E}_{\mathbf{k}\lambda\mu}^*) f_{t'}^{000}(\mathbf{I}) \\
 &+ \sum_{\mathbf{k}\lambda} \frac{1}{i\hbar} \int_{-\infty}^t dt' e^{-i \left(\sum_k \epsilon^0 I_k \right) (t-t')/\hbar} \sum_{r,a} (\mathfrak{S}_{\mathbf{k}\lambda}^*)_{ra} e^{-i \epsilon^0_a (t'-t)/\hbar} (\mathfrak{S}_{\mathbf{k}\lambda})_{ar} e^{-i \epsilon^0_r (t-t')/\hbar} e^{-i \omega_{\mathbf{k}} (t-t')/\hbar} f_{t'}^{000}(\mathbf{I}), \quad (33)
 \end{aligned}$$

which will be discussed later; below we concentrate on the terms of dynamical relevance.

Equation (32) contains a number of self-interaction terms, the Coulombic as well as the transversal photon exchange interactions. These are apparently divergent, and we need appropriate renormalization to obtain a meaningful result.

We first combine two transversal self-interaction terms in Eq. (45) to define a quantity $\Sigma_{1L I_j, s}^{\text{tr}}(t-t')$,

$$\begin{aligned}
 \Sigma_{1L I_j, s}^{\text{tr}}(t-t') &= \sum_{\mathbf{k}\lambda} \left\langle \varphi_j \left| \left[\theta(t-t') \sum_{r \in \Lambda^+} \mathfrak{S}_{\mathbf{k}\lambda} |\varphi_r\rangle e^{-i(\epsilon^0_r + \hbar\omega_{\mathbf{k}})(t-t')/\hbar} \langle \varphi_r | \mathfrak{S}_{\mathbf{k}\lambda}^* \right. \right. \right. \\
 &\quad \left. \left. - \theta(t'-t) \sum_{a \in \Lambda^-} \mathfrak{S}_{\mathbf{k}\lambda}^* |\varphi_a\rangle e^{-i(\epsilon^0_a - \hbar\omega_{\mathbf{k}})(t'-t)/\hbar} \langle \varphi_a | \mathfrak{S}_{\mathbf{k}\lambda} \right] \right| \varphi_s \rangle, \quad (34)
 \end{aligned}$$

and its Fourier transformation

$$\begin{aligned}
 \Sigma_{1L I_j, s}^{\text{tr}}(\omega) &= \frac{q_e^2}{\mathcal{V} \text{ol.}} \sum_{\mathbf{k}} \int d^3 \mathbf{x} \int d^3 \mathbf{y} \varphi_j^\dagger(\mathbf{x}) \alpha^t S^F(\mathbf{x}, \mathbf{y}; \omega - \omega_{\mathbf{k}}) \alpha^s \\
 &\quad \times \left\{ \delta_{ij} - \frac{k_i k_j}{\mathbf{k}^2} \right\} \frac{4\pi c^2 \hbar}{2\omega_{\mathbf{k}}} e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} \varphi_s(\mathbf{y}) \\
 &= i\hbar c q_e^2 \int \frac{d^4 k}{(2\pi)^4} \int d^3 \mathbf{x} \int d^3 \mathbf{y} \varphi_j^\dagger(\mathbf{x}) \alpha^t S^F(\mathbf{x}, \mathbf{y}; \omega - ck^0) \alpha^s \\
 &\quad \times \left\{ \delta_{ij} - \frac{k_i k_j}{\mathbf{k}^2} \right\} \frac{4\pi}{k^2 + i\eta} e^{-i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} \varphi_s(\mathbf{y}), \quad (35)
 \end{aligned}$$

where η represents an infinitesimal positive constant and the mean-field propagator S^F in the MO representation expands as

$$S^F(\mathbf{x}, \mathbf{y}; \omega) = \sum_{r \in \Lambda^+} \frac{\varphi_r(\mathbf{x}) \varphi_r^\dagger(\mathbf{y})}{\hbar\omega - \epsilon^0_r + i\eta} + \sum_{a \in \Lambda^-} \frac{\varphi_a(\mathbf{x}) \varphi_a^\dagger(\mathbf{y})}{\hbar\omega - \epsilon^0_a - i\eta}. \quad (36)$$

We then combine the Coulombic self-interaction contribution as well, to obtain

$$\begin{aligned}
 (\Sigma_{1L}(\omega))_{I_j, s} &= i\hbar c q_e^2 \int \frac{d^4 k}{(2\pi)^4} \int d^3 \mathbf{x} \\
 &\quad \times \int d^3 \mathbf{y} \varphi_j^\dagger(\mathbf{x}) \alpha^t S^F(\mathbf{x}, \mathbf{y}; \omega - ck^0) \alpha^s \\
 &\quad \times \left[\delta_{ij}^0 \delta_{ij}^0 \frac{4\pi}{\mathbf{k}^2} + (1 - \delta_{ij}^0)(1 - \delta_{ij}^0) \left\{ \delta_{ij}^0 - \frac{k_i k_j}{\mathbf{k}^2} \right\} \frac{4\pi}{k^2 + i\eta} \right] \\
 &\quad \times e^{-i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} \varphi_s(\mathbf{y}). \quad (37)
 \end{aligned}$$

Self-energy renormalization using the Coulomb gauge is formulated in Refs. 80 and 81, and we may well use these results. For simplicity of discussion, however, we here assume that the final result should be invariant with respect to the choice of the gauge, and we replace the photon propagator by the Feynman gauge propagator,

$$\begin{aligned}
 (\Sigma_{1L}(\omega))_{I_j, s} &= i\hbar c q_e^2 \int \frac{d^4 k}{(2\pi)^4} \int d^3 \mathbf{x} \int d^3 \mathbf{y} \varphi_j^\dagger(\mathbf{x}) \alpha^t \\
 &\quad \times S^F(\mathbf{x}, \mathbf{y}; \omega - ck^0) \alpha^s \frac{4\pi \eta_{\mu\nu}}{-k^2 - i\eta} e^{-i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} \varphi_s(\mathbf{y}) \\
 &= \int d^3 \mathbf{x} \int d^3 \mathbf{y} \varphi_j^\dagger(\mathbf{x}) \Sigma_{1L}(\mathbf{x}, \mathbf{y}; \omega) \varphi_s(\mathbf{y}), \quad (38)
 \end{aligned}$$

where $\Sigma_{1L}(\mathbf{x}, \mathbf{y}; \omega)$ is the one-loop electronic self-energy constructed from the Hartree-Fock propagator. Apparent divergence in $\Sigma_{1L}(\mathbf{x}, \mathbf{y}; \omega)$ can be renormalized using the standard procedure by subtracting the counter terms, which leaves a finite contribution $\Sigma_{1L}^{\text{fin.}}(\mathbf{x}, \mathbf{y}; \omega)$. Formal expression of the finite part of the single-loop self-energy can be written as

$$\begin{aligned} \Sigma_{1L}^{\text{fin.}}(x, y) &= i\hbar c q_e^2 \alpha^\mu S^F(x, y) \alpha^\nu D_{\mu\nu}(x, y) - \beta \delta m c^2 \delta^4(x - y) - Z_2 \left(c \alpha^\mu i \hbar \partial_\mu - W_{\text{HF}}^{\text{loc}}(\mathbf{x}) - \beta m c^2 \right) \delta^4(x - y) \\ &= i\hbar c q_e^2 \alpha^\mu S_0^F(x, y) \alpha^\nu D_{\mu\nu}(x, y) - \beta \delta m c^2 \delta^4(x - y) - Z_2 \left(c \alpha^\mu i \hbar \partial_\mu - \beta m c^2 \right) \delta^4(x - y) \\ &\quad + i\hbar c \int d^4 \xi \alpha^\mu S_0^F(x, \xi) W_{\text{HF}}^{\text{loc}}(\xi) S_0^F(\xi, y) \alpha^\nu D_{\mu\nu}(x - y) + Z_2 W_{\text{HF}}^{\text{loc}}(\mathbf{x}) \delta^4(x - y) \\ &\quad + i\hbar c \int d^4 \xi \int d^4 \eta \alpha^\mu S_0^F(x, \xi) W_{\text{HF}}^{\text{loc}}(\xi) S^F(\xi, \eta) W_{\text{HF}}^{\text{loc}}(\eta) S_0^F(\eta, y), \end{aligned} \quad (39)$$

where S_0^F is the free-electron propagator and $\delta m c^2$ and Z_2 are the mass and field-renormalization constants.² Detailed discussion of the self-energy, including its numerical calculation procedure, is found in a number of excellent reviews including Refs. 7 and 82.

We next consider terms including \mathcal{J}^μ , which arises from the vacuum polarization. The two terms with $(\mathcal{J}_{-\mathbf{k}}^\nu \varepsilon_{\mathbf{k}\nu})$ or its conjugate but not the both describe scattering of molecular electrons by the current \mathcal{J}^ν , as are schematically shown in Figs. 1(d) and 1(e). In order to get better insight, we introduce the following potential field arising from \mathcal{J}^μ :

$$\mathcal{V}_\mu^{\text{vp}}(\mathbf{x}, t) \alpha^\mu = \frac{1}{i\hbar c} \int d^4 \xi \sum_{\mathbf{k}, \lambda} \varepsilon_{\mathbf{k}, \lambda}(\mathbf{x}) e^{-i\omega_{\mathbf{k}}(t - \xi^0/c)} \varepsilon_{\mathbf{k}, \lambda \mu}^* \mathcal{J}^\mu(\xi) e^{-i\mathbf{k}\xi}. \quad (40)$$

Indeed, the two scattering terms in Eq. (32) essentially reduce to scattering by $\mathcal{V}_\mu^{\text{vp}}$. Apparent complication arises from formal distinction of the time ordering of the source term (\mathcal{J}^μ) and the scattering event ($\varepsilon_{\mathbf{k}, \lambda}$), as well as existence of the upper limit of the integration range in Eq. (32). Assuming approximate stationarity of \mathcal{J}^μ , we therefore should be able to replace those two terms by $\mathcal{V}_\mu^{\text{vp}}$. We further combine the Coulombic contribution to Eq. (40) as




FIG. 1. Schematic view of transversal photon exchange interactions included in Eq. (32) [(a)–(e)] and Eq. (33) [(f) and (g)]. Arrows represent electrons where upward and downward lines indicate positive and negative energy states, respectively. Wavy lines represent transversal photons. Circles attached with a single photon line, or “tadpoles,” which appear in (d)–(f) represent the vacuum expectation values of the current arising from the vacuum polarization. In each diagram, time flows upward and the upper vertices correspond to time t , while the lower ones correspond to t' of Eqs. (32) and (33).

$$\begin{aligned} \mathcal{V}_\mu^{\text{vp}}(\mathbf{x}, t) \alpha^\mu &= \frac{1}{i\hbar c} \int d^4 \xi \sum_{\mathbf{k}, \lambda} \left(\varepsilon_{\mathbf{k}, \lambda}(\mathbf{x}) e^{-i\omega_{\mathbf{k}}(t - \xi^0/c)} \varepsilon_{\mathbf{k}, \lambda \nu}^* \mathcal{J}^\nu(\xi) e^{-i\mathbf{k}\xi} \right. \\ &\quad \left. + e^{i\mathbf{k}\cdot(\mathbf{x} - \xi)} \frac{4\pi}{\mathbf{k}^2} \mathcal{J}^0(\xi) \delta(t - \xi^0/c) \right) \\ &= \int d^4 \xi \int \frac{d^4 k}{(2\pi)^4} \alpha^\mu \left[\delta_\mu^0 \delta_\nu^0 \frac{4\pi q_e^2}{\mathbf{k}^2} + (1 - \delta_\mu^0)(1 - \delta_\nu^0) \right. \\ &\quad \left. \times \left\{ \delta_{\mu\nu} - \frac{k_\mu k_\nu}{\mathbf{k}^2} \right\} \frac{4\pi q_e^2}{k^2 + i\eta} \right] \mathcal{J}^\nu(\xi) e^{-i\mathbf{k}\cdot(\mathbf{x} - \xi)}. \end{aligned} \quad (41)$$

Applying the same argument as Eq. (38), we can replace the square bracket in Eq. (41) by the corresponding expression in the Feynman gauge, $-4\pi q_e^2 \eta_{\mu\nu} / (k^2 + i0)$. We can then combine it into the self-energy as

$$\Sigma^{\text{fin.}}(\mathbf{x}, t; \mathbf{x}', t') = \Sigma_{1L}^{\text{fin.}}(\mathbf{x}, t; \mathbf{x}', t') + \delta^3(\mathbf{x} - \mathbf{x}') \delta(t - t') \mathcal{V}_\mu^{\text{vp}} \alpha^\mu(\mathbf{x}, t). \quad (42)$$

We finally discuss the vacuum polarization term we separated out as Eq. (33). It consists of two types of interaction, the Coulombic and the transversal photon exchange, and each has direct and exchangelike terms [see Figs. 1(f) and 1(g)]. A tricky point is that some of those terms have apparent dependence on the zeroth electronic energy, and it includes convolution with $f_t^{000}(\mathbf{I})$. We can, however, separate it by taking account of time dependence of the amplitude, $f_t(\mathbf{I}) \propto e^{-i\sum_k \varepsilon^0 I_k}$ and extending the time integration to ∞ to rewrite it as

$$\hat{\Delta}^{\text{vp}} f_t^{000}(\mathbf{I}) \approx \Delta E^{\text{vp}} f_t^{000}(\mathbf{I}), \quad (43)$$

where ΔE^{vp} is the vacuum energy shift,

$$\begin{aligned} \Delta E^{\text{vp}} &= -i\hbar \int dt' \int d^3 \mathbf{x}' i\hbar \int \frac{d^4 k}{(2\pi)^4} \left[\delta_\mu^0 \delta_\nu^0 \frac{4\pi q_e^2}{\mathbf{k}^2} + (1 - \delta_\mu^0)(1 - \delta_\nu^0) \right. \\ &\quad \left. \times \left\{ \delta_{\mu\nu} - \frac{k_\mu k_\nu}{\mathbf{k}^2} \right\} \frac{4\pi q_e^2}{k^2 + i\eta} \right] \left(\text{Tr}(\alpha^\mu S^F(x, y) \alpha^\nu S^F(y, x)) \right. \\ &\quad \left. + \text{Tr}(S^F(\mathbf{x}, t - 0; \mathbf{x}, t) \alpha^\mu) \text{Tr}(S^F(\mathbf{y}, t' - 0; \mathbf{y}, t') \alpha^\nu) \right) e^{-i\mathbf{k}\cdot(\mathbf{x} - \mathbf{y})}, \end{aligned} \quad (44)$$

where the square bracket can be replaced by the Feynman gauge expression, $-4\pi q_e^2 \eta_{\mu\nu} / (k^2 + i0)$. Although replacement Eq. (43) still lacks theoretical rigor, Eq. (44) is a formally correct expression for the one-loop vacuum energy shift in QED. Furthermore, as such terms only contribute to a phase factor of the system's wavefunction, we hereafter drop this term for simplicity.

Thus, we can now rewrite Eq. (32) as

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} f_t^{000}(\mathbf{I}) &= \sum_k \epsilon^0 \mathbf{I}_k f_t^{000}(\mathbf{I}) - \sum_j (W_{\text{HF}j}^{\text{loc}})_{I_j, r} f_t^{000}(\mathbf{I}_j^r) + \frac{1}{2} \sum_{j \neq \ell} \langle I_j I_\ell \| r s \rangle f_t^{000}(\mathbf{I}_{j\ell}^{rs}) \\ &+ \sum_{j \neq \ell} \sum_{\mathbf{k}\lambda} \sum_{r,s} (\mathfrak{s}_{\mathbf{k}\lambda})_{I_j r} \int_{-\infty}^t dt' e^{-i \left(\sum_{k \neq j} \epsilon^0 I_k + \epsilon^0_r + \hbar \omega_{\mathbf{k}} \right) (t-t') / \hbar} (\mathfrak{s}_{\mathbf{k}\lambda}^*)_{I_\ell s} f_{t'}^{000}(\mathbf{I}_{j\ell}^{rs}) \\ &+ \sum_{j,s} \iint_{-\infty}^t d\tau'_1 d\tau'_2 \{ \delta(\tau'_1 - (t-0)) + \delta(\tau'_2 - (t-0)) \} e^{-i \sum_{k \neq j} \epsilon^0 I_k (t-\tau'_k) / \hbar} \\ &\times e^{-i \epsilon^0_{I_j} (t-\tau'_1) / \hbar} \sum_{I_j, s}^{\text{fin.}} (\tau'_1, \tau'_2) e^{-i \epsilon^0_{I_\ell} (\tau'_2 - \tau'_1) / \hbar} f_{\tau'_2}^{000}(\mathbf{I}_j^s), \end{aligned} \quad (45)$$

where the self-energy term is defined as

$$\Sigma_{i_j}^{\text{fin.}}(\tau'_1, \tau'_2) \equiv \iint d^3 \mathbf{x} d^3 \mathbf{x}' \varphi_i(\mathbf{x})^\dagger \Sigma^{\text{fin.}}(\mathbf{x}, \tau; \mathbf{x}', \tau') \varphi_j(\mathbf{x}'), \quad (46)$$

and the delta functions in the curly bracket indicate that either τ'_1 or τ'_2 is to be fixed to t (with $t-0$ being an infinitesimal time before t) and $\tau'_<$ represents the smaller one of τ'_1 and τ'_2 . We can also rewrite the real-space expression Eq. (26) as

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{\mathbf{x}\}) &= \left[\sum_j (\hat{h}_j^{\text{mf}} - W_{\text{HF}j}^{\text{loc}}) + V_C \right] \mathcal{F}_t^{000}(\{\mathbf{x}\}) \\ &+ \sum_{j \neq \ell} \sum_{\mathbf{k}\lambda} (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}(\mathbf{x}_j))_{(j)} \int_{-\infty}^t dt' e^{-i \omega_{\mathbf{k}} (t-t')} \int \prod_i d^3 \mathbf{x}'_i S^>(\mathbf{x}_i, t; \mathbf{x}'_i, t') (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{x}_\ell))_{(\ell)} \mathcal{F}_{t'}^{000}(\{\mathbf{x}'\}) \\ &+ \sum_\ell \iint_{-\infty}^t d\tau'_1 d\tau'_2 \iint d^3 \xi'_1 d^3 \xi'_2 \{ \delta(\tau'_1 - (t-0)) + \delta(\tau'_2 - (t-0)) \} \int \prod_k d^3 \mathbf{x}'_k S^F(\mathbf{x}_\ell, t; \xi'_1, \tau'_1) \Sigma^{\text{fin.}}(\xi'_1, \tau'_1; \xi'_2, \tau'_2) \\ &\times S^F(\xi'_2, \tau'_2; \mathbf{x}'_k, \tau'_<) \prod_{k \neq \ell} S^F(\mathbf{x}_k, t; \mathbf{x}'_k, \tau'_<) \mathcal{F}_{\tau'_<}^{000}(\{\mathbf{x}'\}). \end{aligned} \quad (47)$$

The real-time expression of the self-energy operator Eq. (39) represents "fluctuation" (i.e., emission and absorption of virtual photons) induced by the radiation field. Assuming such effects can essentially be included in the form of energy correction, we can approximately rewrite Eq. (45) into a simpler form,

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} f_t^{000}(\mathbf{I}) &= \sum_j (\hat{h}_{\text{mf}} - W_{\text{HF}j}^{\text{loc}})_{I_j, r} f_t^{000}(\mathbf{I}_j^r) + \frac{1}{2} \sum_{j\ell} \langle I_j I_\ell \| r s \rangle f_t^{000}(\mathbf{I}_{j\ell}^{rs}) \\ &+ \sum_{j \neq \ell} \sum_{\mathbf{k}\lambda} \sum_{r,s} (\mathfrak{s}_{\mathbf{k}\lambda})_{I_j r} \int_{-\infty}^t dt' e^{-i \left(\sum_{k \neq j} \epsilon^0 I_k + \epsilon^0_r + \hbar \omega_{\mathbf{k}} \right) (t-t') / \hbar} \\ &\times (\mathfrak{s}_{\mathbf{k}\lambda}^*)_{I_\ell s} f_{t'}^{000}(\mathbf{I}_{j\ell}^{rs}) + \sum_j \bar{\Sigma}_{I_j, r} f_t^{000}(\mathbf{I}_j^r), \end{aligned} \quad (48)$$

where $\bar{\Sigma}_{\ell, m}$ is a time-independent matrix, defined by the Fourier transformation of the self-energy operator as

$$\begin{aligned} \bar{\Sigma}_{\ell, m} &= \frac{1}{2} \int d^3 \mathbf{x} \int d^3 \mathbf{y} \varphi_\ell^\dagger(\mathbf{x}) \left[\Sigma^{\text{fin.}}(\mathbf{x}, \mathbf{y}; \epsilon^0_\ell) \right. \\ &\left. + \Sigma^{\text{fin.}}(\mathbf{x}, \mathbf{y}; \epsilon^0_m) \right] \varphi_m(\mathbf{y}), \end{aligned} \quad (49)$$

which is seen to converge to an appropriate expression⁸³ in the static limit. We thus have removed the divergent terms in Eq. (47). Although our results Eq. (48) or its real-space representation Eq. (47) are aimed at relativistic dynamics, they have interesting applications in the nonrelativistic limit as is discussed in Appendix B.

In this paper, we have limited our discussion to the lowest order expansion. The higher order expansions [i.e., formulation taking account of higher orders of $H_{\text{int}}^{\text{el}}$ appearing in the RHS of Eq. (12)] can be in principle obtained in the same framework although it is not straightforward. We show a sketch of the higher order extensions in Appendix C, and the full expressions will be worked out in future study.

B. Coupling to nuclear dynamics

We next consider inclusion of nuclear degrees of freedom. We start with the formal second-quantized representation. We disregard internal structures of the atomic nuclei and assume that all nuclei in the system are distinguishable nonrelativistic particles. The nuclear Hamiltonian Eq. (8a) can formally be rewritten as

$$\begin{aligned} \mathcal{H}_{\text{nuc}} = & \int d^3\mathbf{R} \sum_a \hat{\chi}_a^\dagger(\mathbf{R}) \left[\frac{1}{2M_a} \left(\frac{\hbar}{i} \nabla - \frac{Z_a|e|}{c} \mathbf{A}^{\text{tr}} \right)^2 \right] \hat{\chi}_a(\mathbf{R}) \\ & + \frac{1}{2} \sum_{a < b} \iint d^3\mathbf{R} d^3\mathbf{R}' \hat{\chi}_a^\dagger(\mathbf{R}) \hat{\chi}_b^\dagger(\mathbf{R}') \frac{Z_a Z_b e^2}{|\mathbf{R} - \mathbf{R}'|} \hat{\chi}_b(\mathbf{R}') \hat{\chi}_a(\mathbf{R}), \end{aligned} \quad (50)$$

with $\hat{\chi}_a$ ($\hat{\chi}_a^\dagger$) representing a formal annihilation (creation) operator of the a th nucleus. We then extend our definition of the TD amplitudes as

$$\begin{aligned} \mathcal{F}_t^{000}(\{\mathbf{x}\}, \mathbf{R}) \equiv & \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) \\ & \times \check{\chi}_{A_1}(\mathbf{R}_1, t) \dots \check{\chi}_{A_{N_n}}(\mathbf{R}_{N_n}, t) | \Psi_t^{\text{int}} \rangle. \end{aligned} \quad (51)$$

A tricky aspect in the present procedure is that our definition of the TD amplitudes requires the zeroth order Hamiltonian in order to specify the interaction representation. We therefore first assume a fictitious potential function $\mathcal{U}(\mathbf{r}, t)$ which mimics the true electron-nucleus Coulombic interaction potential $\hat{U}_{\text{nuc}}(\mathbf{r})$ defined as an operator,

$$\hat{U}_{\text{nuc}}(\mathbf{r}) = \sum_a \int d^3\mathbf{R} \frac{q_e Z_a |e|}{|\mathbf{r} - \mathbf{R}|} \hat{\chi}_a^\dagger(\mathbf{R}) \hat{\chi}_a(\mathbf{R}). \quad (52)$$

Assuming such a space-time function $\mathcal{U}(\mathbf{r}, t)$, the “mean-field” Hamiltonian is redefined as

$$h_{\text{mf}}^{\text{el}}[\mathcal{U}] = c\alpha \cdot \frac{\hbar}{i} \nabla + \beta mc^2 + \mathcal{U}(\mathbf{r}, t) + W_{\text{HF}}^{\text{loc}}(\mathbf{x}), \quad (53)$$

and we can set the zeroth order Hamiltonian \mathcal{H}_0 along with the correction term \mathcal{H}_{int} as

$$\mathcal{H}_0 = \mathcal{H}_{\text{nuc}} + H_{\text{mf}}^{\text{el}}[\mathcal{U}] + H_{\text{rad}}, \quad (54a)$$

$$\mathcal{H}_{\text{int}} = \mathcal{H}_{\text{el-nuc}} + H_{\text{int}}^{\text{el}}, \quad (54b)$$

with

$$H_{\text{mf}}^{\text{el}} = \int d^3\mathbf{r} \psi^\dagger(\mathbf{r}) h_{\text{mf}}^{\text{el}}[\mathcal{U}] \psi(\mathbf{r}) \quad (55)$$

and

$$\mathcal{H}'_{\text{el-nuc}} \equiv \int d^3\mathbf{r} \psi^\dagger(\mathbf{r}) \psi(\mathbf{r}) [\hat{U}_{\text{nuc}}(\mathbf{r}) - \mathcal{U}(\mathbf{r})], \quad (56)$$

where the latter one $\mathcal{H}'_{\text{el-nuc}}$ represents the difference between the fictitious field and the true electron-nucleus potential. Having thus defined the zeroth order Hamiltonian, the extended TD amplitude given by Eq. (51) is now well-defined. We can then derive an equation of motion for $\mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R})$ as

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) = & \langle 0 | [\check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) \check{\chi}_{A_1}(\mathbf{R}_1, t) \dots \check{\chi}_{A_{N_n}}(\mathbf{R}_{N_n}, t), \mathcal{H}_{\text{nuc}} + \mathcal{H}'_{\text{el-nuc}}] | \Psi_t^{\text{int}} \rangle \\ & + \langle 0 | [\check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t), H_{\text{mf}}[\mathcal{U}] + H_{\text{rad}}] \check{\chi}_{A_1}(\mathbf{R}_1, t) \dots \check{\chi}_{A_{N_n}}(\mathbf{R}_{N_n}, t) | \Psi_t^{\text{int}} \rangle \\ & + \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) \check{\chi}_{A_1}(\mathbf{R}_1, t) \dots \check{\chi}_{A_{N_n}}(\mathbf{R}_{N_n}, t) H_{\text{int}}^{\text{el}} | \Psi_t^{\text{int}} \rangle \\ = & \left[\sum_a \frac{-\hbar^2}{2M_a} \nabla_a^2 + \frac{1}{2} \sum_{a \neq b} \frac{Z_a Z_b e^2}{|\mathbf{R}_a - \mathbf{R}_b|} \right] \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) + H_{\text{mf}}[\mathcal{U}] \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) \\ & + \sum_a \sum_{\mathbf{k}\lambda} \frac{Z_a |e| \hbar}{M_a c} \frac{\hbar}{i} \nabla_a \cdot \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{R}_a) \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) a_{\mathbf{k}\lambda} \check{\chi}_{A_1}(\mathbf{R}_1, t) \dots \check{\chi}_{A_{N_n}}(\mathbf{R}_{N_n}, t) | \Psi_t^{\text{int}} \rangle \\ & + \sum_a \sum_{\mathbf{k}\lambda} \frac{(Z_a |e|)^2}{2M_a c^2} \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) (\mathbf{A}^{\text{tr}})^2 \check{\chi}_{A_1}(\mathbf{R}_1, t) \dots \check{\chi}_{A_{N_n}}(\mathbf{R}_{N_n}, t) | \Psi_t^{\text{int}} \rangle \\ & + \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) \check{\chi}_{A_1}(\mathbf{R}_1, t) \dots \check{\chi}_{A_{N_n}}(\mathbf{R}_{N_n}, t) \int d^3\mathbf{r} \psi^\dagger(\mathbf{r}) \psi(\mathbf{r}) (\hat{U}_{\text{nuc}} - \mathcal{U}) | \Psi_t^{\text{int}} \rangle \\ & + \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) \check{\chi}_{A_1}(\mathbf{R}_1, t) \dots \check{\chi}_{A_{N_n}}(\mathbf{R}_{N_n}, t) H_{\text{int}}^{\text{el}} | \Psi_t^{\text{int}} \rangle. \end{aligned} \quad (57)$$

We find that commutation of the nuclear annihilation operators with \mathcal{H}_{nuc} formally yields additional couplings to the $(N_e, 0, 1)$ sector although such radiation-nucleus coupling terms in general should be much smaller in amplitude compared to its electronic counterpart because of large masses of the atomic nuclei and/or slowness of the nuclear motion. Below we only keep the first order coupling terms, which are proportional to $\hbar \nabla_a / i$, whereas we neglect the second order terms, which are proportional to $(\mathbf{A}^{\text{tr}})^2$. Terms arising from $H_{\text{int}}^{\text{el}}$ can be expanded in the same manner as we did in Subsection III A. The remaining terms are expressed in terms of the derivatives of the $(N_e, 0, 0)$ sector amplitude.

We therefore find that the amplitude $\mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R})$ behaves as a “first-quantized nuclear wavefunction” on which H_{nuc} operates as a differential operator. We therefore rewrite the definition of the TD amplitude as

$$\mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) = \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) | \Psi_t^{\text{int}}(\mathbf{R}) \rangle, \quad (58)$$

in which the electronic mean-field Hamiltonian is constructed with the use of the true electron-nucleus interaction, $H_{\text{mf}}(\mathbf{R}) = H_{\text{mf}}[\hat{U}_{\text{nuc}}]$ (with the operator \hat{U}_{nuc} rewritten in the first-quantized operator form), so that we no longer need the fictitious field \mathcal{U} . The above definition can be naturally extended to other sectors as

$$\mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda; \mathbf{R}) = \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) a_{\mathbf{k}\lambda}(t) | \Psi_t^{\text{int}}(\mathbf{R}) \rangle \quad (59)$$

and

$$\mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{R}) = \langle 0 | \check{\psi}(\mathbf{x}_1, t) \dots \check{\psi}(\mathbf{x}_{N_e}, t) a_{\mathbf{k}\lambda}(t) (\check{\psi}_{<}(\mathbf{y}_h, t))_{i_h} (\check{\psi}_{>}(\mathbf{y}_p, t))_{j_p} | \Psi_t^{\text{int}}(\mathbf{R}) \rangle, \quad (60)$$

and so on. The equation of motion for $\mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R})$ reads

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) = & \left[\sum_j \hat{h}^{\text{mf}}(\mathbf{R})_j + \hat{V}_C - \sum_j W_{\text{HF};j}^{\text{loc}}(\mathbf{R}) \right] \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) + \sum_a \frac{1}{2M_a} \left(\frac{\hbar}{i} \nabla_a - i\hbar \mathbf{X}_a \right)^2 \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) + \mathcal{K} \\ & + \frac{1}{2} \sum_{a \neq b} \frac{Z_a Z_b |e|^2}{|\mathbf{R}_a - \mathbf{R}_b|} \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) + \sum_a \sum_{\mathbf{k}\lambda} \frac{Z_a |e|}{M_a c} \left[\frac{\hbar}{i} \nabla_a - i\hbar \mathbf{X}^{(a)} \right] \cdot \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{R}_a) \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda; \mathbf{R}) \\ & + \sum_{\mathbf{k}\lambda} \left(\sum_j (\Lambda^+ \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{x}_j))_{(j)} + \mathcal{J}_{-\mathbf{k}}^{\mu} \boldsymbol{\varepsilon}_{\mathbf{k}\lambda\mu} \right) \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda; \mathbf{R}) + \sum_{\mathbf{k}\lambda} \sum_{i_h j_p} \int d^3 \mathbf{y} (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{y}))_{i_h j_p} \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}; i_h, j_p; \mathbf{k}\lambda; \mathbf{R}). \quad (61) \end{aligned}$$

Since we have switched from the fictitious \mathcal{U} to the true \hat{U}_{nuc} , the electronic wavefunctions depend explicitly on the nuclear coordinate \mathbf{R} . We therefore have included the derivative coupling operator \mathbf{X}_a , which acts on electronic variables in the manner $(\mathbf{X}_a)_{ij} = \langle \varphi_i | \partial / \partial \mathbf{R}_a | \varphi_j \rangle$ and possible mass-polarization terms, \mathcal{K} . The latter one \mathcal{K} represents a set of coordinate-frame dependent differential operators arising from the coupling to the nuclear dynamics; in the center of the mass of the nuclei (CMN) frame, for example, there should be mass-polarization terms as we discuss in [Appendix D](#). For simplicity, however, we here use the Cartesian coordinates in the laboratory frame, where \mathcal{K} vanishes.

We next consider the equation of motion of $\mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda; \mathbf{R})$, which reads

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda; \mathbf{R}) = & H_0^{\text{el,rad}}(\mathbf{R}) \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda; \mathbf{R}) + \left[\sum_a \frac{1}{2M_a} \left(\frac{\hbar}{i} \nabla_a - i\hbar \mathbf{X}_a \right)^2 + \frac{1}{2} \sum_{a \neq b} \frac{Z_a Z_b e^2}{|\mathbf{R}_a - \mathbf{R}_b|} \right] \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda; \mathbf{R}) \\ & + \sum_a \sum_{\mathbf{k}\lambda} \frac{Z_a |e|}{M_a c} \left[\frac{\hbar}{i} \nabla_a - i\hbar \mathbf{X}^{(a)} \right] \cdot \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*(\mathbf{R}_a) \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) + \left(\sum_{\ell} (\Lambda^+ \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*(\mathbf{x}_{\ell}))_{(\ell)} + \mathcal{J}_{\mathbf{k}}^{\nu} \boldsymbol{\varepsilon}_{\mathbf{k}\lambda\nu} \right) \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) \\ & + (\text{terms arising from } H_{\text{int}}^{\text{el}'}) + (\text{terms that do not belong to the } (N_e, 0, 0) \text{ or } (N_e, 0, 1) \text{ sector}), \quad (62) \end{aligned}$$

and that for \mathcal{F}_t^{111} becomes

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{k}\lambda; \mathbf{R}) = & H_0^{\text{el,rad}}(\mathbf{R}) \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{k}\lambda; \mathbf{R}) \\ & + \left[\sum_a \frac{1}{2M_a} \left(\frac{\hbar}{i} \nabla_a - i\hbar \mathbf{X}_a \right)^2 + \frac{1}{2} \sum_{a \neq b} \frac{Z_a Z_b e^2}{|\mathbf{R}_a - \mathbf{R}_b|} \right] \mathcal{F}_t^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda; \mathbf{R}) \\ & + (\Lambda^+ \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*(\mathbf{y}_h) \Lambda^-)_{i_h j_p} \delta^3(\mathbf{y}_h - \mathbf{y}_p) \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) - \sum_{\ell} (\Lambda^+ \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*(\mathbf{x}_{\ell}) \Lambda^-)_{i_{\ell} j_h} \delta^3(\mathbf{x}_{\ell} - \mathbf{y}) \mathcal{F}_t^{000}(\{\mathbf{x}\}; \mathbf{R}) |_{i_{\ell} \rightarrow j_p} \\ & + (\text{terms that do not belong to the } (N_e, 0, 0) \text{ or } (N_e, 1, 1) \text{ sector}) + (\text{terms arising from } H_{\text{int}}^{\text{el}'}). \quad (63) \end{aligned}$$

In addition to those belonging to other sectors than $(N_e, 0, 0)$, $(N_e, 0, 1)$, or $(N_e + 1, 1, 1)$, we neglected all terms arising from nucleus-radiation couplings as these describe the emission of radiation by the nuclear charge current, which should be negligible due to slowness of the nuclear motion ($|\mathbf{R}_a| \ll c$). Switching to the MO representation and solving Eqs. (62) and (63), we obtain a closed equation of $(N_e, 0, 0)$ amplitude,

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} f_t^{000}(\mathbf{I}; \mathbf{R}) &= \sum_a \frac{1}{2M_a} \left(\frac{\hbar}{i} \nabla_a - i\hbar \mathbf{X}_a \right)_{I_j, r}^2 f_t^{000}(\mathbf{I}_j^r; \mathbf{R}) + \frac{1}{2} \sum_{a \neq b} \frac{Z_a Z_b |e|^2}{|\mathbf{R}_a - \mathbf{R}_b|} f_t^{000}(\mathbf{I}; \mathbf{R}) + \sum_j \left(\hat{h}_{\text{mf}}(\mathbf{R}) - W_{\text{HF}j}^{\text{loc}} \right)_{I_j, r} f_t^{000}(\mathbf{I}_j^r; \mathbf{R}) \\
 &+ \frac{1}{2} \sum_{j \neq \ell} \langle I_j I_\ell \| r s \rangle f_t^{000}(\mathbf{I}_j^{rs}; \mathbf{R}) + \sum_{j \neq \ell} \sum_{\mathbf{k}\lambda} \sum_{r,s} \int_{-\infty}^t dt' (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda})_{I_\ell s} e^{-iH_0(t-t')/\hbar} (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*)_{I_j r} f_{t'}^{000}(\mathbf{I}_j^{rs}; \mathbf{R}) \\
 &+ \sum_{j,s} \iint_{-\infty}^t d\tau'_1 d\tau'_2 \{ \delta(\tau'_1 - (t-0)) + \delta(\tau'_2 - (t-0)) \} e^{-iH_0(t-\tau'_2)/\hbar + i\epsilon_{I_j}^0(t-\tau'_2)/\hbar} e^{-i\epsilon_{I_j}^0(t-\tau'_1)/\hbar} \Sigma_{I_j s}^{\text{fin}}(\tau'_1, \tau'_2) e^{-i\epsilon_r^0(\tau'_2 - \tau'_1)/\hbar} f_{\tau'_2}^{000}(\mathbf{I}_j^s; \mathbf{R}) \\
 &+ \sum_{a,j} \sum_{\mathbf{k}\lambda} \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{R}_a) \cdot \frac{Z_a |e|}{M_a} \left[\frac{\hbar}{i} \nabla_a - i\hbar \mathbf{X}_a \right] \int_{-\infty}^t dt' e^{-iH_0(t-t')/\hbar} (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*)_{I_j r} f_{t'}^{000}(\mathbf{I}_j^r; \mathbf{R}) \\
 &+ \sum_{a,j} \sum_{\mathbf{k}\lambda} (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda})_{I_j r} \int_{-\infty}^t dt' e^{-iH_0(t-t')/\hbar} \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*(\mathbf{R}_b) \cdot \frac{Z_b |e|}{M_b} \left[\frac{\hbar}{i} \nabla_b - i\hbar \mathbf{X}_b \right] f_{t'}^{000}(\mathbf{I}_j^r; \mathbf{R}), \tag{64}
 \end{aligned}$$

where we neglected terms that contain nucleus-radiation coupling to the second order. Equation (64) is to be compared to the previous result [Eq. (45)], where the nuclear dynamics is absent. In addition to the nuclear kinematic energy term, which is in the first term in Eq. (64), we have additional terms that arise from electron-nucleus interaction by the transversal photon exchange. Although we admit those terms should be small in general by the factor $|\dot{\mathbf{R}}|/c$, it gives the lowest order relativistic correction to the electron-nucleus interaction due to relativistic electronic motion. We also note that nuclear derivative operators in Eq. (64) accompany the derivative couplings $i\hbar \mathbf{X}$. Since the amplitude is explicitly dependent on the nuclear coordinates, appearance of the derivative coupling terms is a manifestation of the “gauge invariance,” or covariance of the expression with electronic basis set transformations.

Equation (64) represents the main conclusion of this subsection. We also derive the corresponding equation in an external field in Appendix E, where we show that the external field should formally be put into H_0 so that one can treat it in a nonperturbative manner. Our formulation thus defines a Schrödinger-like equation of motion in the equal-time representation (i.e., all coordinates sharing the same timelike coordinate) of a many-body system; hence, the space-time frame is the same as that of the conventional non-relativistic many-electron theory. Also taking account of the fact we are assuming the nuclear motion as nonrelativistic, we expect that there should be no fundamental difficulty in integration of our equation of motion, Eq. (64) using one of existing calculation techniques. Yet, we propose in Appendix F a wavepacket implementation of the nuclear dynamics in an attempt to seek for a best suited numerical algorithm for relativistic electron-nucleus coupled dynamics.

For clarity of later discussion, we summarize the assumptions we made in this formulation. First, we assumed (i) nonrelativistic nuclear motion. The assumption is essential not only for describing the nuclear kinematic term in the Schrödinger-like form in Eq. (64) but also for treating nucleus-radiation coupling as a weak perturbation. For laser-induced dynamics, this assumption puts a formal upper limit of the laser intensity around $\sim 10^{22}$ W/cm², assuming $\lambda = 800$ nm and nuclear mass of order $10^3 m_e$.

Second, we truncated the higher order terms in the Tamm-Dancoff expansion beyond the lowest order, which in turn implicitly assumes (ii) irrelevance of the higher order electron-radiation coupled effects. Such assumption should be valid in models where

smallness of electron-radiation coupling suppresses the higher order effects, but it would break down in unconventional types of dynamics such as those discussed in Refs. 17–19, where higher order QED effects play vital roles. This assumption (ii), in practice, may therefore impose much lower limit than (i) does to the external field intensity acceptable in our formulation. Relevance of higher order effects is, however, not clear just from an order estimate. A desirable strategy is therefore to start from a parameter range $\sim 10^{16}$ W/cm² to $\sim 10^{18}$ W/cm² and critically examine the validity of the lowest order treatment.

We, however, note that the absence of higher order effects in assumption (ii) refers to the “dynamical” electron-radiation coupling such as those arising from physical processes represented by more complex Feynman diagrams. It should not be confused with coupling with “external” radiation fields such as ultrastrong laser fields, which can be included in H_0 and solved by the means of the time-dependent Schrödinger-like equation [Eq. (64), also see Appendix E].

We have also mentioned about the gauge invariance, which we used to simplify the expression of the photon propagator, Coulomb plus transversal photon to the Feynman gauge propagator [see, for example, Eqs. (37) and (38) and related discussions]. We do not consider it as an essential assumption since numerical studies indeed show gauge independence of the final results,^{80,81} let alone difference in the contribution from each individual diagram, and there is also an established technique to directly evaluate the Coulomb gauge propagator.^{80,81}

Another feature of our present approach is that our working equation [Eq. (64)] is a direct extension of the conventional time-dependent Schrödinger equation (TDSE) approach in the sense it reduces to TDSE in the limit of vanishing dynamical electron-radiation coupling. We therefore expect that we can apply it to those models where the conventional TDSE approach provide at least qualitatively reasonable results in order to obtain an insight on how the relativistic effects affect the dynamics.

IV. SUMMARY AND DISCUSSIONS

In this report, we have developed a nonperturbative approach for relativistic electron-nucleus coupled dynamics. We consider that this approach has potential applications to the electron dynamics induced by ultraintense IR laser fields.

Our approach essentially reduces to a set of effective Schrödinger-like equations of equal-time amplitudes of the system; hence, it can be combined with nonperturbative calculation techniques including grid-type time-dependent wavepacket approaches. Possible limitations yet encountered in this approach are rooted to the problems inherent to the Tamm-Dancoff formulation.^{67,70} Nonetheless, the present theory has resolved difficulties faced in application of QED to a marked extent.

The formulation presented here only uses the lowest order expansion, which reproduces two of the essential features in the quantum electrodynamics, retarded and/or energy-dependent interactions among electrons and self-energies arising from the self-interaction of electrons. However, we need higher order expansions to include other types of QED effects including vertex corrections and multiple scatterings. Although we included a sketch of such a formulation in Appendix C, full formulation is to be discussed in another paper.

As we discussed in our closely related publication,¹² the performance in practical calculations should depend strongly on the quality of electronic wavefunctions to be used, which are supposed to take into account an electron-electron correlation. Compared to the path-integral formulation,¹² the present formulation, which derives a Schrödinger-like equation for a many-body wavefunction in the equal-time representation, has better similarity and affinity to the conventional Hamiltonian formalism of nonrelativistic theory, which might work favorably. We therefore can rest on many computational tools⁴⁰ and techniques^{41,42} well developed in quantum chemistry.

Another theoretical interest is to make a close comparison between our obtained equation [Eq. (45)] in the static limit and existing static QED theory such as the eQED approach.^{23,24} We can indeed find rough correspondence to their theory by performing the Fourier transformation to those integral kernels, as we did in Eq. (49). Nevertheless, partly because it is not very straightforward to define a strict static limit of Eq. (45), a detailed comparison is beyond our current scope and left for future study.

Notation

- We use the sign convention (1, -1, -1, -1). Symbol $\eta_{\mu\nu}$ represents the four-dimensional diagonal metric tensor unless specified otherwise; $\eta_{00} = 1$, $\eta_{11} = \eta_{22} = \eta_{33} = -1$. General four-dimensional indices are denoted by Greek letters μ , ν , etc., but Latin letters i , j , k , etc., are also used when we emphasize that they are spacelike components.
- ∂_μ describes the covariant derivatives; $\partial_\mu \doteq (\partial/(c\partial t), \partial/\partial \mathbf{x})$, whereas vectors such as A_μ represents covariant vectors $A_\mu \doteq (A_0, -\mathbf{A})$.
- q_e is the electronic charge, which takes a negative value; $q_e = -|e|$.
- Subscripts i and j attached to Fermionic operators and matrices represent the spinor indices.
- We use the Gauss unit for electromagnetic field; the Coulomb interaction between two charges Q_1 and Q_2 separated by distance r becomes $Q_1 Q_2 / r$, and the fine-structure constant is $e^2 / \hbar c \approx 1/137$.
- We exclusively use spinor matrices $\alpha^\mu \doteq (1, \boldsymbol{\alpha})$ and $\beta = \gamma^0$ instead of γ^μ . Slashed symbols, such as \varkappa , are also defined

as contraction of four-vector and alpha-matrices; $\varkappa \equiv \alpha^\mu \varepsilon_\mu$ [see Eq. (16)].

- Symbol \gtrless attached as a subscript or superscript of an electronic annihilation or creation operator specifies the energy sector; for example, $\psi_>$ and $\psi_<$ represent a positive-energy electron annihilation operator and a negative-energy electron (positron) creation operator, respectively.
- Symbol \mathcal{T} represents the time-ordering operator unless specified otherwise.
- Symbol \mathbf{R} represents the nuclear coordinate, which is a $3N_n$ dimensional vector. The mass and charge of the a th nucleus are denoted by M_a and $Z_a |e|$, respectively.
- Symbol $a_{\mathbf{k}\lambda}$ represents the annihilation operator of a (transversal) photon with the wavevector \mathbf{k} and polarization λ . The associated normalized polarization vector is denoted by $\mathbf{e}_{\mathbf{k}\lambda}^\mu$.

ACKNOWLEDGMENTS

The work has been supported by JSPS KAKENHI, Grant No. JP15H05752.

APPENDIX A: DETAILS OF THE TAMM-DANCOFF EXPANSION

We here describe some details of the Tamm-Dancoff expansion. As a simple example, we show the derivation of Eq. (15), the time evolution equation of an $(N_e, 0, 0)$ sector amplitude.

We first recall the definition of \mathcal{F}_t^{000} in Eq. (11). Since each operator is in the interaction representation, $\tilde{\psi}(\mathbf{x}, t) = e^{\frac{i}{\hbar} H_0 t} \psi(\mathbf{x}) e^{-\frac{i}{\hbar} H_0 t}$, its time-derivative yields commutator with H_0 , as shown in the first term in the RHS of Eq. (12). It then contributes to the mean-field single-particle Dirac Hamiltonian term $H_0 \mathcal{F}_t^{000}(\{\mathbf{x}\})$ in the RHS of Eq. (15). On the other hand, the state vector appearing in the definition of \mathcal{F}_t^{000} [Eq. (11)] is in the interaction representation. Hence, its time derivative yields $i\hbar \frac{\partial}{\partial t} |\Psi_t^{\text{int}}\rangle = H_{\text{int}}(t) |\Psi_t^{\text{int}}\rangle$, which appears in the second term in the RHS of Eq. (12). All Fermionic and Bosonic creation operators appearing in $H_{\text{int}}^{\text{el}}(t)$ are then moved leftward until they come next to the bra-vector $\langle 0|$, where they eliminate the vacuum state. During this operation, the Fermionic (Bosonic) anti-commutation (commutation) rules between equal-time operators apply.

Following the main text, we consider the transversal part, $H_{\text{int}}^{\text{el tr}}$, in the interaction Hamiltonian. We see that $H_{\text{int}}^{\text{el tr}}$ is decomposed as

$$H_{\text{int}}^{\text{el tr}}(t) = \int d^3\xi \left(\psi_>^\dagger(\xi, t) + \psi_<^\dagger(\xi, t) \right) \sum_{\mathbf{k}\lambda} \left(\varkappa_{\mathbf{k}\lambda}(\xi) \hat{a}_{\mathbf{k}\lambda}(t) + \varkappa_{\mathbf{k}\lambda}^*(\xi) \hat{a}_{\mathbf{k}\lambda}^\dagger(t) \right) \left(\psi_>(\xi, t) + \psi_<(\xi, t) \right). \quad (\text{A1})$$

It then follows that, in the equation of \mathcal{F}^{000} , all the “photon creation” operators trivially move to the left end and vanish, whereas all the photon annihilation operators survive. We next see that, in the electronic operators, $\psi_>^\dagger(\xi, t)$ and $\psi_<(\xi, t)$ contain the creation operators. We see that the positron creation operator $\psi_<(\xi, t)$ moves to the left end and vanishes, whereas the electron creation operator $\psi_>^\dagger(\xi, t)$ anticommutes with $\tilde{\psi}(\mathbf{x}_j, t)$ to yield $\Lambda_+ \delta^3(\mathbf{x}_j - \xi)$; hence, there appears $\varkappa_{\mathbf{k}\lambda}(\mathbf{x}_j) \psi_>(\mathbf{x}_j, t)$ in the position of the j th operator, yielding a vector in the 001 sector.

We also see that the remaining set of operators $\int d^3\xi \psi_{\pm}^{\dagger}(\xi, t) \alpha^{\mu} \psi_{\pm}(\xi, t) \sum_{\mathbf{k}\lambda} \varkappa_{\mathbf{k}\lambda}(\xi) \hat{a}_{\mathbf{k}\lambda}(t)$ does not vanish and yields state vectors in the 111 sector.

APPENDIX B: SPIN-ORBIT AND SPIN-SPIN COUPLING IN THE NONRELATIVISTIC LIMIT

Although it is not our major task, the low-energy effective theory with nonrelativistic two-component approximation gives by-products which are certainly useful in many applications.

As is suggested in the textbook of Bethe and Salpeter,⁸⁴ an effective Hamiltonian for interacting particles can be derived from the Tamm-Dancoff expansion. First we note that, in contrast to the remark made in Ref. 84 that (standard) TDA only reproduces the Coulombic term, our Tamm-Dancoff-like expansion does indeed recover those equations that include transversal interaction terms.

Following Ref. 84, we study an N_e -particle system in an external vector field potential $V^{\text{ext}} = \sum_j v_{\mu}^{\text{ext}}(\alpha^{\mu})_{(j)}$,

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{\mathbf{x}_i\}) &= [H_0 + V^{\text{ext}} + V_C] \mathcal{F}_t^{000}(\{\mathbf{x}_i\}) + \sum_i \int \prod_m d^3\mathbf{x}'_m \int d\tau \Sigma(\mathbf{x}_i, t; \mathbf{x}'_i, \tau) \prod_{j \neq i} K_{>}^0(\mathbf{x}_j, t; \mathbf{x}'_j, \tau) \mathcal{F}_{\tau}^{000}(\{\mathbf{x}'_k\}) \\ &+ \frac{c}{2} \int_{-\infty}^t d\tau \sum_{i \neq j} \int \prod_m d^3\mathbf{x}'_m \int \frac{d^4k}{(2\pi)^4} e^{-ik(x_i - x'_j)} \frac{4\pi e^2}{k^2 + i\eta} \left\{ \delta_{\ell, m} - \frac{k_{\ell} k_m}{\mathbf{k}^2} \right\} \\ &\times \left(\alpha^{\ell} K_{>}^0(\mathbf{x}_i, t; \mathbf{x}'_i, \tau) \right)_{(i)} \left(K_{>}^0(\mathbf{x}_j, t; \mathbf{x}'_j, \tau) \alpha^m \right)_{(j)} \prod_{k \neq i, j} K_{>}^0(\mathbf{x}_k, t; \mathbf{x}'_k, \tau) \mathcal{F}_{\tau}^{000}(\{\mathbf{x}'\}), \end{aligned} \quad (\text{B1})$$

which is a slight modification of our real-space equation (47), in which we use, following Ref. 84 Sec. 39, the free-particle Hamiltonian as the zeroth order one; $H_0 = \sum_j h_j^0 + H_{\text{rad}}$ with $h_j^0 = -i\hbar \boldsymbol{\alpha} \cdot \nabla_j + \beta m c^2$. The corresponding positive-energy propagator is denoted by $K_{>}^0$, which reads

$$\begin{aligned} K_{>}^0 &= \theta(t - t') \langle \mathbf{x} | \Lambda^+ e^{-\frac{i}{\hbar} \hat{h}_0(t-t')} \Lambda^+ | \mathbf{x}' \rangle \\ &= \theta(t - t') \sum_{\sigma} \int \frac{d^3\mathbf{p}}{(2\pi\hbar)^3} u_{\mathbf{p}\sigma} e^{i\mathbf{p}(x-x')/\hbar} e^{-i\varepsilon_{\mathbf{p}}(t-t')/\hbar} u_{\mathbf{p}\sigma}^{\dagger}, \end{aligned} \quad (\text{B2})$$

with $u_{\mathbf{p}\sigma} e^{i\mathbf{p}\mathbf{x}/\hbar}$ being the positive energy solution of h^0 with spin-projection σ and $\varepsilon_{\mathbf{p}}$ being its energy, $\varepsilon_{\mathbf{p}} \equiv \sqrt{c^2\mathbf{p}^2 + m^2c^4}$. Assuming the existence of its stationary solutions, we perform Fourier transformation of Eq. (B1) to obtain

$$\begin{aligned} \hbar\omega \mathcal{F}_{\omega}^{000}(\{\mathbf{p}\}) &= \sum_i \varepsilon_{\mathbf{p}_i} \mathcal{F}_{\omega}^{000}(\{\mathbf{p}\}) + \sum_i \left(\Sigma(\mathbf{p}_i, \omega_i^{(1)}) \right)_{(i)} \mathcal{F}_{\omega}^{000}(\{\mathbf{p}\}) \\ &+ \sum_i \int \frac{d^3\mathbf{k}}{(2\pi)^3} v_{\mu}^{\text{ext}}(\mathbf{k}) (\alpha^{\mu})_{(j)} \mathcal{F}_{\omega}^{000}(\mathbf{p}_i - \overset{i}{\underset{j}{\mathbf{k}}}) \\ &+ \frac{1}{2} \sum_{i \neq j} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{4\pi e^2}{\mathbf{k}^2} \mathcal{F}_{\omega}^{000}(\mathbf{p}_i - \overset{i}{\underset{j}{\mathbf{k}}}, \mathbf{p}_j + \overset{j}{\underset{i}{\mathbf{k}}}) \\ &- \frac{1}{2} \sum_{i \neq j} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{4\pi e^2}{\mathbf{k}^2 - (\omega_i^{(2)}(\mathbf{k})/c)^2} \left\{ \delta_{\ell, m} - \frac{k_{\ell} k_m}{\mathbf{k}^2} \right\} \\ &\times \left(\alpha^{\ell} \right)_{(i)} \left(\alpha^m \right)_{(j)} \mathcal{F}_{\omega}^{000}(\mathbf{p}_i - \overset{i}{\underset{j}{\mathbf{k}}}, \mathbf{p}_j + \overset{j}{\underset{i}{\mathbf{k}}}), \end{aligned} \quad (\text{B3})$$

where we introduced frequencies $\omega_i^{(1)} \equiv (\omega - \sum_k \varepsilon_{\mathbf{p}_k}) + \varepsilon_{\mathbf{p}_i}$ and $\omega_i^{(2)}(\mathbf{k}) \equiv (\omega - \sum_k \varepsilon_{\mathbf{p}_k}) + \varepsilon_{\mathbf{p}_i} - \varepsilon_{\mathbf{p}_i - \mathbf{k}}$. Equation (B3) can be seen as an implicit eigenvalue problem, where one solves an eigenvalue ω by ω -dependent matrix in the RHS. We also see that by neglecting the frequency dependent term $(\omega_i^{(2)}(\mathbf{k})/c)^2$, the transversal photon contribution reduces to

$$\begin{aligned} &-\frac{1}{2} \sum_{i \neq j} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{4\pi e^2}{\mathbf{k}^2} \left\{ \delta_{\ell, m} - \frac{k_{\ell} k_m}{\mathbf{k}^2} \right\} (\alpha^{\ell})_{(i)} (\alpha^m)_{(j)} \\ &\times \mathcal{F}_{\omega}^{000}(\mathbf{p}_i - \overset{i}{\underset{j}{\mathbf{k}}}, \mathbf{p}_j + \overset{j}{\underset{i}{\mathbf{k}}}), \end{aligned} \quad (\text{B4})$$

which is equivalent to the momentum representation of the Breit term in Ref. 84 Eq. (38.10). Equation (B3) therefore provides a small extension of Eq. (38.10) in Ref. 84, with an additional self-energy terms and frequency dependency in the transversal photon-exchange interactions.

For low-energy dynamics $|E - mc^2| \ll m_e c^2$, we can apply the same technique as Ref. 84 to reduce it to $2N$ -component representation. Even under such an assumption, however, the frequency dependence of the transversal interaction is not necessarily negligible if the frequency-dependent factor $\omega_i^{(2)}(\mathbf{k})/c$ is not negligible in comparison with a typical value of \mathbf{k} , which is the inverse of the typical length scale of the electronic wavefunction. Assuming inner core electron, such an inverse length scale is of order $\sim (a_B/Z)^{-1}$, with a_B being the Bohr radius and Z being the (effective) atomic number. This implies the frequency condition $\hbar\omega_i^{(2)}(\mathbf{k}) \sim Z\hbar c/(a_B)$, which is usually much smaller than the ‘‘relativistic’’ energy scale $\hbar\omega_i^{(2)}(\mathbf{k}) \sim mc^2$.

As for \mathcal{F}_{ω} , we apply, following Ref. 84, general $4N_e$ -component spinor transformation to $\mathcal{F}_{\omega}^{000}$ as

$$\mathcal{F}_\omega^{000} = \sum_{a_1} \sum_{a_2} \cdots \sum_{C_1} \sum_{C_2} \cdots \left(\prod_j \Xi_{C_j}^{i_j, a_j} \right) \theta_{a_1, a_2, \dots, a_{N_e}}^{C_1, C_2, \dots, C_{N_e}}, \quad (\text{B5})$$

where the indices of type a_j take 0 or 1, whereas those of type C_j take + or -. Symbol $\theta_{a_1, a_2, \dots, a_{N_e}}^{C_1, C_2, \dots, C_{N_e}}$ represents a $2N_e$ -component spinor, which is labeled by an index C_1, C_2, \dots, C_{N_e} and has 2^{N_e} -dimensional spinor index a_1, a_2, \dots, a_{N_e} . We also recall that, in our convention, i_j represents the j th (four-component) spinor index of the amplitude \mathcal{F}_ω^{000} . Four-by-two spinor matrices Ξ_C are given by

$$\Xi_+(\mathbf{p}) \doteq \begin{pmatrix} 1_2 \\ \mathbf{c}\mathbf{p} \cdot \boldsymbol{\sigma} / (mc^2 + \varepsilon_{\mathbf{p}}) \end{pmatrix}, \quad (\text{B6a})$$

$$\Xi_-(\mathbf{p}) \doteq \begin{pmatrix} -\mathbf{c}\mathbf{p} \cdot \boldsymbol{\sigma} / (mc^2 + \varepsilon_{\mathbf{p}}) \\ 1_2 \end{pmatrix}, \quad (\text{B6b})$$

with 1_2 representing the two-dimensional unit matrix. Positive (negative) energy projection $\Lambda_{\mathbf{p}}^+$ ($\Lambda_{\mathbf{p}}^-$) is defined as

$$\Lambda_{\mathbf{p}}^\pm = \frac{\varepsilon_{\mathbf{p}} \pm (\mathbf{c}\mathbf{p} \cdot \boldsymbol{\alpha} + \beta mc^2)}{2\varepsilon_{\mathbf{p}}}, \quad (\text{B7})$$

with which we have

$$\Lambda_{\mathbf{p}}^C \alpha_{CC'}^\mu \Xi_{C'}(\mathbf{p}') = \Xi_C(\mathbf{p}) \alpha_{CC'}^\mu(\mathbf{p}, \mathbf{p}'), \quad (\text{B8})$$

where the explicit expression of $\alpha_{CC'}^\mu(\mathbf{p}, \mathbf{p}')$ are found in Ref. 84 Eq. (16.12) [$\alpha_{CC'}^0(\mathbf{p}, \mathbf{p}')$ in our notation corresponds to $I_{CC}(\mathbf{p}, \mathbf{p}')$ in Ref. 84]. Here, we concentrate on the lowest order expansion with respect to the $|\mathbf{p}|/m_e c$ term of $\alpha_{+,+}^\mu(\mathbf{p}, \mathbf{p}')$, which are, according to Eq. (16.14) in Ref. 84,

$$\alpha_{++}^0(\mathbf{p}, \mathbf{p} + \hbar\mathbf{k}) = 1 + \frac{(\boldsymbol{\sigma} \cdot \mathbf{p})(\boldsymbol{\sigma} \cdot \hbar\mathbf{k})}{(2mc)^2}, \quad (\text{B9a})$$

$$\alpha_{++}(\mathbf{p}, \mathbf{p} + \hbar\mathbf{k}) = \frac{2\mathbf{p} + \hbar\mathbf{k} + i\hbar\mathbf{k} \times \boldsymbol{\sigma}}{2mc}. \quad (\text{B9b})$$

Substitution of expansion Eq. (B5) into Eq. (B3) and application of positive-energy projection operator $\prod_j \Lambda_{\mathbf{p}_j}^+$ retaining only a single component $C_1, C_2, \dots, C_{N_e} = + \dots +$, we obtain

$$\begin{aligned} \prod_j \Lambda_{\mathbf{p}_j}^+ \hbar \Omega \Xi_{++++}(\{\mathbf{p}\}) \theta^{++++}(\{\mathbf{p}\}) &\approx \sum_i \varepsilon_{\mathbf{p}_i} \Xi_{++++}(\{\mathbf{p}\}) \theta^{++++}(\{\mathbf{p}\}) + \sum_i \int \frac{d^3\mathbf{k}}{(2\pi)^3} v_\mu^{\text{ext}}(\mathbf{k}) \Xi_{++++}(\mathbf{p}) (\alpha_{+,+}^\mu)_{(i)} \theta^{++++}(\{\mathbf{p}\}) \\ &+ \frac{1}{2} \sum_{i \neq j} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{4\pi e^2}{\mathbf{k}^2} \Xi_{++++}(\mathbf{p}) (\alpha_{++}^0(\mathbf{p}_i, \mathbf{p}_i - \hbar\mathbf{k}))_{(i)} (\alpha_{++}^0(\mathbf{p}_j, \mathbf{p}_j + \hbar\mathbf{k}))_{(j)} \theta^{++++}(\mathbf{p}_i - \hbar\mathbf{k}, \mathbf{p}_j + \hbar\mathbf{k}) \\ &- \frac{1}{2} \sum_{i \neq j} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{4\pi e^2}{\mathbf{k}^2 - (\omega_i^{(2)}(\mathbf{k})/c)^2} \Xi_{++++}(\mathbf{p}) \sum_{\ell, m} \left\{ \delta_{\ell, m} - \frac{k_\ell k_m}{\mathbf{k}^2} \right\} (\alpha_{++}^\ell(\mathbf{p}_i, \mathbf{p}_i - \hbar\mathbf{k}))_{(i)} \\ &\times (\alpha_{++}^m(\mathbf{p}_j, \mathbf{p}_j + \hbar\mathbf{k}))_{(j)} \theta^{++++}(\mathbf{p}_i - \hbar\mathbf{k}, \mathbf{p}_j + \hbar\mathbf{k}), \end{aligned} \quad (\text{B10})$$

where we have neglected the self-energy term whose contribution is negligible in nonrelativistic limit and summation over spinor indices is made implicit, that is, spinor products are to be understood as $\sum_{a_1} \cdots \sum_{a_{N_e}} \Xi_{++++}^{a_1, \dots, a_{N_e}} \theta_{a_1, \dots, a_{N_e}}^{++++}$. Equation (B10) only differs by the retarded interaction from Ref. 84.

We next expand the retarded interactions. Substitution of Eqs. (B9a) and (B9b) into the last term in the RHS of Eq. (B10) yields the following three terms:

$$V_{ij}^{(1)} = \frac{4\pi e^2}{\mathbf{k}^2 - (\omega_i^{(2)}(\mathbf{k})/c)^2} \frac{1}{(mc)^2} \left\{ \mathbf{p}_i \cdot \mathbf{p}_j - \frac{(\mathbf{p}_i \cdot \mathbf{k})(\mathbf{p}_j \cdot \mathbf{k})}{\mathbf{k}^2} \right\}, \quad (\text{B11a})$$

$$V_{ij}^{(2)} = \frac{4\pi e^2}{\mathbf{k}^2 - (\omega_i^{(2)}(\mathbf{k})/c)^2} \frac{i\hbar}{(mc)^2} \left\{ \mathbf{s}_j \cdot (\mathbf{p}_i \times \mathbf{k}) - \mathbf{s}_i \cdot (\mathbf{p}_j \times \mathbf{k}) \right\}, \quad (\text{B11b})$$

$$V_{ij}^{(3)} = \frac{4\pi e^2}{\mathbf{k}^2 - (\omega_i^{(2)}(\mathbf{k})/c)^2} \frac{\hbar^2}{(mc)^2} \left\{ \mathbf{k}^2 \mathbf{s}_i \cdot \mathbf{s}_j - (\mathbf{s}_i \cdot \mathbf{k})(\mathbf{s}_j \cdot \mathbf{k}) \right\}, \quad (\text{B11c})$$

which are then transformed into real-space representation as

$$V_{ij}^{(1)} = \frac{e^2}{r_{ij}(mc)^2} \left(\mathbf{p}_i \cdot \mathbf{p}_j \left\{ e^{iqr_{ij}} + \frac{1 - e^{iqr_{ij}}}{(qr_{ij})^2} + i \frac{e^{iqr_{ij}}}{qr_{ij}} \right\} - (\mathbf{p}_i \cdot \mathbf{n}_{ij})(\mathbf{p}_j \cdot \mathbf{n}_{ij}) \left\{ e^{iqr_{ij}} + 3i \frac{e^{iqr_{ij}}}{qr_{ij}} + 3 \frac{1 - e^{iqr_{ij}}}{(qr_{ij})^2} \right\} \right), \quad (\text{B12a})$$

$$V_{ij}^{(2)} = \frac{\hbar^2 e^2}{(mc)^2 r_{ij}^3} (\mathbf{s}_i \cdot (\mathbf{p}_j \times \mathbf{n}_{ij}) - \mathbf{s}_j \cdot (\mathbf{p}_i \times \mathbf{n}_{ij})) (1 - iqr_{ij}), \quad (\text{B12b})$$

$$V_{ij}^{(3)} = \frac{\hbar^2 e^2}{(mc)^2} \left((\mathbf{s}_i \cdot \mathbf{s}_j) \frac{8\pi}{3} \delta^3(\mathbf{r}_{ij}) - \left[(\mathbf{s}_i \cdot \mathbf{s}_j) (1 - iqr_{ij} + (qr_{ij})^2) \frac{e^{iqr_{ij}}}{r_{ij}^3} - 3(\mathbf{s}_i \cdot \mathbf{n}_{ij})(\mathbf{s}_j \cdot \mathbf{n}_{ij}) \left(\frac{e^{iqr_{ij}}}{r_{ij}^3} - iq \frac{e^{iqr_{ij}}}{r_{ij}^2} + \frac{q^2 e^{iqr_{ij}}}{3 r_{ij}} \right) \right] \right), \quad (\text{B12c})$$

where $q \equiv \omega_i^{(2)}(\mathbf{k})/c$, $\mathbf{r}_{ij} \equiv \mathbf{r}_i - \mathbf{r}_j$, and $\mathbf{n}_{ij} \equiv \mathbf{r}_{ij}/|\mathbf{r}_{ij}|$, and the primed bracket $[\dots]'$ indicates that the singularity of the expression in the square bracket at the origin has been removed.⁸⁴ We note that these expressions reduce to static expressions in the limit $\omega \rightarrow 0$ as

$$V_{ij}^{(1)} = \frac{e^2}{r_{ij}(mc)^2} (\mathbf{p}_i \cdot \mathbf{p}_j - (\mathbf{p}_i \cdot \mathbf{n}_{ij})(\mathbf{p}_j \cdot \mathbf{n}_{ij})), \quad (\text{B13a})$$

$$V_{ij}^{(2)} = \frac{e^2 \hbar^2}{(mc)^2 r_{ij}^3} (\mathbf{s}_i \cdot (\mathbf{p}_j \times \mathbf{n}_{ij}) - \mathbf{s}_j \cdot (\mathbf{p}_i \times \mathbf{n}_{ij})), \quad (\text{B13b})$$

$$V_{ij}^{(3)} = \frac{e^2 \hbar^2}{(mc)^2} \left((\mathbf{s}_i \cdot \mathbf{s}_j) \frac{8\pi}{3} \delta^3(\mathbf{r}_{ij}) - \left[(\mathbf{s}_i \cdot \mathbf{s}_j) \frac{1}{r_{ij}^3} - 3(\mathbf{s}_i \cdot \mathbf{n}_{ij})(\mathbf{s}_j \cdot \mathbf{n}_{ij}) \right] \right). \quad (\text{B13c})$$

A qualitative difference in our results [Eqs. (B12b) and (B12c)] in contrast to its well-known static counterpart [Eqs. (B13b) and (B13c)] is that the present expressions bear the longer-range terms (with respect to spatial separation r_{ij}) arising from higher order terms of qr_{ij} .

APPENDIX C: HIGHER ORDER EXPANSIONS IN THE ELECTRON-RADIATION COUPLED DYNAMICS

Here, we discuss higher order expansions in the electron-radiation coupled dynamics to be appended to the lowest order

expansion in Subsection III A. We find that a straightforward extension of our discussion made on the lowest order expansion in Subsection III A does not work to a larger number of sectors because of divergences arising from the summation over photon \mathbf{k} vectors. Since such \mathbf{k} vector summation arises from all loop structures in the diagrammatic representation, we first need to separate out these divergent summations. We restart from a formal perturbation expansion of the state vector $|\Psi_t^{\text{int}}\rangle$, which is assumed to correspond to a reference state (eigenstate of H_0) $|\Phi_0\rangle$ in the limit of far past $t \rightarrow -\infty$ as

$$\begin{aligned} |\Psi_t^{\text{int}}\rangle &= U_\eta(t, -\infty)|\Phi_0\rangle/\mathcal{N}_\eta \\ &= \mathcal{T} \exp\left[\frac{1}{i\hbar} \int_{-\infty}^t dt' H_{\text{int}}(t') e^{-\eta|t'|}\right] |\Phi_0\rangle / \mathcal{N}_\eta, \end{aligned} \quad (\text{C1})$$

where η is an infinitesimal positive constant, which is to be later taken the limit $\eta \rightarrow 0$, and the associated normalization factor is defined as $\mathcal{N}_\eta \equiv \sqrt{\langle \Phi_0 | U_\eta(t, -\infty) | \Phi_0 \rangle}$. We then find that the amplitude is written as

$$\begin{aligned} \mathcal{F}_t^{000}(\{x\}) &= \langle 0 | \check{\psi}(x_1) \dots \check{\psi}(x_{N_e}) \left[|\Phi_0\rangle + \frac{1}{i\hbar} \int_{-\infty}^t d\tau_0 H_{\text{int}}(\tau_0) |\Phi_0\rangle + \left(\frac{1}{i\hbar}\right)^2 \int_{-\infty}^t d\tau_0 \int_{-\infty}^{\tau_0} d\tau_1 H_{\text{int}}(\tau_0) H_{\text{int}}(\tau_1) |\Phi_0\rangle \right. \\ &\quad \left. + \left(\frac{1}{i\hbar}\right)^3 \int_{-\infty}^t d\tau_0 \int_{-\infty}^{\tau_0} d\tau_1 \int_{-\infty}^{\tau_1} d\tau_2 H_{\text{int}}(\tau_0) H_{\text{int}}(\tau_1) H_{\text{int}}(\tau_2) U_\eta(\tau_2, -\infty) |\Phi_0\rangle \right] \Big/ \mathcal{N}_\eta, \end{aligned} \quad (\text{C2})$$

whose time derivative leads to an integrodifferential equation,

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{x\}) &= H_0 \mathcal{F}_t^{000}(\{x\}) + \langle 0 | \check{\psi}(x_1) \dots \check{\psi}(x_{N_e}) \left[H_{\text{int}}^{\text{el}}(t) |\Phi_0\rangle + H_{\text{int}}^{\text{el}}(t) \frac{1}{i\hbar} \int_{-\infty}^t d\tau_1 H_{\text{int}}^{\text{el}}(\tau_1) |\Phi_0\rangle + H_{\text{int}}^{\text{el}}(t) \left(\frac{1}{i\hbar}\right)^2 \right. \\ &\quad \left. \times \int_{-\infty}^t d\tau_1 \int_{-\infty}^{\tau_1} d\tau_2 H_{\text{int}}^{\text{el}}(\tau_1) H_{\text{int}}^{\text{el}}(\tau_2) |\Psi_{\tau_2}^{\text{int}}\rangle \right] \Big/ \mathcal{N}_\eta. \end{aligned} \quad (\text{C3})$$

We herein concentrate on the formal theory for simplicity and assume that each integral can be extended to $t \rightarrow -\infty$ and that the state in the far past $t \rightarrow -\infty$ is a radiation-free reference state of N_e electrons with no photons or antiparticles. We also extend our definition of the amplitudes. Only in this section, we allow appearance of multiple electronic timelike coordinates in the amplitude,

$$\mathcal{F}_t^{000}(\{x\}) = \langle 0 | \check{\psi}(x_1) \dots \check{\psi}(x_N) | \Psi_t^{\text{int}} \rangle, \quad (\text{C4})$$

$$\mathcal{F}_t^{001}(\{x\}, \mathbf{k}\lambda) = \langle 0 | \check{\psi}(x_1) \dots \check{\psi}(x_N) a_{\mathbf{k}\lambda}(t) | \Psi_t^{\text{int}} \rangle, \quad (\text{C5})$$

$$\begin{aligned} \mathcal{F}_t^{111}(\{x\}; \mathbf{y}_h, \mathbf{y}_p, \mathbf{i}_h, \mathbf{i}_p; \mathbf{k}\lambda) &= \langle 0 | \check{\psi}(x_1) \dots \check{\psi}(x_N) \left(\check{\psi}_{<}^\dagger(\mathbf{y}_h, t) \right)_{\mathbf{i}_h} \\ &\quad \times \left(\check{\psi}_{>}(\mathbf{y}_p, t) \right)_{\mathbf{i}_p} a_{\mathbf{k}\lambda}(t) | \Psi_t^{\text{int}} \rangle, \end{aligned} \quad (\text{C6})$$

where we require that all timelike variables should not be smaller than t ; $x_j^0/c \geq t$. Those amplitudes can then be calculated by spatial convolution with function $C^>$,

$$\mathcal{F}_t^{nlm}(\{x\}, \dots) = \int \prod_k d^3 \mathbf{x}'_k \prod C^>(x_k; \mathbf{x}'_k, t) \mathcal{F}_t^{nlm}(\{\mathbf{x}\}, \dots), \quad (\text{C7})$$

with nlm being either of 000, 001, or 111, and $\mathcal{F}_t^{nlm}(\{\mathbf{x}\}, \dots)$ in the RHS being the corresponding amplitude with a single timelike variable t .

As we did in the main text, we first replace $H_{\text{int}}^{\text{el}}$ by $H_{\text{int}}^{\text{el tr}}$ and expand the interaction terms. We then find that, in Eq. (C3), the first order correction vanishes, whereas the second order survives. After restoring Coulombic contributions, the second order contribution reads

$$\begin{aligned}
 & \langle 0 | \check{\psi}(x_1) \dots \check{\psi}(x_{N_e}) H_{\text{int}}^{\text{el}}(t) \frac{1}{i\hbar} \int_{-\infty}^t d\tau_1 H_{\text{int}}^{\text{el}}(\tau_1) | \Phi_0 \rangle \\
 &= i\hbar \sum_{i \neq j} (\alpha^\mu)_{(i)} \int_{-\infty}^t d\tau \int d^3\xi (S^F(\mathbf{x}_j, t; \xi, \tau) \alpha^\nu)_{(j)} \mathcal{F}_\tau^{000(0)}(\mathbf{x}_1, t; \dots; \overset{j}{\xi}, \tau; \dots) D_{\mu\nu}(\mathbf{x}_i, t; \xi) \\
 &+ \sum_i \int_{-\infty}^t d\tau \int d^3\xi \Sigma_{1L}(\mathbf{x}_i, t; \xi, \tau) \mathcal{F}_\tau^{000(0)}(\mathbf{x}_1, t; \dots; \overset{i}{\xi}, \tau; \dots) \\
 &+ i\hbar \sum_i \int d^3\mathbf{y} \int d\tau \int d^3\xi \mathcal{S}^\mu(\mathbf{y}, t) D_{\mu\nu}(\mathbf{y}, t; \xi, \tau) (S^F(\mathbf{x}_i, t; \xi, \tau) \alpha^\nu)_{(i)} \mathcal{F}_\tau^{000(0)}(\mathbf{x}_1, t; \dots; \overset{i}{\xi}, \tau; \dots) \\
 &+ \sum_i (\alpha^\mu)_{(i)} \int d\tau \int d^3\xi \mathcal{S}^\nu(\xi, \tau) D_{\mu\nu}(\mathbf{x}_i, t; \xi, \tau) \mathcal{F}_\tau^{000(0)}(\mathbf{x}_1, t; \dots; \overset{i}{\xi}, \tau; \dots) \\
 &+ \int d^3\mathbf{y} \int d\tau \int d^3\xi \mathcal{S}^\mu(\mathbf{y}, t) D_{\mu\nu}(\mathbf{y}, t; \xi, \tau) \mathcal{S}^\nu(\xi, \tau) \mathcal{F}_\tau^{000(0)}(\mathbf{x}_1, t; \dots; \mathbf{x}_{N_e}, t) \\
 &- (i\hbar)^2 \int d^3\mathbf{y} \int d\tau \int d^3\xi \text{Tr}(\alpha^\mu S^F(\mathbf{y}, t; \xi, \tau) \alpha^\nu S^F(\xi, \tau; \mathbf{y}, t)) D_{\mu\nu}(\mathbf{y}, t; \xi, \tau), \tag{C8}
 \end{aligned}$$

with $\mathcal{F}_t^{000(0)}$ defined as

$$\mathcal{F}_t^{000(0)} \equiv \langle 0 | \check{\psi}(x_1) \dots \check{\psi}(x_{N_e}) | \Phi_0 \rangle.$$

The remaining part in Eq. (C3) is expanded as

$$\begin{aligned}
 & \langle 0 | \check{\psi}(x_1) \dots \check{\psi}(x_{N_e}) H_{\text{int}}^{\text{el}}(t) \left(\frac{1}{i\hbar} \right)^2 \int_{-\infty}^t d\tau_1 \int_{-\infty}^{\tau_1} d\tau_2 H_{\text{int}}^{\text{el}}(\tau_1) H_{\text{int}}^{\text{el}}(\tau_2) | \Psi_{\tau_2}^{\text{int}} \rangle \\
 &= \int \prod_j \frac{1}{c} d^4\xi_j \sum_{\mathfrak{D}} \sum_{\ell} \sum_{\mathbf{k}\lambda} \mathcal{V}_\ell^{000,001;\mathfrak{D}}(\{x\}; \{\xi\}; \mathbf{k}\lambda) \mathcal{F}_\ell^{001}(\{x\}; \{\xi\}; \mathbf{k}\lambda) + \int \prod_j \frac{1}{c} d^4\xi_j \sum_{\mathfrak{D}} \sum_{\ell} \sum_{\mathbf{k}\lambda} \int d^3\mathbf{y} \sum_{i_h, j_p} \mathcal{V}_\ell^{000,111;\mathfrak{D}}(\{x\}; \{\xi\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{k}\lambda) \\
 &\times \mathcal{F}_\ell^{111}(\{x\}; \{\xi\}; \mathbf{y}_h, \mathbf{y}_p; i_h, j_p; \mathbf{k}\lambda) + (\text{terms arising from } H_{\text{int}}^{\text{el}}) + (\text{terms that do not belong to sector } 000, 001 \text{ or } 111), \tag{C9}
 \end{aligned}$$

where $\mathcal{V}_\ell^{\mathcal{X};\mathcal{Y};\mathfrak{D}}(\{x\}; \{\xi\}; \mathbf{k}\lambda)$ represents a coefficient (a function of space-time variables) that couple sectors \mathcal{X} and \mathcal{Y} by a process represented by the time-ordered Feynman diagram \mathfrak{D} . The remaining symbols and their meanings depend on their diagrammatic representations such that ℓ represents a set of indices to be summed over, $\{\xi\}$ do a set of integration variables, and the notation $\{\xi\}; \{\xi\}$ indicates that a part of the coordinate set $\{x\} = x_1, \dots, x_{N_e}$ is replaced by

$\{\xi\} = \xi_1, \dots$. In the relevant calculations, we rearrange the Coulombic interaction as follows: all photon lines represent the summation of the Coulombic interaction and the transversal photon exchange, and all the lines representing interelectron interactions (excluding self-interactions) accompany mean-field subtraction term. Several examples of such diagrams are shown below, whose corresponding expressions are

$$\begin{aligned}
 & \int^{ct} \frac{1}{c} d^4\xi_1 \int^{\xi_2} \frac{1}{c} d^4\xi_2 \mathcal{V}_{ij}^{000,001;\mathfrak{D}_1}(\xi_1, \xi_2) \mathcal{F}_{\xi_2}^{001}(x_1, \dots, \overset{i}{\xi}_2, \dots, \overset{j}{\xi}_1, \dots) \\
 &= \int^{ct} \frac{1}{c} d^4\xi_1 \int^{\xi_2} \frac{1}{c} d^4\xi_2 \left[\sum_{\mathbf{k}'\lambda'} (\mathfrak{S}_{\mathbf{k}'\lambda'}(\mathbf{x}_i) S^F(x_i, \xi_2) \mathfrak{S}_{\mathbf{k}\lambda}(\mathbf{x}_i))_{(i)} (S^F(x_j, \xi_1) \mathfrak{S}_{\mathbf{k}'\lambda'}^*(\xi_1))_{(j)} + (S^F(x_i, \xi_2) \mathfrak{S}_{\mathbf{k}\lambda}(\mathbf{x}_i))_{(i)} \right. \\
 &\quad \left. \times c\delta^4(x_j - \xi_1) v_C'(\mathbf{x}_i, \mathbf{x}_j) \right] \mathcal{F}_{\xi_2}^{001}(x_1, \dots, \overset{i}{\xi}_2, \dots, \overset{j}{\xi}_1, \dots; \mathbf{k}\lambda), \tag{C10a}
 \end{aligned}$$

$$\int^{ct} \frac{1}{c} d^4 \xi_2 \mathcal{V}_i^{000,001;\mathcal{D}_2}(\xi_2) \mathcal{F}_{\xi_2}^{001}(x_1, \dots, \overset{i}{\xi_2}, \dots; \mathbf{k}\lambda) = \int^{ct} \frac{1}{c} d^4 \xi_2 \int_{\xi_2^0}^{ct} \frac{1}{c} d^4 \xi_1 \left(\sum_{\mathbf{k}'\lambda'} \mathfrak{s}_{\mathbf{k}'\lambda'}(\mathbf{x}_i) S^F(x_i, \xi_1) \mathfrak{s}_{\mathbf{k}'\lambda'}^*(\xi_1) S^F(\xi_1, \xi_2) \mathfrak{s}_{\mathbf{k}\lambda}(\xi_2) \right. \\ \left. + S^F(x_i, \xi_1) \delta(t - \frac{\xi_1^0}{c}) \frac{1}{|\mathbf{x}_i - \xi_1'} S^F(\xi_1, \xi_2) \mathfrak{s}_{\mathbf{k}\lambda}(\xi_2) \right) \mathcal{F}_{\xi_2}^{001}(x_1, \dots, \overset{i}{\xi_2}, \dots; \mathbf{k}\lambda), \quad (\text{C10b})$$

$$\int^{ct} \frac{1}{c} d^4 \xi_2 \mathcal{V}_i^{000,001;\mathcal{D}_3}(\xi_2) \mathcal{F}_{\xi_2}^{001}(x_1, \dots, x_{N_e}; \mathbf{k}\lambda) = \int^{ct} \frac{1}{c} d^4 \xi_2 \int_{\xi_2^0}^{ct} \frac{1}{c} d^4 \xi_1 \left[\sum_{\mathbf{k}'\lambda'} (\mathfrak{s}_{\mathbf{k}'\lambda'}(\mathbf{x}_i))_{(i)} \text{Tr}[-S^F(\xi_1, \xi_2) \mathfrak{s}_{\mathbf{k}\lambda}(\xi_2) S^F(\xi_2, \xi_1) \mathfrak{s}_{\mathbf{k}'\lambda'}^*(\xi_1)] \right. \\ \left. + \text{Tr}[-S^F(\xi_1, \xi_2) \mathfrak{s}_{\mathbf{k}\lambda}(\xi_2) S^F(\xi_2, \xi_1)] \delta(t - \frac{\xi_1^0}{c}) v'_C(\xi_1, \mathbf{x}_i) \right] \mathcal{F}_{\xi_2}^{001}(x_1, \dots, x_{N_e}; \mathbf{k}\lambda), \quad (\text{C10c})$$

and

$$\int^{ct} \frac{1}{c} d^4 \xi_2 \mathcal{V}_{ij}^{000,001;\mathcal{D}_4}(\xi_2) \mathcal{F}_{\xi_2}^{001}(x_1, \dots, \overset{i}{\xi_2}, \dots) = \int^{ct} \frac{1}{c} d^4 \xi_2 \int_{\xi_2^0}^{ct} \frac{1}{c} d^4 \xi_1 \left[\sum_{\mathbf{k}'\lambda'} (\mathfrak{s}_{\mathbf{k}'\lambda'}(\mathbf{x}_i) S^F(x_i, \xi_1) \mathfrak{s}_{\mathbf{k}\lambda}(\xi_1) S^F(\xi_1, \xi_2) \mathfrak{s}_{\mathbf{k}'\lambda'}^*(\xi_2))_{(i)} \right. \\ \left. + (S^F(x_i, \xi_1) \mathfrak{s}_{\mathbf{k}\lambda}(\xi_1) S^F(\xi_1, \xi_2))_{(i)} \delta(t - \frac{\xi_2^0}{c}) v'_C(\mathbf{x}_i, \xi_2) \right] \mathcal{F}_{\xi_2}^{001}(x_1, \dots, \overset{i}{\xi_2}, \dots), \quad (\text{C10d})$$

where the four-dimensional integral with upper limit is a shorthand notation defined as

$$\int^{ct} \frac{1}{c} d^4 \xi \equiv \frac{1}{c} \int_{-\infty}^{ct} d\xi^0 \int d^3 \xi,$$

whereas the symbol v'_C represents the Coulombic interaction minus the mean-field interaction,

$$v'_C(\mathbf{x}_1, \mathbf{x}_2) \equiv \frac{q_e^2}{|\mathbf{x}_1 - \mathbf{x}_2|} - \frac{W_{\text{HF}}^{\text{loc}}(\mathbf{x}_1) + W_{\text{HF}}^{\text{loc}}(\mathbf{x}_2)}{2}.$$

We see that Eqs. (C10b)–(C10d) are divergent. It is, however, clear that summation over all the different time-ordering with the same topology as Fig. 2(b) makes an expression that is obtained from Eq. (C10b) by reordering the time. We can then apply the standard renormalization procedure for the electronic self-energy Σ to remove formal divergences and obtain a finite result. Application of the same procedure to diagrams Figs. 2(c) and 2(d) [Eqs. (C10c) and (C10d)] should yield the polarization and vertex correction functions, respectively.

We then need expressions for the amplitudes of the $(N_e, 0, 1)$ and $(N_e + 1, 1, 1)$ sectors. Applying the simplest truncation scheme to Eqs. (22) and (23), to truncate sectors other than $(N_e, 0, 0)$, $(N_e, 0, 1)$, and $(N_e + 1, 1, 1)$, we obtain

$$i\hbar \frac{\partial}{\partial t} \mathcal{F}_i^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda) \approx H_0 \mathcal{F}_i^{001}(\{\mathbf{x}\}; \mathbf{k}\lambda) \\ + \left(\sum_{\ell} (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{x}_\ell))_{(\ell)} + \mathcal{J}^{\ell\ell} \mathfrak{e}_{\mathbf{k}\lambda}^* \right) \mathcal{F}^{000}(\{\mathbf{x}\}) \quad (\text{C11})$$

and

$$i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; j_h, j_p; \mathbf{k}\lambda) \\ = H_0 \mathcal{F}_t^{111}(\{\mathbf{x}\}; \mathbf{y}_h, \mathbf{y}_p; j_h, j_p; \mathbf{k}\lambda) + \langle 0 | \psi(\mathbf{x}_1, t) \dots \psi(\mathbf{x}_{N_e}, t) | \Psi_t^{\text{int}} \rangle \\ \times \delta^3(\mathbf{y}_h - \mathbf{y}_p) (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{y}_h) \Lambda^-)_{j_p j_h} - \sum_{\ell} (\Lambda^+ \mathfrak{s}_{\mathbf{k}\lambda}^*(\mathbf{x}_\ell) \Lambda^-)_{i_\ell j_h} \\ \times \delta^3(\mathbf{x}_\ell - \mathbf{y}_h) \mathcal{F}_t^{000}(\{\mathbf{x}\})|_{i_\ell \rightarrow j_p}. \quad (\text{C12})$$

Substitution back of the formal solutions of Eqs. (C11) and (C12) then leads to an effective equation of the $(N_e, 0, 0)$ sector such that

$$i\hbar \frac{\partial}{\partial t} \mathcal{F}_t^{000}(\{\mathbf{x}\}) = H_0 \mathcal{F}_t^{000}(\{\mathbf{x}\}) + \frac{1}{2} \sum_{\mathcal{D}, \mathcal{D}'} \sum_{\chi=001,111} \sum_{\ell, j} \sum_{\mathbf{k}\lambda} \int \frac{1}{c} d^4 \xi_1 \dots \\ \times \int \frac{1}{c} d^4 \xi_1' \dots \mathcal{V}_{\ell}^{000, \chi; \mathcal{D}}(\{\mathbf{x}\}; \xi_1, \dots; \mathbf{k}\lambda) \\ \times \exp\left[\frac{1}{i\hbar} H_{\text{mf}}(t - \xi_2^0)\right] \mathcal{V}_j^{\chi, 000; \mathcal{D}'}(\{\mathbf{x}\}; \xi_1, \dots; \mathbf{k}\lambda) \\ \times \mathcal{F}^{000}(\{\mathbf{x}\}; \xi_1, \dots). \quad (\text{C13})$$




FIG. 2. Examples of time-ordered Feynman diagrams that couple sectors $(N_e, 0, 0)$ and $(N_e, 0, 1)$. Time increases upward. Four panels (a)–(d) correspond to Eqs. (C10a)–(C10d), respectively. Upward (downward) arrows represent electrons (positrons) and wavy lines represent photons.

Here, an additional divergence arises from the summation over $\mathbf{k}\lambda$ if the corresponding photon line makes a loop. But it can be treated in the same manner as Eq. (48). Although Eq. (C13) only shows an outline of the procedure, it is clear that summation over all the diagrams arising from expansion of Eq. (C3) corresponds to the standard Feynman diagrams of the two loop expansion in the standard perturbation theory, in which all formal divergences can be removed. We therefore conceive that there should be no divergence problem although we have yet to calculate all the possible diagrams in MO representation carefully examining whether this expansion works.

APPENDIX D: MASS-POLARIZATION TERMS

Here, we discuss the mass-polarization terms \mathcal{H} we encounter in the derivation of Eq. (61). In the main text, we adopted the lab frame and simply set it zero. On the other hand, if we use the coordinate representation relative to the center of the mass of the nuclei (CMN), Nonzero mass polarization emerges. Hence, we need to consider a general transformation from the lab frame to the CMN frame to derive those terms. Below, the lab frame electronic coordinates are denoted by $\{\mathbf{r}_j\}$ ($j = 1, 2, \dots, N_e$) and the nuclear coordinates in the mass-weighted representation are denoted by $\{\mathbf{Q}_a\}$ ($a = 1, 2, \dots, N_n$), whereas in the CMN frame, electronic coordinates are denoted by $\{\tilde{\mathbf{r}}_j\}$ and the mass-weighted nuclear coordinates are denoted by $\{\Xi_\lambda\}$, among which the CMN coordinate is denoted by $\Xi_\omega \equiv \sum_a \sqrt{M_a} \mathbf{Q}_a / \sqrt{\sum_b M_b}$.

We then consider the following generalized transformation with an orthogonal matrix \mathcal{R} :

$$\tilde{\mathbf{r}}_j = \mathbf{r}_j - \frac{\Xi_\omega}{\sqrt{\sum_b M_b}}, \quad (\text{D1a})$$

$$\Xi_\lambda = \sum_{a'} \mathcal{R}_\lambda^{a'} \mathbf{Q}_{a'}. \quad (\text{D1b})$$

The associated differential operators then transform as

$$\frac{\partial}{\partial \mathbf{r}_j} = \frac{\partial}{\partial \tilde{\mathbf{r}}_j}, \quad (\text{D2a})$$

$$\frac{\partial}{\partial \mathbf{Q}_a} = \sum_{\lambda'} \mathcal{R}_{\lambda'}^a \frac{\partial}{\partial \Xi_{\lambda'}} - \frac{1}{\sqrt{\sum_b M_b}} \mathcal{R}_\omega^a \left(\sum_{j'} \frac{\partial}{\partial \mathbf{r}_{j'}} \right). \quad (\text{D2b})$$

Substitution of these results to the electronic Dirac equation makes no apparent change, whereas that to the kinematic part of the nuclear Schrödinger equation yields

$$\begin{aligned} & \frac{1}{2} \sum_a \left(\frac{\hbar}{i} \frac{\partial}{\partial \mathbf{Q}_a} - \frac{Z_a |e|}{c} \mathbf{A}(\mathbf{R}_a) \right)^2 \\ &= \frac{1}{2} \sum_\lambda \left(\frac{\hbar}{i} \frac{\partial}{\partial \Xi_\lambda} \right)^2 + \frac{1}{2} \sum_a \frac{1}{M_a} \left(\frac{Z_a |e|}{c} \mathbf{A}(\mathbf{R}_a) \right)^2 \\ &+ \frac{1}{2 \sum_b M_b} \left(\sum_j \frac{\hbar}{i} \frac{\partial}{\partial \tilde{\mathbf{r}}_j} \right)^2 - \frac{1}{\sqrt{\sum_b M_b}} \left(\sum_j \frac{\hbar}{i} \frac{\partial}{\partial \tilde{\mathbf{r}}_j} \right) \cdot \frac{\hbar}{i} \frac{\partial}{\partial \Xi_\omega} \\ &+ \sum_\lambda \frac{1}{2} \left\{ \frac{\hbar}{i} \frac{\partial}{\partial \mathbf{Q}_\lambda}, \left(\sum_a \frac{Z_a |e|}{\sqrt{M_a}} \mathbf{A}(\mathbf{R}_a) \mathcal{R}_\lambda^a \right) \right\}_+ \\ &+ \frac{\sum_a Z_a |e| \mathbf{A}(\mathbf{R}_a)}{c \sum_b M_b} \left(\sum_j \frac{\hbar}{i} \frac{\partial}{\partial \tilde{\mathbf{r}}_j} \right), \end{aligned} \quad (\text{D3})$$

with $\{A, B\}_+ \equiv AB + BA$ representing the anticommutator, and \mathbf{R}_a in the RHS are to be understood as a function of new coordinate vectors $\{\Xi_\lambda\}$ in the sense $\mathbf{R}_a = \sum_\lambda (\mathcal{R}^{-1})_{a\lambda} \Xi_\lambda$.

We can then find the mass-polarization term which couples among all the electronic momenta and also charge-polarization terms that add to the electron-radiation coupling term although both terms accompany small factor $m_e/(\sum M_b)$. We also see that the CMN mode formally couple with the electronic momentum operators reflecting the Galilean covariance although we can simply assume that the CMN is fixed at the origin in many of applications.

In dynamical models of molecules, however, the derivative couplings, which approximately scales as $\sqrt{m_e/M_\lambda}$ for the λ th ($\lambda \neq \omega$) nuclear displacement modes around the equilibrium configuration, can be even larger in magnitude than these mass-polarizations.

APPENDIX E: EXTERNAL FIELDS

We here consider chemical dynamics (CD) in an external vector field $\mathcal{A}^{\text{ext}}(\mathbf{r}, t)$. We keep discussions in the main text where we set that \mathbf{A}^{tr} in Eqs. (8a)–(8d) as “internal” or “dynamical” field while adding new terms

$$\begin{aligned} H_{\text{ext}} = & -q_e \int d^3 \mathbf{x} \psi^\dagger(\mathbf{x}) \alpha \psi(\mathbf{x}) \cdot \mathcal{A}^{\text{ext}}(\mathbf{x}, t) + \sum_a \frac{1}{2M_a} \left\{ -2 \frac{Z_a |e|}{c} \mathcal{A}^{\text{ext}}(\mathbf{R}, t) \cdot \left(\frac{\hbar}{i} \nabla_a - \frac{Z_a |e|}{c} \mathbf{A}^{\text{tr}} \right) + \left(\frac{Z_a |e|}{c} \mathcal{A}^{\text{ext}}(\mathbf{R}, t) \right)^2 \right\} \\ & + \frac{1}{4\pi} \int d^3 \mathbf{r} \left\{ -\frac{1}{c} \frac{\partial}{\partial t} \mathcal{A}^{\text{ext}}(\mathbf{r}, t) \cdot 4\pi c \mathbf{\Pi}^{\text{tr}} + (\nabla \times \mathcal{A}^{\text{ext}}) \cdot (\nabla \times \mathbf{A}^{\text{tr}}) \right\}. \end{aligned} \quad (\text{E1})$$

In principle, one can treat these terms in an analogous manner as what we did in Subsection III A. We then need to expand the electronic part in Eq. (E1), $H_{\text{int}}^{\text{el ext}} \equiv -q_e \int d^3 \mathbf{x} \psi^\dagger(\mathbf{x}) \alpha \psi(\mathbf{x}) \cdot \mathcal{A}^{\text{ext}}(\mathbf{x}, t)$, which introduces additional couplings between $(N_e, 0, 0)$ and $(N_e + 1, 1, 0)$ sectors, $(N_e, 0, 1)$ and $(N_e + 1, 1, 1)$ sectors, etc. It then appears, in the lowest order, we need to solve a set of coupled equations of the $(N_e, 0, 0)$, $(N_e, 1, 0)$, and $(N_e, 1, 1)$ sector amplitudes,

which is not a trivial task, since there appears potentially divergent summation over photon wavevectors.

A better insight is obtained by first solving the single-particle propagator in the external field and then by solving the radiation coupling. In such a scheme, one can directly reach the final conclusion just by replacing propagators in Eq. (64) as

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t'} f_t^{000}(\mathbf{I}; \mathbf{R}) &= \sum_a \frac{1}{2M_a} \left(\frac{\hbar}{i} \nabla_a - \frac{Z|e|}{c} \mathcal{A}^{\text{ext}}(\mathbf{R}) - i\hbar \mathbf{X}_a \right)_{I_j, r}^2 f_t^{000}(\mathbf{I}'_j; \mathbf{R}) \\
 &+ \frac{1}{2} \sum_{a \neq b} \frac{Z_a Z_b |e|^2}{|\mathbf{R}_a - \mathbf{R}_b|} f_t^{000}(\mathbf{I}; \mathbf{R}) + \sum_j \left(\hat{h}_{\text{mf}}^{(0)}(\mathbf{R}) - W_{\text{HF}j}^{\text{loc}} - \boldsymbol{\alpha} \cdot \mathcal{A}^{\text{ext}} \right)_{I_j, r} f_t^{000}(\mathbf{I}'_j; \mathbf{R}) \\
 &+ \frac{1}{2} \sum_{j \neq \ell} (I_j I_\ell \| r s) f_t^{000}(\mathbf{I}'_j{}^{\text{rs}}; \mathbf{R}) + \sum_{j, r} \int_{-\infty}^t dt' \Sigma_{\mathcal{A}^{\text{ext}}}^{\text{fin.}}(t, t')_{I_j, r} f_{t'}^{000}(\mathbf{I}'_j; \mathbf{R}) \\
 &+ \sum_{j \neq \ell} \sum_{\mathbf{k}\lambda} \sum_{r, s} \int_{-\infty}^t dt' (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda})_{I_\ell s} e^{-i\mathcal{H}_0^{\mathcal{A}^{\text{ext}}}(t-t')/\hbar} (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*)_{I_j r} f_{t'}^{000}(\mathbf{I}'_j{}^{\text{rs}}; \mathbf{R}) \\
 &+ \sum_{aj} \sum_{\mathbf{k}\lambda} \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}(\mathbf{R}_a) \cdot \frac{Z_a |e|}{M_a} \left[\frac{\hbar}{i} \nabla_a - \frac{Z_a |e|}{c} \mathcal{A}^{\text{ext}}(\mathbf{R}_a) - i\hbar \mathbf{X}_a \right] \int_{-\infty}^t dt' e^{-i\mathcal{H}_0^{\mathcal{A}^{\text{ext}}}(t-t')/\hbar} (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*)_{I_j r} f_{t'}^{000}(\mathbf{I}'_j{}^r; \mathbf{R}) \\
 &+ \sum_{aj} \sum_{\mathbf{k}\lambda} (\boldsymbol{\varepsilon}_{\mathbf{k}\lambda})_{I_j r} \int_{-\infty}^t dt' e^{-i\mathcal{H}_0^{\mathcal{A}^{\text{ext}}}(t-t')/\hbar} \boldsymbol{\varepsilon}_{\mathbf{k}\lambda}^*(\mathbf{R}_b) \cdot \frac{Z_b |e|}{M_b} \left[\frac{\hbar}{i} \nabla_b - \frac{Z_b |e|}{c} \mathcal{A}^{\text{ext}}(\mathbf{R}_b) - i\hbar \mathbf{X}_b \right] f_{t'}^{000}(\mathbf{I}'_j{}^r; \mathbf{R}), \tag{E2}
 \end{aligned}$$

where the zeroth order Hamiltonian is defined as

$$\mathcal{H}_0^{\mathcal{A}^{\text{ext}}} \equiv \mathcal{H}_{\text{nuc}} + H_{\text{mf}}^{\text{el}} + H_{\text{rad}} + H_{\text{ext}} \tag{E3}$$

and

$$\Sigma_{\mathcal{A}^{\text{ext}}}^{\text{fin.}}(t, t')_{j, k} \equiv \iint d^3 \mathbf{r} d^3 \mathbf{r}' \varphi_j(\mathbf{r}) \Sigma_{\mathcal{A}^{\text{ext}}}^{\text{fin.}}(\mathbf{r}, t; \mathbf{r}', t') \varphi_k(\mathbf{r}'), \tag{E4}$$

with the finite part of the self-energy operator in the presence of the external field being $\Sigma_{\mathcal{A}^{\text{ext}}}^{\text{fin.}}(x, y)$. An explicit expression of such self-energy is given in an analogous manner as in Eq. (39),

$$\begin{aligned}
 \Sigma_{\mathcal{A}^{\text{ext}}}^{\text{fin.}}(x, y) &= i\hbar c e^2 \alpha^\mu S_{\mathcal{A}^{\text{ext}}}^{\text{F}}(x, y) \alpha^\nu D_{\mu\nu}(x, y) - \beta \delta m c^2 \delta^4(x - y) \\
 &- Z_2 \left(c \alpha^\mu i \hbar \partial_\mu - W_{\text{HF}}^{\text{loc}}(\mathbf{x}) - q_e \boldsymbol{\alpha} \cdot \mathcal{A}^{\text{ext}} - \beta m c^2 \right) \delta^4(x - y) + \delta^4(x - y) \mathcal{V}_{\mu}^{\text{vp}} \alpha^\mu(\mathbf{x}, t) \\
 &= i\hbar c e^2 \alpha^\mu S^{\text{F}}(x, y) \alpha^\nu D_{\mu\nu}(x, y) - \beta \delta m c^2 \delta^4(x - y) - Z_2 \left(c q_e \alpha^\mu i \hbar \partial_\mu - W_{\text{HF}}^{\text{loc}}(\mathbf{x}) - \beta m c^2 \right) \delta^4(x - y) \\
 &- i\hbar c e^2 \int d^4 \xi \alpha^\mu S^{\text{F}}(x, \xi) \boldsymbol{\alpha} \cdot \mathcal{A}^{\text{ext}} S_0^{\text{F}}(\xi, y) \alpha^\nu D_{\mu\nu}(x - y) - Z_2 \boldsymbol{\alpha} \cdot \mathcal{A}^{\text{ext}}(\mathbf{x}) \delta^4(x - y) \\
 &+ i\hbar c \int d^4 \xi \int d^4 \eta \alpha^\mu S^{\text{F}}(x, \xi) \boldsymbol{\alpha} \cdot \mathcal{A}^{\text{ext}}(\xi) S_{\mathcal{A}^{\text{ext}}}^{\text{F}}(\xi, \eta) \boldsymbol{\alpha} \cdot \mathcal{A}^{\text{ext}}(\eta) S^{\text{F}}(\eta, y) + \delta^4(x - y) \mathcal{V}_{\mu}^{\text{vp}} \alpha^\mu(\mathbf{x}, t), \tag{E5}
 \end{aligned}$$

in which $S_{\mathcal{A}^{\text{ext}}}^{\text{F}}$ and S^{F} being the Hartree-Fock Feynman propagator in the presence and absence of the external field, respectively. The above expansion does not necessarily require that the external field should be included in the mean-field Hamiltonian as long as one can obtain an expression of single-particle propagator. Obviously, however, a set of field-induced states, such as the Volkov states,^{85,86} would be one of the most suitable basis set to be used in dynamics in an external field of relativistic strength.

APPENDIX F: WAVEPACKET FORMULATION OF NUCLEAR DYNAMICS

Here, we append the discussion for the nuclear dynamics since this aspect is inevitable in practical applications of the present theory to molecular problems. Here, we restrict our attention to a mixed quantum-classical formulation or its variant, in which the

nuclear wavepacket is represented by a small number of localized wavepackets instead of a fully quantum-mechanical (delocalized) wavefunction.

The formal equation of motion in the Tamm-Dancoff expansion is given as in Eq. (64). Since the equation has almost the same mathematical form as those in the conventional nonrelativistic chemical dynamics (CD), many calculation techniques so far proposed can be applied for its integration. A crucial difference to the conventional CD is, however, the absence of explicit expressions of the true adiabatic states. Moreover, even if one derives an approximate expression of those states, it should be very different from those in conventional CD but should be a superposition of multiple electronic and radiational states. We therefore consider that it should be more convenient to adopt a “wavepacket picture,” where the electronic state vectors dynamically evolve in time (this contrasts with a more popular approach in which each wavepacket is associated with a single time-independent

state). In such a formulation, it is, in general, not appropriate to use a single wavepacket, but one needs to introduce bifurcation at a point where the character of the electronic state changes drastically. We therefore develop a general (multiple) wavepacket expansion of the nuclear dynamics. We then augment our theory with an appropriate “branching” algorithm in order to take account of bifurcations.

In this section, we use the mass-weighted representation of the nuclear coordinates, and hence, \mathbf{R} here represents mass-weighted coordinates. We start with a formal expansion of the electron-nucleus coupled state,

$$|\Omega_t\rangle = \int d^D \mathbf{R} |\mathbf{R}\rangle \sum_A |\Phi_A : \mathbf{R}\rangle \chi_A(\mathbf{R}, t), \quad (\text{F1})$$

where $|\Phi_A : \mathbf{R}\rangle$ represents a time-independent electronic state at nuclear coordinate \mathbf{R} and $\chi_A(\mathbf{R}, t)$ represents its associated nuclear wavefunction. Recalling Eq. (64) in Sec. III, a formal equation of motion of χ can be rewritten in the following form:

$$i\hbar \chi_A(\mathbf{R}, t) = \sum_B \left[\mathfrak{T}_{\text{nuc}}(\mathbf{R})_{AB} + H_{AB}^{\text{el}}(\mathbf{R}) \right] \chi_B(\mathbf{R}, t), \quad (\text{F2})$$

where $\mathfrak{T}_{\text{nuc}}$ is the (nonrelativistic) kinetic operator for nucleus,

$$\begin{aligned} \mathfrak{T}_{\text{nuc}}(\mathbf{R})_{AB} = & \frac{1}{2} \left(\frac{\hbar}{i} \nabla - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right)^2 \delta_{AB} \\ & - i\hbar \mathbf{X}_{AB} \cdot \left(\frac{\hbar}{i} \nabla - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right) - \frac{\hbar^2}{2} Y_{AB}, \end{aligned} \quad (\text{F3})$$

where $\nabla = \partial/\partial \mathbf{R}$ is the $3N_n$ -dimensional (mass-weighted) gradient, \mathcal{Q}_{nuc} represents a diagonal matrix which consists of reduced charges (the nuclear charges divided by the square-root of nuclear masses), and \mathcal{A}^{ext} represents the external field supervector $\mathcal{A}^{\text{ext}} \doteq (\mathbf{A}^{\text{ext}}(\mathbf{R}_1)^t, \mathbf{A}^{\text{ext}}(\mathbf{R}_2)^t, \dots)$. Since we neglect the radiation from the nuclear charge current, the resultant vector fields are regarded as an external field. Matrices \mathbf{X}_{AB} and Y_{AB} represent the derivative couplings of the first and second order; $\mathbf{X}_{AB} \equiv \langle A | \nabla | B \rangle$ and $Y_{AB} \equiv \langle A | \nabla^2 | B \rangle$. We then expand $\chi_A(\mathbf{R})$ as

$$\chi_A(\mathbf{R}) = \sum_{\sigma} \mathbf{g}_{\sigma} C_A^{\sigma}, \quad (\text{F4})$$

with normalized Gaussian \mathbf{g} , defined as

$$\begin{aligned} \mathbf{g}_{\sigma} = & \exp \left[- \sum_{\ell j} (x^{\ell} - Q_{\sigma}^{\ell}(t)) \frac{\Gamma_{\sigma}^{\ell j}(t)}{2} (x^j - Q_{\sigma}^j(t)) + \sum_{\ell} i K_{\sigma}^{\ell}(t) \right. \\ & \left. \times (x^{\ell} - Q_{\sigma}^{\ell}(t)) + i \Theta_{\sigma}(t) + \frac{1}{4} \ln \left(\det \left(\frac{\Gamma_{\sigma}(t)}{\pi} \right) \right) \right], \end{aligned} \quad (\text{F5})$$

where \mathbf{Q}_{σ} , $\hbar \mathbf{K}_{\sigma}$, Γ_{σ} , and Θ_{σ} represent the central coordinate, momentum, inverse width, and phase of the σ th Gaussian. The inverse width matrix Γ_{σ} is symmetric. For convenience, we also introduce a shorthand notation for nuclear coordinate integral in the bracket form,

$$\langle \sigma | \mathcal{O} | \tau \rangle \equiv \int d^D \mathbf{R} \mathbf{g}_{\sigma}^*(\mathbf{R}) \mathcal{O} \mathbf{g}_{\tau}(\mathbf{R}), \quad (\text{F6})$$

where \mathcal{O} represents an arbitrary function or operator in the nuclear coordinate representation. The overlap matrix is then defined as

$$S_{\sigma\tau} \equiv \langle \sigma | \tau \rangle. \quad (\text{F7})$$

In the RHS of Eq. (F1), coefficients C_A^{σ} are related to the total population of the A th adiabatic state n_A by

$$n_A = \sum_{\sigma\tau} C_A^{\sigma*} S_{\sigma\tau} C_A^{\tau},$$

where the unitarity condition requires $\sum_A n_A = 1$. We then apply the time-dependent variational equation,

$$\langle \delta \Omega_t | \mathcal{H}^{\text{tot}} - i\hbar \partial_t | \Omega_t \rangle = 0, \quad (\text{F8})$$

with \mathcal{H}^{tot} representing the total (electron-nucleus) Hamiltonian and bracket $\langle \dots \rangle$ representing the integration of electronic as well as nuclear coordinates, to derive equations of motion for $C_A^{\sigma}(t)$, and the parameters in the Gaussian functions, \mathbf{Q}_{σ} , \mathbf{K}_{σ} , and Γ_{σ} .

1. Variation of C_A^{σ}

We first decompose the total Hamiltonian \mathcal{H}^{tot} by the electronic basis as

$$\langle A | \mathcal{H}^{\text{tot}} | B \rangle = \mathcal{T} \delta_{AB} + \hat{\mathcal{K}}_{AB}, \quad (\text{F9})$$

where \mathcal{T} is the kinetic operator acting only on the nuclear wavefunction, whereas $\hat{\mathcal{K}}_{AB}$ is the remaining part of the Hamiltonian,

$$\hat{\mathcal{K}}_{AB} = H_{AB}^{\text{el}} - i\hbar \mathbf{X}_{AB} \cdot \left(\frac{\hbar}{i} \nabla - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right) - \frac{\hbar^2}{2} Y_{AB}, \quad (\text{F10})$$

where “hat” on \mathcal{K} is meant to emphasize that it is an operator working on the nuclear wavefunction. We also use notation H_{AB}^{eff} to denote the “electronic part,”

$$H_{AB}^{\text{eff}} = H_{AB}^{\text{el}} - \frac{\hbar^2}{2} Y_{AB}. \quad (\text{F11})$$

We now write down the variational stationary condition for $C_A^{\sigma*}$ as

$$\begin{aligned} 0 = & \sum_{\tau} \sum_B \langle \sigma | \langle A | \mathcal{H}^{\text{tot}} - i\hbar \frac{\partial}{\partial t} | B \rangle | \tau \rangle C_B^{\tau} \\ = & \sum_{\tau} \langle \sigma | f(\Gamma_{\tau}, \dot{\Gamma}_{\tau}; \tau) + \text{Tr} \left(\frac{\hbar^2 \Gamma_{\tau}}{4} \right) - \frac{1}{2} \left(\hbar \mathbf{K}_{\tau} - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right)^2 \\ & + \hbar \dot{\Theta}_{\tau} + \hbar \mathbf{K}_{\sigma} \cdot \mathbf{y}_{\tau} | \tau \rangle C_A^{\tau} \\ & + \sum_{\tau} \sum_B \langle \sigma | H_{AB}^{\text{eff}} - i\hbar \mathbf{X}_{AB} \cdot \left(\frac{\hbar}{i} \nabla - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right) | \tau \rangle C_B^{\tau} \\ & - \sum_{\tau} \langle \sigma | \tau \rangle i\hbar \dot{C}_A^{\tau}, \end{aligned} \quad (\text{F12})$$

where we introduced the notation

$$\mathbf{y}_{\sigma} \equiv \mathbf{R} - \mathbf{Q}_{\sigma}, \quad (\text{F13})$$

and $f(\Gamma_{\tau}, \dot{\Gamma}_{\tau}; \tau)$ represents

$$f(\Gamma_\tau, \dot{\Gamma}_\tau; \tau) \equiv \frac{1}{2} \text{Tr} \left[\left(i\hbar \dot{\Gamma}_\tau - \hbar^2 \Gamma_\tau^2 \right) \left(\mathbf{y}_\tau \mathbf{y}_\tau^\dagger - \frac{\Gamma_\tau^{-1}}{2} \right) \right],$$

with the operator Tr representing the trace over coordinate-like indices. We then obtain the time evolution equation for C_A^σ as

$$\begin{aligned} i\hbar \dot{C}_A^\sigma &= \sum_\tau \sum_B \langle \tilde{\sigma} | \mathcal{H}_{AB}^{\text{tot}} - i\hbar \hat{\partial}_t \delta_{AB} | \tau \rangle C_B^\tau \\ &= \sum_\tau \langle \tilde{\sigma} | f(\Gamma_\tau, \dot{\Gamma}_\tau; \tau) + \text{Tr} \left(\frac{\hbar^2 \Gamma_\tau}{4} \right) - \frac{1}{2} \left(\hbar \mathbf{K}_\tau - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right)^2 \\ &\quad + \hbar \dot{\Theta}_\tau + \hbar \dot{\mathbf{K}}_\sigma \cdot \mathbf{y}_\tau | \tau \rangle C_A^\tau + \sum_\tau \sum_B \langle \tilde{\sigma} | \hat{\mathcal{K}}_{AB} | \tau \rangle C_B^\tau, \end{aligned} \quad (\text{F14})$$

where $\hat{\partial}_t$ represents time derivative whose operation is limited to inside the bracket and bra-vectors with tilde and $\langle \tilde{\sigma} |$ are defined as $\langle \tilde{\sigma} | \equiv \sum_\tau (S^{-1})_{\sigma\tau} \langle \tau |$.

2. Variation of the Gaussian parameters

We first note that variation of the Gaussian parameters ξ , which is one of the four parameters K_σ^ℓ , Q_σ^ℓ , $\Gamma_\sigma^{\ell m}$, and Θ_σ , can be written in the following form:

$$\delta^{(\xi)} \mathbf{g}_\sigma(\mathbf{R}, t) = \delta \xi_\sigma \mathcal{O}_\sigma^{(\xi)} \mathbf{g}_\sigma(\mathbf{R}, t), \quad (\text{F15})$$

where the superscript (ξ) represents the variational parameter and the corresponding $\mathcal{O}_\sigma^{(\xi)}$ is in general a function of nuclear coordinates,

$$\mathcal{O}_\sigma^{(K_\sigma^\ell)} = i(x^\ell - Q_\sigma^\ell), \quad (\text{F16a})$$

$$\mathcal{O}_\sigma^{(Q_\sigma^\ell)} = -\Gamma_\sigma^{\ell m} (x^m - Q_\sigma^m) + iK_\sigma^\ell, \quad (\text{F16b})$$

$$\mathcal{O}_\sigma^{(\Theta_\sigma)} = i, \quad (\text{F16c})$$

$$\mathcal{O}_\sigma^{(\Gamma_\sigma^{\ell m})} = -(x^\ell - Q_\sigma^\ell)(x^m - Q_\sigma^m)/2 + (\Gamma_\sigma^{-1})^{m\ell}/4. \quad (\text{F16d})$$

We then write the variational stationary condition for those parameters,

$$\begin{aligned} 0 &= \sum_\tau \sum_{A,B} \langle \sigma | \mathcal{O}_\sigma C_A^{\sigma*} \langle A | \mathcal{H}^{\text{tot}} - i\hbar \frac{\partial}{\partial t} | B \rangle C_B^\tau | \tau \rangle \\ &= \sum_\tau \langle \sigma | \mathcal{O}_\sigma (\hat{\mathcal{T}} + \hat{\mathcal{K}} - i\hbar \hat{\partial}_t)^{\sigma\tau} | \tau \rangle \\ &\quad - \sum_{\tau\kappa} \sum_{AB} C_A^{\sigma*} \langle \sigma | \mathcal{O}_\sigma | \tau \rangle \langle \tilde{\tau} | \mathcal{H}_{AB}^{\text{tot}} - i\hbar \hat{\partial}_t \delta_{AB} | \kappa \rangle C_B^\kappa, \end{aligned} \quad (\text{F17})$$

where the matrices with Gaussian superscripts σ, τ, \dots represent

$$\mathcal{O}^{\sigma\tau} \equiv \sum_{AB} C_A^{\sigma*} O_{AB} C_B^\tau, \quad (\text{F18})$$

and we also define related quantities for the unit matrix

$$\eta^{\sigma\tau} \equiv \sum_A C_A^{\sigma*} C_A^\tau. \quad (\text{F19})$$

Although we have to find such a solution that eliminates all variation in principle, it would lead to complex expressions for numerical calculations. Here, we rather derive a simpler approximate solution and later confirm that major contributions to variation do vanish.

We first set

$$\hbar \mathbf{K}_\sigma = \dot{\mathbf{Q}}_\sigma \quad (\text{F20})$$

so that all terms containing $\dot{\mathbf{Q}}_\sigma$ vanish at all order. We then concentrate on large contributions, arising from $\sigma = \tau$, or diagonal terms in Eq. (F17), and eliminate variation of type $\mathcal{O} = 1$ and $\mathcal{O} = \mathbf{y}_\sigma$. It can then be shown that all the other residual terms arise from the higher order correlation. Diagonal contribution in the RHS of Eq. (F17) then reads

$$\begin{aligned} \langle \sigma | \mathcal{O}_\sigma \left[\eta^{\sigma\sigma} \left\{ f(\Gamma_\sigma, \dot{\Gamma}_\sigma; \sigma) + \text{Tr} \left(\frac{\hbar^2 \Gamma_\sigma}{4} \right) - \frac{1}{2} \left(\hbar \mathbf{K}_\sigma - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right)^2 \right. \right. \\ \left. \left. + \hbar \dot{\Theta}_\sigma + \hbar \dot{\mathbf{K}}_\sigma \cdot \mathbf{y}_\sigma \right\} + H_{\text{eff}}^{\sigma\sigma} \right] | \sigma \rangle, \end{aligned} \quad (\text{F21})$$

where we have taken account of the fact $\mathbf{X}^{\sigma\sigma} = 0$, which follows from the antisymmetry of \mathbf{X}_{AB} . Requiring Eq. (F21) for $\mathcal{O} = 1$, $\mathcal{O} = y_\sigma^\ell$ and $\mathcal{O} = y_\sigma^\ell y_\sigma^m$ to vanish, we obtain

$$-\hbar \dot{\Theta}_\sigma = \text{Tr} \left(\frac{\hbar^2 \Gamma_\sigma}{4} \right) - \frac{1}{2} (\hbar \mathbf{K}_\sigma)^2 + \langle \sigma | H_{\text{eff}}^{\sigma\sigma} | \sigma \rangle / \eta^{\sigma\sigma}, \quad (\text{F22a})$$

$$\hbar \dot{K}_\sigma^\ell = - \sum_m (2\Gamma_\sigma)^{\ell m} \langle \sigma | \left(\frac{1}{2} \left(\hbar \mathbf{K}_\sigma - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right)^2 + H_{\text{eff}}^{\sigma\sigma} \right) y_\sigma^m | \sigma \rangle / \eta^{\sigma\sigma}, \quad (\text{F22b})$$

$$\begin{aligned} 0 &= \sum_{jk} \langle \sigma | (i\hbar \dot{\Gamma}_\sigma - \hbar^2 \Gamma_\sigma^2)_{jk} \left(\left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{\ell m} \left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{jk} + \left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{\ell k} \left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{jm} \right) \\ &\quad + \langle \sigma | (H_{\text{eff}}^{\sigma\sigma} - \langle \sigma | H_{\text{eff}}^{\sigma\sigma} | \sigma \rangle) y_\sigma^\ell y_\sigma^m | \sigma \rangle / \eta^{\sigma\sigma}. \end{aligned} \quad (\text{F22c})$$

In deriving these equations, we took account of the fact that any odd momenta of Gaussian $\langle \sigma | y_\sigma^{\ell_1} \dots y_\sigma^{\ell_{2n+1}} | \sigma \rangle = 0$ and that

$$\langle \sigma | y_\sigma^j y_\sigma^\ell | \sigma \rangle = \frac{1}{2} (\Gamma_\sigma^{-1})^{\ell j} = \frac{1}{2} (\Gamma_\sigma^{-1})^{j\ell} \quad (\text{F23})$$

and

$$\begin{aligned} \langle \sigma | \prod_{p=1}^4 y_\sigma^{\ell_p} | \sigma \rangle &= (\Gamma_\sigma^{-1}/2)_{\ell_1 \ell_2} (\Gamma_\sigma^{-1}/2)_{\ell_3 \ell_4} + (\Gamma_\sigma^{-1}/2)_{\ell_1 \ell_3} \\ &\quad \times (\Gamma_\sigma^{-1}/2)_{\ell_2 \ell_4} + (\Gamma_\sigma^{-1}/2)_{\ell_1 \ell_4} (\Gamma_\sigma^{-1}/2)_{\ell_2 \ell_3}. \end{aligned} \quad (\text{F24})$$

Equations (F22a)–(F22c) reduce to more familiar forms if we expand $H_{\text{eff}}^{\sigma\sigma}$ around the Gaussian center \mathbf{Q}_σ up to the second order,

$$\begin{aligned} -\hbar \dot{\Theta}_\sigma &= \text{Tr} \left(\frac{\hbar^2 \Gamma_\sigma}{4} \right) - \frac{1}{2} \left(\hbar \mathbf{K}_\sigma - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}}(\mathbf{Q}_\sigma) \right)^2 \\ &\quad + \left(H_{\text{eff}}^{\sigma\sigma}(\mathbf{Q}_\sigma) + \frac{1}{2} \sum_{jk} \frac{\partial^2 H_{\text{eff}}^{\sigma\sigma}}{\partial R^j \partial R^k} \left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{jk} \right) / \eta^{\sigma\sigma}, \end{aligned} \quad (\text{F25a})$$

$$\hbar \dot{K}_\sigma = - \langle \sigma | \nabla \left(\frac{1}{2} \left(\hbar \mathbf{K}_\sigma - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right)^2 + H_{\text{eff}}^{\sigma\sigma} \right) | \sigma \rangle / \eta^{\sigma\sigma}, \quad (\text{F25b})$$

$$\begin{aligned} 0 &= \sum_{jk} \left((i\hbar \dot{\Gamma}_\sigma - \hbar^2 \Gamma_\sigma^2)_{jk} + \left(\frac{1}{2} \frac{\partial^2 H_{\text{eff}}^{\sigma\sigma}}{\partial R^j \partial R^k} \right) / \eta^{\sigma\sigma} \right) \\ &\quad \times \left(\left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{\ell m} \left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{jk} + \left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{\ell k} \left(\frac{\Gamma_\sigma^{-1}}{2} \right)_{jm} \right), \end{aligned} \quad (\text{F25c})$$

where the second one is further rewritten as

$$\begin{aligned} \frac{d}{dt} \left[\hbar \mathbf{K}_\sigma - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right] &= - \langle \sigma | \nabla H_{\text{eff}}^{\sigma\sigma} | \sigma \rangle / \eta^{\sigma\sigma} \\ &\quad - \frac{\mathcal{Q}_{\text{nuc}}}{c} \langle \sigma | \left(\hbar \mathbf{K}_\sigma - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right) \\ &\quad \times (\nabla \times \mathcal{A}^{\text{ext}}) | \sigma \rangle. \end{aligned} \quad (\text{F26})$$

We now fix the time derivatives of the first two set of parameters, $\dot{\Theta}_\sigma$ and $\hbar \mathbf{K}_\sigma$ by Eqs. (F22a) and (F22b), respectively, together with $\dot{\mathbf{Q}}_\sigma$ by Eq. (F20). On the other hand, we formally leave $\dot{\Gamma}$ in

the following expressions as we later employ the frozen Gaussian approximation to fix Γ_σ at a reasonable value, such as that satisfies Eq. (F25c) at the initial point of the dynamics. Substitution of the relation

$$\begin{aligned} \langle \tau | \mathcal{F} - i\hbar \dot{\Theta}_\tau | \sigma \rangle &= \langle \tau | f(\Gamma_\sigma, \dot{\Gamma}_\sigma; \sigma) | \sigma \rangle - \langle \tau | \sigma \rangle \langle \sigma | H_{\text{eff}}^{\sigma\sigma} | \sigma \rangle / \eta^{\sigma\sigma} \\ &\quad - \sum_{\ell, m} \langle \tau | y_\sigma^\ell | \sigma \rangle (2\Gamma_\sigma)_{\ell m} \langle \sigma | H_{\text{eff}}^{\sigma\sigma} y_\sigma^m | \sigma \rangle / \eta^{\sigma\sigma} \end{aligned} \quad (\text{F27})$$

into Eq. (F12) leads to

$$\begin{aligned} 0 &= \sum_\tau \langle \sigma | \mathcal{O}_\sigma \eta^{\sigma\tau} f | \tau \rangle - \sum_\tau \eta^{\sigma\tau} \langle \sigma | \mathcal{O}_\sigma | \tau \rangle \langle \tau | H_{\text{eff}}^{\tau\tau} | \tau \rangle / \eta^{\tau\tau} - \sum_\tau \eta^{\sigma\tau} \langle \sigma | \mathcal{O}_\sigma y_\sigma^\dagger | \tau \rangle (2\Gamma_\tau) \langle \tau | y_\tau H_{\text{eff}}^{\tau\tau} | \tau \rangle / \eta^{\tau\tau} - \sum_\tau \langle \sigma | \mathcal{O}_\sigma H_{\text{eff}}^{\sigma\tau} | \tau \rangle \\ &\quad - \sum_\tau \langle \sigma | \mathcal{O}_\sigma i\hbar \mathbf{X}^{\sigma\tau} \cdot \left(i\hbar \Gamma_\tau \mathbf{y}_\tau + \left(\hbar \mathbf{K}_\tau - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right) \right) | \tau \rangle - \sum_{\tau\lambda} \langle \sigma | \mathcal{O}_\sigma | \tau \rangle \left[\langle \tau | f(\Gamma_\lambda, \dot{\Gamma}_\lambda; \lambda) | \lambda \rangle - \langle \tau | H_{\text{eff}}^{\tau\tau} | \tau \rangle / \eta^{\lambda\lambda} \right. \\ &\quad \left. - \langle \tau | y_\lambda^\dagger | \lambda \rangle (2\Gamma_\lambda) \langle \lambda | y_\lambda H_{\text{eff}}^{\lambda\lambda} | \lambda \rangle / \eta^{\lambda\lambda} \right] \eta^{\sigma\lambda} + \langle \tau | H_{\text{eff}}^{\sigma\lambda} - i\hbar \mathbf{X}^{\sigma\lambda} \cdot \left(\frac{\hbar}{i} \nabla - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right) | \lambda \rangle, \end{aligned} \quad (\text{F28})$$

which shows that the variation indeed vanishes for $\mathcal{O}_\sigma = 1$ and the residual part consists only of the higher correlation for the other \mathcal{O}_σ .

We also substitute Eq. (F27) into Eq. (F12) to obtain a reduced equation for C_A^σ ,

$$\begin{aligned} i\hbar \dot{C}_A^\sigma &= \sum_\tau \langle \tilde{\sigma} | f(\Gamma_\tau, \dot{\Gamma}_\tau; \tau) | \tau \rangle C_A^\tau - \left[\langle \sigma | H_{\text{eff}}^{\sigma\sigma} | \sigma \rangle / \eta^{\sigma\sigma} \right] C_A^\sigma - \sum_\tau \left[\langle \tilde{\sigma} | y_\tau^\dagger | \tau \rangle (2\Gamma_\tau) \langle \tau | H_{\text{eff}}^{\tau\tau} | \tau \rangle / \eta^{\tau\tau} \right] C_A^\tau \\ &\quad + \sum_\tau \sum_B \langle \tilde{\sigma} | H_{AB}^{\text{eff}} - i\hbar \mathbf{X}_{AB} \cdot \left(i\hbar \Gamma_\tau \mathbf{y}_\tau + \left(\hbar \mathbf{K}_\tau - \frac{\mathcal{Q}_{\text{nuc}}}{c} \mathcal{A}^{\text{ext}} \right) \right) | \tau \rangle C_B^\tau, \end{aligned} \quad (\text{F29})$$

which is rewritten in the following form:

$$\begin{aligned} i\hbar \dot{C}_A^\sigma &= \Delta_{\sigma\sigma}^{\text{dag}} C_A^\sigma + \sum_{\tau \neq \sigma} \Delta_{\sigma\tau}^{\text{ofd}} C_A^\tau + \sum_B K_{AB}^{(1)\sigma\sigma} C_B^\sigma \\ &\quad + \sum_\tau K_{AA}^{(2)\sigma\tau} C_A^\tau + \sum_\tau \sum_B K_{AB}^{(3)\sigma\tau} C_B^\tau, \end{aligned} \quad (\text{F30})$$

with

$$\Delta_{\sigma\sigma}^{\text{dag}} = - \langle \sigma | H_{\text{eff}}^{\sigma\sigma} | \sigma \rangle / \eta^{\sigma\sigma} - \langle \tilde{\sigma} | y_\sigma^\dagger | \sigma \rangle (2\Gamma_\sigma) \langle \sigma | H_{\text{eff}}^{\sigma\sigma} y_\sigma | \sigma \rangle / \eta^{\sigma\sigma}, \quad (\text{F31a})$$

$$\begin{aligned} \Delta_{\sigma\tau}^{\text{ofd}} &= \langle \tilde{\sigma} | f(\Gamma_\tau, \dot{\Gamma}_\tau; \tau) | \tau \rangle \\ &\quad - (1 - \delta_{\sigma\tau}) \langle \tilde{\sigma} | y_\tau^\dagger | \tau \rangle (2\Gamma_\tau) \langle \tau | H_{\text{eff}}^{\tau\tau} y_\tau | \tau \rangle / \eta^{\tau\tau}, \end{aligned} \quad (\text{F31b})$$

$$K_{AB}^{(1)\sigma\sigma} = \langle \tilde{\sigma} | H_{AB}^{\text{eff}} - i\hbar \mathbf{X}_{AB} \cdot (i\hbar \Gamma_\tau \mathbf{y}_\tau + \hbar \mathbf{K}_\tau) | \sigma \rangle C_B^\sigma, \quad (\text{F31c})$$

$$K_{AA}^{(2)\sigma\tau} = (1 - \delta_{\sigma\tau}) \langle \tilde{\sigma} | H_{AA}^{\text{eff}} | \sigma \rangle C_A^\tau, \quad (\text{F31d})$$

$$K_{AB}^{(3)\sigma\tau} = (1 - \delta_{\sigma\tau}) (1 - \delta_{AB}) \langle \tilde{\sigma} | H_{AB}^{\text{eff}} - i\hbar \mathbf{X}_{AB} \cdot (i\hbar \Gamma_\tau \mathbf{y}_\tau + \hbar \mathbf{K}_\tau) | \tau \rangle. \quad (\text{F31e})$$

We then see $K_{AB}^{(1)\sigma\sigma}$ describes the electron wavepacket dynamics (including possible nonadiabatic transitions among the wavepacket components) of the σ th wavepacket, and $\Delta_{\sigma\sigma}^{\text{dag}}$ adds or subtracts the nuclear wavepacket contribution of the phase

factor, whereas $K_{AB}^{(3)\sigma\tau}$ describes the nonadiabatic transitions between the σ th and τ th wavepackets. On the other hand, we consider that $K_{AA}^{(2)\sigma\tau}$ and $\Delta_{\sigma\tau}^{\text{ofd}}$ cause “transition to different wavepacket with no electronic state change,” which corresponds to an artifact of our Gaussian expansion scheme [Eq. (F4)]. We therefore expect that the amplitude of these two terms should be much smaller than other terms if our Gaussian propagation scheme is successful.

3. A practical calculation scheme

Having established formal equations of motion, we now consider a practical calculation scheme with branching. We first apply the frozen-Gaussian approximation, to avoid complexity and possible unphysical deformation of Gaussian wavepackets. We set Γ_σ of the initial wavepacket to some reasonable value, in such a manner that Γ^2 equals the Hessian of the electronic Hamiltonian at the initial geometry and set $\dot{\Gamma} = 0$ in subsequent time evolution. We thus start from a single wavepacket, but the final state should be in general a superposition of multiple wavepackets that asymptote to mutually independent adiabatic states. Since our equations of motion do not automatically increase the number of Gaussian wavepackets, we need an additional algorithm that introduces new wavepackets. Following Ref. 55, we call it a “branching” algorithm.

There are several clues to judge path-branching that directly follows from our formulation: (i) the variation residual of Eq. (F17), which does not exactly vanish except for $\mathcal{O} = 1$, or (ii) $K_{AA}^{(2)\sigma r}$ and $\Delta_{\sigma r}^{\text{off}}$ in Eq. (F30), which should be small by physical requirements. One can therefore introduce a new wavepacket such that it reduces those inconvenient quantities. Such algorithm should be most desirable for fully consistent realization of our Gaussian wavepacket formulation; however, we here seek for a simpler algorithm that directly realizes branchings.

An alternative approach is to judge from the nuclear kinematics. In case we know the asymptotic adiabatic states, we can (a) introduce branching so that each wavepacket should consist of a single adiabatic state in the final asymptotic region. Even if we do not know those states *a priori*, we can also use more intuitive quantity, “force matrix,” introduced in Refs. 54 and 55. In the present formulation, it appears as the first (nonradiative) term in the RHS of Eq. (F26),

$$\mathbf{F}_{\sigma\sigma} = -\langle\sigma|\nabla H_{\text{eff}}^{\sigma\sigma}|\sigma\rangle/\eta^{\sigma\sigma} = \sum_{A,B} C_A^{\sigma*} \mathbf{F}_{AB}^{\sigma\sigma} C_B^{\sigma} / \sum_{A'} C_{A'}^{\sigma*} C_{A'}^{\sigma}, \quad (\text{F32})$$

where we find a state-dependent force, or a variant of the force matrix,

$$\mathbf{F}_{AB}^{\sigma\sigma} = -\langle\sigma|\nabla H_{AB}|\sigma\rangle. \quad (\text{F33})$$

One can then (b) use the eigenvectors of the force matrix to decompose the wavepacket into most rapidly departing parts.⁵⁵ In either scheme, (a) or (b), we can construct branching algorithm following the idea proposed in Ref. 55: (1) Set a relative population threshold $n_{\text{thr}}^{\text{rel}}$ and relative adiabatic coupling thresholds $\xi_{\text{thr}}^{\text{rel}}$. (2) Start checking wavepackets which have a mixture of distinct adiabatic states more than $n_{\text{thr}}^{\text{rel}}$ when the strength of the nonadiabatic coupling decreases and passes through the thresholds $\xi_{\text{thr}}^{\text{rel}}$ (relative to the peak) downward, and divide the wavepacket by projecting onto either (a) the adiabatic state vectors or (b) the eigenstates of the force matrix.

At this point, our discussion on the branching algorithm cannot be further extended since branching does not directly follow from the variational principle [Eq. (F8)]. The best algorithm is therefore only be judged through numerical applications by comparing accuracy, convergence, computational costs, etc., which we leave to our future study.

REFERENCES

- V. B. Berestetskii, E. M. Lifshitz, and L. P. Pitaevskii, *Quantum Electrodynamics* (Butterworth-Heinemann, Oxford, 1982).
- S. Weinberg, *Quantum Field Theory* (Cambridge University Press, 2004).
- M. Reiher and A. Wolf, *Relativistic Quantum Chemistry* (Wiley, Weinheim, 2015).
- W. Liu, *Handbook of Relativistic Quantum Chemistry* (Springer, Berlin, 2017).
- P. J. Mohr, *Ann. Phys.* **88**, 26 (1974).
- J. Sucher, *Phys. Rev. A* **22**, 348 (1980).
- P. J. Mohr, G. Plunien, and G. Soff, *Phys. Rep.* **293**, 227 (1998).
- S. J. Rose, I. P. Grant, and N. C. Pyper, *J. Phys. B: At. Mol. Phys.* **11**, 1171 (1978).
- E. J. Baerends, W. H. E. Schwarz, P. Schwerdtfeger, and J. G. Snijders, *J. Phys. B: At. Mol. Phys.* **23**, 3225 (1990).
- G. A. Mourou, T. Tajima, and S. V. Bulanov, *Rev. Mod. Phys.* **78**, 309 (2006).
- A. Di Piazza, C. Muller, K. Z. Hatsagortsyan, and C. H. Keitel, *Rev. Mod. Phys.* **84**, 1177 (2012).
- K. Hanasaki and K. Takatsuka submitted for publication in *Phys. Rev. A*.
- M. Aoyama, K. Yamakawa, Y. Akahane, J. Ma, N. Inoue, H. Ueda, and H. Kiriya, *Opt. Lett.* **28**, 1594 (2003).
- C. Danson, D. Hillier, N. Hopps, and D. Neely, *High Power Laser Sci. Eng.* **3**, e3 (2015).
- M. D. Perry, D. Pennington, B. C. Stuart, G. Tiethohl, J. A. Britten, C. Brown, S. Hermann, B. Golick, M. Kartz, J. Miller, H. T. Powell, M. Vergino, and V. Yanovsky, *Opt. Lett.* **24**, 160 (1999).
- C. I. Moore, A. Ting, S. J. McNaught, J. Qiu, H. R. Burris, and P. Sprangle, *Phys. Rev. Lett.* **82**, 1688 (1999).
- J. M. Cole, K. T. Behm, E. Gerstmayr, T. G. Blackburn, J. C. Wood, C. D. Baird, M. J. Duff, C. Harvey, A. Ilderton, A. S. Joglekar, K. Krushelnick, S. Kuschel, M. Marklund, P. McKenna, C. D. Murphy, K. Poder, C. P. Ridgers, G. M. Samarin, G. Sarri, D. R. Symes, A. G. R. Thomas, J. Warwick, M. Zepf, Z. Najmudin, and S. P. D. Mangles, *Phys. Rev. X* **8**, 011020 (2018).
- L. L. Ji, A. Pukhov, I. Yu. Kostyukov, B. F. Shen, and K. Akli, *Phys. Rev. Lett.* **112**, 145003 (2014).
- W. Yan, C. Fruhling, G. Golovin, D. Haden, J. Luo, P. Zhang, B. Zhao, J. Zhang, C. Liu, M. Chen, S. Chen, S. Banerjee, and D. Umstadter, *Nat. Commun.* **11**, 514 (2017).
- S. Palaniyappan, R. Mitchell, R. Sauer, I. Ghebregziabher, S. White, M. Decamp, and B. Walker, *Phys. Rev. Lett.* **100**, 183001 (2008).
- P. Grugan, N. Ekanayake, S. White, S. Luo, P. Ruan, and B. Walker, *J. Phys. B: At. Mol. Phys.* **51**, 165601 (2018).
- P. Pyykko and J.-P. Desclaux, *Acc. Chem. Res.* **12**, 276 (1979); P. Pyykko, *Chem. Rev.* **88**, 563 (1988).
- W. Liu, *Phys. Rep.* **537**, 59 (2014).
- W. Liu, *Nat. Sci. Rev.* **3**, 204 (2016).
- P. Pyykko and Y. F. Zhao, *Angew. Chem., Int. Ed. Engl.* **30**, 604 (1991).
- P. Pyykko, *Angew. Chem., Int. Ed. Engl.* **43**, 4412 (2004).
- P. Schwerdtfeger, M. Dolg, W. H. E. Schwarz, G. A. Bowmaker, and P. D. W. Boyd, *J. Chem. Phys.* **91**, 1762 (1989).
- R. Fukuda, M. Hada, and H. Nakatsuji, *J. Chem. Phys.* **118**, 1015 (2003).
- M. Douglas and N. Kroll, *Ann. Phys.* **82**, 89 (1974); G. Jansen and B. Hess, *Phys. Rev. A* **39**, 6016 (1989).
- T. Nakajima and K. Hirao, *J. Chem. Phys.* **113**, 7786 (2000); *Chem. Rev.* **112**, 385 (2012).
- T. Tsuchiya, M. Abe, T. Nakajima, and K. Hirao, *J. Chem. Phys.* **115**, 4463 (2001).
- W. Liu, *J. Chem. Phys.* **123**, 241102 (2005); W. Liu and D. Peng, *ibid.* **125**, 044102 (2006).
- W. Liu, *Mol. Phys.* **108**, 1679 (2010).
- T. Saue, *ChemPhysChem* **12**, 3077 (2011).
- D. Peng and M. Reiher, *Theor. Chem. Acc.* **131**, 1081 (2012).
- J. D. Talman, *Phys. Rev. Lett.* **57**, 1091 (1986).
- H. Nakatsuji and H. Nakashima, *Phys. Rev. Lett.* **95**, 050407 (2005); H. Nakatsuji, *Acc. Chem. Res.* **45**, 1480 (2012).
- H. J. Aa Jensen, K. G. Dyall, T. Saue, and K. Faegri, Jr., *J. Chem. Phys.* **104**, 4083 (1996).
- P. G. Szalay, T. Müller, G. Gidofalvi, H. Lischka, and R. Shepard, *Chem. Rev.* **112**, 108 (2012).
- K. Faegri, Jr., *Theor. Chem. Acc.* **105**, 252 (2001).
- T. Yanai, T. Nakajima, Y. Ishikawa, and K. Hirao, *J. Chem. Phys.* **114**, 6526 (2001).
- T. Saue and H. J. Aa Jensen, *J. Chem. Phys.* **111**, 6211 (1999).
- L. Visschera, O. Visser, P. J. C. Aerts, H. Merenga, and W. C. Nieuwpoort, *Comput. Phys. Commun.* **81**, 120 (1994).
- T. Yanai, H. Nakano, T. Nakajima, T. Tsuneda, S. Hirata, Y. Kawashima, Y. Nakao, M. Kamiya, H. Sekino, and K. Hirao, in *Computational Science-ICCS 2003*, Lecture Notes in Computer Science Vol. 2660, edited by P. M. A. Sloot *et al.* (Springer, 2003), p. 84.

- ⁴⁵L. D. Landau, *Phys. Z. Sowjetunion* **2**, 46 (1932).
- ⁴⁶C. Zener, *Proc. R. Soc. A* **137**, 696 (1932).
- ⁴⁷E. C. G. Stueckelberg, *Helv. Phys. Acta* **5**, 369 (1932).
- ⁴⁸W. Domcke, D. Yarkony, and H. Köppel, *Conical Intersections* (World Scientific, New Jersey, 2004).
- ⁴⁹M. Baer, *Beyond Born-Oppenheimer: Electronic Nonadiabatic Coupling and Applications* (Wiley, New Jersey, 2006).
- ⁵⁰H. Nakamura, *Nonadiabatic Transition: Concepts, Basic Theories and Applications* (World Scientific, Singapore, 2012).
- ⁵¹A. H. Zewail, *Femtochemistry: Ultrafast Dynamics of the Chemical Bond, I and II* (World Scientific, Singapore, 1994).
- ⁵²Y. Arasaki, K. Takatsuka, K. Wang, and V. McKoy, *Phys. Rev. Lett.* **90**, 248303 (2003); Y. Arasaki and K. Takatsuka, *Phys. Chem. Chem. Phys.* **13**, 4756 (2011).
- ⁵³Y. Arasaki and K. Takatsuka, *Phys. Chem. Chem. Phys.* **12**, 1239 (2010); Y. Arasaki, K. Wang, V. McKoy, and K. Takatsuka, *ibid.* **13**, 8681 (2011).
- ⁵⁴K. Takatsuka, *J. Phys. Chem. A* **111**, 10196 (2007).
- ⁵⁵T. Yonehara and K. Takatsuka, *J. Chem. Phys.* **129**, 134109 (2008).
- ⁵⁶K. Hanasaki and K. Takatsuka, *Phys. Rev. A* **81**, 052514 (2010).
- ⁵⁷T. Yonehara, K. Hanasaki, and K. Takatsuka, *Chem. Rev.* **112**, 499 (2012).
- ⁵⁸K. Takatsuka, T. Yonehara, K. Hanasaki, and Y. Arasaki, *Chemical Theory Beyond the Born-Oppenheimer Paradigm* (World Scientific, Singapore, 2015).
- ⁵⁹F. Hangelberg, *Electron Dynamics in Molecular Interactions* (Imperial College Press, London, 2014).
- ⁶⁰I. Tamm, *J. Phys. (USSR)* **9**, 449 (1945).
- ⁶¹S. M. Dancoff, *Phys. Rev.* **78**, 382 (1950).
- ⁶²S. Tomonaga, *Prog. Theor. Phys.* **1**, 27 (1946).
- ⁶³J. Schwinger, *Phys. Rev.* **74**, 1439 (1948).
- ⁶⁴J. Schwinger, *Phys. Rev.* **75**, 651 (1949).
- ⁶⁵R. P. Feynman, *Phys. Rev.* **76**, 749 (1949).
- ⁶⁶R. P. Feynman, *Phys. Rev.* **76**, 769 (1949).
- ⁶⁷F. J. Dyson, *Phys. Rev.* **90**, 994 (1953); F. J. Dyson, M. Ross, E. E. Salpeter, S. S. Schweber, M. K. Sundaresan, W. M. Vischer, and H. A. Bethe, *ibid.* **95**, 1644 (1954).
- ⁶⁸M. Cini, *Nuovo Cimento* **10**, 526 (1953); **10**, 614 (1953).
- ⁶⁹W. M. Visscher, *Phys. Rev.* **96**, 788 (1954).
- ⁷⁰R. J. Perry, A. Harindranath, and K. G. Wilson, *Phys. Rev. Lett.* **65**, 2959 (1990); R. J. Perry and A. Harindranath, *Phys. Rev. D* **43**, 4051 (1991).
- ⁷¹A. Harindranath and R. J. Perry, *Phys. Rev. D* **43**, 492 (1991).
- ⁷²D. Andrae, *Phys. Rep.* **336**, 413 (2000).
- ⁷³J. C. Slater, *Phys. Rev.* **81**, 385 (1951).
- ⁷⁴W. Kohn and L. J. Sham, *Phys. Rev.* **140**, A1133 (1965).
- ⁷⁵R. T. Sharp and G. K. Horton, *Phys. Rev.* **90**, 317 (1953).
- ⁷⁶J. D. Talman and W. F. Shadwick, *Phys. Rev. A* **14**, 36 (1976).
- ⁷⁷W. Heitler, *The Quantum Theory of Radiation*, 3rd ed. (Clarendon Press, Oxford, 1954), p. 121.
- ⁷⁸J. Sakurai, *Advanced Quantum Mechanics* (Addison-Wesley, Boston, 1987), p. 134.
- ⁷⁹An explicit expression of the Coulombic self-interaction can formally be written as $V_{SI} = \frac{e^2}{2} \iint d^3r d^3r' \delta^3(\mathbf{r} - \mathbf{r}') \psi^\dagger(\mathbf{r}) \psi(\mathbf{r}) / |\mathbf{r} - \mathbf{r}'|$. See also Eq. (32).
- ⁸⁰D. Hedendahl and J. Holmberg, *Phys. Rev. A* **85**, 012514 (2012).
- ⁸¹J. Holmberg, S. Salomonson, and I. Lindgren, *Phys. Rev. A* **92**, 012509 (2015).
- ⁸²N. J. Snyderman, *Ann. Phys.* **211**, 43 (1991).
- ⁸³One can show that the one-loop self-energy contribution to the effective Hamiltonian in the Hermitian form, shown as Eq. (289) in [I. Lindgren, S. Salomonson, and B. Åsén, *Phys. Rep.* **389**, 161 (2004)], $\mathcal{H}_{\text{eff}}^S = \mathcal{S}^{-1/2} \mathcal{K} \mathcal{S}^{-1/2}$ gives an off-diagonal energy correction similar to the Eq. (49) (in the sense that the expression includes the zeroth order energies of the both initial and final states). We also find that Ref. 23 used the same form of expression as radiation-correction to the orbital energy although the self-energy expression might differ in details.
- ⁸⁴H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One- and Two-Electron Atoms* (Springer Berlin, 1957).
- ⁸⁵D. M. Volkov, *Z. Phys.* **94**, 250 (1935); *Zh. Eksp. Teor. Fiz.* **7**, 1286 (1937).
- ⁸⁶V. I. Ritus, *J. Sov. Laser Res.* **6**, 497 (1985).