TITLE:

Scapular kinematic alterations during arm elevation with decrease in pectoralis minor stiffness after stretching in healthy individuals

AUTHOR(S):

Umehara, Jun; Nakamura, Masatoshi; Nishishita, Satoru; Tanaka, Hiroki; Kusano, Ken; Ichihashi, Noriaki

CITATION:

Umehara, Jun ...[et al]. Scapular kinematic alterations during arm elevation with decrease in pectoralis minor stiffness after stretching in healthy individuals. Journal of Shoulder and Elbow Surgery 2018, 27(7): 1214-1220

ISSUE DATE:

2018-07

URL:

<http://hdl.handle.net/2433/244286>

RIGHT:

© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Scapular kinematics and pectoralis minor stiffness

Title page

Scapular kinematic alterations during arm elevation with decrease in pectoralis minor stiffness
after stretching in healthy individuals

Jun Umehara, MS.^{1)*}, Masatoshi Nakamura, PhD.²⁾, Satoru Nishishita MS.¹⁾, Hiroki Tanaka
MS.³⁾, Ken Kusano, MS.⁴⁾, Noriaki Ichihashi, PhD.¹⁾

1) Human Health Sciences, Graduate School of Medicine, Kyoto University, Kyoto, Japan

2) Institute for Human Movement and Medical Sciences, Niigata University of Health and
Welfare, Niigata, Japan

3) Rehabilitation Unit, Kyoto University Hospital, Kyoto, Japan

4) ASICS Corporation, Institution of Sports Science, Kobe, Japan

***Corresponding author**

Jun Umehara

Human Health Sciences, Graduate School of Medicine, Kyoto University,

53 Shogoin-Kawahawa-cho, Kyoto 606-8507, Japan

Phone: +81-75-751-3935; Fax: +81-75-751-3909

E-mail: umehara.jun.77z@st.kyoto-u.ac.jp

Financial biases: None

Ethics committee approval

The study design was approved by the ethic committee of Kyoto University Graduate School
and the Faculty of Medicine (R0233).

Scapular kinematics and pectoralis minor stiffness

26 **Comments**

27 Please publish the Figure 3 in color.

28

29 **Acknowledgement**

30 This study was supported by the Grant – in – Aid for Scientific Research (B) 15H03043.

31

32

Scapular kinematics and pectoralis minor stiffness

Abstract

Background: Pectoralis minor tightness may be seen in individuals with scapular dyskinesis, and stretching is used for the treatment of altered scapular motion in sports and clinical fields. However, few researchers have reported on the effects of the pectoralis minor stiffness on scapular motion during arm elevation. The purpose of this study is to investigate whether acute decrease of pectoralis minor stiffness after stretching changes the scapular motion during arm elevation.

Methods: Fifteen dominant and 15 non-dominant upper limbs in healthy men were allocated as control and interventional limbs, respectively. In the intervention limb group, the shoulder was passively and horizontally abducted at 150° of elevation for five minutes to stretch the pectoralis minor muscle. Before and after stretching, three-dimensional scapular motion during abduction and scaption was examined using an electromagnetic sensor. Pectoralis minor stiffness was measured using ultrasonic shear wave elastography before and immediately after stretching, and after arm elevation.

Results: In the interventional limb, the pectoralis minor stiffness decreased by 3.2 kPa immediately after stretching and 2.5 kPa after arm elevation. The maximal changes in scapular kinematics after stretching were 4.8° of external rotation and 3.3° of posterior tilt in abduction, and 4.5° of external rotation and 3.7° of posterior tilt in scaption. No changes in upward rotation in abduction or scaption were seen.

Conclusion: Stretching for pectoralis minor muscle increases external rotation and posterior tilt of the scapula during arm elevation.

Level of evidence: Basic Science, Kinesiology Study.

Keywords: Shoulder; Physical therapy; Biomechanics; Stretching; Muscle stiffness; Elastography; Pectoralis minor muscle;

Scapular kinematics and pectoralis minor stiffness

Introduction

The shoulder joint consists of the scapula, humerus, and clavicle, and is one of the largest and most complex joints in humans. The coordinated movement of these bones is important for optimal shoulder motion. Early authors investigating scapula motion in healthy individuals defined scapulohumeral rhythm,¹⁰ and it has been established that the scapula rotates upward, externally, and tilts posteriorly during arm elevation in healthy individuals.^{11,19,22} Additional researchers reported that scapular motion of the patients with impingement syndrome or glenohumeral instability was decreased in external and upward rotation and posterior tilt as compared with that of healthy individuals.^{2,17,21} Scapular dyskinesis has been defined as the set of abnormal motions and positions of scapula,¹⁴ and the evaluation and treatment for scapular dyskinesis may be essential for shoulder rehabilitation.

The onset scapular dyskinesis was related to the tightness of soft tissue surrounding the scapula.^{7,13} The tightness of the pectoralis minor muscle (PMi),^{7,13} the short head of the biceps brachii,¹³ the levator scapula,⁷ or the rhomboid⁷ has been speculated to cause scapular dyskinesis. Of these shoulder muscles, the PMi is the only muscle whose relationship between tightness and scapular dyskinesis has been verified by experimental study. Borstad et al.⁴ examined three-dimensional (3D) scapular motion during elevation in healthy individuals with and without a shortened PMi and showed that a decrease in external rotation and posterior tilt are seen in individuals with a shortened PMi. The altered scapular kinematics, which is found in subjects with shortened PMi, seen in this previous study⁴ was similar to that observed in many patients with shoulder disease.^{15,17} Therefore, PMi tension may be important factor in scapular dyskinesis.

Stretching is applied as an approach to scapular dyskinesis caused by the PMi tightness. Borstad et al.³ recommended a unilateral corner stretch as one self-stretch method for the PMi. Umehara et al.³¹ also showed that shoulder horizontal abduction at an elevation of 150° was the

Scapular kinematics and pectoralis minor stiffness

most effective stretching technique for the PMi. Considering that there is a correlation between PMi stiffness and scapular dyskinesis, it is obvious that investigating not only the stretching maneuver but also the change in the PMi stiffness and scapular motion after stretching is important. However, little is examined on this relationship.

The purpose of the present study was to investigate whether the acute decrease in PMi stiffness after stretching alters the 3D scapular motion during arm elevation. Borstad et al.⁴ reported a decrease in external rotation and posterior tilt of scapula in individuals with a shortened PMi as compared with in healthy individuals. Therefore, we hypothesized that the decrease in PMi stiffness after stretching augments the external rotation and posterior tilt of scapula during arm elevation.

Scapular kinematics and pectoralis minor stiffness

95 **Materials and Methods**

96 **Participants**

97 This study was a controlled experimental study. Twenty men (age, 25.4 ± 3.1 years; height,
98 171.5 ± 5.3 cm; weight, 67.6 ± 8.5 kg) participated in this study. Dominant and non-dominant
99 upper limbs were allocated as control and interventional limbs respectively. The subjects were
100 randomly recruited from the students at our institution. Upon selection, the subjects orally
101 confirmed that they do not meet the exclusion criteria, which included female gender,
102 designation as an athlete or performing any extensive exercise, a history of orthopedic or
103 nervous system disease in upper limb. Considering that the low body mass index minimized
104 skin motion artifacts in the measurement of scapular motion during arm elevation, we also
105 excluded the subject with body mass index >25 , calculated using the height and weight. Prior
106 to the experiment, four men—one with a daily extensive exercise regimen, one with a history
107 of shoulder pain, and two with a high body mass index—were excluded. The aim and
108 procedures of the study were explained to all subjects, and informed consent was obtained. The
109 study protocol was approved by the ethics committee of our institution, and conformed the
110 principle of the Declaration of Helsinki.

111

112 **Experimental Procedures**

113 The participants, while sitting on a wooden stool, performed shoulder abduction (elevation in
114 the coronal plane) and scaption (elevation in the scapular plane) before and after the PMi
115 stretching. The stretching procedure of the PMi is described in detail in our previous study
116 (Figure 1).³¹ The participants underwent stretching to the point of discomfort (but not pain) for
117 five minutes (30 seconds, 10 repetitions, 10-second intervals). Each elevation plane was marked
118 on the floor using sections of elastic tape. In the starting posture, keeping the upper limb aside
119 the body with the elbow fully extended, the palm facing the body, and the eyes looking straight

Scapular kinematics and pectoralis minor stiffness

forward on the target at eye height, the participant was asked to raise their arm to full elevation in four seconds and then lower it to starting position in four seconds three times consecutively to the rhythm of a metronome with 60 BPM. The participant underwent sufficient familiarization to the abduction and scaption before the assessment.

Instrumentation

Scapular Kinematics

The 3D motion of the shoulder complex during arm elevation before and after stretching was measured using an electromagnetic tracking device (Liberty; Polhemus, Colchester, VT, USA) at 120 Hz. This system consists of a transmitter, five sensors, and a digitizing stylus operated by an electronic unit. The transmitter was fixed on a rigid wooden board at a height of 40 cm from the floor and 30 cm behind the subjects. An electromagnetic field was generated by the transmitter, and was sensed by these sensors and the stylus. This electromagnetic field represented the global coordinate system, with the X-axis pointing forward, the Y-axis pointing upward, the Z-axis pointing to the right, and the origin located at the transmitter. Next, the sensors were attached to the bony landmarks of the subjects with adhesive tape. The thoracic sensor was placed on the sternum just inferior to the jugular notch, the humeral sensor was placed on the middle point of the humerus with a thermoplastic cuff, and the scapular sensor was placed on the flat surface of acromion. Based on these sensors' placement, the local coordinate system of the thorax, humerus, and scapula were established by digitizing each bony landmark. All definitions of the local coordinate system were in accordance with the shoulder standardization proposal of the International Society of Biomechanics,³³ and the glenohumeral rotation center in the humeral segment was defined with reference to the previous study.²³

The rotation of the distal coordinate system was described with respect to the proximal coordinate system, according to the Euler angle of the International Society of Biomechanics.³³

Scapular kinematics and pectoralis minor stiffness

To describe the joint motion in correspondence with human kinesiology, the motion of the scapula around the Ys-axis was defined as internal rotation (positive) and external rotation (negative); the motion around the Xs-axis was defined as downward rotation (positive) and upward rotation (negative); the motion around Zs-axis was defined as posterior tilt (positive) and anterior tilt (negative); and the motion of the humerus around Xh-axis was defined as elevation (positive) (Figure 2). These motions were calculated using MATLAB (The Math Works, Natick, MA, USA). The scapular rotation was measured in every 10° of humeral elevation relative to the thorax, from 30° to 120° of humeral elevation. These angles were selected because the previous study¹² reported that there was little influence of the artifact of soft tissue on measuring the scapular motion in humeral elevations of less than 120° using a surface method. The elevation was examined three times, and the mean value was used for analysis.

Muscle Stiffness

The PMi stiffness was measured before stretching, immediately after stretching, and after arm elevation using ultrasonic shear wave elastography (Aixplorer, SuperSonic Imagine, Aix-en-Provence, France) with an ultrasound transducer (SL15-4: 4 to 15 MHz linear probe) (Figure 3). The ultrasonic shear wave elastography monitors the propagation of shear waves generated in tissue using acoustic radiation forces, and is able to evaluate the tissue elasticity of individual muscles.²⁸ The shear elastic modulus of the muscle represents muscle stiffness, and has been used as a quantitative indicator of the stretching effect in many previous studies.^{24,29,34} The shear elastic modulus (G) was calculated from the shear wave propagation speed (V) generated by the transducer using the formula of $G = \rho V^2$, in which ρ is the muscle density (1,000 kg/m³). The validity of applying the shear wave elastography to evaluate the skeletal muscle was reported in a previous study.⁸ The stiffness was measured three times in each session, and the

Scapular kinematics and pectoralis minor stiffness

mean value was used for analysis. All calculation of the stiffness was blinded by anonymizing the ultrasonic image, and a region of interest of the shear modulus was carefully chosen as large as possible with exclusion of subcutaneous adipose tissues and aponeuroses.

For the reliability study, the PMi stiffness was measured in ten healthy men (age, 24.9±1.5 years; height, 171.7±6.5.cm; weight, 70.4±7.8 kg) prior to this study. Following the completion of the measurement method mentioned above, the stiffness was measured three times with sufficient rest interval. The intra-observer reliability of the ultrasonic measurement was confirmed using the intraclass correlation coefficient (1,3) ($ICC_{1,3}$) with 95% confidence interval (95% CI). $ICC_{1,3}$ was 0.99 (95% CI: 0.97-0.99). A previous study investigating the reliability coefficient reported that a range from 0.81 to 1.00 was “almost perfect” reproducibility.¹⁶ The measurement of PMi stiffness in our study, therefore, was considered to be reproducible.

Data analysis

Only those subjects whose decrease in muscle stiffness remained until after arm elevation were analyzed in this study. We focused on the effect of the decrease in PMi stiffness on the scapular motion but not the PMi stretching, so that we could examine the direct relationship between the PMi stiffness and scapular motion. Statistical analysis was performed with IBM SPSS Statistical software (version 22; IBM, Armonk, NY, USA).

Regarding the stiffness in raw data before stretching, a paired *t*-test was performed to compare the interventional and the control limbs. The amount of change in the stiffness was calculated by subtracting the value of the stiffness before stretching from that present immediately after stretching or after elevation. For the change in stiffness, a two-way analysis of variance (ANOVA) with repeated measures on two factors [limb (two levels, interventional limb; control limb) × time (two levels, immediately after stretching - before stretching; after

Scapular kinematics and pectoralis minor stiffness

arm elevation - before stretching)] was used to demonstrate that the stretching decreased the PMi stiffness. If a significant main effect was found, then a Bonferroni post hoc test was performed. A confidence level of .05 was used in all statistics tests. Cohen's *d* values were also reported as the effect size, with the values of 0.2, 0.5, and 0.8 considered to elicit small, moderate, and large effects, respectively.⁶

The amount of change in the scapular motion was calculated by subtracting the value of scapular motion before stretching from that of it after stretching. For the change in each scapular motion (i.e. internal/external rotation; downward/upward rotation; posterior/anterior tilt), a two-way ANOVA with repeated measures on two factors [limb (two levels, interventional limb; control limb) $\times$ angle (ten levels, every 10° from 30° to 120°)] was used to determine the effects of the change in PMi stiffness on the scapular motion during elevation. If a significant interaction was found, then a paired *t*-test for post hoc test was performed to compare the interventional limb with the control limb with respect to each angle. If a significant main effect of the side was also found, then a Bonferroni post hoc test was performed to compare the interventional limb with the control limb. The significant main effect of the angle was ignored because the present study was interested in the comparison between the interventional limb and the control limb.

Scapular kinematics and pectoralis minor stiffness

Results

In the dominant limb, the decrease in the PMi stiffness immediately after stretching and after arm elevation occurred in fifteen men (age, 24.9 ± 3.3 years; height, 171.9 ± 5.9 cm; weight, 67.2 ± 8.4 kg); therefore, the results of these fifteen men (thirty shoulders) are shown below.

The paired *t*-test showed no significant differences between the interventional limb and the control limb in stiffness before stretching ($P = .063$, 95%IC: $-0.24 - 7.98$). For the amount of change in the stiffness, a two-way ANOVA indicated a significant main effect of the limb but not the time, with no significant interaction between the limb and the time (Table 1).

The raw value and the amount of change in scapular motion for abduction are shown in Table 2. For the amount of change in the internal/external rotation of scapula, a two-way ANOVA showed a significant interaction between the limb and the angle ($F = 4.519$, $P = .029$). Then, a post hoc test indicated that the change in the interventional limb was significantly greater than that in the control limb from 40° to 120° abduction ($P = .001 - .014$), and an increase in external rotation in interventional limb was found after stretching. For the amount of change in the downward/upward rotation of the scapula, there were no significant interactions between the limb and the angle ($F = 0.345$, $P = .726$), and no significant main effect in the limb ($F = 0.129$, $P = .725$). For the amount of change in the posterior/anterior tilt of scapula, a two-way ANOVA showed no significant interaction between the limb and the angle ($F = 0.891$, $P = .378$), but a significant main effect in the limb was seen ($F = 4.966$, $P = .043$). A post hoc test indicated that the change in the interventional limb was significantly greater than that in the control limb ($P = .043$), and that the posterior tilt in the interventional limb increased after stretching throughout the full evaluated range.

The raw value and the amount of change in scapular motion for scaption are shown in Table 3. For the amount of change in the internal/external rotation of scapula, a two-way ANOVA showed a significant interaction between the limb and the angle ($F = 6.655$, $P = .004$).

Scapular kinematics and pectoralis minor stiffness

238 Then, a post hoc test indicated that the amount of change from 40° to 120° in the interventional
239 limb was significantly greater than that in the control limb ($P = .001 - .034$), and that the external
240 rotation in the interventional limb increased after stretching. For the amount of change in the
241 downward/upward rotation of scapula, there were no significant interactions between the limb
242 and the angle ($F = 0.295$, $P = .750$), and no significant main effect in the limb ($F = 0.006$, P
243 $= .940$). For the amount of change in the posterior/anterior tilt of scapula, a two-way ANOVA
244 showed a significant interaction between the limb and the angle ($F = 4.397$, $P = .032$). Then, a
245 post hoc test indicated that the amount of change from 50° to 120° in the interventional limb
246 was significantly greater than in the control limb ($P = .006 - .035$), and the posterior tilt in the
247 interventional limb increased after stretching.

248

Scapular kinematics and pectoralis minor stiffness

249 Discussion

250 The present study investigated the effects of PMi stiffness on 3D scapular motion during arm
251 elevation, and found a decrease in PMi stiffness and an increase in external rotation and
252 posterior tilt of the scapula after stretching. These results indicate that the alteration in scapular
253 motion in combination with decrease in PMi stiffness occurred after stretching, and accorded
254 with our hypothesis. To the best of our knowledge, this is the first study to demonstrate that an
255 acute decrease in PMi stiffness after stretching changes the 3D scapular motion during arm
256 elevation.

257 The decrease in the PMi stiffness occurred immediately after stretching and lasted until
258 after arm elevation. The previous study²⁴ showed a positive correlation between the rate of
259 change in the shear elastic modulus and the rate of change in muscle stiffness, and therefore,
260 the decrease seen in shear elastic modulus after stretching indicates a decrease in muscle
261 stiffness (a so-called increase in the muscle flexibility).^{1,9,30,34} Therefore, the decrease in the
262 PMi stiffness immediately after stretching and after arm elevation confirmed that the stretching
263 used in the current study was sufficient in decreasing PMi stiffness until the end of the
264 evaluation.

265 To the best of our knowledge, there is only one study that investigated the relationship
266 between the acute change in the flexibility of PMi and 3D scapular motion. Williams et al.³²
267 measured the PMi length (coracoid process to forth rib) and the scapular kinematics before and
268 after two types of stretching, focused stretch or gross stretch, for one minute (30 seconds, two
269 repetitions, 30-second intervals). They³² concluded that there are no changes in the scapular
270 kinematics after either form of stretching, which was inconsistent with our results, which noted
271 that a change in scapular motion occurred after PMi stretching. This discrepancy between the
272 previous study and our results could be attributed to the duration time and the index of
273 stretching effect. Among various studies on stretching duration, there is a previous study²⁵ that

Scapular kinematics and pectoralis minor stiffness

examined the minimum time required for stretching to change the passive property. Nakamura et al.²⁵ concluded that stretching for more than two minutes was recommended to decrease the passive property of the gastrocnemius muscle. Therefore, though it is necessary to consider the difference in the muscles studied, it is possible that the stretching duration of the PMi was insufficient to elicit a change in scapular kinematics in Williams's study. Furthermore, there is also a possibility the length of the PMi may not be sufficient to represent PMi flexibility due to the bias of skins, soft tissues, and/or posture. In contrast, we measured the PMi stiffness quantitatively using the shear elastic modulus measured by ultrasonic shear wave elastography. Therefore, our study might be more valid than their study in investigating the relation of the PMi stiffness and the scapular motion.

These results mean that the decrease in the PMi stiffness after stretching induced the external rotation and posterior tilt of scapula. Generally, the scapula rotates upward and externally and tilts posteriorly during arm elevation in healthy individuals.^{12,18,22} The PMi tightness might cause the internal and downward rotation and anterior tilt of scapula from an anatomic perspective.^{5,20,26} Borstad et al.⁴ showed that the external rotation and posterior tilt in individuals with shortened PMi muscles were decreased when compared with individuals with long PMi during arm elevation. Therefore, considering these studies, our results suggested that the scapular external rotation and posterior tilt increased in abduction and scaption with a decrease in PMi stiffness. For the posterior tilt in the interventional limb group after stretching, there were differences in the amount of change between abduction and scaption. These differences in behavior, which the increase in the posterior tilt and occurred from 30° to 120° in abduction and from 50° to 120° in scaption, might depend on the difference in the plane of elevation. Compared with scaption, abduction needs a slightly greater posterior tilt of the scapula, due to the difference of the plane although there are no significant differences.¹⁹ Therefore, it is rational that the posterior tilt of the scapula may arise from the early phase of

Scapular kinematics and pectoralis minor stiffness

abduction by the decrease in PMi stiffness.

The maximum amount of change in external rotation and the posterior tilt were 4.8° and 3.3° in abduction, and 4.5° and 3.7° in scaption. Ludewig and Cook¹⁷ reported that individuals with shoulder impingement had increased internal rotation, decreased upward rotation, and decreased posterior tilt during arm elevation. The previous¹⁷ study suggested that 4° to 6° of change in scapular motion is important in narrowing of the subacromial space and the occurrence of impingement, because individuals with impingement syndrome showed increased internal rotation of 5.2° , decreased upward rotation of 4.1° , and increased anterior tilt of 5.8° as compared with healthy individuals. We believe that our findings are of clinical relevance because the change in scapular motion after stretching shown in the present study approximately correspond with the range of changes seen in scapular motion between healthy individuals and those with impingement syndrome indicated in Ludewig and Cook's study. The change in the scapular motion after the stretching observed in the current study may be of clinical significance; however, further research is warranted to validate these theories within pathologic subjects.

However, when interpreting our findings, one should note the following: first, the subjects were all healthy men, as prescribed by the exclusion criteria. Therefore, it is unclear whether the findings can be generalized to individuals with impingement syndrome. Second, the stiffness of only the PMi was measured among the shoulder girdle muscles. Therefore, this study does not exactly promise that only a decrease in the PMi stiffness changed the scapular kinematics, and it is possible that other muscles such as the pectoralis major muscle, the subscapularis muscle, or glenohumeral ligaments and capsules were also stretched and thus had an effect. This is the limitation of a in-vivo study. Third, the current study investigated only the acute effect of the PMi stretching on the scapular motion, so its long-term effect is unknown. The recent study examining the effects of self-stretching of the PMi for six weeks on the

Scapular kinematics and pectoralis minor stiffness

324 scapular kinematics concluded that stretching did not change PMi length and scapular
325 kinematics in individuals with and without shoulder pain.²⁷ Therefore, future study should
326 evaluate the long-term effects of the therapist-applied PMi stretching on muscle stiffness and
327 scapular motion.
328

Scapular kinematics and pectoralis minor stiffness

329 **Conclusion**

330 We investigated the effects of PMi stiffness after stretching on the change in scapular motion
331 during arm elevation. Our results indicated a decrease in PMi stiffness, increased external
332 rotation and posterior tilt of the scapula occurred during arm elevation after stretching. These
333 findings might be relevant knowledge for the approach to scapular dyskinesis and in further
334 studies.

335

Scapular kinematics and pectoralis minor stiffness

References

1. Akagi R, Takahashi H. Acute effect of static stretching on hardness of the gastrocnemius muscle. *Med Sci Sports Exerc* 2013;45(7):1348-1354. doi:10.1249/MSS.0b013e3182850e17.
2. Borstad JD, Ludewig PM. Comparison of scapular kinematics between elevation and lowering of the arm in the scapular plane. *Clin Biomech* 2002;17:650-659. doi:10.1016/S0268-0033(02)00136-5.
3. Borstad JD, Ludewig PM. Comparison of three stretches for the pectoralis minor muscle. *J Shoulder Elbow Surg* 2006;15(3):324-330. doi:10.1016/j.jse.2005.08.011.
4. Borstad JD, Ludewig PM. The effect of long versus short pectoralis minor resting length on scapular kinematics in healthy individuals. *J Orthop Sports Phys Ther* 2005;35(4):227-238. doi:10.2519/jospt.2005.1669.
5. Burkhart SS, Morgan CD, Ben Kibler W. The disabled throwing shoulder: Spectrum of pathology Part I: Pathoanatomy and biomechanics. *Arthrosc - J Arthrosc Relat Surg* 2003;19(4):404-420. doi:10.1053/jars.2003.50128.
6. Cohen J. Statistical Power Analysis. *Current Directions in Psychological Science* 1992; 1(3):98-101
7. Cools AMJ, Struyf F, De Mey K, Maenhout A, Castelein B, Cagnie B. Rehabilitation of scapular dyskinesis: from the office worker to the elite overhead athlete. *Br J Sports Med* 2014;48(8):692-697. doi:10.1136/bjsports-2013-092148.
8. Eby SF, Song P, Chen S, Chen Q, Greenleaf JF, An KN. Validation of shear wave elastography in skeletal muscle. *J Biomech* 2013;46(14):2381-2387. doi:10.1016/j.jbiomech.2013.07.033
9. Ichihashi N, Umegaki H, Ikezoe T, Nakamura M, Nishishita S, Fujita K, et al. The effects of a 4-week static stretching programme on the individual muscles

Scapular kinematics and pectoralis minor stiffness

- 361 comprising the hamstrings. J Sports Sci 2016;34(23):2155-2159.
- 362 doi:10.1080/02640414.2016.1172725.
- 363 10. Inman VT, Saunders FRCS, Abbott LC. Observations on the function of the shoulder
- 364 joint. J Bone Jt Surg Am 1944;26(1):1-30.
- 365 11. Johnson GR, Stuart PR, Mitchell S. A method for the measurement of three-
- 366 dimensional scapular movement. Clin Biomech 1993;8(5):269-273. doi:10.1016/0268-
- 367 0033(93)90037-I.
- 368 12. Karduna AR, McClure PW, Michener LA, Sennett B. Dynamic measurements of
- 369 three-dimensional scapular kinematics: a validation study. J Biomech Eng
- 370 2001;123(2):184-190. doi:10.1115/1.1351892.
- 371 13. Kibler WB, Ludewig PM, McClure PW, Michener LA, Bak K, Sciascia AD. Clinical
- 372 implications of scapular dyskinesis in shoulder injury: the 2013 consensus statement
- 373 from the “Scapular Summit”. Br J Sports Med 2013;47(14):877-885.
- 374 doi:10.1136/bjsports-2013-092425.
- 375 14. Kibler WB. The Role of the Scapula in Athletic Shoulder Function. Am J Sports Med
- 376 1998;26(2):325-337. doi:10.1177/03635465980260022801
- 377 15. Kijima T, Matsuki K, Ochiai N, Yamaguchi T, Sasaki Y, Hashimoto E, et al. In vivo
- 378 3-dimensional analysis of scapular and glenohumeral kinematics: comparison of
- 379 symptomatic or asymptomatic shoulders with rotator cuff tears and healthy
- 380 shoulders. J Shoulder Elb Surg 2015;24(11):1817-1826. doi:10.1016/j.jse.2015.06.003.
- 381 16. Landis JR, Koch GG. The measurement of observer agreement for categorical data.
- 382 Biometrics 1977;33(1):159-174.
- 383 17. Ludewig PM, Cook TM. Alterations in shoulder kinematics and associated muscle
- 384 activity in people with symptoms of shoulder impingement. Phys Ther
- 385 2000;80(3):276-291.

Scapular kinematics and pectoralis minor stiffness

- 386 18. Ludewig PM, Cook TM, Nawoczenski D a. Three-dimensional scapular orientation
387 and muscle activity at selected positions of humeral elevation. J Orthop Sports Phys
388 Ther 1996;24(2):57-65. doi:10.2519/jospt.1996.24.2.57.
- 389 19. Ludewig PM, Phadke V, Braman JP, Hassett DR, Cieminski CJ, LaPrade RF. Motion
390 of the shoulder complex during multiplanar humeral elevation. J Bone Joint Surg Am
391 2009;91(2):378-389. doi:10.2106/JBJS.G.01483.
- 392 20. Ludewig PM, Reynolds JF. The association of scapular kinematics and glenohumeral
393 joint pathologies. J Orthop Sports Phys Ther 2009;39(2):90-104.
394 doi:10.2519/jospt.2009.2808.
- 395 21. Matias R, Pascoal AG. The unstable shoulder in arm elevation: A three-dimensional
396 and electromyographic study in subjects with glenohumeral instability. Clin Biomech
397 2006;21:S52-S58. doi:10.1016/j.clinbiomech.2005.09.014.
- 398 22. McClure PW, Michener LA, Sennett BJ, Karduna AR. Direct 3-dimensional
399 measurement of scapular kinematics during dynamic movements in vivo. J Shoulder
400 Elb Surg 2001;10(3):269- 277. doi:10.1067/mse.2001.112954.
- 401 23. Murray IA. Determining upper limb kinematics and dynamics during everyday tasks.
402 PhD Thesis, Cent Rehabil Eng Stud Univ Newcastle upon Tyne 1999;1-470.
403 <https://theses.ncl.ac.uk/dspace/bitstream/10443/185/1/murray99.pdf>.
- 404 24. Nakamura M, Ikezoe T, Kobayashi T, Umegaki H, Takeno Y, Nishishita S, et al.
405 Acute effects of static stretching on muscle hardness of the medial gastrocnemius
406 muscle belly in humans: An ultrasonic shear-wave elastography study. Ultrasound
407 Med Biol 2014;40(9):1991-1997. doi:10.1016/j.ultrasmedbio.2014.03.024.
- 408 25. Nakamura M, Ikezoe T, Takeno Y, Ichihashi N. Time course of changes in passive
409 properties of the gastrocnemius muscle-tendon unit during 5 min of static stretching.
410 Man Ther 2013;18(3):211-215. doi:10.1016/j.math.2012.09.010.

Scapular kinematics and pectoralis minor stiffness

- 411 26. Novak CB, Mackinnon SE. Repetitive use and static postures: A source of nerve
412 compression and pain. *J Hand Ther* 1997;10(2):151-159. doi:10.1016/S0894-
413 1130(97)80069-5.
- 414 27. Rosa DP, Borstad JD, Pogetti LS, Camargo PR. Effects of a stretching protocol for
415 the pectoralis minor on muscle length, function, and scapular kinematics in
416 individuals with and without shoulder pain. *J Hand Ther* 2017;30(1):20-29.
417 doi:10.1016/j.jht.2016.06.006.
- 418 28. Shiina T, Nightingale KR, Palmeri ML, et al. WFUMB guidelines and
419 recommendations for clinical use of ultrasound elastography: Part 1: basic principles
420 and terminology. *Ultrasound Med Biol* 2015;41(5):1126-1147.
421 doi:10.1016/j.ultrasmedbio.2015.03.009.
- 422 29. Taniguchi K, Shinohara M, Nozaki S, Katayose M. Acute decrease in the stiffness of
423 resting muscle belly due to static stretching. *Scand J Med Sci Sports* 2015;25(1):32-
424 40. doi:10.1111/sms.12146.
- 425 30. Umehara J, Hasegawa S, Nakamura M, Nishishita S, Umegaki H, Tanaka H, et al.
426 Effect of scapular stabilization during cross-body stretch on the hardness of
427 infraspinatus, teres minor, and deltoid muscles: An ultrasonic shear wave
428 elastography study. *Musculoskelet Sci Pract* 2017;27:91-96.
429 doi:10.1016/j.math.2016.10.004.
- 430 31. Umehara J, Nakamura M, Fujita K, Kusano K, Nishishita S, Araki K, et al. Shoulder
431 horizontal abduction stretching effectively increases shear elastic modulus of
432 pectoralis minor muscle. *J Shoulder Elb Surg* 2017; 26(7):1159-1165.
433 doi:10.1016/j.jse.2016.12.074.
- 434 32. Williams JG, Laudner KG, McLoda T. The acute effects of two passive stretch
435 maneuvers on pectoralis minor length and scapular kinematics among collegiate

Scapular kinematics and pectoralis minor stiffness

- 436 swimmers. *Int J Sports Phys Ther* 2013;8(1):25-33.
- 437 33. Wu G, van der Helm FC, Veeger HE, Makhsous M, Van Roy P, Anglin C, et al. ISB
438 recommendation on definitions of joint coordinate systems of various joints for the
439 reporting of human joint motion—Part II: shoulder, elbow, wrist and hand. *J*
440 *Biomech* 2005;38(5):981-992. doi:10.1016/j.jbiomech.2004.05.042.
- 441 34. Yamauchi T, Hasegawa S, Nakamura M, Nishishita S, Yanase K, Fujita K, et al.
442 Effects of two stretching methods on shoulder range of motion and muscle stiffness
443 in baseball players with posterior shoulder tightness: a randomized controlled trial. *J*
444 *shoulder Elb Surg* 2016;25(9):1395-1403. doi:10.1016/j.jse.2016.04.025.
- 445

Scapular kinematics and pectoralis minor stiffness

Figure and Table Legends

Figure 1; Stretching of the pectoralis minor muscle. As directed, the subject sat on the wooden stool and the interventional limb was brought to maximal horizontal abduction and external rotation at an arm elevation of 150° with the elbow in 90° flexion, and was subsequently maximally externally rotated by the investigator. During the stretching, the participants was instructed to remain relaxed. The investigator operated the upper limb of the subject using one hand and held the trunk using the other hand.

Figure 2; The definition of coordinate systems and motions relative to the thorax for the scapula. The scapula are seen in the posterior view of the right shoulder.

Figure 3; Posture and measurement site of the pectoralis minor stiffness are shown in A. The participant sat on a wooden stool with their arm relaxed on a platform in a position with 90° of shoulder abduction and 90° of elbow flexion. The measurement site was defined as the midpoint between the coracoid process and the fourth rib-sternum junction. The probe was placed parallel to the muscle fascicle of the PMi on the ultrasonic image as B. The participant was instructed to hold their breath during measurement to prevent elongation of the PMi due to the motion of rib cage. PMA, pectoralis major muscle; PMi, pectoralis minor muscle; IM, intercostal muscle.

Table 1; Pectoralis minor muscle stiffness (kPa) pre-assessment and post assessment.

Table 2; Raw value and amount of change in scapular motion for abduction.

Table 3; Raw value and amount of change in scapular motion for scaption.

Scapular kinematics and pectoralis minor stiffness

471 Figure 1

472

473

Scapular kinematics and pectoralis minor stiffness

474 Figure 2

Scapular kinematics and pectoralis minor stiffness

478 Figure 3

479

480

Scapular kinematics and pectoralis minor stiffness

Table 1

Pectoralis minor stiffness (kPa)		Interventional limb	Control limb	Statistical significance
Before-stretching	Raw value	12.7±3.6	11.4±3.8	
	Change	-	-	
	ES (<i>d</i>)		-	
Immediately after-stretching	Raw value	9.4±2.2	11.0±3.4	Interaction F = 1.935; P = .186
	Change	-3.2±2.0	-0.4±2.4	
	ES (<i>d</i>)		1.2	Main effect Limb: F = 14.140; P = .002 Time: F = 0.869; P = .367
After-arm elevation	Raw value	10.2±2.5	10.8±3.6	
	Change	-2.5±1.9	-0.6±2.0	
	ES (<i>d</i>)		0.9	

Change, amount of change between before and after stretching, and before and after arm elevation. Value ± standard deviation; ES, effect size.

Scapular kinematics and pectoralis minor stiffness

Table 2

		Internal/External rotation(°)			Downward/Upward rotation(°)			Posterior/Anterior tilt(°)		
		IN	CON	ES	IN	CON	ES	IN	CON	ES
E l e v a t i o n a n g l e (o)	Before	12.1±4.5	17.3±6.6		-5.5±4.8	-5.2±3.9		-4.9±5.9	-2.9±4.7	
	30 After	10.2±5.4	16.4±7.7	0.5	-6.9±5.5	-5.7±3.8	0.4	-3.9±6.8	-2.7±5.0	0.6
	Change	-1.9±2.2	-1.0±1.8		-1.4±2.8	-0.5±1.1		0.9±1.5	0.2±0.7	
	Before	10.5±4.0	15.7±6.4		-9.8±4.8	-9.6±4.1		-3.3±6.2	-1.7±4.9	
	40 After	8.3±4.9	14.7±7.1	0.6	-10.9±6.0	-10.5±3.9	0.1	-2.3±7.1	-1.7±5.2	0.7
	Change	-2.3±2.4*	-1.0±1.6		-1.1±3.0	-0.9±1.2		1.0±1.6	0.0±0.8	
	Before	9.6±4.4	14.4±6.6		-14.0±5.2	-14.3±4.4		-1.6±6.2	0.0±4.9	
	50 After	6.7±5.7	13.5±7.4	0.8	-15.4±6.7	-15.2±4.6	0.1	-0.3±7.3	0.0±5.3	0.8
	Change	-2.9±3.0†	-0.9±1.5		-1.4±3.5	-0.9±1.2		1.3±1.8	0.02±0.9	
	Before	9.1±4.6	13.3±6.4		-18.5±5.4	-18.9±4.6		0.3±6.6	2.0±5.0	
	60 After	5.9±6.0	12.1±7.0	0.8	-19.6±6.9	-19.7±4.7	0.1	1.9±7.7	2.2±5.4	0.8
	Change	-3.3±3.0†	-1.2±1.3		-1.1±3.6	-0.8±1.5		1.6±2.0	0.2±1.2	
	Before	9.1±5.1	12.4±6.7		-21.2±12.2	-23.4±4.7		2.4±6.9	4.0±5.3	
	70 After	5.7±6.5	11.2±7.0	0.8	-23.7±7.5	-23.9±5.2	0.1	4.2±8.0	4.5±6.0	0.6
	Change	-3.4±3.2†	-1.2±1.3		-0.9±4.1	-0.5±1.5		1.8±2.4	0.5±1.3	
	Before	9.4±5.4	12.1±6.9		-22.8±5.6	-27.7±5.3		4.5±6.7	6.3±5.5	
	80 After	6.0±7.0	11.1±7.5	0.7	-27.6±7.1	-28.1±5.8	0.1	6.7±7.8	7.1±6.1	0.6
	Change	-3.4±3.6†	-1.0±1.4		-0.7±3.9	-0.4±1.6		2.2±2.7	0.9±1.3	
	Before	9.8±6.0	12.3±7.5		-30.9±5.4	-31.6±5.7		6.4±6.5	8.2±5.4	
	90 After	6.4±7.4	11.2±8.1	0.7	-31.4±7.3	-31.7±6.2	0.1	9.1±7.7	9.3±6.1	0.6
	Change	-3.4±3.7*	-1.0±1.4		-0.6±4.2	-0.05±1.6		2.7±3.0	1.1±1.4	
	Before	10.6±6.5	12.8±8.4		-34.6±5.5	-35.0±6.1		8.0±6.1	9.8±5.4	
	100 After	6.7±7.9	11.6±9.1	0.8	-35.0±7.8	-35.1±6.8	0.1	11.0±7.4	11.0±6.3	0.6
	Change	-3.9±3.9†	-1.2±1.4		-0.4±4.2	-0.1±1.6		2.9±3.4	1.2±1.3	
	Before	11.8±7.2	13.9±9.4		-38.0±5.8	-38.1±6.5		9.3±5.7	10.8±5.5	
	110 After	7.5±8.5	12.6±9.9	0.9	-38.4±4.4	-38.1±7.1	0.1	12.6±7.1	12.0±6.0	0.7
	Change	-4.4±4.1†	-1.3±1.6		-0.3±4.3	-0.03±1.6		3.3±3.4	1.1±1.3	
	Before	14.0±7.8	15.9±10.5		-41.1±6.1	-40.9±6.7		9.9±5.7	11.1±5.8	
	120 After	9.1±9.2	14.9±11.3	1.0	-41.4±9.0	-41.0±7.2	<0.1	13.2±7.4	12.2±6.5	0.7
	Change	-4.8±4.4‡	-1.0±1.8		-0.3±4.5	-0.07±1.4		3.3±3.6	1.2±1.4	

IN, interventional limb; CON, control limb; ES, effect size of amount of change; Before, raw value before stretching; After, raw value after stretching; Change, amount of change between before and after stretching. Values are expressed as mean ± standard deviation. The *asterisk* indicates that the change in scapular motion in the interventional limb is significantly ($P < .05$) greater than it in the control limb; *single dagger* indicates that it in the interventional limb is significantly ($P < .01$) greater than it in the control limb; and *double dagger*

Scapular kinematics and pectoralis minor stiffness

496 indicates that it in the interventional limb is significantly ($P < .001$) greater than it in the control limb.

497

498

Scapular kinematics and pectoralis minor stiffness

Table 3

		Internal/External rotation (°)			Downward/Upward rotation (°)			Posterior/Anterior tilt (°)		
		IN	CON	ES	IN	CON	ES	IN	CON	ES
Eleva- tion (°)	Before	21.3±3.8	26.1±5.6		-3.7±6.1	-3.7±3.4		-4.5±5.6	-2.9±5.0	
	30 After	20.7±4.0	25.8±6.7	0.3	-3.6±7.0	-3.4±3.7	0.1	-3.6±6.3	-2.4±5.6	0.3
	Change	-1.1±2.9	-0.3±2.1		0.1±2.5	0.3±0.9		0.9±1.4	0.5±0.9	
	Before	21.0±3.6	25.6±5.7		-7.5±6.1	-7.9±3.8		-3.5±5.8	-1.8±5.0	
	40 After	19.6±6.7	25.3±6.8	0.5	-7.6±7.1	-7.7±4.0	0.1	-2.7±6.4	-1.3±5.7	0.3
	Change	-1.4±2.9*	-0.3±1.8		-0.1±2.8	0.3±0.9		0.8±1.5	0.5±1.0	
	Before	20.9±3.5	25.0±5.5		-12.1±6.0	-12.6±4.1		-2.2±5.8	-0.6±5.2	
	50 After	19.2±4.0	24.7±6.8	0.5	-12.3±7.0	-12.6±4.4	0.1	-0.9±6.5	-0.1±5.8	0.7
	Change	-1.6±2.9*	-0.3±2.0		-0.1±3.0	0.1±1.2		1.2±1.3*	0.4±1.0	
	Before	20.8±3.5	24.4±5.7		-17.0±6.0	-17.6±4.1		-0.8±6.1	0.9±5.0	
	60 After	18.9±4.4	24.2±6.9	0.6	-17.3±7.3	-17.9±4.5	0.2	0.5±6.8	1.3±5.8	0.8
	Change	-1.9±3.2†	-0.2±2.2		-0.3±3.5	-0.3±1.5		1.3±1.4*	0.3±1.3	
	Before	20.8±3.5	23.5±5.8		-21.8±5.9	-22.4±4.3		0.6±6.3	2.6±5.2	
	70 After	18.5±4.8	23.0±7.3	0.6	-22.4±7.4	-22.8±5.0	0.1	2.0±6.9	3.1±5.8	0.7
	Change	-2.3±3.4†	-0.5±2.3		-0.6±3.8	-0.4±1.6		1.5±1.6*	0.4±1.0	
	Before	20.3±4.0	22.8±6.2		-26.2±11.1	-26.9±4.8		2.2±6.5	4.4±5.3	
	80 After	17.7±5.5	22.4±7.4	0.7	-26.9±7.7	-27.6±5.5	< 0.1	4.3±7.3	4.9±5.9	1.0
	Change	-2.6±3.4†	-0.4±2.3		-0.7±3.9	-0.7±1.6		2.1±1.8*	0.5±1.2	
	Before	20.1±4.5	22.2±6.6		-30.1±6.0	-30.9±5.2		3.8±6.7	6.2±5.4	
	90 After	17.0±6.6	21.6±7.7	0.7	-31.0±8.0	-31.7±6.1	< 0.1	6.6±7.7	6.8±6.0	1.1
	Change	-3.0±3.9†	-0.6±2.3		-0.9±4.0	-0.8±1.4		2.8±2.2†	0.7±1.3	
	Before	20.1±5.0	21.7±7.1		-33.6±6.1	-34.2±5.6		5.5±6.7	7.9±5.5	
	100 After	16.8±6.9	21.2±8.0	0.8	-34.6±8.3	-35.3±6.8	< 0.1	8.8±8.1	8.9±6.2	1.0
	Change	-3.3±4.1†	-0.5±2.3		-1.0±4.3	-1.0±1.5		3.3±2.6*	1.0±1.5	
	Before	20.1±5.8	21.4±7.7		-37.1±6.3	-37.3±6.1		7.1±6.6	9.2±5.7	
	110 After	16.3±8.0	20.5±8.8	0.8	-38.1±8.4	-38.3±7.1	< 0.1	10.9±8.4	10.5±6.6	0.9
	Change	-3.8±4.3†	-0.8±2.4		-1.0±4.4	-1.0±1.3		3.8±3.0*	1.3±1.6	
	Before	20.4±6.6	21.5±8.9		-40.4±6.6	-40.1±6.5		8.3±6.4	10.1±6.1	
	120 After	15.9±8.6	20.1±10.2	0.7	-41.0±8.9	-41.1±7.6	0.1	12.0±8.3	11.5±7.4	0.8
	Change	-4.5±4.9†	-1.4±2.5		-0.6±4.7	-1.0±1.4		3.7±3.1*	1.5±1.8	

IN, interventional limb; CON, control limb; ES, effect size of amount of change; Before, raw value before stretching; After, raw value after stretching; Change, amount of change between before and after stretching. Values are expressed as mean ± standard deviation. The *asterisk* indicates that the change in scapular motion in the interventional limb is significantly ($P < .05$) greater than it in the control limb; the *single dagger*

Scapular kinematics and pectoralis minor stiffness

506 indicates that it in the interventional limb is significantly ($P < .01$) greater than it in the control limb.
507
508