

Research



Cite this article: Toju H, Tanaka Y. 2019
Consortia of anti-nematode fungi and bacteria in
the rhizosphere of soybean plants attacked by
root-knot nematodes. *R. Soc. open sci.* **6**: 181693.
<http://dx.doi.org/10.1098/rsos.181693>

Received: 10 October 2018
Accepted: 21 February 2019

Subject Category:

Biology (whole organism)

Subject Areas:

ecology/environmental science/microbiology

Keywords:

disease-suppressive soil, *glycine max*,
meloidogyne, nematophagous fungi,
phytopathogenic pathogens and pests,
sustainable agriculture

Author for correspondence:

Hirokazu Toju
e-mail: toju.hirokazu.4c@kyoto-u.ac.jp

Electronic supplementary material is available
online at <https://dx.doi.org/10.6084/m9.figshare.c.4443020>.

Consortia of anti-nematode fungi and bacteria in the rhizosphere of soybean plants attacked by root-knot nematodes

Hirokazu Toju^{1,2} and Yu Tanaka^{2,3}

¹Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2133, Japan

²Precursory Research for Embryonic Science and Technology (PRESTO), Japan Science and Technology Agency, Kawaguchi, Saitama 332-0012, Japan

³Graduate School of Agriculture, Kyoto University, Kitashirakawa-oiwake-cho, Sakyo, Kyoto 606-8502, Japan

HT, 0000-0002-3362-3285

Cyst and root-knot nematodes are major risk factors of agroecosystem management, often causing devastating impacts on crop production. The use of microbes that parasitize or prey on nematodes has been considered as a promising approach for suppressing phytopathogenic nematode populations. However, effects and persistence of those biological control agents often vary substantially depending on regions, soil characteristics and agricultural practices: more insights into microbial community processes are required to develop reproducible control of nematode populations. By performing high-throughput sequencing profiling of bacteria and fungi, we examined how root and soil microbiomes differ between benign and nematode-infected plant individuals in a soybean field in Japan. Results indicated that various taxonomic groups of bacteria and fungi occurred preferentially on the soybean individuals infected by root-knot nematodes or those uninfected by nematodes. Based on a network analysis of potential microbe–microbe associations, we further found that several fungal taxa potentially preying on nematodes (*Dactylellina* (Orbiliiales), *Rhizophydium* (Rhizophydiales), *Clonostachys* (Hypocreales), *Pochonia* (Hypocreales) and *Purpureocillium* (Hypocreales)) co-occurred in the soybean rhizosphere at a small spatial scale. This study suggests how ‘consortia’ of anti-nematode microbes can derive from indigenous (resident) microbiomes, providing basic information for managing anti-nematode microbial communities in agroecosystems.

1. Introduction

Plant pathogenic nematodes, such as cyst and root-knot nematodes, are major threats to crop production worldwide [1,2]. Soybean fields, in particular, are often damaged by such phytopathogenic nematodes, resulting in substantial yield loss [3,4]. A number of chemical nematicides and biological control agents (e.g. nematophagous fungi in the genera *Purpureocillium* and *Clonostachys*) have been used to suppress nematode populations in farmlands [5,6]. However, once cyst and root-knot nematodes appear in a farmland, they often persist in the soil for a long time [7], causing high financial costs in agricultural management. Finding ways to suppress pathogenic nematode populations in agroecosystems is a key to reducing risk and management costs in production of soybean and other crop plants.

To reduce damage by cyst and root-knot nematodes, a number of studies have evaluated effects of crop varieties/species, crop rotations, fertilizer inputs and tillage intensity on nematode density in farmland soil [1,8–10]. However, the results of those studies varied considerably depending on regions, soil characteristics and complicated interactions among multiple factors (e.g. interactions between organic matter inputs and tillage frequency) [11]. Thus, it remains an important challenge to understand the mechanisms by which phytopathogenic nematode populations are suppressed in some farmland soils but not in others [12]. New lines of information are required for building general schemes for making agroecosystems robust against the emergence of pest nematodes.

Based on the technological advances in high-throughput DNA sequencing, more and more studies have examined structures of microbial communities (microbiomes) in order to evaluate biotic environmental conditions in the endosphere and rhizosphere of plants [13–16]. Recent studies have uncovered microbiome compositions of ‘disease-suppressive soils’, in which pests and pathogens damaging crop plants have been suppressed for long periods of time [17–19]. Some studies have further discussed how some microbes within such disease-suppressive microbiomes contribute to the health and growth of crop plant species [17,20,21]. In one of the studies, soil microbiome compositions were compared among soybean fields that differed in the density of cyst nematodes [12]. The study then revealed that bacteria and fungi potentially having negative impacts on nematode populations (e.g. *Purpureocillium* and *Pochonia*) were more abundant in the long term than in short-term monoculture fields of soybeans [12]. Such among-farmland comparisons have provided invaluable insights into ecosystem functions of indigenous (native) microbiomes. Nonetheless, the potential relationship between cropping system management and community processes of anti-nematode microbes remains obscured because the farmlands compared in those studies could vary in climatic and edaphic factors. Moreover, because incidence of cyst and root-knot nematodes generally varies at small spatial scales [22], there can be spatial heterogeneity in abundance and community compositions of anti-nematode bacteria and fungi within a farmland. Studies focusing on fine-scale assembly of anti-nematode microbes are required for developing agroecosystem management protocols for controlling phytopathogenic nematodes.

We conducted an Illumina sequencing analysis of bacteria and fungi in a soybean (*Glycine max*) field and then examined how root and rhizosphere microbiome structures varied among host plant individuals that differed in damage by root-knot nematodes (*Meloidogyne* sp.). Based on the data of microbiomes at a small spatial scale, we statistically explored microbial species/taxa that occurred preferentially in the roots or rhizosphere soil of nematode-infected soybean individuals. We further investigated the structure of networks depicting co-abundance patterns of microbial species/taxa within the soybean field, thereby examining whether multiple anti-nematode bacteria and fungi form consortia (assemblages) on/around the plant individuals infected by root-knot nematodes. Our results suggest that various taxonomic groups of anti-nematode bacteria and fungi are present within indigenous microbiomes. This study also suggests that microbiome assembly at fine spatial scales is a key to managing populations and communities of such functional microbes.

2. Methods

2.1. Sampling

Fieldwork was conducted at the soybean field on the Hokubu Campus of Kyoto University, Japan (35.033°N, 135.784°E). In the field, the soybean strain ‘Sachiyutaka’ was sown at 15 cm intervals in two lines (electronic supplementary material, figure S1) on 4 July 2016 (basal fertilizer, N:P₂O₅:K₂O = 3:10:10 g m⁻²). In the lines, 69 and 62 individuals (‘set 1’ and ‘set 2’, respectively),




Figure 1. Study site and soybeans. (a) Soybean field in which sampling was conducted. (b) Soybean states. Soybean individuals were classified into three categories: those heavily attacked by root-knot nematodes ('no leaf'; left), those exhibited normal growth ('green'; right) and those showing intermediate characters ('yellow'; middle). (c) Relationship between soybean states and biomass. Dry mass significantly differed among 'no leaf', 'yellow' and 'green' soybean individuals (ANOVA; $F_2 = 20.5$, $p < 0.0001$). (d) Spatial distribution of 'no leaf', 'yellow' and 'green' soybean individuals. Sampling sets 1 and 2 are shown separately.

respectively, were sampled at every other position (i.e. 30 cm intervals; figure 1) on 7 October 2016. The sampled soybean individuals were classified into three categories: normal individuals with green leaves (green), individuals with yellow leaves (yellow) and those with no leaves (no leaf) (figure 1a–c). Among them, 'green' individuals exhibited normal growth, while 'no leaf' individuals were heavily infected by root-knot nematodes: 'yellow' individuals showed intermediate characters. In total, 97 'green', 19 'yellow' and 15 'no leaf' individuals were sampled (figure 1d).

For each individual, two segments of 5-cm terminal roots and rhizosphere soil were collected from around 10 cm below the soil surface. The root and soil samples were transferred into a cool box in the field and then stored at -80°C until DNA extraction in the laboratory. The above-ground bodies of the individuals were placed in drying ovens at 80°C for 72 h to measure dry mass. The dry mass data indicated that 'green', 'yellow' and 'no leaf' soybean individuals differed significantly in their biomass (figure 1c).

2.2. DNA extraction, PCR and sequencing

The root segments of each individual were transferred to a 15 ml tube and washed in 70% ethanol by vortexing for 10 s. The samples were then transferred to a new 15 ml tube and then washed again in 70% ethanol by sonication (42 Hz) for 5 min. After an additional sonication wash in a new tube, one of the two root segments was dried and placed in a 1.2 ml tube for each soybean individual. DNA extraction was then performed with a cetyltrimethylammonium bromide method [23] after pulverizing the roots with 4 mm zirconium balls at 25 Hz for 3 min using a TissueLyser II (Qiagen).

For DNA extraction from the rhizosphere soil, the ISOIL for Beads Beating kit (Nippon Gene) was used as instructed by the manufacturer. For each sample, 0.5 g of soil was placed into a 2 ml microtube of the ISOIL kit. To increase the yield of DNA, 10 mg of skim milk powder (Wako, 198–10605) was added to each sample [24].

For each of the root and soil samples, the 16S rRNA V4 region of the prokaryotes and the internal transcribed spacer 1 (ITS1) region of fungi were amplified. The PCR of the 16S rRNA region was performed with the forward primer 515f [25] fused with 3–6-mer Ns for improved Illumina sequencing quality [26] and the forward Illumina sequencing primer (5'- TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG- (3–6-mer Ns) - (515f) -3') and the reverse primer 806rB [27] fused with 3–6-mer Ns and the reverse sequencing primer (5'- GTC TCG TGG GCT CGG AGA TGT GTA

TAA GAG ACA G (3–6-mer Ns) - (806rB) -3') (0.2 μ M each). To prevent the amplification of mitochondrial and chloroplast 16S rRNA sequences, specific peptide nucleic acids (mPNA and pPNA; [26]) (0.25 μ M each) were added to the reaction mix of KOD FX Neo (Toyobo). The temperature profile of the PCR was 94°C for 2 min, followed by 35 cycles at 98°C for 10 s, 78°C for 10 s, 50°C for 30 s, 68°C for 50 s and a final extension at 68°C for 5 min. To prevent generation of chimaeric sequences, the ramp rate through the thermal cycles was set to 1°C s⁻¹ [28]. Illumina sequencing adaptors were then added to respective samples in the supplemental PCR using the forward fusion primers consisting of the P5 Illumina adaptor, 8-mer indexes for sample identification [29] and a partial sequence of the sequencing primer (5'- AAT GAT ACG GCG ACC ACC GAG ATC TAC AC - (8-mer index) - TCG TCG GCA GCG TC -3') and the reverse fusion primers consisting of the P7 adaptor, 8-mer indexes and a partial sequence of the sequencing primer (5'- CAA GCA GAA GAC GGC ATA CGA GAT - (8-mer index) - GTC TCG TGG GCT CGG -3'). KOD FX Neo was used with a temperature profile of 94°C for 2 min, followed by eight cycles at 98°C for 10 s, 55°C for 30 s, 68°C for 50 s (ramp rate = 1°C s⁻¹) and a final extension at 68°C for 5 min. The PCR amplicons of the 131 soybean individuals were then pooled after a purification/equalization process with the AMPureXP Kit (Beckman Coulter). Primer dimers, which were shorter than 200 bp, were removed from the pooled library by supplemental purification with AMPureXP: the ratio of AMPureXP reagent to the pooled library was set to 0.6 (v/v) in this process.

The PCR of fungal ITS1 region was performed with the forward primer ITS1F_KYO1 [30] fused with 3–6-mer Ns for improved Illumina sequencing quality [26] and the forward Illumina sequencing primer (5'- TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG- (3–6-mer Ns) - (ITS1F_KYO1) -3') and the reverse primer ITS2_KYO2 [30] fused with 3–6-mer Ns and the reverse sequencing primer (5'- GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA G (3–6-mer Ns) - (ITS2_KYO2) -3'). The buffer and polymerase system of KOD FX Neo was used with a temperature profile of 94°C for 2 min, followed by 35 cycles at 98°C for 10 s, 50°C for 30 s, 68°C for 50 s, and a final extension at 68°C for 5 min. Illumina sequencing adaptors and 8-mer index sequences were then added in the second PCR as described above. The amplicons were purified and pooled as described above.

The sequencing libraries of the prokaryote 16S and fungal ITS regions were processed in an Illumina MiSeq sequencer (run center: KYOTO-HE; 15% PhiX spike-in). Because the quality of forward sequences is generally higher than that of reverse sequences in Illumina sequencing, we optimized the MiSeq run setting in order to use only forward sequences. Specifically, the run length was set at 271 forward (R1) and 31 reverse (R4) cycles in order to enhance forward sequencing data: the reverse sequences were used only for discriminating between 16S and ITS1 sequences based on the sequences of primer positions.

2.3. Bioinformatics

The raw sequencing data were converted into FASTQ files using the program bcl2fastq 1.8.4 distributed by Illumina. The output FASTQ files were demultiplexed with the program Claident v0.2.2017.05.22 [31,32], by which sequencing reads whose 8-mer index positions included nucleotides with low (less than 30) quality scores were removed. The sequencing data were deposited to DNA Data Bank of Japan (DDBJ) (DRA006845). Only forward sequences were used in the following analyses after removing low-quality 3'-ends using Claident. Noisy reads [31] were subsequently discarded and then denoised dataset consisting of 2 041 573 16S and 1 325 199 ITS1 reads were obtained.

For each dataset of 16S and ITS1 regions, filtered reads were clustered with a cut-off sequencing similarity of 97% using the program VSEARCH [33] as implemented in Claident. The operational taxonomic units (OTUs) representing less than 10 sequencing reads were subsequently discarded. The molecular identification of the remaining OTUs was performed based on the combination of the query-centric auto-*k*-nearest neighbour (QCauto) method [32] and the lowest common ancestor (LCA) algorithm [34] as implemented in Claident. Note that taxonomic identification results based on the combination of the QCauto search and the LCA taxonomic assignment are comparable to, or sometimes more accurate than, those with the alternative approaches [32,35,36]. In total, 5351 prokaryote (bacterial or archaeal) OTUs and 1039 fungal OTUs were obtained for the 16S and ITS1 regions, respectively (electronic supplementary material, data S1). The UNIX codes used in the above bioinformatic pipeline are available as electronic supplementary material, data S2.

For each combination of target region (16S or ITS1) and sample type (root or soil), we obtained a *sample* × OTU matrix, in which a cell entry depicted the number of sequencing reads of an OTU in a sample (electronic supplementary material, data S3). The cell entries whose read counts represented less than 0.1% of the total read count of each sample were removed to minimize effects of

PCR/sequencing errors [37]. The filtered matrix was then rarefied to 1000 reads per sample using the ‘rarefy’ function of the vegan 2.4–1 package [38] of R 3.4.3 [39]. Samples with less than 1000 reads were discarded in this process: the numbers of samples in the rarefied sample \times OTU matrices were 119, 128, 117 and 128 for root prokaryote, root fungal, soil prokaryote and soil fungal matrices, respectively (electronic supplementary material, data S4).

2.4. Prokaryote and fungal community structure

Relationship between the number of sequencing reads and that of detected OTUs was examined for each dataset (root prokaryote, root fungal, soil prokaryote or soil fungal dataset) with the ‘rarecurve’ function of the R vegan package. Likewise, relationship between the number of samples and that of OTUs was examined with the vegan ‘specaccum’ function. For each dataset, difference in OTU compositions among ‘green’, ‘yellow’ and ‘no leaf’ soybean individuals was examined by the permutational analysis of variance (PERMANOVA; [40]) with the vegan ‘adonis’ function (10 000 permutations). To control effects of sampling positions (lines) on the community structure, the information of sampling sets (set 1 or set 2) was included as an explanatory variable in the PERMANOVA. The variation in OTU compositions was visualized with non-metric multidimensional scaling (NMDS) using the vegan ‘metaMDS’ function. To examine the potential relationship between root/soil microbial community structure and plant biomass, an additional PERMANOVA was performed for each dataset. The information of sampling sets was included in the models. To explore signs of spatial autocorrelation in the community data, a Mantel’s correlogram analysis was performed with the vegan ‘mantel.correlog’ function. The ‘Bray–Curtis’ metric of β -diversity was used in the PERMANOVA, NMDS and Mantel’s correlogram analyses.

2.5. Screening of host-state-specific OTUs

To explore prokaryote/fungal OTUs that preferentially occurred on/around ‘green’, ‘yellow’ or ‘no leaf’ soybean individuals, a randomization test was performed by shuffling the plant state labels in each of the root prokaryote, root fungal, soil prokaryote and soil fungal data matrices (100 000 permutations). We then evaluated preference of a prokaryote/fungal OTU (i) for a plant state (j) (‘green’, ‘yellow’ or ‘no leaf’) as follows:

$$\text{Preference}(i, j) = \frac{[N_{\text{observed}}(i, j) - \text{Mean}(N_{\text{randomized}}(i, j))]}{\text{s.d.}(N_{\text{randomized}}(i, j))},$$

where $N_{\text{observed}}(i, j)$ denoted the mean number of the sequencing reads of OTU i among state j soybean samples in the original data, and the $\text{Mean}(N_{\text{randomized}}(i, j))$ and $\text{s.d.}(N_{\text{randomized}}(i, j))$ were the mean and standard deviation of the number of sequencing reads for the focal OTU–plant state combination across randomized matrices. Regarding this standardized preference index, values larger than three generally represent strong preferences (false discovery rate (FDR) < 0.05 ; see results of a previous study [35]): hence, we listed OTUs whose preference values exceeded three.

2.6. Microbe–microbe networks

To examine how prokaryote and fungal OTUs co-occurred in root or soil samples, a co-abundance network analysis was performed based on the sparse inverse covariance estimation for ecological association inference (Spiec-Easi) method [41]. In each of the root and soil sample analyses, the input data matrix was prepared by merging the sample \times OTU matrices of prokaryotes and fungi. As inferences of co-abundance patterns were unavailable for rare OTUs, only the OTUs detected from 30 or more samples were retained in the input matrices. For each of the root and soil data matrices, a co-abundance analysis was performed with the ‘spiec.easi’ function of the R ‘SpiecEasi’ package [41]. The networks depicting the co-abundance patterns were drawn using the R ‘igraph’ package [42].

3. Results

3.1. Prokaryotes and fungal community structure

On average, 107.9 (s.d. = 18.0), 25.4 (s.d. = 8.9), 172.5 (s.d. = 17.3) and 78.3 (s.d. = 10.5) OTUs per sample were observed, respectively, from the root prokaryote, root fungal, soil prokaryote and soil fungal dataset




Figure 2. Prokaryote and fungal community structure. (a) Phylum-level compositions of prokaryotes in the root and soil datasets. Mean proportions of sequencing reads are shown for respective taxa. The numbers of the samples from which sequencing data were successfully obtained are shown in the parentheses. (b) Order-level compositions of fungi in the root and soil datasets.

after filtering and rarefaction steps (electronic supplementary material, figure S2). The total number of OTUs observed was 1387, 346, 1191 and 769 for the root prokaryote, root fungal, soil prokaryote and soil fungal datasets, respectively (electronic supplementary material, figure S3).

In the soybean field, the prokaryote community on roots was dominated by the bacterial classes Proteobacteria, Actinobacteria, Chloroflexi and Bacteroidetes, while that of rhizosphere soil consisted mainly of Proteobacteria, Actinobacteria and Acidobacteria, and the archaeal lineage Thaumarchaeota (figure 2a). The fungal community of roots was dominated by the fungal orders Hypocreales, Sordariales, Pleosporales, while that of soil consisted mainly of Hypocreales, Agaricales, Eurotiales, Mortierellales and Filobasidiales (figure 2b). Regarding the order-level compositions of fungi in the rhizosphere soil, the proportion of Orbiliales reads was much higher in ‘yellow’ (3.62%) and ‘no leaf’ (4.82%) soybean individuals than in ‘green’ ones (0.89%) (figure 2). The genus level compositions of the samples are shown in electronic supplementary material, figure S4.

In each dataset (i.e. root prokaryote, root fungal, soil prokaryote or soil fungal data), microbial community structure varied among ‘green’, ‘yellow’ or ‘no leaf’ soybean individuals, although the effects of sampling sets on the community structure were much stronger (figure 3). Even within each sampling set, spatial autocorrelations of bacterial/fungal community structure were observed (electronic supplementary material, figure S5). Significant relationships between microbial community structure and soybean biomass were observed in the soil prokaryote and soil fungal datasets but not in the root prokaryote and root fungal datasets (table 1).

3.2. Screening of host-state-specific OTUs

In the root microbiome, only an unidentified fungal OTU showed a strong preference for ‘green’ soybean individuals, while 18 bacterial and four fungal OTUs occurred preferentially on ‘no leaf’ host individuals (table 2; electronic supplementary material, figure S6). The list of the bacteria showing preferences for ‘no leaf’ soybean individuals included OTUs whose 16S rRNA sequences were allied to those of *Dyella*, *Herbaspirillum*, *Labrys*, *Phenylobacterium*, *Gemmata*, *Chitinophaga*, *Pedobacter*, *Niastella* and *Streptomyces* (table 2). The four fungal OTUs showing preferences for ‘no leaf’ hosts were unidentified basidiomycetes (table 2).

In the rhizosphere soil microbiome, seven prokaryote OTUs, including those belonging to Chloroflexi (e.g. *Sphaerobacteraceae* sp.) and Proteobacteria (*Kofleriaceae* sp.), occurred preferentially on ‘green’ host individuals (table 3). Likewise, five fungal OTUs, including those allied to basidiomycete yeasts in the genera *Solicoccozyma* and *Saitozyma*, showed preferences for ‘green’ soybean individuals (table 3). Results also revealed that 26 bacterial and 11 fungal OTUs had biased distributions in the rhizosphere of ‘no leaf’ soybean individuals (table 3). The list of microbes showing preferences for ‘no leaf’ hosts included OTUs allied to bacteria in the genera *Pseudomonas*, *Nevskia*, *Cellvibrio*, *Massilia*, *Duganella*, *Novosphingobium*, *Mucilaginibacter* and *Flavobacterium* and OTUs allied to fungi in the genera *Burgoa*, *Clonostachys*, *Plectosphaerella*, *Xylaria*, *Dactylellina*, *Talaromyces*, *Cladosporium*, *Alternaria* and *Peniophora* (table 3). The list of microbes that preferentially occurred on ‘no leaf’ hosts involved OTUs with high sequence similarity to the nematophagous fungi, *Clonostachys rosea* (Hypocreales) and *Dactylellina* sp.




Figure 3. Diversity of microbiome structures among samples. (a) NMDS of the root prokaryote dataset. The results of the PERMANOVA, in which sampling set (set 1 or set 2) and plant state (green, yellow or no leaf) were included as explanatory variables, are shown. (b) NMDS of the root fungal dataset. (c) NMDS of the soil prokaryote dataset. (d) NMDS of the soil fungal dataset.

Table 1. Relationship between prokaryote/fungal community structure and the biomass of soybean individuals. For each dataset (i.e. root prokaryote, root fungal, soil prokaryote or soil fungal data), a PERMANOVA model of community structure was constructed. The information of the sampling set (set 1 or set 2) and the dry mass of host soybean individuals were included as explanatory variables.

variable	d.f.	F_{model}	p
root prokaryotes			
sampling set	1	10.4	<0.0001
dry mass	1	1.3	0.1139
root fungi			
sampling set	1	14.0	<0.0001
dry mass	1	0.6	0.8267
soil prokaryotes			
sampling set	1	15.4	<0.0001
dry mass	1	3.1	0.002
soil fungi			
sampling set	1	36.7	<0.0001
dry mass	1	2.2	0.0145

(Orbiliiales) (table 3). The reads of the *Clonostachys* (F_0257) and *Dactyloellina* (F_0163) OTUs, respectively, represented 9.5% and 3.5% of the sequencing reads of 'no leaf' samples (electronic supplementary material, data S5). The indices of preferences for 'yellow' soybean individuals are shown in electronic supplementary material, data S5.

Table 2. Prokaryote and fungal OTUs showing strong preferences for host states in the root microbiome datasets. The prokaryote/fungal OTUs that showed strong preferences for ‘green’ or ‘no leaf’ soybean individuals (preference value ≥ 3) are shown. The taxonomic assignment results based on the Qcauto–LCA pipeline are shown with the top-hit results of NCBI BLAST searches. The OTU code starting with P (P_XXXX) and F (F_XXXX) are prokaryotes and fungi, respectively.

OTU	phylum	class	order	family	genus	NCBI top hit	accession	cover (%)	Identity ()
green									
F_0437	Ascomycota	—	—	—	—	<i>Knufia</i> sp.	KP235641.1	83	98
no leaf									
P_3453	Proteobacteria	Gammaproteobacteria	Xanthomonadales	Rhodanobacteraceae	—	<i>Dyella marenis</i>	LN890104.1	100	99
P_3207	Proteobacteria	Gammaproteobacteria	Legionellales	Coxiellaceae	<i>Aquicella</i>	<i>Aquicella siphonis</i>	NR_025764.1	100	94
P_2827	Proteobacteria	Betaproteobacteria	—	—	—	<i>Duganella zoogloeoides</i>	KT983992.1	100	100
P_2733	Proteobacteria	Betaproteobacteria	Burkholderiales	Oxalobacteraceae	<i>Herbaspirillum</i>	<i>Herbaspirillum chlorophenolicum</i>	MG571754.1	100	100
P_2590	Proteobacteria	Alphaproteobacteria	—	—	—	<i>Crocococcus mobilis</i>	NR_152701.1	100	88
P_2481	Proteobacteria	Alphaproteobacteria	Rickettsiales	Rickettsiaceae	—	<i>Rickettsia japonica</i>	KU586263.1	100	91
P_2279	Proteobacteria	Alphaproteobacteria	Rhizobiales	Xanthobacteraceae	<i>Labrys</i>	<i>Labrys monachus</i>	KT694157.1	100	100
P_2042	Proteobacteria	Alphaproteobacteria	Caulobacterales	Caulobacteraceae	<i>Phenylobacterium</i>	<i>Phenylobacterium</i> sp.	JX458410.1	100	99
P_3664	Proteobacteria	—	—	—	—	<i>Desulfofrigus oceanense</i>	AB568590.1	97	93
P_3658	Proteobacteria	—	—	—	—	<i>Rudaea</i> sp.	KM253197.1	100	85
P_1748	Planctomycetes	Planctomycetia	Planctomycetales	Gemmataceae	<i>Gemmata</i>	<i>Gemmata</i> sp.	GQ889445.1	100	99
P_1278	Chloroflexi	Thermomicrobia	—	—	—	<i>Sphaerobacter thermophilus</i>	AJ871227.1	100	92
P_1058	Bacteroidetes	—	—	—	—	<i>Chitinophaga polysaccharaea</i>	MG322237.1	100	92
P_1049	Bacteroidetes	—	—	—	—	<i>Pedobacter terrae</i>	MG819444.1	100	98
P_0994	Bacteroidetes	—	—	—	—	<i>Chitinophaga terrae</i>	LN890054.1	100	95

(Continued.)

Table 2. (Continued.)

OTU	phylum	class	order	family	genus	NCBI top hit	accession	cover (%)	identity ()
P_0887	Bacteroidetes	Chitinophagia	Chitinophagales	Chitinophagaceae	<i>Mastella</i>	<i>Mastella koreensis</i>	NR_074595.1	100	100
P_0498	Actinobacteria	Actinobacteria	Streptomyetales	Streptomycetaceae	—	<i>Streptomyces albiaxialis</i>	KP170480.1	100	98
P_0444	Actinobacteria	Actinobacteria	Streptomycetales	Streptomycetaceae	<i>Streptomyces</i>	<i>Streptomyces olivaceoviridis</i>	KP823723.1	100	98
F_0796	Basidiomycota	—	—	—	—	<i>Classicalaceae</i> sp.	KY548838.1	92	84
F_0792	Basidiomycota	—	—	—	—	<i>Classicalaceae</i> sp.	KY548838.1	92	83
F_0790	Basidiomycota	—	—	—	—	<i>Classicalaceae</i> sp.	KY548838.1	91	83
F_0786	Basidiomycota	—	—	—	—	<i>Classicalaceae</i> sp.	KY548838.1	90	84

Table 3. Prokaryote and fungal OTUs showing strong preferences for host states in the soil microbiome datasets. The prokaryote/fungal OTUs that showed strong preferences for ‘green’ or ‘no leaf’ soybean individuals (preference value ≥ 3) are shown. The taxonomic assignment results based on the Qcauto–LCA pipeline are shown with the top-hit results of NCBI BLAST searches. The OTU code starting with P (P_XXXX) and F (F_XXXX) are prokaryotes and fungi, respectively.

OTU	phylum	class	order	family	genus	NCBI top hit	accession	cover (%)	identity (%)
green									
P_0697	Actinobacteria	—	—	—	—	<i>Gaiella oculata</i>	NR_118138.1	100	91
P_1264	Chloroflexi	Thermomicrobia	Sphaerobacterales	Sphaerobacteraceae	<i>Sphaerobacter</i>	<i>Shewanella fodinae</i>	FM887036.1	98	84
P_1281	Chloroflexi	Thermomicrobia	—	—	—	<i>Thermomicrobium carboxidum</i>	NR_134218.1	100	87
P_2949	Proteobacteria	Deltaproteobacteria	Myxococcales	Kofleriaceae	<i>Hallangium</i>	<i>Koferia flava</i>	HF937255.1	100	91
P_3762	—	—	—	—	—	<i>Planctomycetales bacterium</i>	AY673390.1	98	94
P_3715	—	—	—	—	—	<i>Brochothrix thermosphacta</i>	MG807446.1	99	86
P_0032	—	—	—	—	—	<i>Nitrosocosmicus exaquare</i>	CP017922.1	100	99
Fungi									
F_0477	Ascomycota	—	—	—	—	No significant match	—	—	—
F_0141	Ascomycota	Eurotiomycetes	—	—	—	<i>Penicillium clavigerum</i>	NR_121317.1	100	81
F_0700	Basidiomycota	Tremellomycetes	Filobasidiales	Pisikurozymaceae	<i>Solicozozyma</i>	<i>Solicozozyma terreus</i>	KY102958.1	100	100
F_0734	Basidiomycota	Tremellomycetes	Tremellales	Trimorphomycetaceae	<i>Saitozyma</i>	<i>Saitozyma podzolica</i>	KY102943.1	82	99
F_0738	Basidiomycota	Tremellomycetes	Tremellales	Trimorphomycetaceae	<i>Saitozyma</i>	<i>Saitozyma podzolica</i>	KY102943.1	84	99

(Continued.)

Table 3. (Continued.)

OTU	phylum	class	order	family	genus	NCBI top hit	accession	cover (%)	Identity (%)
no leaf									
P_3294	Proteobacteria	Gammaproteobacteria	Pseudomonadales	Pseudomonadaceae	<i>Pseudomonas</i>	<i>Pseudomonas psychrotolerans</i>	KY623077.1	100	100
P_3256	Proteobacteria	Gammaproteobacteria	Nevskiales	Sinobacteraceae	<i>Nevskia</i>	<i>Nevskia persephonica</i>	J0710442.1	97	99
P_3189	Proteobacteria	Gammaproteobacteria	Cellvibrionales	Cellvibrionaceae	<i>Cellvibrio</i>	<i>Cellvibrio mixtus</i>	KC329916.1	100	100
P_3308	Proteobacteria	Gammaproteobacteria	—	—	—	<i>Steroidobacter</i> sp.	KP185148.1	100	95
P_3093	Proteobacteria	Deltaproteobacteria	Myxococcales	—	—	<i>Sorangineae bacterium</i>	JF719608.1	100	94
P_3004	Proteobacteria	Deltaproteobacteria	Myxococcales	Polyangiaceae	<i>Byssovorax</i>	<i>Polyangium spumosum</i>	KX572839.2	100	97
P_3114	Proteobacteria	Deltaproteobacteria	—	—	—	<i>Stigmatella hybrida</i>	KX572784.2	100	91
P_2747	Proteobacteria	Betaproteobacteria	Burkholderiales	Oxalobacteraceae	—	<i>Massilia kyonggiensis</i>	NR_126273.1	100	100
P_2827	Proteobacteria	Betaproteobacteria	—	—	—	<i>Duganella radidis</i>	LC191531.1	100	100
P_2552	Proteobacteria	Alphaproteobacteria	Sphingomonadales	Sphingomonadaceae	—	<i>Novosphingobium sediminicola</i>	KX987160.1	100	100
P_1637	Gemmatimonadetes	Gemmatimonadetes	Gemmatimonadales	Gemmatimonadaceae	<i>Gemmatimonas</i>	<i>Gemmatimonas aurantiaca</i>	KF228166.1	100	93
P_1544	Gemmatimonadetes	Gemmatimonadetes	Gemmatimonadales	Gemmatimonadaceae	<i>Gemmatimonas</i>	<i>Gemmatimonas</i> sp.	LN876485.1	100	89
P_0962	Bacteroidetes	Sphingobacteriia	Sphingobacteriales	Sphingobacteriaceae	<i>Mucilaginibacter</i>	<i>Mucilaginibacter gofawali</i>	AP017313.1	100	99
P_0892	Bacteroidetes	Chitinophagia	Chitinophagales	Chitinophagaceae	—	<i>Ferruginibacter profundus</i>	NR_148259.1	100	88

(Continued.)

Table 3. (Continued.)

OTU	phylum	class	order	family	genus	NCBI top hit	accession	cover (%)	Identity (%)
P_1095	Bacteroidetes	—	—	—	—	<i>Flavisolibacter</i>	NR_041500.1	100	95
P_1051	Bacteroidetes	—	—	—	—	<i>ginsengisoli</i> <i>Flavobacterium</i>	KP875419.1	100	100
P_1008	Bacteroidetes	—	—	—	—	<i>lindanitolerans</i> <i>Solitalea canadensis</i>	CP003349.1	100	88
P_0652	Actinobacteria	Thermoleophila	Solirubrobacterales	Solirubrobacteraceae	<i>Solirubrobacter</i>	<i>Solirubrobacter</i>	NR_133858.1	99	92
P_5169	—	—	—	—	—	<i>phytolaccae</i> <i>Desulfotomaculum</i>	NR_074579.1	97	85
P_5087	—	—	—	—	—	<i>nigrificans</i> <i>Stenotrophobacter</i>	NR_146022.1	99	97
P_4649	—	—	—	—	—	<i>roseus</i> <i>Alkalilimnicola ehrlichii</i>	NR_074775.1	99	81
P_4607	—	—	—	—	—	<i>Verrucomicrobia</i>	JF488114.1	100	92
P_4606	—	—	—	—	—	<i>Ruminococcus</i>	KX155563.1	99	83
P_4595	—	—	—	—	—	<i>flavofaciens</i> <i>Moorella thermoacetica</i>	NR_043076.1	97	84
P_3783	—	—	—	—	—	<i>Fimbriimonas</i>	CP007139.1	100	88
P_3739	—	—	—	—	—	<i>ginsengisoli</i> <i>Solibacter usitatus</i>	GQ287461.1	100	88
F_0866	Mucoromycota	Glomeromycetes	—	—	—	<i>Acaulospora delicata</i>	JF439203.1	45	95
F_0620	Basidiomycota	Agaricomycetes	Polyporales	—	<i>Burgoa</i>	<i>Burgoa anomala</i>	AB972783.1	100	100

(Continued.)

Table 3. (Continued.)

OTU	phylum	class	order	family	genus	NCBI top hit	accession	cover (%)	Identity (%)
F_0785	Basidiomycota	—	—	—	—	<i>Radulomyces copeiandii</i>	MG722738.1	87	99
F_0257	Ascomycota	Sordariomycetes	Hypocreales	Bionectriaceae	<i>Clonostachys</i>	<i>Clonostachys rosea</i>	KY320599.1	100	100
F_0237	Ascomycota	Sordariomycetes	Glomerellales	Plectosphaerellaceae	—	<i>Plectosphaerella plurivora</i>	KU204617.1	98	99
F_0413	Ascomycota	Sordariomycetes	—	—	—	<i>Xylariales</i> sp.	KY031690.1	100	100
F_0163	Ascomycota	Orbiliomycetes	Orbiliales	Orbiliaceae	<i>Dactylellina</i>	<i>Dactylellina</i> aff. <i>ellipospora</i>	KT215204.1	100	99
F_0131	Ascomycota	Eurotiomycetes	Eurotiales	—	—	<i>Talaromyces verruculosus</i>	KC937053.1	100	98
F_0003	Ascomycota	Dothideomycetes	Capnodiales	Cladosporiaceae	<i>Cladosporium</i>	<i>Cladosporium cladosporioides</i>	MG946764.1	100	100
F_0482	Ascomycota	—	—	—	—	<i>Alternaria alternata</i>	KY367499.2	100	100
F_0973	—	—	—	—	—	<i>Peniophora incarnata</i>	EU918698.1	100	98




Figure 4. Microbe–microbe co-abundance networks. (a) Positive co-abundance network of the root microbiome data. Pairs of OTUs linked by a blue line frequently co-occurred in the same soybean samples. (b) Negative co-abundance network of the root microbiome data. Pairs of OTUs linked by a red line rarely co-occurred in the same soybean samples. (c) Positive co-abundance network of the soil microbiome data. (d) Negative co-abundance network of the soil microbiome data.

3.3. Microbe–microbe networks

The structures of microbe–microbe networks (figure 4) were more complicated in the soil microbiome data (figure 4c,d) than in the root microbiome data (figure 4a,b). Within the network representing co-abundance of microbes across root samples, the *Clonostachys* OTU (F_0257) had a significant link with a *Streptomyces* OTU, while *Dactylellina* was absent from the root microbiome network data (figure 4a). Within the positive co-abundance network of the rhizosphere soil microbiome (figure 4c), the *Clonostachys* (F_0257) and *Dactylellina* (F_0163) nematophagous fungal OTUs were connected with each other (table 4). In addition, the *Clonostachys* OTU was linked with two bacterial OTUs (*Ralstonia* and *Rhizobiales*) and fungal OTUs in the genera *Calonectria* and *Purpureocillium* (table 4). Likewise, the *Dactylellina* OTU was connected also with two Alphaproteobacterial OTUs and a bacterial OTU allied to *Nitrospira japonica* as well as fungal OTUs in the genera *Rhizophydium*, *Pochonia* and *Purpureocillium* (table 4).

4. Discussion

Based on Illumina sequencing, we compared root-associated/rhizosphere microbial communities between soybean individuals infected by root-knot nematodes and those showing no symptoms. The

Table 4. Prokaryote/fungal OTUs linked to nematophagous fungi in the microbe–microbe networks. For each of the microbe–microbe co-abundance networks (figure 4a,c), the prokaryote/fungal OTUs that showed positive co-abundance patterns with *Clonostachys* (F_0257) and *Dactylellina* (F_0163) nematophagous fungal OTUs are listed. The taxonomic assignment results based on the Qcauto–LCA pipeline are shown with the top-hit results of NCBI BLAST searches. The OTU code starting with P (P_XXXX) and F (F_XXXX) are prokaryotes and fungi, respectively.

OTU	phylum	class	order	family	genus	NCBI top hit	accession	Cover (%)	Identity (%)
root: OTUs linked to <i>Clonostachys rosea</i> (F_0257)									
P_0510	Actinobacteria	Actinobacteria	Streptomycetales	Streptomycetaceae	—	<i>Streptomyces nigrogriseolus</i>	MG984076.1	100	98
soil: OTUs linked to <i>Clonostachys rosea</i> (F_0257)									
P_2689	Proteobacteria	Betaproteobacteria	Burkholderiales	Burkholderiaceae	<i>Ralstonia</i>	<i>Ralstonia pickettii</i>	MF179868.1	100	100
P_2243	Proteobacteria	Alphaproteobacteria	Rhizobiales	—	—	<i>Pedomicrobium americanum</i>	NR_104908.1	100	90
F_0163	Ascomycota	Orbiliomycetes	Orbiliales	Orbiliaceae	<i>Dactylellina</i>	<i>Dactylellina</i> aff. <i>ellipospora</i>	KT215204.1	100	99
F_0278	Ascomycota	Sordariomycetes	Hypocreales	Nectriaceae	<i>Calonectria</i>	<i>Calonectria zuluensis</i>	NR_137728.1	97	100
F_0310	Ascomycota	Sordariomycetes	Hypocreales	Ophiocordycipitaceae	—	<i>Purpureocillium lilacinum</i>	KP691502.1	100	100
soil: OTUs linked to <i>Dactylellina</i> sp. (F_0163)									
P_2443	Proteobacteria	Alphaproteobacteria	Rhodospirillales	—	—	<i>Azospirillum brasilense</i>	KY010284.1	100	92
P_2589	Proteobacteria	Alphaproteobacteria	—	—	—	<i>Elstera litoralis</i>	KR856497.1	100	92
P_3774	—	—	—	—	—	<i>Nitrospira japonica</i>	LT828648.1	100	100
F_0812	Chytridiomycota	Chytridiomycetes	Rhizophydiales	Rhizophydiaceae	<i>Rhizophyidium</i>	<i>Rhizophyidium</i> sp.	AY349124.1	99	100
F_0278	Ascomycota	Sordariomycetes	Hypocreales	Nectriaceae	<i>Calonectria</i>	<i>Calonectria zuluensis</i>	NR_137728.1	97	100
F_0265	Ascomycota	Sordariomycetes	Hypocreales	Claviicipitaceae	<i>Pochonia</i>	<i>Pochonia chlamydosporia</i>	KY977543.1	100	100
F_0257	Ascomycota	Sordariomycetes	Hypocreales	Bionectriaceae	<i>Clonostachys</i>	<i>Clonostachys rosea</i>	KY320599.1	100	100
F_0310	Ascomycota	Sordariomycetes	Hypocreales	Ophiocordycipitaceae	—	<i>Purpureocillium lilacinum</i>	KP691502.1	100	100

results indicated that, in both soybean roots and rhizosphere soil, prokaryote and fungal community structures significantly varied depending on host plant states (figures 2 and 3). We further performed statistical analyses for screening prokaryote and fungal OTUs preferentially associated with infected and benign soybean host individuals (tables 2 and 3; figure 4). The results are based on purely descriptive data and hence they, in principle, are not direct evidences of interactions among plants, nematodes and microbiomes: i.e. causal relationship among those agents remains unknown. As this study provided only 'snap-shot' information of microbiome structure at the end of a growing season, we need to conduct further studies uncovering temporal microbiome dynamics throughout the growing season of soybeans. Nonetheless, as detailed below, the statistical analyses suggest assembly of diverse anti-nematode bacteria and fungi from indigenous microbial communities in the soybean field, providing a basis for exploring ways to reduce damage by root-knot nematodes with those indigenous functional microbes.

Within the root microbiome analysed, various taxonomic groups of bacteria preferentially occurred on 'no leaf' soybean samples (table 2). Among them, the genus *Streptomyces* is known to involve some species that suppress nematode populations, potentially used as biological control agents for root-knot nematodes [43–46]. By contrast, *Herbaspirillum*, *Rickettsia*, *Chitinophaga* and *Pedobacter* have been reported as symbionts of nematodes, potentially playing beneficial roles for host nematodes [47–49]. Results of these statistical analyses should be interpreted with caution, as they are likely to highlight not only prospective microbes potentially parasitizing on pests/pathogens, but also microbes that can form mutualistic interactions with disease agents.

Within the soybean rhizosphere soil microbiome, diverse taxonomic groups of not only bacteria, but also fungi preferentially occurred around 'no leaf' soybean individuals (table 3). Among them, *Pseudomonas* has been known to suppress root-knot nematode populations [50,51] potentially by producing hydrogen cyanide [52] or extracellular protease [53], but interactions with root-knot nematodes have not yet been examined for other bacteria preferentially found in the rhizosphere of 'no leaf' soybean individuals. Meanwhile, the list of the fungal OTUs frequently observed in the rhizosphere of 'no leaf' soybeans included some fungi whose ability to suppress nematode populations had been well documented (table 3). *Clonostachys rosea*, for example, has been known as a prospective biological control agent of plant- and animal-pathogenic nematodes [54,55]. An observational study based on green fluorescent protein imaging has indicated that the conidia of the fungus adhere to nematode cuticle and their germ tubes penetrate nematode bodies, eventually killing the invertebrate hosts [56]. The fungus is also known to produce a subtilisin-like extracellular protease, which plays an important role during the penetration of nematode cuticles [57]. Our analysis also highlighted a nematophagous fungus in the genus *Dactylellina* (teleomorph = *Orbilina*), which could capture juveniles of nematodes with hyphal traps [58]. Species in the genus and many other fungi in the order Orbiliales produce characteristic trap structures with their hyphae to prey on nematodes [59–61], often nominated as prospective biological control agents [62–64].

An additional analysis focusing on *Clonostachys* and *Dactylellina* highlighted bacteria and fungi that frequently co-occurred with the nematophagous fungi (figure 4). In the root microbiome, *Clonostachys* and a *Streptomyces* OTU showed positively correlated distributions across soybean samples (table 4). In the rhizosphere microbiome, *Clonostachys* and *Dactylellina* showed significant co-abundance patterns (table 4). Moreover, in the soil, the two nematophagous fungi co-occurred frequently with other taxonomic groups of nematophagous fungi such as *Purpureocillium*, *Pochonia* and *Rhizophydium* (table 4 and figure 5). Among them, fungi in the genus *Purpureocillium* (Hypocreales: Ophiocordycipitaceae) have been known to suppress plant parasitic nematodes, insect pests and oomycete phytopathogens [65–68]. Another Hypocreales genus, *Pochonia* (previously placed in the genus *Verticillium*; teleomorph = *Metacordyceps*; Clavicipitaceae) has been known as nematophagous as well and they can kill eggs and females of root-knot (*Meloidogyne* spp.) and cyst (*Globodera* spp.) nematodes [69–72]. Species in the chytrid genus *Rhizophydium* include species that use nematodes as parasites or saprophytes [73,74]. They are known to explore host nematodes in the form of zoospores [73]. All these results suggest that indigenous anti-nematode or nematophagous microbes can form consortia in soil ecosystems of soybean fields. It is important to note that the members of the consortia do not necessarily interact with each other directly: i.e. they may merely share habitat preferences [36,37,75]. However, the inferred structure of microbe–microbe networks helps us understand overall consequences of ecological processes in microbiomes [15].

Along with the consortia of anti-nematode microbes, an OTU in the genus *Calonectria*, which causes leaf blight, wilt and root rot of various plant species [76,77], was frequently observed (table 4). The phytopathogenic fungus might have attacked soybean individuals weakened by root-knot nematodes.




Figure 5. Spatial distribution of nematophagous fungal OTUs. (a) Sampling set 1. For each soybean individual, the proportions of sequencing reads representing nematophagous fungal OTUs are shown. (b) Sampling set 2.

Alternatively, *Calonectria* may have infected host soybeans earlier than root-knot nematodes, followed by the emergence of nematodes and their exploiters (i.e. anti-nematode microbes). Given that fungi can interact with each other both antagonistically and mutualistically in the soil [78,79], direct interactions between *Calonectria* and nematophagous fungi in the genera *Clonostachys*, *Dactylellina*, *Purpureocillium*, *Pochonia* and *Rhizophydium* are of particular interest. Studies examining potential interactions involving soybeans, root-knot nematodes, anti-nematode bacteria/fungi and *Calonectria* will help us understand ecological processes that structure consortia of nematophagous fungi.

Although this study did not evaluate potential effects of background environmental conditions (e.g. soil pH and inorganic nitrogen concentration) on microbiome structure, management of edaphic conditions are expected to have great impacts on dynamics of anti-nematode microbiomes. A number of studies have explored ways to suppress nematode populations by optimizing cropping systems [1]. Crop rotation, in which planting of a crop variety and that of nematode-resistant varieties/species are rotated, has been recognized as an effective technique for regulating root-knot and cyst nematode populations [8,80,81]. By contrast, long-term continual cropping in soybean monoculture fields can increase anti-nematode bacteria and fungi (e.g. *Pseudomonas*, *Purpureocillium* and *Pochonia*), potentially resulting in lowered densities of cyst nematodes [12]. Tillage regimes [9–11] and introduction of organic matter (e.g. alfalfa leaves or crop residue) [82–84] have great impacts on nematode densities in farmlands, but their effects vary considerably among studies [1]. In addition, because nematode-infected plant individuals can show highly aggregated distributions at a small spatial scale within a farmland (figure 1d), tillage can promote the spread of plant damaging nematodes [22]. Frequent tillage may have negative impacts on populations of nematophagous fungi as a consequence of hyphal fragmentation (cf. [85]), but such destructive effects on fungal communities have not yet been tested intensively. Given that microbiome structures were not taken into account in most previous studies evaluating effects of cropping systems on nematode suppression (but see [12,21]), more insights into the relationship between agroecosystem management and indigenous (native) microbiome dynamics are required for building reproducible ways to develop disease-suppressive soil.

We herein found that consortia of anti-nematode bacteria and fungi could develop at a small spatial scale within a field of soybeans infected by root-knot nematodes. Given the diversity of those anti-nematode microbes observed in this study, multiple biological control agents are potentially available *in situ* without introducing exogenous ones depending on base compositions and conditions of indigenous microbiomes. In this respect, design of cropping systems (e.g. crop rotations, tillage frequencies, and inputs of fertilizer or organic matter) is of particular importance in activating and maximizing ecosystem functions that stem from resident microbial diversity [15]. Because those indigenous microbes, in general, have adapted to local biotic and abiotic environments, their populations are expected to persist more stably than exogenous microbes artificially introduced to a target agroecosystem (see [19] for reviews of the success/failure of microbial introduction). Elucidating the relationship between cropping systems and microbiome processes is the key to designing disease-suppressive agroecosystems.

Data accessibility. Data are available from the electronic supplementary material, data S1–S5 and DNA DDBJ (DRA006845).

Authors' contributions. H.T. conceived and designed the work. H.T. and Y.T. performed fieldwork. H.T. conducted molecular experiment and analysed the data. H.T. wrote the manuscript with Y.T. All authors gave final approval for publication.

Competing interests. The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be constructed as conflict of interest.

Funding. This work was financially supported by JSPS KAKENHI Grant no. (15KT0032) and JST PRESTO (JPMJPR16Q6) to H.T.

Acknowledgements. We thank Tatsuhiko Shiraiwa for his support in fieldwork and Mizuki Shinoda, Ko Mizushima, Sarasa Amma and Hiroki Kawai for their support in molecular experiments. We are also grateful to Ryoji Shinya for his advice on the biology of nematodes. We are also grateful to anonymous reviewers for their productive comments.

References

- Barker KR, Koenning SR. 1998 Developing sustainable systems for nematode management. *Annu. Rev. Phytopathol.* **36**, 165–205. (doi:10.1146/annurev.phyto.36.1.165)
- Abad P *et al.* 2008 Genome sequence of the metazoan plant-parasitic nematode *Meloidogyne incognita*. *Nat. Biotech.* **26**, 909. (doi:10.1038/nbt.1482)
- Wrather JA, Anderson T, Arsyad D, Gai J, Ploper L, Porta-Puglia A, Ram H, Yorinori J. 1997 Soybean disease loss estimates for the top 10 soybean producing countries in 1994. *Plant Dis.* **81**, 107–110. (doi:10.1094/PDIS.1997.81.1.107)
- Wrather JA, Koenning SR. 2006 Estimates of disease effects on soybean yields in the United States 2003 to 2005. *J. Nematol.* **38**, 173–180.
- Li J, Zou C, Xu J, Ji X, Niu X, Yang J, Huang X, Zhang K-Q. 2015 Molecular mechanisms of nematode-nematophagous microbe interactions: basis for biological control of plant-parasitic nematodes. *Annu. Rev. Phytopathol.* **53**, 67–95. (doi:10.1146/annurev-phyto-080614-120336)
- Schmitt D, Corbin F, Nelson L. 1983 Population dynamics of *Heterodera glycines* and soybean response in soils treated with selected nematocides and herbicides. *J. Nematol.* **15**, 432–437.
- Meyer SL, Roberts DP. 2002 Combinations of biocontrol agents for management of plant-parasitic nematodes and soilborne plant-pathogenic fungi. *J. Nematol.* **34**, 1–8.
- Nusbaum C, Ferris H. 1973 The role of cropping systems in nematode population management. *Annu. Rev. Phytopathol.* **11**, 423–440. (doi:10.1146/annurev.phy.11.090173.002231)
- Okada H, Harada H. 2007 Effects of tillage and fertilizer on nematode communities in a Japanese soybean field. *Appl. Soil Ecol.* **35**, 582–598. (doi:10.1016/j.apsoil.2006.09.008)
- Thomas S. 1978 Population densities of nematodes under seven tillage regimes. *J. Nematol.* **10**, 24.
- Donald P, Tyler D, Boykin D. 2009 Short-and long-term tillage effects on *Heterodera glycines* reproduction in soybean monoculture in west Tennessee. *Soil Tillage Res.* **104**, 126–133. (doi:10.1016/j.still.2009.02.002)
- Hamid MI, Hussain M, Wu Y, Zhang X, Xiang M, Liu X. 2017 Successive soybean-monoculture cropping assembles rhizosphere microbial communities for the soil suppression of soybean cyst nematode. *FEMS Microbiol. Ecol.* **93**, fiw222. (doi:10.1093/femsec/fiw222)
- Edwards J, Johnson C, Santos-Medellin C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA, Sundaresan V. 2015 Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc. Natl Acad. Sci. USA* **112**, E911–E920. (doi:10.1073/pnas.1414592112)
- Lundberg DS *et al.* 2012 Defining the core *Arabidopsis thaliana* root microbiome. *Nature* **488**, 86–90. (doi:10.1038/nature11237)
- Toju H *et al.* 2018 Core microbiomes for sustainable agroecosystems. *Nat. Plants* **4**, 247–257. (doi:10.1038/s41477-018-0139-4)
- Schlaeppli K, Bulgarelli D. 2015 The plant microbiome at work. *Mol. Plant-Microbe Int.* **28**, 212–217. (doi:10.1094/MPMI-10-14-0334-FI)
- Mendes R *et al.* 2011 Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science* **332**, 1097–1100. (doi:10.1126/science.1203980)
- Berendsen RL, Pieterse CM, Bakker PA. 2012 The rhizosphere microbiome and plant health. *Trends Plant Sci.* **17**, 478–486. (doi:10.1016/j.tplants.2012.04.001)
- Mendes R, Garbeva P, Raaijmakers JM. 2013 The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS Microbiol. Rev.* **37**, 634–663. (doi:10.1111/1574-6976.12028)
- Cha J-Y *et al.* 2016 Microbial and biochemical basis of a *Fusarium* wilt-suppressive soil. *ISME J.* **10**, 119. (doi:10.1038/ismej.2015.95)
- Hussain M, Hamid MI, Tian J, Hu J, Zhang X, Chen J, Xiang M, Liu X. 2018 Bacterial community assemblages in the rhizosphere soil, root endosphere and cyst of soybean cyst nematode-suppressive soil challenged with nematodes. *FEMS Microbiol. Ecol.* **94**, fiy142.
- Gavassoni WL, Tylka GL, Munkvold GP. 2001 Relationships between tillage and spatial patterns of *Heterodera glycines*. *Phytopathology* **91**, 534–545. (doi:10.1094/PHYTO.2001.91.6.534)
- Sato H, Murakami N. 2008 Reproductive isolation among cryptic species in the ectomycorrhizal genus *Strobilomyces*: population-level CAPS marker-based genetic analysis. *Mol. Phylog. Evol.* **48**, 326–334. (doi:10.1016/j.ympev.2008.01.033)
- Takada-Hoshino Y, Matsumoto N. 2004 An improved DNA extraction method using skim milk from soils that strongly adsorb DNA. *Microbes Env.* **19**, 13–19. (doi:10.1264/jsm.2.19.13)
- Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N, Knight R. 2011 Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc. Natl Acad. Sci. USA* **108**, 4516–4522. (doi:10.1073/pnas.1000080107)
- Lundberg DS, Yourstone S, Mieczkowski P, Jones CD, Dangl JL. 2013 Practical innovations for high-throughput amplicon sequencing. *Nat. Methods* **10**, 999–1002. (doi:10.1038/nmeth.2634)
- Apprill A, McNally S, Parsons R, Weber L. 2015 Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquat. Microb. Ecol.* **75**, 129–137. (doi:10.3354/ame01753)
- Stevens JL, Jackson RL, Olson JB. 2013 Slowing PCR ramp speed reduces chimera formation from environmental samples. *J. Microbiol. Methods* **93**, 203–205. (doi:10.1016/j.mimet.2013.03.013)
- Hamady M, Walker JJ, Harris JK, Gold NJ, Knight R. 2008 Error-correcting barcoded primers for pyrosequencing hundreds of samples in multiplex. *Nat. Methods* **5**, 235–237. (doi:10.1038/nmeth.1184)
- Toju H, Tanabe AS, Yamamoto S, Sato H. 2012 High-coverage ITS primers for the DNA-based identification of ascomycetes and basidiomycetes in environmental samples. *PLoS ONE* **7**, e40863. (doi:10.1371/journal.pone.0040863)
- Tanabe AS. 2018 Claident v0.2.2018.05.29, a software distributed by author at <http://www.fifthdimension.jp/>.
- Tanabe AS, Toju H. 2013 Two new computational methods for universal DNA barcoding: a benchmark using barcode sequences of bacteria, archaea, animals, fungi, and land plants. *PLoS ONE* **8**, e76910. (doi:10.1371/journal.pone.0076910)

33. Rognes T, Mahé F, Flouri T, Quince C, Nichols B. 2014 Vsearch: program. See <https://github.com/torognes/vsearch>.
34. Huson DH, Auch AF, Qi J, Schuster SC. 2007 MEGAN analysis of metagenomic data. *Genome Res.* **17**, 377–386. (doi:10.1101/gr.5969107)
35. Toju H, Tanabe A, Ishii H. 2016 Ericaceous plant–fungus network in a harsh alpine–subalpine environment. *Mol. Ecol.* **25**, 3242–3257. (doi:10.1111/mec.13680)
36. Toju H, Yamamoto S, Tanabe AS, Hayakawa T, Ishii HS. 2016 Network modules and hubs in plant–root fungal biome. *J. R. Soc. Interface* **13**, 20151097. (doi:10.1098/rsif.2015.1097)
37. Peay KG, Russo SE, McGuire KL, Lim Z, Chan JP, Tan S, Davies SJ. 2015 Lack of host specificity leads to independent assortment of dipterocarps and ectomycorrhizal fungi across a soil fertility gradient. *Ecol. Lett.* **18**, 807–816. (doi:10.1111/ele.12459)
38. Oksanen J *et al.* 2012 Vegan: community ecology package. R package version 2.0-3. See <http://CRAN.R-project.org/package=vegan>.
39. R-Core-Team. 2018 R 3.5.1: A language and environment for statistical computing. See <http://www.R-project.org/>.
40. Anderson MJ. 2001 A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* **26**, 32–46. (doi:10.1111/j.1442-9993.2001.01070.pp.x)
41. Kurtz ZD, Mueller CL, Mirdali ER, Littman DR, Blaser MJ, Bonneau RA. 2015 Sparse and compositionally robust inference of microbial ecological networks. *PLoS Comp. Biol.* **11**, e1004226. (doi:10.1371/journal.pcbi.1004226)
42. Csardi G, Nepusz T. 2006 The igraph software package for complex network research. *Int. J. Complex Syst.* **1695**, 1–9.
43. Esnard J, Potter TL, Zuckerman BM. 1995 *Streptomyces costaricanus* sp. nov., isolated from nematode-suppressive soil. *Int. J. Syst. Evol. Microbiol.* **45**, 775–779. (doi:10.1099/00207173-45-4-775)
44. Chubachi K, Furukawa M, Fukuda S, Takahashi S, Matsumura S, Itagawa H, Shimizu T, Nakagawa A. 1999 Control of root-knot nematodes by *Streptomyces*: screening of root-knot nematode-controlling actinomycetes and evaluation of their usefulness in a pot test. *Nematol. Res.* **29**, 42–45. (doi:10.3725/jjn1993.29.2_42)
45. Siddiqui Z, Mahmood I. 1999 Role of bacteria in the management of plant parasitic nematodes: a review. *Biores. Tech.* **69**, 167–179. (doi:10.1016/S0960-8524(98)00122-9)
46. Samac DA, Kinkel LL. 2001 Suppression of the root-lesion nematode (*Pratylenchus penetrans*) in alfalfa (*Medicago sativa*) by *Streptomyces* spp. *Plant Soil* **235**, 35–44. (doi:10.1023/A:1011820002779)
47. Baquiran J-P, Thater B, Sedky S, De Ley P, Crowley D, Orwin PM. 2013 Culture-independent investigation of the microbiome associated with the nematode *Acrobeloides maximus*. *PLoS ONE* **8**, e67425. (doi:10.1371/journal.pone.0067425)
48. Cheng X-Y, Tian X-L, Wang Y-S, Lin R-M, Mao Z-C, Chen N, Xie B-Y. 2013 Metagenomic analysis of the pinewood nematode microbiome reveals a symbiotic relationship critical for xenobiotics degradation. *Sci. Rep.* **3**, 1869. (doi:10.1038/srep01869)
49. Tian X, Cheng X, Mao Z, Chen G, Yang J, Xie B. 2011 Composition of bacterial communities associated with a plant–parasitic nematode *Bursaphelenchus mucronatus*. *Curr. Biol.* **62**, 117–125. (doi:10.1007/s00284-010-9681-7)
50. Siddiqui IA, Ehteshamul-Haque S. 2001 Suppression of the root rot–root knot disease complex by *Pseudomonas aeruginosa* in tomato: the influence of inoculum density, nematode populations, moisture and other plant-associated bacteria. *Plant Soil* **237**, 81–89. (doi:10.1023/A:1013313103032)
51. Siddiqui I, Shaikat S. 2004 Systemic resistance in tomato induced by biocontrol bacteria against the root-knot nematode, *Meloidogyne javanica* is independent of salicylic acid production. *J. Phytopathol.* **152**, 48–54. (doi:10.1046/j.1439-0434.2003.00800.x)
52. Siddiqui IA, Shaikat SS, Sheikh IH, Khan A. 2006 Role of cyanide production by *Pseudomonas fluorescens* CHA0 in the suppression of root-knot nematode, *Meloidogyne javanica* in tomato. *World J. Microbiol. Biotechnol.* **22**, 641–650. (doi:10.1007/s11274-005-9084-2)
53. Siddiqui IA, Haas D, Heeb S. 2005 Extracellular protease of *Pseudomonas fluorescens* CHA0, a biocontrol factor with activity against the root-knot nematode *Meloidogyne incognita*. *Appl. Environ. Microbiol.* **71**, 5646–5649. (doi:10.1128/AEM.71.9.5646-5649.2005)
54. Zou CG, Tu HH, Liu XY, Tao N, Zhang KQ. 2010 PacC in the nematophagous fungus *Clonostachys rosea* controls virulence to nematodes. *Environ. Microbiol.* **12**, 1868–1877. (doi:10.1111/j.1462-2920.2010.02191.x)
55. Baloyi M, Laing M, Yobo K. 2011 Isolation and in vitro screening of *Bacillus thuringiensis* and *Clonostachys rosea* as biological control agents against sheep nematodes. *African J. Agr. Res.* **6**, 5047–5054. (doi:10.5897/AJAR11.261)
56. Zhang L, Yang J, Niu Q, Zhao X, Ye F, Liang L, Zhang K-Q. 2008 Investigation on the infection mechanism of the fungus *Clonostachys rosea* against nematodes using the green fluorescent protein. *Appl. Microbiol. Biotech.* **78**, 983–990. (doi:10.1007/s00253-008-0012-9)
57. Zou CG, Tao N, Liu WJ, Yang JK, Huang XW, Liu XY, Tu HH, Gan ZW, Zhang KQ. 2010 Regulation of subtilisin-like protease prC expression by nematode cuticle in the nematophagous fungus *Clonostachys rosea*. *Environ. Microbiol.* **12**, 3243–3252. (doi:10.1111/j.1462-2920.2010.02296.x)
58. Xie H, Aminuzzaman F, Xu L, Lai Y, Li F, Liu X. 2010 Trap induction and trapping in eight nematode-trapping fungi (Orbiliaceae) as affected by juvenile stage of *Caenorhabditis elegans*. *Mycopathologia* **169**, 467–473.
59. Liu B, Liu X-Z, Zhuang W-Y. 2005 *Orbilbia querci* sp. nov. and its knob-forming nematophagous anamorph. *FEMS Microbiol. Lett.* **245**, 99–105. (doi:10.1016/j.femsle.2005.02.027)
60. Yang Y, Yang E, An Z, Liu X. 2007 Evolution of nematode-trapping cells of predatory fungi of the Orbiliaceae based on evidence from rRNA-encoding DNA and multiprotein sequences. *Proc. Natl. Acad. Sci. USA* **104**, 8379–8384. (doi:10.1073/pnas.0702770104)
61. Yu H, Duan J, Wang B, Jiang X. 2012 The function of snodprot in the cerato-platanin family from *Dactylellina cionopaga* in nematophagous fungi. *Biosci. Biotech. Biochem.* **76**, 1835–1842. (doi:10.1271/bbb.120173)
62. Yang J, Liang L, Zhang Y, Li J, Zhang L, Ye F, Gan Z, Zhang K-Q. 2007 Purification and cloning of a novel serine protease from the nematode-trapping fungus *Dactylellina varietas* and its potential roles in infection against nematodes. *Appl. Microbiol. Biotech.* **75**, 557–565. (doi:10.1007/s00253-007-0839-6)
63. Moosavi MR, Zare R. 2012 Fungi as biological control agents of plant-parasitic nematodes. In *Plant defence: biological control* (eds J Méillon, K Ramawat), pp. 67–107. New York, NY: Springer.
64. Timper P. 2011 Utilization of biological control for managing plant-parasitic nematodes. In *Biological control of plant-parasitic nematodes* (eds J Méillon, K Ramawat), pp. 259–289. New York, NY: Springer.
65. Wang G, Liu Z, Lin R, Li E, Mao Z, Ling J, Yang Y, Yin W-B, Xie B. 2016 Biosynthesis of antibiotic leucinostatin in bio-control fungus *Purpureocillium lilacinum* and their inhibition on *Phytophthora* revealed by genome mining. *PLoS Pathog.* **12**, e1005685. (doi:10.1371/journal.ppat.1005685)
66. Lopez-Lima D, Carrion G, Núñez-Sánchez ÁE. 2014 Isolation of fungi associated with *Criconeimoides* sp. and their potential use in the biological control of ectoparasitic and semiendoparasitic nematodes in sugar cane. *Austral. J. Crop Sci.* **8**, 389.
67. Goffré, D, Folgarait P. 2015 *Purpureocillium lilacinum*, potential agent for biological control of the leaf-cutting ant *Acromyrmex lundi*. *J. Invert. Pathol.* **130**, 107–115. (doi:10.1016/j.jip.2015.07.008)
68. Lopez DC, Sword GA. 2015 The endophytic fungal entomopathogens *Beauveria bassiana* and *Purpureocillium lilacinum* enhance the growth of cultivated cotton (*Gossypium hirsutum*) and negatively affect survival of the cotton bollworm (*Helicoverpa zea*). *Biol. Control* **89**, 53–60. (doi:10.1016/j.biocontrol.2015.03.010)
69. Siddiqui I, Atkins S, Kerry B. 2009 Relationship between saprotrophic growth in soil of different biotypes of *Pochonia chlamydosporia* and the infection of nematode eggs. *Annu. Appl. Biol.* **155**, 131–141. (doi:10.1111/j.1744-7348.2009.00328.x)
70. Niu X-M, Wang Y-L, Chu Y-S, Hse X-H, Li N, Wei L-X, Mo M-H, Zhang K-Q. 2009 Nematodetoxic aurovertin-type metabolites from a root-knot nematode parasitic fungus *Pochonia chlamydosporia*. *J. Agric. Food Chem.* **58**, 828–834. (doi:10.1021/jf903259n)
71. Tobin J, Haydock P, Hare M, Woods S, Crump D. 2008 Effect of the fungus *Pochonia chlamydosporia* and fosthiazate on the multiplication rate of potato cyst nematodes (*Globodera pallida* and *G. rostochiensis*) in potato crops grown under UK field conditions. *Biol. Control* **46**, 194–201. (doi:10.1016/j.biocontrol.2008.03.014)

72. Atkins SD, Hidalgo-Diaz L, Clark IM, Morton CO, De Oca NM, Kerry BR. 2003 Approaches for monitoring the release of *Pochonia chlamydosporia* var. *catenulata*, a biocontrol agent of root-knot nematodes. *Mycol. Res.* **107**, 206–212. (doi:10.1017/S095375620300724X)
73. Esser R. 1983 Fungi that utilize zoospores to parasitize nematodes. *Nematol. Circular* No.101, Department of Agriculture and Consumer Service, Division of Plant Industry.
74. Barr D. 1970 Two varieties of *Rhizophyidum sphaerocarum* (Chytridiales). *Can. J. Bot.* **48**, 1067–1071.
75. Dickie IA. 2007 Host preference, niches and fungal diversity. *New Phytol.* **174**, 230–233. (doi:10.1111/j.1469-8137.2007.02055.x)
76. Kuruppu P, Schneider R, Russin J. 2004 Factors affecting soybean root colonization by *Calonectria ilicicola* and development of red crown rot following delayed planting. *Plant Disease* **88**, 613–619. (doi:10.1094/PDIS.2004.88.6.613)
77. Vitale A, Crous P, Lombard L, Polizzi G. 2013 *Calonectria* diseases on ornamental plants in Europe and the Mediterranean basin: an overview. *J. Plant Pathol.* **95**, 463–476. (doi:10.4454/JPP.V95I3.007)
78. Verma M, Brar SK, Tyagi R, Surampalli R, Valero J. 2007 Antagonistic fungi, *Trichoderma* spp.: panoply of biological control. *Biochem. Engin. J.* **37**, 1–20. (doi:10.1016/j.bej.2007.05.012)
79. Toju H, Kishida O, Katayama N, Takagi K. 2016 Networks depicting the fine-scale co-occurrences of fungi in soil horizons. *PLoS ONE* **11**, e0165987. (doi:10.1371/journal.pone.0165987)
80. Koenning S, Schmitt D, Barker K. 1993 Effects of cropping systems on population density of *Heterodera glycines* and soybean yield. *Plant Disease* **77**, 780–786. (doi:10.1094/Pd-77-0780)
81. Chen S. 2007 Tillage and crop sequence effects on *Heterodera glycines* and soybean yields. *Agronomy J.* **99**, 797–807.
82. Jaffee B. 2004 Do organic amendments enhance the nematode-trapping fungi *Dactylellina haptotyla* and *Arthrobotrys oligospora*? *J. Nematol.* **36**, 267.
83. Jaffee B. 2006 Interactions among a soil organic amendment, nematodes, and the nematode-trapping fungus *Dactylellina candidum*. *Phytopathology* **96**, 1388–1396. (doi:10.1094/PHYTO-96-1388)
84. Hershman D, Bachi P. 1995 Effect of wheat residue and tillage on *Heterodera glycines* and yield of doublecrop soybean in Kentucky. *Plant Disease* **79**, 631–633. (doi:10.1094/Pd-79-0631)
85. Verbruggen E, Toby Kiers E. 2010 Evolutionary ecology of mycorrhizal functional diversity in agricultural systems. *Evol. Appl.* **3**, 547–560. (doi:10.1111/j.1752-4571.2010.00145.x)