

ABSTRACTS (PH D THESIS)

Generation of transgenic rice with altered lignin composition and comparative characterization of their biomass utilization properties**(Graduate School of Agriculture, Laboratory of Metabolic Science of Forest Plants and Microorganisms, RISH, Kyoto University)****Yuri Takeda**

Lignocellulose, a complex bio-composite composed of lignin, cellulose, and hemicelluloses, constitutes the secondary cell walls of vascular plants and accounts for the highest proportion of terrestrial biomass on Earth. Hence, sustainable production and utilization of lignocellulosic biomass are crucial. Trees and grass biomass crops represent major lignocellulose feedstocks. While the annual biomass production of typical tree species is estimated to be less than 20 t ha⁻¹ year⁻¹, some large-sized grass biomass crops, such as *Sorghum*, sugarcane, and *Erianthus*, produce several-fold higher biomass yields compared with those of typical trees [1]. In addition, delignification of grass lignocellulose is generally easier than that of wood lignocellulose [1]. These characteristics render grass biomass crops potential materials for biochemical and thermochemical platforms and/or combustion use, although trees are indispensable for wood-based materials and for pulp and paper production.

Lignin is mainly composed of three different types of aromatic component, *p*-hydroxyphenyl (H), guaiacyl (G), and syringyl (S) units. Lignin aromatic composition, i.e., H/G/S lignin unit ratio, has been considered to be an important structural trait that affects the physico-chemical properties of lignocellulosic biomass. However, our knowledge regarding the relationship between the lignin aromatic composition and biomass utilization characteristics is yet considerably limited especially in monocotyledonous grass species. The main objectives of this PhD research were therefore to generate novel transgenic rice plants that produce lignocellulose with distinct lignin aromatic composition and then, through comparative characterization of their lignocellulose properties, to elucidate the effect of lignin composition on the utilization properties of grass lignocellulose as a source of biofuels and biomaterials.

First, transgenic rice lines augmented with G and S lignin units were successfully obtained through manipulating coniferaldehyde 5-hydroxylase (*CAld5H*) gene expression. Among three *CAld5H* genes identified in rice, *OsCAld5H1* appeared to be predominantly expressed in lignin-producing rice vegetative tissues. RNAi-mediated down-regulation of *OsCAld5H1* resulted in altered lignins largely enriched in G units, whereas up-regulation of *OsCAld5H1* under the control of a modified ubiquitin promoter resulted in lignins substantially enriched in S units, as demonstrated by a series of wet-chemical and NMR structural analyses. The results collectively demonstrate that *OsCAld5H1* expression is a key factor controlling the S/G lignin composition in rice cell walls [2].

To further investigate the effect of *OsCAld5H1*-suppression on rice lignin structure, the author characterized loss-of-function mutants of *OsCAld5H1* generated by the CRISPR/Cas9 genome editing system. A series of cell wall analysis demonstrated that, although lignins in the mutant were predictably enriched in G units, all tested mutant lines produced considerable amounts of S units. Intriguingly, lignin γ -*p*-coumaroylation analysis by the DFRC method revealed that the enrichment of G lignin units in the *OsCAld5H1*-knockout mutants was limited to the non- γ -*p*-coumaroylated lignin units, whereas grass-specific γ -*p*-coumaroylated lignin units were apparently unaffected. These data suggested that CAld5H is mainly involved in the production of non- γ -*p*-coumaroylated S lignin units, common in both eudicots and grasses, but not in the production of grass-specific γ -*p*-coumaroylated S units in rice [3].

Second, transgenic rice lines augmented with H lignin units were successfully obtained through manipulating *p*-coumaroyl ester 3-hydroxylase (*C3'H*) gene expression. *C3'H*-knockdown lines generated via RNAi-driven gene silencing, with about 0.5% of the residual expression levels, reached maturity and set seeds, whereas *C3'H*-knockout rice mutants generated via the CRISPR/Cas9-mediated targeted mutagenesis were severely dwarfed and sterile. Cell wall analysis of the mature *C3'H*-knockdown RNAi lines revealed that their lignins were largely enriched in H units while being substantially reduced in the normally dominant G and S units. Interestingly, however, lignin γ -*p*-coumaroylation analysis by the DFRC

 ABSTRACTS (PH D THESIS)

method revealed that the enrichment of H units was limited to the non-acylated lignin units, with grass-specific γ -*p*-coumaroylated lignin units remaining apparently unchanged. These data suggested that, similar to what the author has shown for CA1d5H, C3'H is mainly involved in the production of non- γ -*p*-coumaroylated lignin units, but not in the production of grass-specific γ -*p*-coumaroylated lignin units in rice. Suppression of C3'H also resulted in a substantial reduction in wall cross-linking ferulates. These results demonstrate that C3'H expression is an important determinant not only of lignin content and composition but also of the degree of cell wall cross-linking in grasses [4].

The author further characterized the CRISPR/Cas9-mediated C3'H-knockout mutants. In line with their severely impaired growth phenotype, the C3'H-knockout lines appeared to show irregular vasculature and ectopic lignification in culm and root. Transcriptomic analysis data suggested that, along with altered expressions of the phenylpropanoid biosynthetic genes, gene expressions related to various biological processes were perturbed in the C3'H-knockout lines. Lignin analysis of the mutant cell walls demonstrated that, as observed in the C3'H-knockdown lines, H lignin units were substantially increased but the augmentation of H lignin units in the mutants were limited to non γ -*p*-coumaroylated lignin units, with grass-specific γ -*p*-coumaroylated lignin units being mostly unaffected, further supporting the author's notion that there is C3'H-independent metabolic pathway(s) responsible for the production of grass-specific γ -*p*-coumaroylated lignin units.

Lastly, the author used the above-described transgenic rice lines to study the impact of the altered lignin composition on the chemical reactivity, enzymatic saccharification efficiency and calorific value of rice lignocellulose. It was suggested that H and G lignins are more susceptible to acid-induced condensation reactions than S lignins. The H-lignin-enriched transgenic rice was shown to display significantly enhanced biomass saccharification efficiency without any pretreatment or with alkali and acid pretreatments, whereas the S-lignin-enriched transgenic rice showed further enhanced saccharification efficiency after liquid hot water pretreatment. While no significant differences in the biomass heating values were observed between the transgenic rice materials tested, analysis of synthetic lignins comprising only H, G or S units suggested that increasing H or G units may be beneficial to increase the heating value of biomass [5].

Acknowledgements

The author thanks Professor Dr. Toshiyuki Takano and Dr. Masahiro Sakamoto, Graduated school of Agriculture, Kyoto University, Dr. Taichi Koshiba and Dr. Shinya Murakami, RISH, Kyoto University, Professor Dr. John Ralph and Dr. Steven D. Karlen, University of Wisconsin, Professor Dr. Keishi Osakabe, Dr. Yuriko Osakabe and Dr. Takefumi Hattori, Tokushima University, Professor Dr. Hironori Kaji and Ms. Ayaka Maeno, Institute for Chemical Research (ICR), Kyoto University, and Dr. Kazuaki Katagiri, Mr. Shimpei Yamaguchi and Mr. Takuya Ehiro, Osaka Research Institute of Industrial Science and Technology, for their scientific advices and technical supports. The author also thanks Dr. Masaharu Kuroda, National Agriculture and Food Research Organization, Professor Dr. Ko Shimamoto and Dr. Daisuke Miki, Nara Institute of Science and Technology, and Professor Dr. Seiichi Toki, Dr. Masaki Endo and Mr. Masafumi Mikami, NARO, for providing transformation vectors. The author acknowledges the Japan Society for the Promotion of Science (JSPS) for financial supports (#17J09654). Part of this study was conducted using the facilities in the DASH/FBAS at RISH, Kyoto University, and the NMR spectrometer in the JURC at ICR, Kyoto University.

Reference

[1] Umezawa (2018) *Phytochem. Rev.* 17:1305-1327; [2] Takeda *et al.* (2017) *Planta* 246:337-349; [3] Takeda *et al.* (2018) *Plant J.* 95:796-811; [4] Takeda *et al.* (2019) *Plant J.* 97:543-554; [5] Takeda *et al.* (2019) *J. Wood Sci.* 65:6.