

1 **Predominant accumulation of a 3-hydroxy- γ -decalactone in the male**
2 **rectal gland complex of the Japanese orange fly, *Bactrocera tsuneonis***

3

4 Hajime Ono^{a,1}, Masataka Nakahira^{a,1}, Satoshi Ohno^{a,1}, Jun Otake^a, Tomoya
5 Kanno^a, Isao Tokushima^a, Yoshimitsu Higashiura^b, Ichiro Nishi^b and Ritsuo
6 Nishida^a

7

8 ^aGraduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan

9 ^bYamaguchi Prefectural Agriculture and Forestry General Technology Center, Yamaguchi
10 742-2805, Japan

11 ¹These authors contributed equally to this work

12

13

14

15

16

17

18

19

20 CONTACT

21 Hajime Ono onoono@kais.kyoto-u.ac.jp; Ritsuo Nishida ritz@kais.kyoto-u.ac.jp

22

23 Running Head: A γ -decalactone from the Japanese orange fly

24 ABSTRACT

25 The Japanese orange fly, *Bactrocera tsuneonis*, infests various citrus crops. While male
26 pheromone components accumulated in the rectal glands are well-characterized for
27 *Bactrocera*, but information regarding the chemical factors involved in the life cycles of
28 *B. tsuneonis* remains scarce. Herein, several volatile chemicals including a γ -decalactone,
29 (3*R*,4*R*)-3-hydroxy-4-decanolide [(3*R*,4*R*)-HD], were identified as major components,
30 along with acetamide and spiroketals as minor components in the rectal gland complexes
31 of male *B. tsuneonis* flies. The lactone (3*R*,4*R*)-HD was also identified in female rectal
32 gland complexes. The amount of this compound in mature males was significantly higher
33 than those observed in females and immature males. The lactone (3*R*,4*R*)-HD was
34 detected in flies fed with sucrose only, indicating that this lactone is not derived from
35 dietary sources during adulthood, but biosynthesized *in vivo*. The predominant
36 accumulation of (3*R*,4*R*)-HD in mature males also suggests a possible role in reproductive
37 behavior.

38

39 KEYWORDS

40 (3*R*,4*R*)-3-hydroxy-4-decanolide; *Bactrocera tsuneonis*; Tephritidae; rectal gland
41 complex; pest fruit fly

42

43 Many dacine fruit fly species comprising a major subfamily of Tephritidae are destructive
44 pests of fruits and vegetables [1]. The Japanese orange fly, *Bactrocera tsuneonis* (Miyake),
45 is an important pest of citrus fruits, in particular mandarin orange [2,3]. *B. tsuneonis* is a
46 univoltine and oligophagous species widely distributed in southwestern China, Taiwan,
47 and Japan. In Japan, the distribution of *B. tsuneonis* was restricted in the southern island
48 Kyushu, but it has recently spread into the adjacent areas of western Honshu and the
49 Shikoku islands. Hence, an effective lure to monitor *B. tsuneonis* population is urgently
50 needed. However, little is known about the semiochemical factors of the life cycles of *B.*
51 *tsuneonis*.

52 In most dacine tephritid fly species, males furnish a glandular complex in the
53 rectum, known as the rectal gland, that is considered to be a secretory organ of male sex
54 pheromones [4]. Various compounds, including aliphatic and aromatic volatiles, have
55 been identified from the rectal glands [5]. In many dacine species, males acquire
56 pheromonal components from plant secondary metabolites during adulthood [6]. For
57 example, males of the Oriental fruit fly, *Bactrocera dorsalis*, are strongly attracted to a
58 specific phenylpropanoid, methyl eugenol (ME), which is an essential oil component of
59 various plants, and subsequently feed voraciously on the compound. [7,8]. The ingested
60 ME is then biotransformed into two sex pheromone components *in vivo*, and these
61 metabolites are subsequently sequestered in the male rectal gland to attract conspecific
62 females [9,10]. Because of its robust attractiveness to male flies, ME has been used as a
63 lure for pest management programs [11,12]. In the olive fly, multiple female sex
64 pheromones are secreted by the female rectal gland [13]. Thus, the identification of
65 chemical substances in the rectal glands of tephritid fruit flies is very important to
66 understand the biological significance of these compounds in their life cycles and to

67 develop effective lures for pest managements [6].

68 To characterize the semiochemicals involved in *B. tsuneonis* life cycles, we
69 analyzed the volatile chemical composition of their rectal tissues. We identified a
70 predominant and unique hydroxy γ -lactone, along with an acetamide and a series of
71 spiroketals in both males and females. The stereochemistry of the γ -lactone was
72 determined and the rectal constituents were quantified in the context of maturation and
73 sex differences. We also determined whether the γ -lactone was synthesized from dietary
74 sources during adulthood.

75

76 Results and discussion

77 *Identification of rectal gland components*

78 A rectal gland complex was dissected from a mature male fly. An ethanolic extract of the
79 tissue was analyzed by gas chromatography-mass spectrometry (GC-MS) (Figure 1). A
80 major component (**1**) and several characteristic minor components, **2-5**, as well as general
81 insect wax components including higher hydrocarbons were identified. The EI-MS and
82 CI-MS of compound **1** afford major ions of $[M-H_2O]^+$ at m/z 168 and $[M+H]^+$ at m/z 187,
83 respectively, for a possible molecular formula of $C_{10}H_{18}O_3$. We dissected rectal gland
84 complexes from 117 laboratory-eclosed males and extracted their contents to isolate **1** for
85 further analyses. Approximately 0.7 mg of **1** was obtained by a silica gel column
86 chromatography. The ^{13}C -NMR, DEPT, and HMQC spectra revealed 10 carbon signals.
87 These signals were assigned to one methyl carbon (δ 14.2), six methylene carbons (δ 39.6,
88 31.8, 29.3, 28.4, 25.7 and 22.7), two methine carbons (δ 84.7 and 69.3), and one carbonyl
89 carbon (δ 175.3) from an ester/lactone group. The 1H -NMR spectrum revealed
90 characteristic signals derived from two protons adjacent to oxygens (δ 4.49 and 4.37) and

91 one pair of geminal protons (δ 2.80 and 2.56). The ^1H - ^1H COSY spectrum indicated a
92 connectivity of H2-H3-H4-H5-H6 and HMBC spectrum provided the following
93 correlations: from H2 to C1, C3 and C4; from H3 to C1 and C4; from H4 to C3; and from
94 H10 to C8 and C9. Therefore, we assigned the structure of **1** to be a γ -decalactone, 3-
95 hydroxy-4-decanolide (HD). Because **1** has four possible stereoisomers, including two
96 pairs of enantiomers due to the two chiral centers at C-3 and C-4, we synthesized a
97 racemic mixture of *erythro*-lactones, and two optical isomers of *threo*-HD to determine
98 the unambiguous structure of **1**. We synthesized racemic *erythro*-lactones—(3*R*,4*S*)-HD
99 and (3*S*,4*R*)-HD—from (*E*)-3-decenoic acid via epoxidation and subsequent lactonization
100 using amberlyst-15 [15] (Figure S1A). In addition, (3*R*,4*R*)- and (3*S*,4*S*)-HD were
101 synthesized via Sharpless asymmetric dihydroxylation of methyl (*E*)-3-decenoate using
102 AD-mix- β and AD-mix- α , respectively [16,17] (Figure S1B). Comparison of the ^1H -
103 NMR and ^{13}C -NMR spectra of **1** with those of the racemic *erythro*-lactones and each
104 enantiomeric *threo*-lactone indicated that the configuration of **1** was a *threo*-HD, either
105 (3*R*,4*R*)-HD or (3*S*,4*S*)-HD (Figure S5 and S10). Because the optical rotation of **1** ($[\alpha]_{\text{D}}^{23}$
106 +61.5) corresponded to that of (3*R*,4*R*)-HD ($[\alpha]_{\text{D}}^{22}$ +40.1), rather than that of (3*S*,4*S*)-HD
107 ($[\alpha]_{\text{D}}^{21}$ -35.1), the absolute structure was confirmed as (3*R*,4*R*)-HD. To the best of our
108 knowledge, this is the first identification of **1** from insect species, as this compound has
109 been reported only from culture media as a fermented yeast product (*Yarrowia lipolytica*)
110 derived from an artificially added lipid [18].

111 We identified the minor components of the rectal volatiles to be *N*-(3-
112 methylbutyl)acetamide (**2**), (*E,E*)-2,8-dimethyl-1,7-dioxaspiro[5.5]undecane (**3**), and
113 (*E,Z*)-2,8-dimethyl-1,7-dioxaspiro[5.5]undecane (**4**) by comparing their retention times
114 and mass spectra with those of the corresponding synthetic and authentic samples. In

115 addition, **5** was putatively identified as 2-methyl-8-ethyl-1,7-dioxaspiro[5.5]undecane by
116 comparing its mass spectrum with a mass spectrum library (Wiley7Nist05.L). The mass
117 spectral data are provided in Table S1 and Figure S11. These compounds were previously
118 identified from the male rectal glands of several tephritid species [5].

119

120 *Quantification of rectal volatile components*

121 In the Dacini genera, male pheromones are accumulated in the rectal gland and play a
122 vital role in mating behavior [6]. Therefore, the major components in the rectal tissues, **1**
123 and **2**, were compared between sexes to determine whether these compounds were
124 distributed in a sexually dimorphic pattern. Female rectal gland complexes were dissected
125 as previously described [14] and subsequently extracted by ethanol. The amounts of **1** and
126 **2** were compared between wild males and females captured in a mandarin orange orchard.
127 While **1** and **2** were detected in both wild males and females, the amount of **1** in males
128 was significantly higher than that in females (Figure 2). The mean **1** content in the male
129 samples was approximately 16 µg/gland, but the quantities varied among individuals
130 (ranging from 0.87 to 33.2 µg/gland). The quantities of **2** were smaller than those of **1**,
131 with mean of 496 and 135 ng/gland in males and females, respectively. While the amount
132 of **2** in males was higher than that in females, no significant difference was detected
133 between the sexes. The variation in **1** content among wild males can be ascribed to age.
134 Indeed, morphological changes in the rectal gland are observed with age, and the storage
135 and secretion in the reservoir of the gland increase with male sexual maturation [4,19].
136 Therefore, we quantified **1** and **2** in flies eclosed in a laboratory at different ages: within
137 three days after eclosion (0–3 d AE) for sexually immature flies; as well as 7–10 d AE
138 and 21–24 d AE for mature flies. A small amount of **1** was detected in males at 0–3 d AE

139 and significant increases were observed in both mature males at 7–10 and 21–24 d AE
140 (Figure 3). The mean **1** contents were 0.30, 6.73 and 4.00 $\mu\text{g/gland}$ in males at 0–3, 7–10,
141 and 21–24 d AE, respectively, indicating that **1** increased by > 20 times with maturation.
142 Similarly, significant enhancement of **2** was observed in females at 7–10 d AE. It should
143 be noted that the quantities of this compound varied widely in the 7–10 d AE females, i.e.,
144 more than 1.9 $\mu\text{g/gland}$ of **1** was detected in 7 females, while $< 0.4 \mu\text{g/gland}$ was detected
145 in 6 females ($n = 13$). These results indicate that **1** accumulates in males as a function of
146 maturation. Small amounts of **2** were detected in both immature males and females. While
147 the various quantities of **2** (ranging from 0 to 1.57 $\mu\text{g/gland}$) were observed in all male
148 groups, a significant enhancement in its content was observed in males at 7–10 d AE.
149 Similarly, an enhancement of **2** was observed in females at 7–10 d AE, but two distinct
150 patterns were observed, i.e., 7 females contained $> 670 \text{ ng/gland}$, but 6 females contained
151 $< 130 \text{ ng/gland}$ ($n = 13$). This dimorphic pattern was observed for both **1** and **2**, so we
152 examined relationship of the contents of these compounds among individual females. The
153 correlation between the contents of **1** and **2** showed that some females contain significant
154 amounts of these compounds, while others contain only small quantities (Figure S2).

155 Because the male sex pheromones of dacine species are accumulated or are
156 synthesized from dietary pheromone or pheromone precursors, sequestered pheromones
157 can only be detected from male rectal glands after feeding the pheromone or its precursor
158 directly [20]. We further examined whether *B. tsuneonis* biosynthesized **1** and **2** from
159 dietary sources or *de novo*. The flies were fed exclusively with sucrose solution to prevent
160 ingestion of complex food as a dietary source for components in the rectal tissues. Even
161 the flies were fed with only sucrose after eclosion, **1** and **2** were detected in both males
162 and females at 7–10 d AE (Figure 4). The amounts of **1** in flies fed with only sucrose were

163 similar to those in flies fed with normal food in both males and females. The amounts of
164 **2** in flies fed with only sucrose were smaller than those in flies fed with normal food, but
165 no significant difference was observed between feeding groups. The contents of **1** and **2**
166 in males and females regardless of their dietary ingredients indicate that these compounds
167 are not directly derived from ingested food but are synthesized *in vivo*.

168 In preliminary indoor behavioral experiments, we examined the attractiveness of
169 sexually mature male and female flies to an extract of the rectal gland complex, **1**, **2**, and
170 a synthetic mixture of **1** and **2**, in a small screen cage. However, neither mature male nor
171 female flies were attracted to the tested samples. Further, we conducted a field trap
172 experiment with synthetic mixtures of **1**, **2**, **3**, and **4** in various combinations/doses in
173 orange orchards in Yamaguchi Prefecture during the outbreak season of the adult flies.
174 However, no *B. tsuneonis* adults were captured by those traps (unpublished data).
175 Although a physiological function of these volatile components in the rectal gland
176 complex could not be determined, **1** was accumulated in mature males, similar to the male
177 sex pheromones of other *Bactrocera* fruit flies. The distinct pattern in the contents in
178 mature females suggest that this compound may play a role in reproduction of *B.*
179 *tsuneonis*, such as an indicator of sexual maturation and chemical signal during mating
180 events. Elucidation of the roles of these components in the rectal tissues could provide a
181 clue to control this hardly controllable pest fruit fly.

182

183 Experimental

184 *Insects*

185 Last instar larvae of *B. tsuneonis* immediately before pupariation were obtained from
186 mandarin fruits in the local citrus orchards of Suo-Oshima Island, Yamaguchi, and kept

187 in a laboratory in the Yamaguchi Prefectural Agriculture and Forestry General
188 Technology Center, Yamaguchi, Japan. The emerged pupae were kept indoors under
189 ambient temperatures from November to June of the next year. The adults eclosed at the
190 first half of June were provided with water and a diet of four parts sucrose and one part
191 dry yeast AY-65 (Asahi Food & Healthcare, Ltd., Tokyo, Japan) at 25 °C and subjected
192 to a 16 h light/8 h dark cycle. For analysis of the rectal glands from the sugar-only feed
193 group, male and female flies were fed with 2 % sucrose solution after eclosion. For
194 analysis of the rectal gland volatiles of wild fruit flies, males and females were captured
195 in August 2018 in the local citrus orchards of Suo-Oshima Island, Yamaguchi.

196

197 *Instruments*

198 GC-Mass spectra were measured using an Agilent 5975 inert XL MSD mass spectrometer
199 coupled with an Agilent 6890 gas chromatograph equipped with a capillary column (HP-
200 5MS, 29 m × 0.25 mm, 0.25 µm film thickness, helium as a carrier gas) programmed
201 from 60 °C (2 min holding) to 290 °C at a rate of 10 °C/min. GC quantification was
202 performed using an HP-6850 gas chromatograph equipped with an Agilent HP-5MS 5%
203 phenyl methyl siloxane-coated capillary column (15 m × 0.25 mm, 0.25 µm film
204 thickness) with a flame ionization detector using the same program for GC-MS analyses
205 for the GC oven. The ¹H-NMR and ¹³C-NMR spectra were measured using a Bruker
206 Avance III 500 spectrometer with TMS as an internal standard. The optical rotations were
207 measured using a JASCO P-1010 spectropolarimeter.

208

209 *Rectal sample preparation*

210 The rectal gland complexes were dissected from adult flies. Contents of the tissue were

211 extracted with 250 μ L of ethanol per gland for GC quantification or GC-MS analysis.

212

213 *Extraction and purification of compound 1 from male rectal glands*

214 Rectal gland complexes were dissected from 117 males eclosed indoor and extracted with

215 ethanol. The combined extract (17 mg) was subjected to chromatography on a silica gel

216 (500 mg) and eluted with 20% methyl acetate in hexane to isolate the major rectal volatile,

217 compound **1** (yield: approximately 700 μ g). $[\alpha]_D^{23} +61.5$ ($c = 0.065$, CH₃OH). EI-MS: m/z

218 (%) 168 (3, [M-H₂O]⁺), 158 (1), 139 (12), 126 (7), 115 (40), 97 (85), 83 (19), 69 (23), 55

219 (100), 43 (64). CI-MS (CH₄): m/z (%) 187 (24, [M+H]⁺), 169 (74), 151 (63), 127 (100),

220 109 (55). ¹H-NMR (CDCl₃): δ 4.49 (1H, m, H-3), 4.37 (1H, m, H-4), 2.80 (1H, dd, $J =$

221 17.6, 5.6 Hz, H-2), 2.56 (1H, dd, $J = 17.6, 1.0$ Hz, H-2), 1.88 (1H, m, H-5), 1.71 (1H, m,

222 H-5), 1.55–1.25 (8H, m, H-6, H-7, H-8 and H-9), 0.90 (3H, t, $J = 7.0$ Hz, H-10). ¹³C-

223 NMR (CDCl₃): δ 175.3 (C-1), 84.7 (C-4), 69.3 (C-3), 39.6 (C-2), 31.8 (C-8 or C-9), 29.3

224 (C-7), 28.4 (C-5), 25.7 (C-6), 22.7 (C-8 or C-9), 14.2 (C-10).

225

226 *Synthesis of a racemic mixture of erythro-lactones*

227 First, mCPBA (6.65 g) was added to a solution of (*E*)-methyl dec-3-enoate (2.76 g) in

228 CH₂Cl₂ (10 mL), and the mixture was stirred at room temperature for 20 h. The reaction

229 was quenched by addition of a 5% aqueous Na₂SO₃ solution. After filtration and

230 evaporation, the reaction mixture was extracted using diethyl ether, washed with water

231 and brine, and subsequently dried. Epoxy ester (2.04 g, yield: 67%) was obtained as a

232 racemic mixture. Amberlyst-15 (1.0 g) was added to a solution of epoxy ester (2.0 g) in

233 benzene (50 mL), and the mixture was stirred at room temperature for 18 h. After filtration

234 and evaporation, the reaction mixture (1.35 g) was subjected to silica gel chromatography

235 eluted with 50% methyl acetate in hexane to afford a racemic mixture of *erythro*-
236 lactones—(3*R*,4*S*)-HD and (3*S*,4*R*)-HD—(720 mg, yield: 39%) with a minor *threo*-
237 lactone (Figure S1). The racemic *erythro*-lactone was further purified via HPLC on a
238 silica gel column (YMC-Pack SIL, 5 μ m, 10 \times 300 mm, YMC Co., Ltd., Japan, flow rate
239 of 2.5 mL/min with 60% ethyl acetate in hexane; yield: 21 mg). The spectral data are
240 provided in Table S2.

241

242 *Synthesis of each enantiomeric threo-lactone*

243 First, (*E*)-3-decenoic acid was esterified with methanol containing 1% H₂SO₄ to yield
244 methyl (*E*)-3-decenoate. AD-mix- α or - β (0.7 g) was added to 50% aqueous *tert*-BuOH
245 (5 mL) and stirred at room temperature until two clear phases appeared. CH₃SO₂NH₂ (49
246 mg) was added to this solution and subsequently cooled to 0 °C. Methyl (*E*)-3-decenoate
247 (90 mg) was added to the solution and the reaction mixture was stirred at 0 °C for 24 h.
248 The reaction was quenched by addition of NaHSO₃ and extracted with EtOAc. The extract
249 was washed with 2 N NaOH and brine, and subsequently dried. (3*R*,4*R*)-3-HD (85 mg,
250 yield: 82%) and (3*S*,4*S*)-3-HD (82 mg, yield: 80%) were obtained from the reaction with
251 AD-mix- β and AD-mix- α , respectively. (Figure S1B). The spectral data are provided in
252 Table S2.

253

254 *Synthesis of the minor components of the rectal gland*

255 First, *N*-(3-methylbutyl)acetamide (**2**) was obtained by acetylation of 3-
256 methylbutylamine using acetic anhydride. Subsequently, (*E,E*)- and (*E,Z*)-2,8-dimethyl-
257 1,7-dioxaspiro[5.5]undecanes (**3** and **4**) were synthesized as previously described [21]
258 and the corresponding mass spectral data are shown in Table S3.

259

260 *Quantification of rectal volatiles*

261 A 1 or 5 μL portion of each rectal tissue extract (1 gland/250 μL) was subjected to GC
262 quantifications with 1-pentadecanol (Wako Pure Chemical Industries, Japan) as an
263 internal standard. The contents of the relevant compounds in the rectal tissues were
264 determined by comparing FID intensities with those of standard samples with known
265 concentrations.

266

267

268 **Author contributions**

269 HO and RN conceived and designed research. HO, MN, SO, JO, TK, IT, YH, IN and RN
270 conducted experiments and analyzed data. HO and RN wrote the manuscript. All authors
271 read and approved the manuscript.

272

273 **Acknowledgements**

274 This work was supported by the Project of the NARO Biooriented Technology Research
275 Advancement Institution (the special scheme project on regional developing strategy) to
276 HO and YH; Matsushima Horticultural Development Foundation to HO; and JSPS
277 KAKENHI Grant Number 18H02214 to HO.

278

279 **Disclosure statement**

280 No potential conflict of interest was reported by the authors.

281

282 **References**

- 283 [1] Fletcher BS. The biology of dacine fruit flies. *Annu Rev Entomol* 1987;32:115–
284 144.
- 285 [2] Yasumatsu K, Nagatomi A. Studies on the control of *Dacus (Tetradacus)*
286 *tsuneonis* Miyake (Diptera: Trypetidae). 1. Some fundamental and biological
287 investigations essential for its control. *Kyushu Univ Fac Agr Sci Bull*
288 1959;17:129–146.
- 289 [3] Weems Jr. HV. Japanese orange fly (*Dacus tsuneonis* Miyake) (Diptera:
290 Tephritidae). *Entomol Circ* 1967;56:1–2.
- 291 [4] Fletcher BS. Storage and release of a sex pheromone by the Queensland fruit fly,
292 *Dacus tryoni* (Diptera: Trypetidae). *Nature* 1968;219:631–632.
- 293 [5] Fletcher MT, Kitching W. Chemistry of fruit flies. *Chem Rev* 1995;95:789–828.
- 294 [6] Tan KH, Nishida R, Jang EB, et al. Pheromones, male lures, and trapping of
295 tephritid fruit flies. In: Shelly T, Epsky N, Jang EB, Reyes-Flores, Vargas R,
296 editors. *Trapping and the detection, control, and regulation of tephritid fruit flies*.
297 Dordrecht: Springer; 2014. 15–74.
- 298 [7] Nishida R, Tan K, Serit M, et al. Accumulation of phenylpropanoids in the rectal
299 glands of male Oriental fruit fly, *Dacus dorsalis*. *Experientia* 1988;44:534–536.
- 300 [8] Tan KH, Nishida R. Methyl eugenol: Its occurrence, distribution, and role in
301 nature, especially in relation to insect behavior and pollination. *J Insect Sci*
302 2012;12:1–60.
- 303 [9] Tan, KH, Nishida R. Sex pheromone and mating competition after methyl
304 eugenol consumption in the *Bactrocera dorsalis* complex. In: McPherson BA,
305 Steck GJ, editors. *Fruit fly pests: a world assessment of their biology and*
306 *management*. Delray Beach (FL): St. Lucie Press; 1996. 147–153.
- 307 [10] Hee AK, Tan K. Male sex pheromonal components derived from methyl eugenol
308 in the hemolymph of the fruit fly *Bactrocera papayae*. *J Chem Ecol*
309 2004;30:2127–2138.
- 310 [11] Steiner LF, Mitchell WC, Harris EJ, et al. Oriental fruit fly eradication by male
311 annihilation. *J Econ Entomol* 1965;58:961–964.
- 312 [12] Koyama J, Teruya T, Tanaka K. Eradication of the oriental fruit fly (Diptera:
313 Tephritidae) from the Okinawa Islands by a male annihilation method. *J Econ*
314 *Entomol* 1984;77:468–472.
- 315 [13] Baker R, Herbert R, Howse PE, et al. Identification and synthesis of the major
316 sex pheromone of the olive fly (*Dacus oleae*). *J Chem Soc, Chem Commun*
317 1980;52–53.
- 318 [14] Mazomenos G.E., Haniotakis BE. A multicomponent female sex pheromone of

- 319 *Dacus oleae* Gmelin: Isolation and bioassay. J Chem Ecol 1981;7:437–444.
- 320 [15] Jefford CW, Wang Y. A simple, general distereoselective synthesis of 5-
321 hydroxyalkylbutan-4-olides. J Chem Soc 1987;1513–1514.
- 322 [16] Wang ZM, Zhang XL, Sharpless KB, et al. A general approach to γ -lactones via
323 osmium-catalyzed asymmetric dihydroxylation. Synthesis of (-)- and (+)-
324 muricatacin. Tetrahedron Lett 1992;33:6407–6410.
- 325 [17] Garbe LA, Morgenthal K, Kuscher K, et al. Metabolism of deuterated erythro-
326 dihydroxy fatty acids in *Saccharomyces cerevisiae*: Enantioselective formation
327 and characterization of hydroxylactones. Helv Chim Acta 2008;91:993–1007.
- 328 [18] García EE, Aguedo M, Gomes N, et al. Production of 3-hydroxy- γ -decalactone,
329 the precursor of two decenolides with flavouring properties, by the yeast
330 *Yarrowia lipolytica*. J Mol Catal B 2009;57:22–26.
- 331 [19] Fletcher BS. The structure and function of the sex pheromone glands of the male
332 Queensland fruit fly, *Dacus tryoni*. J Insect Physiol 1969;15:1309–1322.
- 333 [20] Nishida R, Iwahashi O, Tan KH. Accumulation of *Dendrobium superbum*
334 (Orchidaceae) fragrance in the rectal glands by males of the melon fly, *Dacus*
335 *cucurbitae*. J Chem Ecol 1993;19:713–722.
- 336 [21] Schwartz BD, Hayes PY, Kitching W, et al. Flexible synthesis of
337 enantiomerically pure homopropargylic alcohols. J Org Chem 2005;3054–3065.
338
339

340 Figure captions

341

342 Figure 1. Representative total ion chromatogram of an ethanol extract from a male rectal
343 gland of *Bactrocera tsuneonis*. The structures of the main components are also shown. **1**:
344 (3*R*,4*S*)-3-Hydroxy-4-decanolide, **2**: *N*-(3-Methylbutyl)acetamide, **3**: (*E,E*)- and (*E,Z*)-
345 2,8-Dimethyl-1,7-dioxaspiro[5.5]undecane, **4**: (*E,Z*)-2,8-Dimethyl-1,7-
346 dioxaspiro[5.5]undecane, **5**: 2-Methyl-8-ethyl-1,7-dioxaspiro[5.5]undecane.

347

348 Figure 2. The amounts of compounds **1** and **2** in the wild male and female rectal gland
349 complexes of *Bactrocera tsuneonis* where each value is plotted as a dot ($n = 5-7$). The
350 box plot shows 25–75% (box), median (band inside), and minima to maxima (whiskers).
351 Welch's *t*-test: $p < 0.01$.

352

353 Figure 3. The amounts of compounds **1** and **2** in immature (0–3-d AE) and mature (7–10
354 or 21–24-d AE) flies of *Bactrocera tsuneonis* where each value is plotted as a dot ($n = 6-$
355 20). Boxes with different letters are significantly different at $p < 0.05$ as determined by
356 Steel-Dwass test.

357

358 Figure 4. The amounts of compounds **1** and **2** in mature flies of *Bactrocera tsuneonis* 7–
359 10-d AE fed with normal food or sucrose solution only where each value is plotted as a
360 dot ($n = 6-15$). The box plot shows 25–75% (box), median (band inside), and minima to
361 maxima (whiskers). Boxes with different letters are significantly different at $p < 0.05$ as
362 determined by Steel-Dwass test.

Abundance









Fig. 1









Fig. 2




Fig. 3








Fig. 4



A



B







Fig. S1

Fig. S2




CI-MS


Fig. S3



A



B



C




Fig. S5





Fig. S6





Fig. S7




Fig. S8







Fig. S9




A (3*R*,4*R*)-3-Hydroxy-4-decanolide




B (3*R*,4*S*)- and (3*S*,4*R*)-3-Hydroxy-4-decanolide




A




C



B



D



SUPPLEMENTARY MATERIAL

Predominant accumulation of a 3-hydroxy- γ -decalactone in the male rectal gland complex of the Japanese orange fly, *Bactrocera tsuneonis*

Hajime Ono^{a,1*}, Masataka Nakahira^{a,1}, Satoshi Ohno^{a,1}, Jun Otake^a, Tomoya Kanno^a, Isao Tokushima^a, Yoshimitsu Higashiura^b, Ichiro Nishi^b and Ritsuo Nishida^{a*}

^aGraduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan

^bYamaguchi Prefectural Agriculture and Forestry General Technology Center, Yamaguchi 742-2805, Japan

*Corresponding authors. E-mails: onoono@kais.kyoto-u.ac.jp; ritz@kais.kyoto-u.ac.jp

¹These authors contributed equally to this work

Contents:

Table S1. EI-MS spectral data of the extract of the male rectal gland complexes.

Table S2. Spectral data of the synthesized 3-hydroxy- γ -decalactones.

Table S3. EI-MS spectral data of the synthesized minor components in the rectal gland.

Figure S1. Synthetic routes for 3-hydroxy- γ -decalactone isomers. (A) racemic (\pm)-*erythro*- γ -lactone. (B) (+)- and (-)-*threo*- γ -lactones: (3*R*,4*R*)-3-hydroxy-4-decanolide and (3*S*,4*S*)-3-hydroxy-4-decanolide.

Figure S2. Correlation diagram between amounts of **1** ((3*R*,4*S*)-3-hydroxy-4-decanolide) and **2** (*N*-(3-methylbutyl)acetamide) in individual females from 7- to 10-days after eclosion. Data are derived from Figure 2B.

Figure S3. CI-Mass spectra of (3*R*,4*R*)-3-hydroxy-4-decanolide (**1**) isolated from *Bactrocera tsuneonis* (A), and synthesized **1** (B).

Figure S4. EI-Mass spectra. (A) (3*R*,4*R*)-3-hydroxy-4-decanolide (**1**) isolated from *Bactrocera tsuneonis*. (B) synthesized **1**. (C) racemic (3*R*,4*S*)- and (3*S*,4*R*)-3-hydroxy-4-decanolide.

Figure S5. ¹H NMR spectrum of **1** isolated from *Bactrocera tsuneonis*.

Figure S6. ¹³C NMR spectrum of **1** isolated from *Bactrocera tsuneonis*.

Figure S7. ¹H-¹H COSY spectrum of **1** isolated from *Bactrocera tsuneonis*.

Figure S8. HMQC spectrum of **1** isolated from *Bactrocera tsuneonis*.

Figure S9. HMBC spectrum of **1** isolated from *Bactrocera tsuneonis*.

Figure S10. ¹H NMR spectra of synthesized 3-hydroxy-4-decanolide. (A) (3*R*,4*R*)-3-hydroxy-4-decanolide. (B) racemic (3*R*,4*S*)- and (3*S*,4*R*)-3-hydroxy-4-decanolide.

Figure S11. EI-Mass spectra of the minor components (**2-5**) contained in the rectal gland complexes of *Bactrocera tsuneonis*. (A) *N*-(3-methylbutyl)acetamide (**2**). (B) (*E,E*)-2,8-dimethyl-1,7-dioxaspiro[5.5]undecane (**3**). (C) (*E,Z*)-2,8-dimethyl-1,7-dioxaspiro[5.5]undecane (**4**). (D) 2-methyl-8-ethyl-1,7-dioxaspiro[5.5]undecane (**5**).

Table S1. EI-MS spectral data of the extract of the male rectal gland complexes.

Compound	<i>Rt</i> (min)	Mass fragments [<i>m/z</i> (%)]
1	16.7	168 (2, [M-H ₂ O] ⁺), 158 (1), 139 (8), 126 (5), 115 (39), 97 (84), 83 (15), 69 (23), 55 (100), 43 (60).
2	9.31	129 (8, M ⁺), 114 (17), 86 (33), 73 (98), 72 (87), 60 (38), 55 (22), 43 (100).
3	9.49	184 (11, M ⁺), 169 (3), 140 (17), 125 (11), 115 (94), 112 (100), 97 (51), 83 (14), 69 (27), 55(26), 43 (23).
4	10.6	184 (6, M ⁺), 169 (3), 140 (6), 125 (8), 115 (100), 112 (47), 97 (57), 83 (9), 69 (34), 55(23), 43 (16).
5	10.8	198 (15, M ⁺), 183 (2), 169 (24), 154 (12), 140 (20), 129 (56), 126 (45), 115 (100), 112 (89), 97 (48), 83 (55), 69 (39), 55 (40), 44 (48).

Table S2. Spectral data of the synthesized 3-hydroxy- γ -decalactones.

A mixture of (3*R*,4*S*)- and (3*S*,4*R*)-3-hydroxy-4-decanolide

EI-MS: m/z (%) 168 (0.2, [M-H₂O]⁺), 158 (2), 139 (2), 126 (3), 115 (53), 97 (100), 83 (10), 69 (18), 55 (81), 43 (40).

¹H-NMR (CDCl₃): δ 4.37 (1H, m, H-4), 4.27 (1H, m, H-3), 2.82 (1H, dd, $J = 18.0, 6.7$ Hz, H-2), 2.51 (1H, dd, $J = 18.0, 3.6$ Hz, H-2), 1.68-1.54 (2H, m, H-5), 1.50-1.24 (8H, m, H-6, H-7, H-8 and H-9), 0.89 (3H, t, $J = 6.9$ Hz, H-10).

¹³C-NMR (CDCl₃): δ 175.9 (C-1), 88.3 (C-3), 71.6 (C-4), 37.7 (C-2), 33.1, 31.6, 29.0, 25.2, 22.5, 14.0 (C-10).

(3*R*,4*R*)-3-Hydroxy-4-decanolide

$[\alpha]_D^{25} +40.1$ ($c = 1.1$, CH₃OH, 22°C)

EI-MS: m/z (%) 168 (0.3, [M-H₂O]⁺), 158 (2), 139 (3), 126 (3), 115 (53), 97 (100), 83 (11), 69 (22), 55 (99), 43 (51).

CI-MS: m/z (%) 187 (31, [M+H]⁺), 169 (82), 151 (63), 127 (100), 109 (51).

¹H-NMR (CDCl₃): δ 4.48 (1H, m, H-3), 4.37 (1H, m, H-4), 2.80 (1H, dd, $J = 17.7, 5.5$ Hz, H-2), 2.56 (1H, dd, $J = 17.7, 0.5$ Hz, H-2), 1.87 (1H, m, H-5), 1.72 (1H, m, H-5), 1.55-1.25 (8H, m, H-6, H-7, H-8 and H-9), 0.89 (3H, t, $J = 7.0$ Hz, H-10).

¹³C-NMR (CDCl₃): δ 175.5 (C-1), 84.8 (C-4), 69.1 (C-3), 39.5 (C-2), 31.6 (C-8 or C-9), 29.1 (C-7), 28.3 (C-5), 25.5 (C-6), 22.6 (C-8 or C-9), 14.0 (C-10).

(3*S*,4*S*)-3-Hydroxy-4-decanolide

$[\alpha]_D^{25} -35.1$ ($c = 1.1$, CH₃OH, 21°C)

Table S3. EI-MS spectral data of the synthesized minor components in the rectal gland.

Compound	Mass fragments [m/z (%)]
<i>N</i> -(3-Methylbutyl)acetamide (2)	129 (12, M ⁺), 114 (22), 86 (35), 73 (100), 72 (85), 60 (37), 55 (20), 43 (84).
(<i>E,E</i>)-2,8-Dimethyl-1,7-dioxaspiro[5.5]undecane (3)	184 (13, M ⁺), 169 (3), 140 (22), 125 (12), 115 (96), 112 (100), 97 (55), 83 (17), 69 (31), 55 (29), 43 (24).
(<i>E,Z</i>)-2,8-Dimethyl-1,7-dioxaspiro[5.5]undecane (4)	184 (7, M ⁺), 169 (3), 140 (6), 125 (7), 115 (100), 112 (45), 97 (54), 83 (9), 69 (35), 55(23), 43 (17).




Figure S1. Synthetic routes for 3-hydroxy- γ -decalactone isomers. (A) racemic (\pm)-*erythro*- γ -lactone. (B) (+)- and (-)-*threo*- γ -lactones: (3*R*,4*R*)-3-hydroxy-4-decanolide and (3*S*,4*S*)-3-hydroxy-4-decanolide.




Figure S2. Correlation diagram between amounts of **1** ((3*R*,4*S*)-3-hydroxy-4-decanolide) and **2** (*N*-(3-methylbutyl)acetamide) in individual females from 7- to 10-days after eclosion. Data are derived from Figure 2B.

CI-MS





Figure S3. CI-Mass spectra of (3R,4R)-3-hydroxy-4-decanolide (**1**) isolated from *Bactrocera tsuneonis* (A), and synthesized **1** (B).

EI-MS




Figure S4. EI-Mass spectra. (A) (3R,4R)-3-hydroxy-4-decanolide (**1**) isolated from *Bactrocera tsuneonis*. (B) synthesized **1**. (C) racemic (3R,4S)- and (3S,4R)-3-hydroxy-4-decanolide.




Figure S5. ¹H NMR spectrum of **1** isolated from *Bactrocera tsuneonis*.




Figure S6. ¹³C NMR spectrum of **1** isolated from *Bactrocera tsuneonis*.




Figure S7. ^1H - ^1H COSY spectrum of **1** isolated from *Bactrocera tsuneonis*.




Figure S8. HMQC spectrum of **1** isolated from *Bactrocera tsuneonis*.




Figure S9. HMBC spectrum of **1** isolated from *Bactrocera tsuneonis*.





Figure S10. ^1H NMR spectra of synthesized 3-hydroxy-4-decanolide. (A) (3*R*,4*R*)-3-hydroxy-4-decanolide. (B) racemic (3*R*,4*S*)- and (3*S*,4*R*)-3-hydroxy-4-decanolide.




Figure S11. EI-Mass spectra of the minor components (**2-5**) contained in the rectal gland complexes of *Bactrocera tsuneonis*. (A) *N*-(3-methylbutyl)acetamide (**2**). (B) (*E,E*)-2,8-dimethyl-1,7-dioxaspiro[5.5]undecane (**3**). (C) (*E,Z*)-2,8-dimethyl-1,7-dioxaspiro[5.5]undecane (**4**). (D) 2-methyl-8-ethyl-1,7-dioxaspiro[5.5]undecane (**5**).