

Truncation error estimates of approximate operators in a generalized particle method

Yusuke Imoto

Received: date / Accepted: date

Abstract To facilitate the numerical analysis of particle methods, we derive truncation error estimates for the approximate operators in a generalized particle method. Here, a generalized particle method is defined as a meshfree numerical method that typically includes other conventional particle methods, such as smoothed particle hydrodynamics or moving particle semi-implicit methods. A new regularity of discrete parameters is proposed via two new indicators based on the Voronoi decomposition of the domain along with two hypotheses of reference weight functions. Then, truncation error estimates are derived for an interpolant, approximate gradient operator, and approximate Laplace operator in the generalized particle method. The convergence rates for these estimates are determined based on the frequency with which they appear in the regularity and hypotheses. Finally, the estimates are computed numerically, and the results are shown to be in good agreement with the theoretical results.

Keywords generalized particle method · truncation error estimate · approximate operator · smoothed particle hydrodynamics method · moving particle semi-implicit method

1 Introduction

Particle methods, such as the smoothed particle hydrodynamics (SPH) [10, 18, 19] and moving particle semi-implicit (MPS) methods [15, 16, 29], are numerical methods for solving partial differential equations that are based on points called particles distributed in a domain. In such methods, an interpolant and several approximate differential operators are defined in terms of linear combinations of weighted interactions between neighboring particles. When such methods are applied to partial differential equations, the equations are effectively discretized in space. As the discretization procedure does not require mesh generation in the

Y. Imoto
Institute for Advanced Study, Kyoto University, Yoshida Konoe-cho, Sakyo-ku, Kyoto, Japan
Tel.: +81-75-753-9441
E-mail: imoto.yusuke.4e@kyoto-u.ac.jp

domain, particle methods can be applied to moving boundary problems, such as the deformation and destruction of structures [5, 22] and flow problems associated with free surfaces [21, 23].

The accuracy of particle methods has been widely researched. From an engineering perspective, many studies have been conducted into the convergence of such methods in practical applications, such as Amicarelli [1, 2], Fulk [9], and Quinlan et al. [25]. On the other hand, few studies in the literature have presented numerical analyses of these methods from a mathematical perspective. In the 1980s, Mas-Gallic and Raviart [20] and Raviart [26] provided error estimates for particle methods when applied to parabolic and hyperbolic partial differential equations on unbounded domains. In the 2000s, Ben Moussa and Via [4] and Ben Moussa [3] provided error estimates of nonlinear conservation laws on bounded domains. In their work, the time integrations of the particle positions and volumes were obtained by solving the differential equations with respect to advection fields. However, as their method is only applicable to problems described by solvable differential equations, it cannot be used with other problems, such as those involving the Navier–Stokes equations.

Sometime later, Ishijima and Kimura [13] developed a truncation error estimate for an approximate gradient operator in the MPS method. By introducing a regularity for particle distributions based on an indicator called the equivolume partition radius, they determined the conditions that depend solely on the space distributions of the particles. However, a practical limitation is that the indicator cannot be computed.

In previous works, we established truncation error estimates for an interpolant, approximate gradient operator, and approximate Laplace operator of a generalized particle method in which the particle volumes were given as Voronoi volumes [11, 12]. A generalized particle method is a numerical method that typically includes conventional particle methods, such as the SPH and MPS methods. In previous studies, we derived truncation error estimates by introducing a regularity using an indicator known as the covering radius, which is used in the numerical analysis of meshfree methods based on moving least-square methods and radial basis functions [17, 27, 30]. Although the formulations and conditions in those works are computable, they are difficult to deploy in practical computations as the computational costs associated with particle volumes based on Voronoi decomposition are high.

The focus of the current work was to analyze particle methods under more practical conditions by extending our results to cases with commonly used particle volumes. We also introduce another indicator of particle volumes, which we refer to as a Voronoi deviation, that represents the deviation between particle volumes and Voronoi volumes. Then, utilizing the Voronoi deviation, we extend the regularity and introduce two hypotheses of reference weight functions. Using the regularity and hypotheses, we derive truncation error estimates of the interpolant, approximate gradient operator, and approximate gradient operator of the generalized particle method. Finally, we numerically analyze our estimates and compare the results to those from the theory.

The remainder of this paper is organized as follows. The interpolant and approximate operators of the generalized particle method are introduced in Section 2. A regularity describing the family of discrete parameters is discussed in Section 3, after which we propose our primary theorem with respect to the truncation error

estimates and provide some corollaries. Then, the primary theorem is proven in Section 4, numerical results are detailed in Section 5, and some concluding remarks are outlined in Section 6.

In the remainder of this section, we describe some notation and define some relevant function spaces. Let \mathbb{R}^+ , \mathbb{R}_0^+ , and \mathbb{N}_0 be the set of positive real numbers, the set of nonnegative real numbers, and the set of nonnegative integers, respectively. Let d be the dimension of a space. Let \mathbb{A}^d be the set of all d -dimensional multi-indices. For $x = (x_1, x_2, \dots, x_d)^\top \in \mathbb{R}^d$ and $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d)^\top \in \mathbb{A}^d$, x^α is defined as $x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_d^{\alpha_d}$. If there is no ambiguity, the symbol $|\cdot|$ is used to denote the following: $|x|$ denotes the Euclidean norm for $x \in \mathbb{R}^d$; $|S|$ denotes the volume of S for $S \subset \mathbb{R}^d$; $|\alpha|$ denotes $|\alpha| := \alpha_1 + \alpha_2 + \dots + \alpha_d$ for $\alpha \in \mathbb{A}^d$. For $S \subset \mathbb{R}^d$, let $\text{diam}(S)$ be $\text{diam}(S) := \sup\{|x - y|; x, y \in S\}$. For $S \subset \mathbb{R}^d$, let $C(\bar{S})$ be the space of real continuous functions defined in \bar{S} with the norm $\|\cdot\|_{C(\bar{S})}$ defined as

$$\|v\|_{C(\bar{S})} := \max_{x \in \bar{S}} |v(x)|.$$

For $S \subset \mathbb{R}^d$ and $\ell \in \mathbb{N}$, let $C^\ell(\bar{S})$ be the space of functions in $C(\bar{S})$ with derivatives up to the ℓ th order with its seminorm $|\cdot|_{C^\ell(\bar{S})}$ and norm $\|\cdot\|_{C^\ell(\bar{S})}$ defined as

$$\begin{aligned} |v|_{C^\ell(\bar{S})} &:= \max_{\alpha \in \mathbb{A}^d, |\alpha|=\ell} \|D^\alpha v\|_{C(\bar{S})}, \\ \|v\|_{C^\ell(\bar{S})} &:= \max_{j=0,1,\dots,\ell} |v|_{C^j(\bar{S})}, \end{aligned}$$

respectively. Here $D^\alpha v := \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_d^{\alpha_d} v$ with multi-index $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d)$.

2 Approximate operators in a generalized particle method

Let Ω be a bounded domain in \mathbb{R}^d . Let H be a fixed positive number. For Ω and H , we define extended domain Ω_H as

$$\Omega_H := \left\{ x \in \mathbb{R}^d \mid \exists y \in \Omega \text{ s.t. } |x - y| < H \right\}.$$

For $N \in \mathbb{N}$, we define a particle distribution \mathcal{X}_N and particle volume set \mathcal{V}_N as

$$\begin{aligned} \mathcal{X}_N &:= \{x_i \in \Omega_H; i = 1, 2, \dots, N, \quad x_i \neq x_j (i \neq j)\}, \\ \mathcal{V}_N &:= \left\{ V_i \in \mathbb{R}^+; i = 1, 2, \dots, N, \quad \sum_{i=1}^N V_i = |\Omega_H| \right\}, \end{aligned}$$

respectively. We refer to $x_i \in \mathcal{X}_N$ and $V_i \in \mathcal{V}_N$ as a particle and particle volume, respectively. An example of the particle distribution \mathcal{X}_N in $\Omega_H (\subset \mathbb{R}^2)$ is shown in Figure 1.

We define an admissible reference weight function set \mathcal{W} as

$$\mathcal{W} := \left\{ w \in C(\mathbb{R}_0^+); \text{supp}(w) = [0, 1], \int_{\mathbb{R}^d} w(|x|) dx = 1, \text{absolutely continuous} \right\},$$

we refer to $w \in \mathcal{W}$ as a reference weight function, and we define the influence radius $h_N \in \mathbb{R}$ as satisfying $0 < h_N < H$ and $h_N \rightarrow 0 (N \rightarrow \infty)$. If there is




Fig. 1 Particle distribution X_N in $\Omega_H \subset \mathbb{R}^2$.

no ambiguity, we denote h_N as h . For reference weight function w and influence radius h , we define the weight function $w_h \in C(\mathbb{R}_0^+)$ as

$$w_h(r) := \frac{1}{h^d} w\left(\frac{r}{h}\right). \quad (1)$$

Note that the weight function w_h satisfies

$$\text{supp}(w_h) = [0, h], \quad \int_{\mathbb{R}^d} w_h(|x|) dx = 1,$$

and is absolutely continuous.

For $v \in C(\overline{\Omega}_H)$, we define interpolant Π_h , approximate gradient operator ∇_h , and approximate Laplace operator Δ_h as

$$\Pi_h v(x) := \sum_{i \in \Lambda_0(x, h)} V_i v(x_i) w_h(|x_i - x|), \quad (2)$$

$$\nabla_h v(x) := d \sum_{i \in \Lambda(x, h)} V_i \frac{v(x_i) - v(x)}{|x_i - x|} \frac{x_i - x}{|x_i - x|} w_h(|x_i - x|), \quad (3)$$

$$\Delta_h v(x) := 2d \sum_{i \in \Lambda(x, h)} V_i \frac{v(x_i) - v(x)}{|x_i - x|^2} w_h(|x_i - x|), \quad (4)$$

respectively. Here, for $x \in \mathbb{R}^d$ and $r \in \mathbb{R}^+ \cup \{\infty\}$, $\Lambda_0(x, r)$ and $\Lambda(x, r)$ are index sets of particles defined as

$$\begin{aligned} \Lambda_0(x, r) &:= \{i = 1, 2, \dots, N; 0 \leq |x - x_i| < r\}, \\ \Lambda(x, r) &:= \{i = 1, 2, \dots, N; 0 < |x - x_i| < r\}, \end{aligned}$$

respectively.

As discussed later in Appendix A, the approximate operators (2), (3), and (4) indicate a wider class of approximate operators of particle methods than those in the SPH and MPS methods. Therefore, we refer to the approximate operators (2), (3), and (4) as generalized approximate operators and to a particle method that uses them as a generalized particle method.




Fig. 2 Example of the Voronoi decomposition of Ω_H associated with the particle distribution \mathcal{X}_N .

3 Truncation error estimates of approximate operators

We first introduce a regularity of discrete parameters. Let $\{\sigma_i\}$ be the Voronoi decomposition of Ω_H associated with the particle distribution \mathcal{X}_N , where σ_i is the Voronoi region defined as

$$\sigma_i := \{x \in \Omega_H; |x_i - x| < |x_j - x|, \forall x_j \in \mathcal{X}_N (j \neq i)\}, \quad i = 1, 2, \dots, N.$$

We define a particle volume decomposition $\Xi = \{\xi_i\}$ as a decomposition of Ω_H satisfying

$$|\xi_i| = V_i, \quad \bigcup_{i=1}^N \bar{\xi}_i = \bar{\Omega}_H \quad (i = 1, 2, \dots, N), \quad \xi_i \cap \xi_j = \emptyset \quad (i \neq j).$$

An example of the Voronoi decomposition of Ω_H associated with the particle distribution \mathcal{X}_N is shown in Figure 2. We define a covering radius r_N for particle distribution \mathcal{X}_N as

$$r_N := \max_{i=1,2,\dots,N} \sup_{x \in \sigma_i} |x_i - x|. \quad (5)$$

Moreover, we define a Voronoi deviation d_N for the particle distribution \mathcal{X}_N and the particle volume set \mathcal{V}_N as

$$d_N := \inf_{\Xi} d_{\Xi} \quad (6)$$

with

$$d_{\Xi} := \max_{i=1,2,\dots,N} \left\{ \sum_{j=1}^N \frac{|\sigma_i \cap \xi_j| + |\xi_i \cap \sigma_j|}{|\sigma_i|} |x_i - x_j| \right\}.$$

Then, we define a regularity for a family consisting of a particle distribution \mathcal{X}_N , particle volume set \mathcal{V}_N , and influence radius h as follows:

Definition 1 A family $\{(\mathcal{X}_N, \mathcal{V}_N, h_N)\}_{N \rightarrow \infty}$ is said to be regular with order m ($m \geq 1$) if there exists a positive constant c_0 such that

$$h_N^m \geq c_0(r_N + d_N), \quad \forall N \in \mathbb{N}. \quad (7)$$




Fig. 3 Two examples of covering radii r_N for particle distributions with same number of particles. The covering radius r_N for the uniform particle distribution (left) is smaller than that for the non-uniform particle distribution (right).

Remark 1 As shown in Figure 3, the covering radius r_N becomes large in the case of a particle distribution with both dense and sparse regions. Therefore, the covering radius r_N can be considered as an indicator representing the uniformness of particle distribution \mathcal{X}_N .

Remark 2 A Voronoi deviation d_N equals zero if and only if the particle volumes are given as the Voronoi volume ($V_i = |\sigma_i|$). Moreover, the Voronoi deviation d_N becomes large if the particle volumes are given as values far from the Voronoi volumes. Therefore, the Voronoi deviation d_N can be regarded as an indicator of the deviation between the particle volume set and the Voronoi volume set.

Remark 3 For a given family $\{(\mathcal{X}_N, \mathcal{V}_N, h_N)\}_{N \rightarrow \infty}$ and given constant m ($m \geq 1$), it is possible to determine whether or not the family is regular with order m as the covering radius r_N and Voronoi deviation d_N are absolutely computable, as shown in Appendix B.

Next, we introduce two hypotheses of reference weight function w :

Hypothesis 1 For $n \in \mathbb{N}$, the reference weight function w satisfies for all $\alpha \in \mathbb{A}^d$ with $1 \leq |\alpha| \leq n$,

$$\int_{\mathbb{R}^d} x^\alpha w(|x|) dx = 0.$$

Hypothesis 2 For $k \in \mathbb{N}_0$, the reference weight function w satisfies

$$\max \left\{ \sup_{r \in (0,1)} |w^{(k+1)}(r)|, \sup_{r \in (0,1)} |(w^{(k)})'(r)| \right\} < \infty,$$

where for $j \in \mathbb{N}_0$, $w^{(j)}(r) : (0, \infty) \rightarrow \mathbb{R}$ is defined as

$$w^{(j)}(r) := \begin{cases} \lim_{s \downarrow 0} \frac{w(s)}{s^j}, & r = 0, \\ \frac{w(r)}{r^j}, & r > 0 \end{cases} \quad (8)$$

and $(w^{(k)})'$ is $dw^{(k)}/dr$.

Remark 4 All reference functions $w \in \mathcal{W}$ satisfy Hypothesis 1 with $n = 1$. Moreover, for all $n \in \mathbb{N}$ and $k \in \mathbb{N}$, reference weight functions satisfying Hypothesis 1 with n and Hypothesis 2 with k can be constructed as shown in Appendix C.

We now state a theorem that defines truncation error estimates of approximate operators in the generalized particle method with a continuous norm:

Theorem 3 *Suppose that a family $\{(\mathcal{X}_N, \mathcal{V}_N, h_N)\}_{N \rightarrow \infty}$ is regular with order $m (m \geq 1)$ and that reference weight function w satisfies Hypothesis 1 with n . Then, there exists a positive constant c independent of N such that*

$$\|v - \Pi_h v\|_{C(\bar{\Omega})} \leq c h^{\min\{m-1, n+1\}} \|v\|_{C^{n+1}(\bar{\Omega}_H)}, \quad v \in C^{n+1}(\bar{\Omega}_H). \quad (9)$$

In addition, if $w \in \mathcal{W}$ satisfies Hypothesis 2 with $k = 0$, then we have

$$\|\nabla v - \nabla_h v\|_{C(\bar{\Omega})} \leq c h^{\min\{m-1, n+1\}} \|v\|_{C^{n+2}(\bar{\Omega}_H)}, \quad v \in C^{n+2}(\bar{\Omega}_H), \quad (10)$$

and if $w \in \mathcal{W}$ satisfies Hypothesis 2 with $k = 1$, then we have

$$\|\Delta v - \Delta_h v\|_{C(\bar{\Omega})} \leq c h^{\min\{m-2, n+1\}} \|v\|_{C^{n+3}(\bar{\Omega}_H)}, \quad v \in C^{n+3}(\bar{\Omega}_H). \quad (11)$$

The proof of Theorem 3 is presented in the next section. As shown in the corollaries in Appendix A, the approximate operators commonly used in the SPH and MPS methods are valid for Theorem 3 under appropriate settings.

4 Proof of truncation error estimates

The following notation will be used in the subsequent proof of Theorem 3. Hereafter, let c be a generic positive constant independent of N (allowed dependence on the fixed positive parameter H). For $\alpha \in \mathbb{A}^d$, set I_α as

$$I_\alpha(x) := \sum_{i \in \mathcal{A}_0(x, h)} V_i (x_i - x)^\alpha w_h(|x_i - x|) - \int_{\mathbb{R}^d} y^\alpha w_h(|y|) dy, \quad x \in \bar{\Omega}.$$

For $\alpha \in \mathbb{A}^d$ and $\ell \in \mathbb{N}$, set $I_{\alpha, \ell}$ as

$$I_{\alpha, \ell}(x) := \sum_{i \in \mathcal{A}(x, h)} V_i \frac{(x_i - x)^\alpha}{|x_i - x|^\ell} w_h(|x_i - x|) - \int_{\mathbb{R}^d} \frac{y^\alpha}{|y|^\ell} w_h(|y|) dy, \quad x \in \bar{\Omega}.$$

For $\ell \in \mathbb{N}$, set J_ℓ as

$$J_\ell(x) := \sum_{i \in \mathcal{A}_0(x, h)} V_i |x_i - x|^\ell |w_h(|x_i - x|)|, \quad x \in \bar{\Omega}.$$

We now present the following lemma.

Lemma 1 *Suppose that $w \in \mathcal{W}$ satisfies Hypothesis 1 with n . Then, there exists a positive constant c independent of N such that*

$$\begin{aligned} & \|v - \Pi_h v\|_{C(\bar{\Omega})} \\ & \leq c \left(\sum_{0 \leq |\alpha| \leq n} \|I_\alpha\|_{C(\bar{\Omega})} + \|J_{n+1}\|_{C(\bar{\Omega})} \right) \|v\|_{C^{n+1}(\bar{\Omega}_H)}, \\ & v \in C^{n+1}(\bar{\Omega}_H), \end{aligned} \quad (12)$$

$$\begin{aligned} & \|\nabla v - \nabla_h v\|_{C(\bar{\Omega})} \\ & \leq c \left(\sum_{2 \leq |\alpha| \leq n+2} \|I_{\alpha,2}\|_{C(\bar{\Omega})} + \|J_{n+1}\|_{C(\bar{\Omega})} \right) \|v\|_{C^{n+2}(\bar{\Omega}_H)}, \\ & v \in C^{n+2}(\bar{\Omega}_H), \end{aligned} \quad (13)$$

$$\begin{aligned} & \|\Delta v - \Delta_h v\|_{C(\bar{\Omega})} \\ & \leq c \left(\sum_{1 \leq |\alpha| \leq n+3} \|I_{\alpha,2}\|_{C(\bar{\Omega})} + \|J_{n+1}\|_{C(\bar{\Omega})} \right) \|v\|_{C^{n+3}(\bar{\Omega}_H)}, \\ & v \in C^{n+3}(\bar{\Omega}_H). \end{aligned} \quad (14)$$

Proof First, we prove (12). We fix $x \in \bar{\Omega}$. Then, let $B(x, r)$ be the open ball in \mathbb{R}^d with center x and radius r , i.e.,

$$B(x, r) := \{y \in \mathbb{R}^d; |y - x| < r\}.$$

From $h < H$, we have $B(x, h) \subset \Omega_H$. Then, for all $v \in C^{\ell+1}(\bar{\Omega}_H)$ ($\ell \in \mathbb{N}$) and $x_i \in B(x, h)$, we obtain the Taylor expansion of v as

$$\begin{aligned} v(x_i) &= \sum_{0 \leq |\alpha| \leq \ell} \frac{D^\alpha v(x)}{\alpha!} (x_i - x)^\alpha + \sum_{|\alpha| = \ell+1} (x_i - x)^\alpha R_\alpha(x_i, x), \\ R_\alpha(x_i, x) &:= \frac{|\alpha|}{\alpha!} \int_0^1 (1-t)^{|\alpha|-1} D^\alpha v(tx + (1-t)x_i) dt. \end{aligned} \quad (15)$$

From (2) and (15) with $\ell = n$, we have

$$\begin{aligned} \Pi_h v(x) &= \sum_{0 \leq |\alpha| \leq n} \frac{D^\alpha v(x)}{\alpha!} \sum_{i \in \Lambda_0(x, h)} V_i(x_i - x)^\alpha w_h(|x_i - x|) \\ &+ \sum_{|\alpha| = n+1} \sum_{i \in \Lambda_0(x, h)} R_\alpha(x_i, x) V_i(x_i - x)^\alpha w_h(|x_i - x|). \end{aligned}$$

Moreover, by Hypothesis 1, we have

$$\begin{aligned} \Pi_h v(x) - v(x) &= \sum_{0 \leq |\alpha| \leq n} \frac{D^\alpha v(x)}{\alpha!} I_\alpha(x) \\ &+ \sum_{|\alpha|=n+1} \sum_{i \in \Lambda_0(x,h)} R_\alpha(x_i, x) V_i(x_i - x)^\alpha w_h(|x_i - x|). \end{aligned} \quad (16)$$

Because

$$|R_\alpha(y, z)| \leq \frac{1}{\alpha!} |v|_{C^{|\alpha|}(\overline{\Omega}_H)}, \quad y \in \overline{\Omega}, \quad z \in B(y, h), \quad \alpha \in \mathbb{A}^d, \quad (17)$$

we have

$$\begin{aligned} \left| \sum_{|\alpha|=n+1} \sum_{i \in \Lambda_0(x,h)} R_\alpha(x_i, x) V_i(x_i - x)^\alpha w_h(|x_i - x|) \right| \\ \leq c |J_{n+1}(x)| |v|_{C^{n+1}(\overline{\Omega}_H)}. \end{aligned} \quad (18)$$

Moreover, we have

$$\left| \sum_{0 \leq |\alpha| \leq n} \frac{D^\alpha v(x)}{\alpha!} I_\alpha(x) \right| \leq c \|v\|_{C^n(\overline{\Omega})} \sum_{0 \leq |\alpha| \leq n} |I_\alpha(x)|. \quad (19)$$

Therefore, from (16), (18), and (19), we obtain (12).

Next, we prove (13). From (3) and (15) with $\ell = n + 1$, we have

$$\begin{aligned} \nabla_h v(x) &= d \sum_{1 \leq |\alpha| \leq n+1} \frac{D^\alpha v(x)}{\alpha!} \sum_{i \in \Lambda(x,h)} V_i \frac{(x_i - x)(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|) \\ &+ d \sum_{|\alpha|=n+2} \sum_{i \in \Lambda(x,h)} R_\alpha(x_i, x) V_i \frac{(x_i - x)(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|). \end{aligned}$$

Because for $\beta \in \mathbb{A}^d$ with $|\beta| = 2$,

$$d \int_{\mathbb{R}^d} \frac{y^\beta}{|y|^2} w_h(|y|) dy = \begin{cases} 1, & \text{all elements of } \beta \text{ are even,} \\ 0, & \text{otherwise,} \end{cases} \quad (20)$$

we have

$$d \sum_{|\alpha|=1} \frac{D^\alpha v(x)}{\alpha!} \int_{\mathbb{R}^d} \frac{yy^\alpha}{|y|^2} w_h(|y|) dy = \nabla v(x). \quad (21)$$

Hypothesis 1 with n yields

$$\int_{\mathbb{R}^d} \frac{yy^\alpha}{|y|^2} w_h(|y|) dy = 0 \quad \alpha \in \mathbb{A}^d \text{ with } 2 \leq |\alpha| \leq n + 1. \quad (22)$$

From (21) and (22), we have

$$\begin{aligned} \nabla_h v(x) - \nabla v(x) &= -d \sum_{1 \leq |\alpha| \leq n+1} \frac{D^\alpha v(x)}{\alpha!} \int_{\mathbb{R}^d} \frac{yy^\alpha}{|y|^2} w_h(|y|) dy \\ &\quad + d \sum_{1 \leq |\alpha| \leq n+1} \frac{D^\alpha v(x)}{\alpha!} \sum_{i \in \Lambda(x,h)} V_i \frac{(x_i - x)(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|) \\ &\quad + d \sum_{|\alpha|=n+2} \sum_{i \in \Lambda(x,h)} R_\alpha(x_i, x) V_i \frac{(x_i - x)(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|). \end{aligned} \quad (23)$$

From (17), we have

$$\begin{aligned} \left| \sum_{|\alpha|=n+2} \sum_{i \in \Lambda(x,h)} R_\alpha(x_i, x) V_i \frac{(x_i - x)(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|) \right| \\ \leq c |J_{n+1}(x)| |v|_{C^{n+2}(\overline{\Omega}_H)}. \end{aligned} \quad (24)$$

Moreover, we have

$$\begin{aligned} \sum_{1 \leq |\alpha| \leq n+1} \left| \sum_{i \in \Lambda(x,h)} V_i \frac{(x_i - x)(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|) - \int_{\mathbb{R}^d} \frac{yy^\alpha}{|y|^2} w_h(|y|) dy \right| \\ \leq c \sum_{2 \leq |\alpha| \leq n+2} |I_{\alpha,2}(x)|. \end{aligned} \quad (25)$$

Therefore, from (23), (24), and (25), we obtain (13).

Finally, we prove (14). From (4) and (15) with $\ell = n + 2$, we have

$$\begin{aligned} \Delta_h v(x) &= 2d \sum_{1 \leq |\alpha| \leq n+2} \frac{D^\alpha v(x)}{\alpha!} \sum_{i \in \Lambda(x,h)} V_i \frac{(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|) \\ &\quad + 2d \sum_{|\alpha|=n+3} \sum_{i \in \Lambda(x,h)} R_\alpha(x_i, x) V_i \frac{(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|). \end{aligned}$$

From (20), we have

$$2d \sum_{|\alpha|=2} \frac{D^\alpha v(x)}{\alpha!} \int_{\mathbb{R}^d} \frac{y^\alpha}{|y|^2} w_h(|y|) dy = \Delta v(x).$$

Hypothesis 1 with n yields

$$\int_{\mathbb{R}^d} \frac{y^\alpha}{|y|^2} w_h(|y|) dy = 0, \quad \alpha \in \mathbb{A}^d \text{ with } |\alpha| = 1 \text{ or } 3 \leq |\alpha| \leq n + 2.$$

Therefore, we have

$$\begin{aligned} \Delta_h v(x) - \Delta v(x) &= 2d \sum_{1 \leq |\alpha| \leq n+2} \frac{D^\alpha v(x)}{\alpha!} I_{\alpha,2}(x) \\ &\quad + 2d \sum_{|\alpha|=n+3} \sum_{i \in \Lambda(x,h)} R_\alpha(x_i, x) V_i \frac{(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|). \end{aligned} \quad (26)$$

From (17), we have

$$\left| \sum_{|\alpha|=n+3} \sum_{i \in \Lambda(x,h)} R_\alpha(x_i, x) V_i \frac{(x_i - x)^\alpha}{|x_i - x|^2} w_h(|x_i - x|) \right| \leq c |J_{n+1}(x)| |v|_{C^{n+3}(\overline{\mathcal{O}}_H)}. \quad (27)$$

Moreover, we have

$$\left| \sum_{1 \leq |\alpha| \leq n+2} \frac{D^\alpha v(x)}{\alpha!} I_{\alpha,2}(x) \right| \leq c \|v\|_{C^{n+2}(\overline{\mathcal{O}})} \sum_{1 \leq |\alpha| \leq n+2} |I_{\alpha,2}(x)|. \quad (28)$$

Therefore, from (26), (27), and (28), we obtain (14).

Next, we show estimates of I_α , $I_{\alpha,\ell}$, and J_ℓ .

Lemma 2 *There exists a positive constant c independent of N such that*

$$\|I_\alpha\|_{C(\overline{\mathcal{O}})} \leq c \left(1 + 2\frac{r_N}{h}\right)^d \left(\frac{r_N + d_N}{h}\right), \quad \alpha \in \mathbb{A}^d. \quad (29)$$

Proof We arbitrarily fix $x \in \overline{\mathcal{O}}$, $\alpha \in \mathbb{A}^d$, and particle volume decomposition $\Xi = \{\xi_i \mid i = 1, 2, \dots, N\}$ and split I_α into

$$I_\alpha(x) = E_1(x) + E_2(x) + E_3(x)$$

with

$$\begin{aligned} E_1(x) &:= \sum_{i \in \Lambda_0(x,h)} V_i (x_i - x)^\alpha w_h(|x_i - x|) \\ &\quad - \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| (x_i - x)^\alpha w_h(|x_j - x|), \\ E_2(x) &:= \sum_{i=1}^N \sum_{j=1}^N (x_i - x)^\alpha \int_{\sigma_j \cap \xi_i} \{w_h(|x_j - x|) - w_h(|y - x|)\} dy, \\ E_3(x) &:= \sum_{i=1}^N \sum_{j=1}^N (x_i - x)^\alpha \int_{\sigma_j \cap \xi_i} w_h(|y - x|) dy - \int_{\mathbb{R}^d} y^\alpha w_h(|y|) dy. \end{aligned}$$

Then, we estimate E_1 , E_2 , and E_3 .

First, we estimate E_1 . Because

$$\sum_{j=1}^N |\sigma_j \cap \xi_i| = V_i, \quad i = 1, 2, \dots, N, \quad (30)$$

we can rewrite E_1 as

$$E_1 = \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| (x_i - x)^\alpha \{w_h(|x_i - x|) - w_h(|x_j - x|)\}.$$

From

$$|(y-x)^\alpha| \leq \text{diam}(\Omega_H)^{|\alpha|}, \quad y \in \Omega_H, \quad (31)$$

we obtain

$$|E_1(x)| \leq c \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| |w_h(|x_i - x|) - w_h(|x_j - x|)|. \quad (32)$$

From

$$|w_h(|y-x|) - w_h(|z-x|)| = 0, \quad \forall y, z \in \mathbb{R}^d \setminus B(x, h),$$

we have

$$\begin{aligned} & \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| |w_h(|x_i - x|) - w_h(|x_j - x|)| \\ & \leq \sum_{i \in \Lambda_0(x, h)} \sum_{j=1}^N |\sigma_j \cap \xi_i| |w_h(|x_i - x|) - w_h(|x_j - x|)| \\ & \quad + \sum_{i=1}^N \sum_{j \in \Lambda_0(x, h)} |\sigma_j \cap \xi_i| |w_h(|x_i - x|) - w_h(|x_j - x|)| \\ & = \sum_{i \in \Lambda_0(x, h)} \sum_{j=1}^N (|\sigma_i \cap \xi_j| + |\sigma_j \cap \xi_i|) |w_h(|x_i - x|) - w_h(|x_j - x|)|. \end{aligned} \quad (33)$$

Because w_h is absolutely continuous, we have

$$\begin{aligned} & |w_h(|y-x|) - w_h(|z-x|)| \\ & = \left| \{(y-x) - (z-x)\} \int_0^1 w'_h(t|y-x| + (1-t)|z-x|) dt \right| \\ & \leq |y-z| \left| \int_0^1 w'_h(t|y-x| + (1-t)|z-x|) dt \right| \\ & \leq |y-z| \int_0^h |w'_h(r)| dr \\ & \leq \frac{|y-z|}{h^{d+1}} \int_0^1 |w'(r)| dr, \end{aligned} \quad (34)$$

for all $y, z \in \mathbb{R}^d$. Here, w' and w'_h are dw/dr and dw_h/dr , respectively. Moreover, we have

$$\sum_{i \in \Lambda_0(x, r)} |\sigma_i| \leq |B(x, 1)| (r + r_N)^d, \quad \forall r \in \mathbb{R}_0^+. \quad (35)$$

From (33), (34), and (35), we have

$$\begin{aligned}
& \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| |w_h(|x_i - x|) - w_h(|x_j - x|)| \\
&= \frac{c}{h^{d+1}} \sum_{i \in \Lambda_0(x, h)} \sum_{j=1}^N (|\sigma_i \cap \xi_j| + |\sigma_j \cap \xi_i|) |x_i - x_j| \\
&\leq \frac{c}{h^{d+1}} \sum_{i \in \Lambda_0(x, h)} |\sigma_i| \sum_{j=1}^N \frac{|\sigma_i \cap \xi_j| + |\sigma_j \cap \xi_i|}{|\sigma_i|} |x_i - x_j| \\
&\leq c \frac{d\Xi}{h^{d+1}} \sum_{i \in \Lambda_0(x, h)} |\sigma_i| \\
&\leq c \left(1 + \frac{r_N}{h}\right)^d \frac{d\Xi}{h}. \tag{36}
\end{aligned}$$

Therefore, from (32) and (36), we obtain

$$|E_1(x)| \leq c \left(1 + \frac{r_N}{h}\right)^d \frac{d\Xi}{h}.$$

Next, we estimate E_2 . Because $\text{supp}(w_h) = [0, h]$ and $\sigma_j \subset B(x_j, r_N)$, we have

$$\int_{\sigma_j \cap \xi_i} |w_h(|x_j - x|) - w_h(|y - x|)| dy = 0, \quad i = 1, 2, \dots, N, \quad j \notin \Lambda_0(x, h + r_N). \tag{37}$$

From (37), we have

$$\begin{aligned}
& \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |w_h(|x_j - x|) - w_h(|y - x|)| dy \\
&= \sum_{i=1}^N \sum_{j \in \Lambda_0(x, h + r_N)} \int_{\sigma_j \cap \xi_i} |w_h(|x_j - x|) - w_h(|y - x|)| dy \\
&= \sum_{j \in \Lambda_0(x, h + r_N)} \int_{\sigma_j} |w_h(|x_j - x|) - w_h(|y - x|)| dy.
\end{aligned}$$

Moreover, from (34) and (35), we have

$$\begin{aligned}
\sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |w_h(|x_j - x|) - w_h(|y - x|)| dy &\leq \frac{c}{h^{d+1}} \sum_{j \in \Lambda_0(x, h + r_N)} \int_{\sigma_j} |x_j - y| dy \\
&\leq c \frac{r_N}{h^{d+1}} \sum_{j \in \Lambda_0(x, h + r_N)} |\sigma_j| \\
&\leq c \left(1 + 2\frac{r_N}{h}\right)^d \frac{r_N}{h}. \tag{38}
\end{aligned}$$

Therefore, from (31) and (38), we obtain

$$\begin{aligned} |E_2(x)| &\leq \sum_{i=1}^N \sum_{j=1}^N |(x_i - x)^\alpha| \int_{\sigma_j \cap \xi_i} |w_h(|x_j - x|) - w_h(|y - x|)| dy \\ &\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |w_h(|x_j - x|) - w_h(|y - x|)| dy \\ &\leq c \left(1 + 2 \frac{r_N}{h}\right)^d \frac{r_N}{h}. \end{aligned}$$

Finally, we estimate E_3 . Because

$$\int_{\mathbb{R}^d} y^\alpha w_h(|y|) dy = \int_{\Omega_H} (y - x)^\alpha w_h(|y - x|) dy,$$

we can rewrite E_3 as

$$E_3(x) = \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} \{(x_i - x)^\alpha - (y - x)^\alpha\} w_h(|y - x|) dy.$$

Because $E_3 = 0$ when $|\alpha| = 0$, we estimate when $|\alpha| \geq 1$. Let β_k ($k = 1, 2, \dots, |\alpha|$) be d -dimensional multi-indices with satisfying

$$\sum_{k=1}^{|\alpha|} \beta_k = \alpha, \quad |\beta_k| = 1 \quad (k = 1, 2, \dots, |\alpha|).$$

Then, we have, for all $y, z \in \mathbb{R}^d$,

$$\begin{aligned} |y^\alpha - z^\alpha| &\leq \left| y^\alpha - y^{\alpha - \beta_1} z^{\beta_1} \right| + \left| y^{\alpha - \beta_1} z^{\beta_1} - z^\alpha \right| \\ &\leq |y - z| |y|^{\alpha - 1} + \left| y^{\alpha - \beta_1} - z^{\alpha - \beta_1} \right| |z| \\ &\leq |y - z| |y|^{\alpha - 1} + |y - z| |y|^{\alpha - 2} |z| + \left| y^{\alpha - \beta_1 - \beta_2} - z^{\alpha - \beta_1 - \beta_2} \right| |z|^2 \\ &\quad \vdots \\ &\leq |y - z| \sum_{k=1}^{|\alpha|} |y|^{\alpha - k} |z|^{k-1}. \end{aligned} \tag{39}$$

From (31) and (39), we obtain

$$\begin{aligned} |E_3(x)| &\leq \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |(x_i - x)^\alpha - (y - x)^\alpha| |w_h(|y - x|)| dy \\ &\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |y - x_i| |w_h(|y - x|)| dy. \end{aligned} \tag{40}$$

By $\text{supp}(w_h) = [0, h]$ and $\sigma_j \subset B(x_j, r_N)$, if $j \notin \Lambda_0(x, h + r_N)$, then

$$\int_{\sigma_j \cap \xi_i} |y - x_i| |w_h(|y - x|)| dy = 0, \quad i = 1, 2, \dots, N. \tag{41}$$

Moreover, from $w \in \mathcal{W} \subset C(\mathbb{R}_0^+)$, we have

$$|w_h(|y-x|)| = \frac{1}{h^d} \left| w\left(\frac{|y-x|}{h}\right) \right| \leq \frac{1}{h^d} \|w\|_{C(\mathbb{R}_0^+)}, \quad \forall y \in \Omega_H. \quad (42)$$

From (35), (41), and (42), we have

$$\begin{aligned} & \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |y-x_i| |w_h(|y-x|)| dy \\ &= \sum_{i=1}^N \sum_{j \in \Lambda_0(x, h+r_N)} \int_{\sigma_j \cap \xi_i} |y-x_i| |w_h(|y-x|)| dy \\ &\leq \frac{c}{h^d} \sum_{i=1}^N \sum_{j \in \Lambda_0(x, h+r_N)} \int_{\sigma_j \cap \xi_i} |y-x_i| dy \\ &\leq \frac{c}{h^d} \sum_{i=1}^N \sum_{j \in \Lambda_0(x, h+r_N)} \int_{\sigma_j \cap \xi_i} (|y-x_j| + |x_j-x_i|) dy \\ &\leq \frac{c}{h^d} \left(r_N \sum_{j \in \Lambda_0(x, h+r_N)} |\sigma_j| + \sum_{j \in \Lambda_0(x, h+r_N)} \sum_{i=1}^N |\sigma_j \cap \xi_i| |x_j-x_i| \right) \\ &\leq \frac{c}{h^d} \left(\sum_{j \in \Lambda_0(x, h+r_N)} |\sigma_j| \right) \left\{ r_N + \max_{j=1,2,\dots,N} \left(\sum_{i=1}^N \frac{|\sigma_i \cap \xi_j| + |\sigma_j \cap \xi_i|}{|\sigma_j|} |x_j-x_i| \right) \right\} \\ &\leq c \left(1 + 2 \frac{r_N}{h} \right)^d (r_N + d\Xi). \end{aligned} \quad (43)$$

Therefore, from (40), (43), and $h \leq H$, we obtain

$$\begin{aligned} |E_3(x)| &\leq c \left(1 + 2 \frac{r_N}{h} \right)^d (r_N + d\Xi) \\ &\leq c \left(1 + 2 \frac{r_N}{h} \right)^d \frac{r_N + d\Xi}{h}. \end{aligned}$$

From the estimates of E_1 , E_2 , and E_3 , we obtain

$$\|I_\alpha\|_{C(\bar{\Omega})} \leq c \left(1 + 2 \frac{r_N}{h} \right)^d \frac{r_N + d\Xi}{h}.$$

Because Ξ is arbitrary, we establish (29).

Lemma 3 *Suppose that a reference weight function w satisfies Hypothesis 2 with k . Then, there exists a positive constant c independent of N such that for all $\alpha \in \mathbb{A}^d$ and $\ell \in \mathbb{N}$ with $1 \leq \ell - k \leq |\alpha|$,*

$$\|I_{\alpha,\ell}\|_{C(\bar{\Omega})} \leq c \left(1 + 2 \frac{r_N}{h} \right)^d \frac{r_N + d_N}{h^{k+1}}. \quad (44)$$

Proof We arbitrarily fix $x \in \bar{\Omega}$, $\alpha \in \mathbb{A}^d$, particle volume decomposition $\Xi = \{\xi_i \mid i = 1, 2, \dots, N\}$, and $\ell \in \mathbb{N}$ with $1 \leq \ell - k \leq |\alpha|$ and split $I_{\alpha,\ell}$ into

$$I_{\alpha,\ell}(x) = E_4(x) + E_5(x) + E_6(x)$$

with

$$\begin{aligned}
E_4(x) &:= \sum_{i \in \Lambda(x, h)} V_i \frac{(x_i - x)^\alpha}{|x_i - x|^\ell} w_h(|x_i - x|) \\
&\quad - \sum_{i \in \Lambda(x, \infty)} \sum_{j \in \Lambda(x, \infty)} |\sigma_j \cap \xi_i| \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} \frac{w_h(|x_j - x|)}{|x_j - x|^k}, \\
E_5(x) &:= \sum_{i \in \Lambda(x, \infty)} \sum_{j \in \Lambda(x, \infty)} |\sigma_j \cap \xi_i| \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} \frac{w_h(|x_j - x|)}{|x_j - x|^k} \\
&\quad - \sum_{i \in \Lambda(x, \infty)} \sum_{j=1}^N \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} \int_{\sigma_j \cap \xi_i} \frac{w_h(|y - x|)}{|y - x|^k} dy, \\
E_6(x) &:= \sum_{i \in \Lambda(x, \infty)} \sum_{j=1}^N \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} \int_{\sigma_j \cap \xi_i} \frac{w_h(|y - x|)}{|y - x|^k} dy \\
&\quad - \int_{\mathbb{R}^d} \frac{y^\alpha}{|y|^\ell} w_h(|y|) dy.
\end{aligned}$$

Then, we estimate E_4 , E_5 , and E_6 .

First, we estimate E_4 and set $w^{(k)}$ as (8) and $w_h^{(k)}$ as

$$w_h^{(k)}(r) := \frac{1}{h^{d+k}} w^{(k)}\left(\frac{r}{h}\right), \quad r \in \mathbb{R}_0^+.$$

Then, from (30), we can rewrite E_4 as

$$E_4(x) = \sum_{i \in \Lambda(x, \infty)} \sum_{j=1}^N |\sigma_j \cap \xi_i| \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} \{w_h^{(k)}(|x_i - x|) - w_h^{(k)}(|x_j - x|)\}.$$

Because

$$\left| \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} \right| \leq |x_i - x|^{|\alpha| - \ell + k} \leq \text{diam}(\Omega_H)^{|\alpha| - \ell + k}, \quad i \in \Lambda(x, \infty), \quad (45)$$

we obtain

$$|E_4(x)| \leq c \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| \left| w_h^{(k)}(|x_i - x|) - w_h^{(k)}(|x_j - x|) \right|.$$

From $\text{supp}(w_h^{(k)}) = [0, h]$, we have

$$w_h^{(k)}(|x_i - x|) - w_h^{(k)}(|x_j - x|) = 0, \quad i, j \notin \Lambda(x, h).$$

Thus, we obtain

$$\begin{aligned}
|E_4(x)| &\leq c \left(\sum_{i \in \Lambda(x, h)} \sum_{j=1}^N |\sigma_j \cap \xi_i| \left| w_h^{(k)}(|x_i - x|) - w_h^{(k)}(|x_j - x|) \right| \right. \\
&\quad \left. + \sum_{i=1}^N \sum_{j \in \Lambda(x, h)} |\sigma_j \cap \xi_i| \left| w_h^{(k)}(|x_i - x|) - w_h^{(k)}(|x_j - x|) \right| \right). \quad (46)
\end{aligned}$$

Using an argument similar to (34), if w satisfies Hypothesis 2 with k , then for all $y, z \in \mathbb{R}^d$,

$$|w_h^{(k)}(|y-x|) - w_h^{(k)}(|z-x|)| \leq \frac{|y-z|}{h^{d+k+1}} \int_0^1 |(w^{(k)})'(r)| dr. \quad (47)$$

From (46) and (47), we obtain

$$\begin{aligned} |E_4(x)| &\leq \frac{c}{h^{d+k+1}} \sum_{i \in \Lambda(x, h)} \sum_{j=1}^N (|\sigma_i \cap \xi_j| + |\sigma_j \cap \xi_i|) |x_i - x_j| \\ &\leq \frac{c}{h^{d+k+1}} \sum_{i \in \Lambda(x, h)} |\sigma_i| \sum_{j=1}^N \frac{|\sigma_i \cap \xi_j| + |\sigma_j \cap \xi_i|}{|\sigma_i|} |x_i - x_j| \\ &\leq c \left(1 + \frac{r_N}{h}\right)^d \frac{d\Xi}{h^{k+1}}. \end{aligned}$$

Next, we estimate E_5 . By using $w_h^{(k)}$, we can rewrite E_5 as

$$E_5(x) = \sum_{i \in \Lambda(x, \infty)} \sum_{j=1}^N \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} \int_{\sigma_j \cap \xi_i} \left\{ w_h^{(k)}(|x_j - x|) - w_h^{(k)}(|y - x|) \right\} dy.$$

From (45), we obtain

$$\begin{aligned} |E_5(x)| &\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} \left| w_h^{(k)}(|x_j - x|) - w_h^{(k)}(|y - x|) \right| dy \\ &\leq c \sum_{j=1}^N \int_{\sigma_j} \left| w_h^{(k)}(|x_j - x|) - w_h^{(k)}(|y - x|) \right| dy. \end{aligned}$$

By $\text{supp}(w_h^{(k)}) = [0, h]$ and $\sigma_j \subset B(x_j, r_N)$, we have

$$\int_{\sigma_j} \left| w_h^{(k)}(|x_j - x|) - w_h^{(k)}(|y - x|) \right| dy = 0, \quad j \notin \Lambda(x, h + r_N). \quad (48)$$

From (47) and (48), we obtain

$$\begin{aligned} |E_5(x)| &\leq c \sum_{j \in \Lambda_0(x, h+r_N)} \int_{\sigma_j} \left| w_h^{(k)}(|x_j - x|) - w_h^{(k)}(|y - x|) \right| dy \\ &\leq \frac{c}{h^{d+k+1}} \sum_{j \in \Lambda_0(x, h+r_N)} \int_{\sigma_j} |x_j - y| dy \\ &\leq c \frac{r_N}{h^{d+k+1}} \sum_{j \in \Lambda_0(x, h+r_N)} |\sigma_j| \\ &\leq c \left(1 + 2\frac{r_N}{h}\right)^d \frac{r_N}{h^{k+1}}. \end{aligned}$$

Finally, we estimate E_6 . Using $w_h^{(k)}$, we can rewrite E_6 as

$$\begin{aligned} E_6(x) &= \sum_{i \in \Lambda(x, \infty)} \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} \left\{ \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} - \frac{(y - x)^\alpha}{|y - x|^{\ell-k}} \right\} w_h^{(k)}(|y - x|) dy \\ &\quad - \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i^*(x)} \frac{(y - x)^\alpha}{|y - x|^{\ell-k}} w_h^{(k)}(|y - x|) dy, \end{aligned}$$

where $\xi_i^*(x)$ is

$$\xi_i^*(x) = \begin{cases} \xi_i, & x = x_i, \\ \emptyset, & \text{otherwise.} \end{cases}$$

For $\alpha \in \mathbb{A}^d$, let β_j ($j = 1, 2, \dots, |\alpha|$) be d -dimensional multi-indices satisfying

$$|\beta_j| = 1 \quad \text{and} \quad \sum_{j=1}^{|\alpha|} \beta_j = \alpha.$$

Let β_j^* ($j = 0, 1, \dots, |\alpha|$) be d -dimensional multi-indices defined as

$$\beta_j^* := \begin{cases} 0, & j = 0, \\ \sum_{\ell=1}^j \beta_\ell, & j = 1, 2, \dots, |\alpha|. \end{cases}$$

For all $y, z \in \mathbb{R}^d \setminus \{0\}$, when $|\alpha| = \ell - k$, we have

$$\begin{aligned} \left| \frac{y^\alpha}{|y|^{\ell-k}} - \frac{z^\alpha}{|z|^{\ell-k}} \right| &\leq \sum_{j=0}^{\ell-k-1} \left| \frac{y^{\beta_{|\alpha|-j}^*} z^{\beta_j^*}}{|y|^{\ell-k-j} |z|^j} - \frac{y^{\beta_{|\alpha|-j-1}^*} z^{\beta_{j+1}^*}}{|y|^{\ell-k-j-1} |z|^{j+1}} \right| \\ &\leq \sum_{j=0}^{\ell-k-1} \left| \frac{y^{\beta_{|\alpha|-j}^*} z^{\beta_j^*} - y^{\beta_{|\alpha|-j-1}^*} z^{\beta_{j+1}^*}}{|y|^{\ell-k-j} |z|^j} \right| \\ &\quad + \sum_{j=0}^{\ell-k-1} \left| \frac{y^{\beta_{|\alpha|-j-1}^*} z^{\beta_{j+1}^*}}{|y|^{\ell-k-j} |z|^j} - \frac{y^{\beta_{|\alpha|-j-1}^*} z^{\beta_{j+1}^*}}{|y|^{\ell-k-j-1} |z|^{j+1}} \right| \\ &\leq 2(\ell - k) \frac{|y - z|}{|y|}. \end{aligned} \tag{49}$$

Moreover, from (39) and (49), when $|\alpha| > \ell - k$, we have

$$\begin{aligned} \left| \frac{y^\alpha}{|y|^{\ell-k}} - \frac{z^\alpha}{|z|^{\ell-k}} \right| &\leq \left| \frac{y^\alpha}{|y|^{\ell-k}} - \frac{y^{\beta_{|\alpha|-\ell+k}^*} z^{\beta_{\ell-k}^*}}{|z|^{\ell-k}} \right| \\ &\quad + \left| \frac{y^{\beta_{|\alpha|-\ell+k}^*} z^{\beta_{\ell-k}^*}}{|z|^{\ell-k}} - \frac{z^\alpha}{|z|^{\ell-k}} \right| \\ &\leq |y|^{|\alpha|-\ell+k} \left| \frac{y^{\beta_{\ell-k}^*}}{|y|^{\ell-k}} - \frac{z^{\beta_{\ell-k}^*}}{|z|^{\ell-k}} \right| + \left| y^{\beta_{|\alpha|-\ell+k}^*} - z^{\beta_{|\alpha|-\ell+k}^*} \right| \\ &\leq 2(\ell - k) |y - z| |y|^{|\alpha|-\ell+k-1} \\ &\quad + |y - z| \sum_{j=0}^{|\alpha|-\ell+k-1} |y|^j |z|^{|\alpha|-\ell+k-1-j}. \end{aligned}$$

Therefore, when $|\alpha| \geq \ell - k$, we have for all $y \in \Omega_H \setminus \{x\}$ and $i \in \Lambda(x, \infty)$,

$$\left| \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} - \frac{(y - x)^\alpha}{|y - x|^{\ell-k}} \right| \leq c \frac{|y - x_i|}{|y - x|}. \quad (50)$$

From (50), we obtain

$$\begin{aligned} |E_6(x)| &\leq \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} \left| \frac{(x_i - x)^\alpha}{|x_i - x|^{\ell-k}} - \frac{(y - x)^\alpha}{|y - x|^{\ell-k}} \right| w_h^{(k)}(|y - x|) dy \\ &\quad + \left| \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i^*(x)} \frac{(y - x)^\alpha}{|y - x|^{\ell-k}} w_h^{(k)}(|y - x|) dy \right| \\ &\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |y - x_i| w_h^{(k+1)}(|y - x|) dy \\ &\quad + \left| \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i^*(x)} \frac{(y - x)^\alpha}{|y - x|^{\ell-k}} w_h^{(k)}(|y - x|) dy \right|. \end{aligned}$$

Because $|\alpha| \geq \ell - k$, we have

$$\begin{aligned} &\left| \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i^*(x)} \frac{(y - x)^\alpha}{|y - x|^{\ell-k}} w_h^{(k)}(|y - x|) dy \right| \\ &\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i^*(x)} \left| \frac{(y - x)^\alpha}{|y - x|^{\ell-k-1}} \right| w_h^{(k+1)}(|y - x|) dy \\ &\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i^*(x)} |y - x| w_h^{(k+1)}(|y - x|) dy. \end{aligned}$$

Therefore, we have

$$|E_6(x)| \leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |y - x_i| w_h^{(k+1)}(|y - x|) dy.$$

Because for all $y \in \Omega_H$,

$$\left| w_h^{(k+1)}(|y - x|) \right| = \frac{1}{h^{d+k+1}} \left| w^{(k+1)} \left(\frac{|y - x|}{h} \right) \right| \leq \frac{1}{h^{d+k+1}} \|w^{(k+1)}\|_{C(\mathbb{R}_0^+)},$$

by the same procedure as (43), we have

$$\sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |y - x_i| w_h^{(k+1)}(|y - x|) dy \leq c \left(1 + 2 \frac{r_N}{h}\right)^d \frac{r_N + d_\Xi}{h^{k+1}}.$$

Therefore, we obtain

$$|E_6(x)| \leq c \left(1 + 2 \frac{r_N}{h}\right)^d \frac{r_N + d_\Xi}{h^{k+1}}.$$

From the estimates of E_4 , E_5 , and E_6 , we obtain

$$\|I_{\alpha, \ell}\|_{C(\bar{\Omega})} \leq c \left(1 + 2\frac{r_N}{h}\right)^d \frac{r_N + d_{\Xi}}{h^{k+1}}.$$

Because Ξ is arbitrary, we establish (44).

Lemma 4 *There exists a positive constant c independent of N such that*

$$\|J_{\ell}\|_{C(\bar{\Omega})} \leq c \left\{ \left(1 + 2\frac{r_N}{h}\right)^d \frac{r_N + d_N}{h} + h^{\ell} \right\}, \quad \ell \in \mathbb{N}. \quad (51)$$

Proof We arbitrarily fix $x \in \bar{\Omega}$ and particle volume decomposition $\Xi = \{\xi_i \mid i = 1, 2, \dots, N\}$, and split J_{ℓ} into

$$J_{\ell}(x) = E_7(x) + E_8(x) + E_9(x) + E_{10}(x)$$

with

$$\begin{aligned} E_7(x) &:= J_{\ell}(x) - \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| |x_i - x|^{\ell} |w_h(|x_j - x|)|, \\ E_8(x) &:= \sum_{i=1}^N \sum_{j=1}^N |x_i - x|^{\ell} \int_{\sigma_j \cap \xi_i} \{|w_h(|x_j - x|)| - |w_h(|y - x|)|\} dy, \\ E_9(x) &:= \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} \{|x_i - x|^{\ell} - |y - x|^{\ell}\} |w_h(|y - x|)| dy, \\ E_{10}(x) &:= \int_{\mathbb{R}^d} |y - x|^{\ell} |w_h(|y - x|)| dy. \end{aligned}$$

Then, we estimate E_7 , E_8 , E_9 , and E_{10} .

From (30), we can rewrite E_7 as

$$E_7(x) = \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| |x_i - x|^{\ell} \{|w_h(|x_i - x|)| - |w_h(|x_j - x|)|\}.$$

For all $y \in \Omega_H$, we have

$$|y - x|^{\ell} \leq \text{diam}(\Omega_H)^{\ell}. \quad (52)$$

From (36) and (52), we obtain

$$\begin{aligned} |E_7(x)| &\leq \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| |x_i - x|^{\ell} \left| |w_h(|x_i - x|)| - |w_h(|x_j - x|)| \right| \\ &\leq c \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| \left| |w_h(|x_i - x|)| - |w_h(|x_j - x|)| \right| \\ &\leq c \sum_{i=1}^N \sum_{j=1}^N |\sigma_j \cap \xi_i| |w_h(|x_i - x|) - w_h(|x_j - x|)| \\ &\leq c \left(1 + \frac{r_N}{h}\right)^d \frac{d_{\Xi}}{h}. \end{aligned}$$

From (38) and (52), we obtain

$$\begin{aligned}
|E_8(x)| &\leq \sum_{i=1}^N \sum_{j=1}^N |x_i - x|^\ell \int_{\sigma_j \cap \xi_i} \left| |w_h(|x_j - x|)| - |w_h(|y - x|)| \right| dy \\
&\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} \left| |w_h(|x_j - x|)| - |w_h(|y - x|)| \right| dy \\
&\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |w_h(|x_j - x|) - w_h(|y - x|)| dy \\
&\leq c \left(1 + 2 \frac{r_N}{h}\right)^d \frac{r_N}{h}.
\end{aligned}$$

For all $x_i \in \mathcal{X}_N$ and $y \in \Omega_H$, we have

$$\begin{aligned}
\left| |x_i - x|^\ell - |y - x|^\ell \right| &= |(x_i - x) - (y - x)| \sum_{k=1}^{\ell} (x_i - x)^{k-1} (y - x)^{\ell-k} \\
&\leq \ell \operatorname{diam}(\Omega_H)^{\ell-1} |y - x_i|.
\end{aligned} \tag{53}$$

From (43), (53), and $h < H$, we obtain

$$\begin{aligned}
|E_9(x)| &\leq \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} \left| |x_i - x|^\ell - |y - x|^\ell \right| |w_h(|y - x|)| dy \\
&\leq c \sum_{i=1}^N \sum_{j=1}^N \int_{\sigma_j \cap \xi_i} |y - x_i| |w_h(|y - x|)| dy \\
&\leq c \left(1 + 2 \frac{r_N}{h}\right)^d (r_N + d\Xi) \\
&\leq c \left(1 + 2 \frac{r_N}{h}\right)^d \frac{r_N + d\Xi}{h}.
\end{aligned}$$

From (1), we obtain

$$|E_{10}(x)| \leq \int_{\mathbb{R}^d} |y|^\ell |w_h(|y|)| dy = h^\ell \int_{\mathbb{R}^d} |y|^\ell |w(|y|)| dy.$$

From the estimates of E_7 , E_8 , E_9 , and E_{10} , we obtain

$$\|J_\ell\|_{C(\overline{\Omega})} \leq c \left\{ \left(1 + 2 \frac{r_N}{h}\right)^d \frac{r_N + d\Xi}{h} + h^\ell \right\}.$$

Because Ξ is arbitrary, we establish (51).

Using the lemmas defined above, we now prove Theorem 3.

Proof (Theorem 3) By Lemmas 1, 2, and 4, we have for all $v \in C^{n+1}(\overline{\Omega}_H)$

$$\|v - \Pi_h v\|_{C(\overline{\Omega})} \leq c \left\{ \left(1 + 2 \frac{r_N}{h_N}\right)^d \frac{r_N + d_N}{h_N} + h_N^{n+1} \right\} \|v\|_{C^{n+1}(\overline{\Omega}_H)}. \tag{54}$$




Fig. 4 Particle distribution \mathcal{X}_N with $\Delta x = 2^{-5}$ ($N = 1,521$). The gray area represents Ω .

Moreover, by Lemmas 1, 3, and 4, when w satisfies Hypothesis 2 with $k = 0$, we have for all $v \in C^{n+2}(\overline{\Omega}_H)$

$$\|\nabla v - \nabla_h v\|_{C(\overline{\Omega})} \leq c \left\{ \left(1 + 2\frac{r_N}{h_N}\right)^d \frac{r_N + d_N}{h_N} + h_N^{n+1} \right\} \|v\|_{C^{n+2}(\overline{\Omega}_H)}, \quad (55)$$

and when w satisfies Hypothesis 2 with $k = 1$ for all $v \in C^{n+3}(\overline{\Omega}_H)$,

$$\|\Delta v - \Delta_h v\|_{C(\overline{\Omega})} \leq c \left\{ \left(1 + 2\frac{r_N}{h_N}\right)^d \frac{r_N + d_N}{h_N^2} + h_N^{n+1} \right\} \|v\|_{C^{n+3}(\overline{\Omega}_H)}. \quad (56)$$

Because the family $\{(\mathcal{X}_N, \mathcal{V}_N, h_N)\}_{N \rightarrow \infty}$ is regular, by applying (7) to (54), (55), and (56), we obtain (9), (10), and (11), respectively. We now conclude the proof of Theorem 3.

5 Numerical results

Set $\Omega = (0, 1)^2$ and $H = 0.1$. Then, $\Omega_H = (-0.1, 1.1)^2$. We now compute the truncation errors of $v : \Omega_H \rightarrow \mathbb{R}$, which are defined as $v(x, y) = \sin(2\pi(x + y))$. Particle distribution \mathcal{X}_N is set as

$$\mathcal{X}_N = \left\{ \left((i + \eta_{ij}^{(1)})\Delta x, (j + \eta_{ij}^{(2)})\Delta x \right) \in \Omega_H; i, j \in \mathbb{Z} \right\},$$

where Δx is taken by $2^{-5}, 2^{-6}, \dots, 2^{-12}$ and $\eta_{ij}^{(k)}$ ($i, j \in \mathbb{Z}, k = 1, 2$) are random numbers satisfying $|\eta_{ij}^{(k)}| < 1/4$. Particle distribution \mathcal{X}_N with $\Delta x = 2^{-5}$ is shown in Figure 4. Particle volume set \mathcal{V}_N is defined as

$$\mathcal{V}_N = \left\{ V_i = \frac{|\Omega_H|}{N} \mid i = 1, 2, \dots, N \right\}.$$

For $m = 1, 3, 5$, the influence radius h_N is set as

$$h_N = 2.6 \times 2^{5/m-5} \Delta x^{1/m}.$$

Note that if $\Delta x = 2^{-5}$, then $h = 2.6 \times 2^{-5}$ for all m . Using the discrete parameters above, the covering radius r_N satisfies $r_N \leq \sqrt{2}(1 + 1/4)\Delta x/2$. Moreover, the Voronoi deviation d_N satisfies $d_N \leq 64(1 + \sqrt{2})\Delta x/\pi$. Therefore, the family $\{(\mathcal{X}_N, \mathcal{V}_N, h_N)\}$ is regular with order m .

For the interpolant, we consider the following three cases of reference weight functions:

$$\begin{aligned}
 (\text{II1}) \quad w(r) &:= \frac{3}{\pi} \begin{cases} 1-r, & 0 \leq r < 1, \\ 0, & 1 \leq r, \end{cases} \\
 (\text{II2}) \quad w(r) &:= \frac{40}{7\pi} \begin{cases} 1-6r^2+6r^3, & 0 \leq r < \frac{1}{2}, \\ 2(1-r)^3, & \frac{1}{2} \leq r < 1, \\ 0, & 1 \leq r, \end{cases} \\
 (\text{II3}) \quad w(r) &:= \frac{5}{\pi} \begin{cases} (1-r)(2-3r), & 0 \leq r < 1, \\ 0, & 1 \leq r. \end{cases}
 \end{aligned}$$

(II1) is the lowest-order polynomial function belonging to \mathcal{W} . (II2) is the cubic B-spline commonly used in the SPH method and belonging to \mathcal{W} . (II3) is the lowest-order polynomial function belonging to \mathcal{W} that satisfies Hypothesis 1 with $n = 3$.

For the approximate gradient operator, we consider the following three cases of reference weight functions:

$$\begin{aligned}
 (\nabla 1) \quad w(r) &:= \frac{6}{\pi} \begin{cases} r(1-r), & 0 \leq r < 1, \\ 0, & 1 \leq r, \end{cases} \\
 (\nabla 2) \quad w(r) &:= \frac{40}{7\pi} \begin{cases} 6r^2-9r^3, & 0 \leq r < \frac{1}{2}, \\ 3r(1-r)^2, & \frac{1}{2} \leq r < 1, \\ 0, & 1 \leq r, \end{cases} \\
 (\nabla 3) \quad w(r) &:= \frac{15}{2\pi} \begin{cases} r(1-r)(5-7r), & 0 \leq r < 1, \\ 0, & 1 \leq r. \end{cases}
 \end{aligned}$$

($\nabla 1$) is the lowest-order polynomial function belonging to \mathcal{W} that satisfies Hypothesis 2 with $k = 0$. ($\nabla 2$) is chosen so that the approximate gradient operator (3) with ($\nabla 2$) coincides with that in the SPH method with the cubic B-spline (see Appendix A). ($\nabla 3$) is the lowest-order polynomial function belonging to \mathcal{W} that satisfies Hypothesis 1 with $n = 3$ and Hypothesis 2 with $k = 0$.

For the approximate Laplace operator, we consider the following three cases of reference weight functions:

$$\begin{aligned}
 (\Delta 1) \quad w(r) &:= \frac{10}{\pi} \begin{cases} r^2(1-r), & 0 \leq r < 1, \\ 0, & 1 \leq r, \end{cases} \\
 (\Delta 2) \quad w(r) &:= \frac{40}{7\pi} \begin{cases} 6r^2 - 9r^3, & 0 \leq r < \frac{1}{2}, \\ 3r(1-r)^2, & \frac{1}{2} \leq r < 1, \\ 0, & 1 \leq r, \end{cases} \\
 (\Delta 3) \quad w(r) &:= \frac{30}{\pi} \begin{cases} r^2(1-r)(3-4r), & 0 \leq r < 1, \\ 0, & 1 \leq r. \end{cases}
 \end{aligned}$$

($\Delta 1$) is the lowest-order polynomial function belonging to \mathcal{W} that satisfies Hypothesis 2 with $k = 1$. ($\Delta 2$) is chosen so that approximate Laplace operator (4) with ($\Delta 2$) coincides with that in the SPH method with the cubic B-spline (see Appendix A). ($\Delta 3$) is the lowest-order polynomial function belonging to \mathcal{W} that satisfies Hypothesis 1 with $n = 3$ and Hypothesis 2 with $k = 1$.

The above settings were used in the computation of the following relative errors

$$\frac{\|v - \Pi_h v\|_{\ell^\infty(\Omega)}}{\|v\|_{C(\bar{\Omega})}}, \quad \frac{\|\nabla v - \nabla_h v\|_{\ell^\infty(\Omega)}}{\|\nabla v\|_{C(\bar{\Omega})}}, \quad \frac{\|\Delta v - \Delta_h v\|_{\ell^\infty(\Omega)}}{\|\Delta v\|_{C(\bar{\Omega})}}.$$

Here, the discrete norm $\|\cdot\|_{\ell^\infty(\Omega)}$ is defined as

$$\|v\|_{\ell^\infty(\Omega)} := \max_{i \in \Lambda(\Omega)} |v(x_i)|.$$

Figure 5 shows graphs of the relative errors of (a) interpolant Π_h , (b) approximate gradient operator ∇_h , and (c) approximate Laplace operator Δ_h versus the influence radius h_N with regular orders $m = 1, 3, 5$. In Figure 5, the slopes of the triangles show the theoretical convergence rates obtained via Theorem 3. Table 1 lists the numerical and theoretical convergence rates obtained from the cases of $\Delta x = 2^{-11}$ and 2^{-12} , where the theoretical convergence rates correspond to Theorem 3. In the case of $m = 1$, as the settings could not be applied to Theorem 3, only numerical results without convergence were obtained. In contrast, the settings in cases $m = 3$ and 5 could be applied Theorem 3; thus, the numerical results with convergence were obtained. Moreover, the approximate operators with reference weight functions satisfying Hypothesis 1 with $n = 3$ became higher convergence orders in the cases where $m = 5$ as per Theorem 3.

6 Conclusions

We analyzed truncation errors in a generalized particle method, which is a wider class of particle methods that includes commonly used methods such as the SPH and MPS methods. In our analysis, we introduced two indicators: the first was the covering radius, which represents the maximum radius of the Voronoi region associated with the particle distribution, while the second was the Voronoi deviation,




Fig. 5 Graphs of the relative errors of (a) the interpolant, (b) approximate gradient operator, and (c) approximate Laplace operator versus the influence radius with regular orders $m = 1, 3, 5$.

which indicates the deviation between particle volumes and Voronoi volumes. With the covering radius and Voronoi deviation, we introduced a regularity of a family of discrete parameters, which includes the particle distribution, particle volume set, and influence radius associated with the number of particles. Moreover, we introduced two hypotheses of reference weight functions. With the regularity and hypotheses of reference weight functions, we established truncation error estimates

Table 1 Numerical and theoretical convergence rates of (a) the interpolant, (b) approximate gradient operator, and (c) approximate Laplace operator with regular orders $m = 1, 3, 5$. The numerical convergence rates were obtained for the cases of $\Delta x = 2^{-11}$ and 2^{-12} .

	$m = 1$		$m = 3$		$m = 5$	
	Numer.	Theor.	Numer.	Theor.	Numer.	Theor.
(a) Interpolant						
($\Pi 1$)	-0.10	N/A	2.02	2	2.00	2
($\Pi 2$)	0.05	N/A	2.13	2	2.01	2
($\Pi 3$)	0.00	N/A	4.20	2	7.41	4
(b) Gradient						
($\nabla 1$)	-0.05	N/A	2.11	2	2.02	2
($\nabla 2$)	-0.02	N/A	2.08	2	2.03	2
($\nabla 3$)	-0.06	N/A	3.56	2	7.69	4
(c) Laplacian						
($\Delta 1$)	-1.14	N/A	2.91	1	2.05	2
($\Delta 2$)	-1.09	N/A	2.39	1	2.23	2
($\Delta 3$)	-1.02	N/A	2.76	1	7.50	3

for the continuous norm. The convergence rates are dependent on the regular order and order of the reference weight functions appearing in a hypothesis. Moreover, as it was possible to validate the conditions by calculation, we showed the numerical convergence orders were in good agreement with the theoretical ones.

In a forthcoming paper, we plan to establish error estimates of the generalized particle method for the Poisson and heat equations.

Acknowledgments

We are grateful to the referee for the constructive comments provided. We would like to thank Drs. Daisuke Tagami and Hayato Waki for their helpful comments over the course of this work. This research was supported by JSPS KAKENHI Grant Number 17K17585 and the JSPS A3 Foresight Program.

A Description of conventional particle methods by the generalized particle method

This appendix provides a description of conventional particle methods, such as the smoothed particle hydrodynamics (SPH) [18, 24] and the moving particle semi-implicit (MPS) methods [16], in the context of the generalized particle method. In the SPH method, upon using the reference weight function $w^{\text{SPH}} \in \mathcal{W}$ and parameters $m_i, \rho_i \in \mathbb{R}^+$ ($i = 1, 2, \dots, N$), for $v \in$

$C(\overline{\Omega}_H)$, the approximate operators are defined as

$$\Pi_h^{\text{SPH}}v(x) := \sum_{i=1}^N \frac{m_i}{\rho_i} v(x_i) w_h^{\text{SPH}}(|x - x_i|), \quad x \in \Omega_H, \quad (57)$$

$$\nabla_h^{\text{SPH}}v(x) := \sum_{i=1}^N \frac{m_i}{\rho_i} \{v(x) - v(x_i)\} \nabla w_h^{\text{SPH}}(|x - x_i|), \quad x \in \Omega_H, \quad (58)$$

$$\Delta_h^{\text{SPH}}v(x) := 2 \sum_{i \in \Lambda(x,h)} \frac{m_i}{\rho_i} \frac{v(x) - v(x_i)}{|x - x_i|} \frac{x - x_i}{|x - x_i|} \cdot \nabla w_h^{\text{SPH}}(|x - x_i|), \quad x \in \Omega_H. \quad (59)$$

By setting $w = w^{\text{SPH}}$ and $\mathcal{V}_N = \{V_i = m_i/\rho_i; i = 1, 2, \dots, N\}$, the generalized interpolant (2) coincides with (57). Moreover, because

$$-\int_{\mathbb{R}^d} \frac{x}{d} \cdot \nabla w^{\text{SPH}}(|x|) dx = \int_{\mathbb{R}^d} w^{\text{SPH}}(|x|) dx = 1,$$

by setting

$$w(r) = -d^{-1}r(w^{\text{SPH}})'(r),$$

and $\mathcal{V}_N = \{V_i = m_i/\rho_i; i = 1, 2, \dots, N\}$, (3) and (4) coincide with (58) and (59), respectively.

From Theorem 3, we obtain the following corollary that is a truncation error estimate of approximate operators (58) and (59).

Corollary 1 *Suppose that parameters ρ_i, m_i satisfy*

$$\sum_{i=1}^N \frac{m_i}{\rho_i} = |\Omega_H|,$$

and that $\{(\mathcal{X}_N, \mathcal{V}_N, h_N)\}_{N \rightarrow \infty}$ is regular with order m , where $\mathcal{V}_N = \{\rho_i/m_i; i = 1, 2, \dots, N\}$. Moreover, suppose that w^{SPH} satisfies the following conditions;

$$w^{\text{SPH}} \in C^2(\mathbb{R}_0^+), \quad (w^{\text{SPH}})'(r) < 0 \quad (0 < r < 1), \quad \lim_{s \downarrow 0} \left| \frac{1}{s} (w^{\text{SPH}})'(s) \right| < \infty. \quad (60)$$

Then, there exists a positive constant c independent of N such that

$$\begin{aligned} \|v - \Pi_h^{\text{SPH}}v\|_{C(\overline{\Omega})} &\leq ch^{\min\{2, m-1\}} \|v\|_{C^2(\overline{\Omega}_H)}, & v \in C^2(\overline{\Omega}_H), \\ \|\nabla v - \nabla_h^{\text{SPH}}v\|_{C(\overline{\Omega})} &\leq ch^{\min\{2, m-1\}} \|v\|_{C^3(\overline{\Omega}_H)}, & v \in C^3(\overline{\Omega}_H), \\ \|\Delta v - \Delta_h^{\text{SPH}}v\|_{C(\overline{\Omega})} &\leq ch^{\min\{2, m-2\}} \|v\|_{C^4(\overline{\Omega}_H)}, & v \in C^4(\overline{\Omega}_H). \end{aligned}$$

Remark 5 Note that representative reference weight functions employed in the SPH method, such as the cubic B-spline, quintic B-spline, and Wendland function (5-order positive definite function) [8, 18], satisfy (60).

In the MPS method [14], upon using reference weight function $w^{\text{MPS}} \in \mathcal{W}$ and parameters $\hat{n}, \hat{\lambda} \in \mathbb{R}^+$ for $v \in C^1(\overline{\Omega}_H)$, approximate differential operators can be defined as

$$\nabla_h^{\text{MPS}}v(x) := \frac{d}{\hat{n}} \sum_{i \neq j} \frac{v(x_i) - v(x)}{|x - x_i|} \frac{x_i - x}{|x - x_i|} w_h^{\text{MPS}}(|x - x_j|), \quad x \in \Omega_H, \quad (61)$$

$$\Delta_h^{\text{MPS}}v(x) := \frac{2d}{\hat{n}\hat{\lambda}} \sum_{i \neq j} \{v(x_i) - v(x)\} w_h^{\text{MPS}}(|x - x_j|), \quad x \in \Omega_H. \quad (62)$$

Note that an interpolant is not defined in the MPS method. By setting $w = w^{\text{MPS}}$ and $\mathcal{V}_N = \{V_i = \hat{n}^{-1}; i = 1, 2, \dots, N\}$, the approximate gradient operator (3) coincides with (61). Moreover, by setting $w(r) = \hat{\lambda}^{-1}r^2 w^{\text{MPS}}(r)$ and $\mathcal{V}_N = \{V_i = \hat{n}^{-1}; i = 1, 2, \dots, N\}$, approximate Laplace operator (4) coincides with (62).

Corollary 2 *Suppose that*

$$\hat{n} = \frac{N}{|\Omega_H|}, \quad \hat{\lambda} = \int_{\mathbb{R}^d} |x|^2 w^{\text{MPS}}(|x|) dx, \quad w^{\text{MPS}} \in \mathcal{W}.$$

Moreover, suppose that $\{(\mathcal{X}_N, \mathcal{V}_N, h_N)\}_{N \rightarrow \infty}$ is regular with order m , where $\mathcal{V}_N = \{V_i = \hat{n}^{-1}; i = 1, 2, \dots, N\}$. Then, there exists a positive constant c independent of N such that

$$\|\Delta v - \Delta_h^{\text{MPS}} v\|_{C(\bar{\Omega})} \leq c h^{\min\{2, m-2\}} \|v\|_{C^4(\bar{\Omega}_H)}, \quad v \in C^4(\bar{\Omega}_H).$$

Furthermore, when w^{MPS} satisfies Hypothesis 2 with $k = 0$,

$$\|\nabla v - \nabla_h^{\text{MPS}} v\|_{C(\bar{\Omega})} \leq c h^{\min\{2, m-1\}} \|v\|_{C^3(\bar{\Omega}_H)}, \quad v \in C^3(\bar{\Omega}_H).$$

Remark 6 Note that the reference weight function, which is commonly used in the MPS method and defined as

$$w^{\text{MPS}}(r) := \begin{cases} \frac{1}{r} - 1, & 0 \leq r < 1, \\ 0, & 1 \leq r, \end{cases}$$

does not satisfy $w^{\text{MPS}} \in \mathcal{W}$. In contrast, the continuous reference weight function as introduced in [28] satisfies $w^{\text{MPS}} \in \mathcal{W}$. However, as far as we know, no reference weight functions that also satisfy Hypothesis 2 with $k = 0$ are proposed in the MPS method.

B Computational procedure of the indicators

This appendix introduces the procedures for computing the indicators introduced in this paper, namely, the covering radius (5) and Voronoi deviation (6).

The covering radius r_N can be computed as follows. As per the methods used to construct Voronoi decompositions, such as the increment method [6], we first draw the boundaries of the Voronoi region in Ω_H . Next, for each particle, we compute the maximum distance from particle x_i to the boundary of its Voronoi region σ_i (i.e., $\max_{y \in \bar{\sigma}_i} |x_i - y|$). Finally, we obtain the covering radius r_N by computing the maximum of these distances.

Next, we consider the Voronoi deviation d_N . Let $\zeta \in \mathbb{R}^{3N}$ be

$$\zeta := (|\sigma_1|, |\sigma_2|, \dots, |\sigma_N|, V_1, V_2, \dots, V_N, 0, 0, \dots, 0)^T.$$

Using parameters $q, s_i, a_{ij} \in \mathbb{R}^+$ ($i, j = 1, 2, \dots, N$), we set $z \in \mathbb{R}^{N^2+N+1}$ as

$$z := (a_{11}, a_{12}, \dots, a_{NN}, s_1, s_2, \dots, s_N, q)^T.$$

Moreover, we set $M \in \mathbb{R}^{3N \times (N^2+N+1)}$ so that equation $Mz = \zeta$ represents

$$\sum_{j=1}^N a_{ij} = |\sigma_i|, \quad \sum_{j=1}^N a_{ji} = V_i, \quad i = 1, 2, \dots, N$$

and

$$q = s_i + \sum_{j=1}^N \frac{a_{ij} + a_{ji}}{|\sigma_i|} |x_i - x_j|, \quad i = 1, 2, \dots, N.$$

Then, by considering a_{ij} to be $|\sigma_i \cap \xi_j|$, we find that the minimum value of q with condition $Mz = \zeta$ coincides with the Voronoi deviation d_N . We therefore consider the linear problem:

$$\text{Minimize } b^T z \quad \text{subject to } Mz = \zeta, \quad z \geq 0. \quad (63)$$

Here, $b := (0, 0, \dots, 0, 1)^T \in \mathbb{R}^{N^2+N+1}$. The solution $b^T z$ of (63) is equivalent to the Voronoi deviation d_N . Because $Mz = \zeta$ is unique for $(\mathcal{X}_N, \mathcal{V}_N, h_N)$, the linear problem is computable via numerical methods for linear programming problems, such as the simplex method [7].

C Construction of reference weight functions

For all $n \in \mathbb{N}$ ($n \geq 2$), it is possible to construct a reference weight function satisfying Hypothesis 1 with n as the condition of Hypothesis 1 can be rewritten to include a finite number of conditions

$$\int_0^1 r^{d+2j-1} w(r) dr = 0, \quad j = 1, 2, \dots, \lfloor n/2 \rfloor.$$

Here, the Gauss symbol $\lfloor a \rfloor$ denotes the largest integer that is less than or equal to a . For example, function w is set as the p th polynomial function:

$$w(r) := \begin{cases} \gamma_d \left(1 + \sum_{\ell=1}^p a_\ell r^\ell \right), & 0 \leq r < 1, \\ 0, & r \geq 1. \end{cases}$$

Then, if coefficients a_ℓ satisfy the linear equations

$$\begin{aligned} \gamma_d \left(1 + \sum_{\ell=1}^p \frac{a_\ell}{\ell + d} \right) &= 1, \\ \sum_{\ell=1}^p a_\ell &= 0, \\ \sum_{\ell=1}^p \ell a_\ell &= 0, \\ 1 + \sum_{\ell=1}^p \frac{d+2j}{d+\ell+2j} a_\ell &= 0, \quad j = 1, 2, \dots, \lfloor n/2 \rfloor, \end{aligned}$$

then w satisfies $w \in \mathcal{W}$ and Hypothesis 1 with n . Therefore, to construct reference functions with Hypothesis 1 with n represented by polynomial functions, the degree of the polynomial functions must be at least $\lfloor n/2 \rfloor + 2$.

Moreover, for all $k \in \mathbb{N}_0$, reference weight functions satisfying Hypothesis 2 with k can be constructed based on the following proposition.

Proposition 1 *Assume that reference the weight function w defined in \mathbb{R}_0^+ satisfies $w \in C^1(\mathbb{R}_0^+)$ and is represented by a polynomial function in $[0, s]$ for $s \in (0, 1]$. Let p_0 be the minimum degree of w in $[0, s]$. Then, if $p_0 - k \geq 1$, w satisfies Hypothesis 2 with k .*

Proof From the assumption, w can be represented by

$$w(r) = \sum_{\ell=p_0}^p a_\ell r^\ell, \quad 0 \leq r < s,$$

where $p \in \mathbb{N}$ and $a_\ell \in \mathbb{R}$ ($\ell = p_0, p_0 + 1, \dots, p$). Set $w^{(k)}$ as (8). Since

$$\begin{aligned} \sup_{r \in (0, s)} |w^{(k+1)}(r)| &\leq \sum_{\ell=p_0}^p |a_\ell| s^{\ell-k-1} < \infty, \\ \sup_{r \in (s, \infty)} |w^{(k+1)}(r)| &\leq \frac{\|w\|_{C([s, 1])}}{s^{k+1}} < \infty \end{aligned}$$

and

$$\begin{aligned} \sup_{r \in (0, s)} \left| \frac{d}{dr} w^{(k)}(r) \right| &\leq \sum_{\ell=p_0}^p (\ell - k) |a_\ell| s^{\ell-k-1} < \infty, \\ \sup_{r \in (s, \infty)} \left| \frac{d}{dr} w^{(k)}(r) \right| &\leq \frac{k \|w\|_{C^1([s, 1])}}{s^{k+1}} < \infty, \end{aligned}$$

if $p_0 - k \geq 1$, we have w satisfies Hypothesis 2 with k .

This proposition means that the regularity of the reference functions around zero is important.

References

1. Amicarella, A., Marongiu, J.C., Leboeuf, F., Leduc, J., Caro, J.: SPH truncation error in estimating a 3D function. *Comput. & Fluids* **44**(1), 279–296 (2011)
2. Amicarella, A., Marongiu, J.C., Leboeuf, F., Leduc, J., Neuhauser, M., Fang, L., Caro, J.: SPH truncation error in estimating a 3D derivative. *Int. J. Numer. Methods Eng.* **87**(7), 677–700 (2011)
3. Ben Moussa, B.: On the convergence of SPH method for scalar conservation laws with boundary conditions. *Methods Appl. Anal.* **13**(1), 29–62 (2006)
4. Ben Moussa, B., Vila, J.: Convergence of SPH method for scalar nonlinear conservation laws. *SIAM J. Numer. Anal.* **37**(3), 863–887 (2000)
5. Benz, W., Asphaug, E.: Simulations of brittle solids using smooth particle hydrodynamics. *Comput. Phys. Commun.* **87**(1), 253–265 (1995)
6. Boissonnat, J.D., Yvinec, M.: *Algorithmic geometry*. Cambridge Univ. Press, Cambridge (1998)
7. Dantzig, G.B.: *Linear programming and extensions*. Princeton Landmarks in Mathematics. Princeton Univ. Press, Princeton (1998)
8. Dehnen, W., Aly, H.: Improving convergence in smoothed particle hydrodynamics simulations without pairing instability. *Monthly Not. Roy. Astronom. Soc.* **425**(2), 1068–1082 (2012)
9. Fulk, D.A., Quinn, D.W.: An analysis of 1-d smoothed particle hydrodynamics kernels. *J. Comput. Phys.* **126**(1), 165–180 (1996)
10. Gingold, R.A., Monaghan, J.J.: Smoothed particle hydrodynamics-theory and application to non-spherical stars. *Monthly Not. Roy. Astronom. Soc.* **181**, 375–389 (1977)
11. Imoto, Y., Tagami, D.: A truncation error estimate of the interpolants of a particle method based on the Voronoi decomposition. *JSIAM Letters* **8**, 29–32 (2016)
12. Imoto, Y., Tagami, D.: Truncation error estimates of approximate differential operators of a particle method based on the Voronoi decomposition. *JSIAM Letters* **9**, 69–72 (2017)
13. Ishijima, K., Kimura, M.: Truncation error analysis of finite difference formulae in meshfree particle methods (*in Japanese*). *Trans. Japan Soc. Ind. Appl. Math.* **20**, 165–182 (2010).
14. Khayyer, A., Gotoh, H.: Enhancement of stability and accuracy of the moving particle semi-implicit method. *J. Comput. Phys.* **230**(8), 3093–3118 (2011)
15. Koshizuka, S.: A particle method for incompressible viscous flow with fluid fragmentation. *J. Comput. Fluid Dyn.* **4**, 29–46 (1995)
16. Koshizuka, S., Oka, Y.: Moving-particle semi-implicit method for fragmentation of incompressible fluid. *Nuclear Sci. Eng.* **123**(3), 421–434 (1996)
17. Levin, D.: The approximation power of moving least-squares. *Math. Comp.* **67**(224), 1517–1531 (1998)
18. Liu, M., Liu, G.: Smoothed particle hydrodynamics (SPH): an overview and recent developments. *Arch. Comput. Methods Eng.* **17**(1), 25–76 (2010)
19. Lucy, L.B.: A numerical approach to the testing of the fission hypothesis. *Astronom. J.* **82**, 1013–1024 (1977)
20. Mas-Gallic, S., Raviart, P.: A particle method for first-order symmetric systems. *Numer. Math.* **51**(3), 323–352 (1987)
21. Monaghan, J.J.: Simulating free surface flows with SPH. *J. Comput. Phys.* **110**(2), 399–406 (1994)
22. Monaghan, J.J., Lattanzio, J.C.: A simulation of the collapse and fragmentation of cooling molecular clouds. *Astrophys. J.* **375**, 177–189 (1991)
23. Murotani, K., Koshizuka, S., Tamai, T., Shibata, K., Mitsume, N., Yoshimura, S., Tanaka, S., Hasegawa, K., Nagai, E., Fujisawa, T.: Development of hierarchical domain decomposition explicit MPS method and application to large-scale tsunami analysis with floating objects. *J. Adv. Simulation Sci. Eng.* **1**(1), 16–35 (2014)
24. Price, D.J.: Smoothed particle hydrodynamics and magnetohydrodynamics. *J. Comput. Phys.* **231**(3), 759–794 (2012)
25. Quinlan, N.J., Basa, M., Lastiwka, M.: Truncation error in mesh-free particle methods. *Int. J. Numer. Methods Eng.* **66**(13), 2064–2085 (2006)

26. Raviart, P.A.: An analysis of particle methods. Numerical methods in fluid dynamics (Como, 1983), *Lect. Notes Math.*, vol. 1127, pp. 243–324. Springer, Berlin (1985)
27. Schaback, R.: Error estimates and condition numbers for radial basis function interpolation. *Adv. Comput. Math.* **3**(3), 251–264 (1995)
28. Shakibaeinia, A., Jin, Y.C.: A weakly compressible MPS method for modeling of open-boundary free-surface flow. *Int. J. Numer. Meth. Fl.* **63**(10), 1208–1232 (2010)
29. Shakibaeinia, A., Jin, Y.C.: MPS mesh-free particle method for multiphase flows. *Comput. Methods Appl. Mech. Eng.* **229**, 13–26 (2012)
30. Wendland, H.: Scattered Data Approximation, *Cambridge Monographs on Applied and Computational Mathematics*, vol. 17. Cambridge Univ. Press, Cambridge (2005)