

Derived Categories in Representation Theory

Jun-ichi Miyachi

Department of Mathematics, Tokyo Gakugei University,
Koganei-shi, Tokyo, 184-8501, Japan
e-mail : miyachi@u-gakugei.ac.jp

We survey recent methods of derived categories in the representation theory of algebras.

1 Triangulated Categories and Brown Representability

Definition 1.1 A triangulated category \mathcal{C} is an additive category together with


- (1) an autofunctor $T : \mathcal{C} \xrightarrow{\sim} \mathcal{C}$ (i.e. there is T^{-1} such that $T \circ T^{-1} = T^{-1} \circ T = 1_{\mathcal{C}}$) called the translation, and
- (2) a collection \mathcal{T} of sextuples (X, Y, Z, u, v, w) :

$$X \xrightarrow{u} Y \xrightarrow{v} Z \xrightarrow{w} T(X)$$

called (distinguished) triangles. These data are subject to the following four axioms:

- (TR1) (1) Every sextuple (X, Y, Z, u, v, w) which is isomorphic to a (distinguished) triangle is a (distinguished) triangle.
- (2) Every morphism $u : X \rightarrow Y$ is embedded in a (distinguished) triangle

$$X \xrightarrow{u} Y \xrightarrow{v} Z \xrightarrow{w} T(X)$$



- (3) For any $X \in \mathcal{C}$,

$$X \xrightarrow{1} X \rightarrow 0 \rightarrow T(X)$$

is a (distinguished) triangle

(TR2) A sextuple

$$X \xrightarrow{u} Y \xrightarrow{v} Z \xrightarrow{w} T(X)$$

is a (distinguished) triangle if and only if

$$Y \xrightarrow{v} Z \xrightarrow{w} T(X) \xrightarrow{-T(u)} T(Y)$$

is a (distinguished) triangle.

(TR3) For any (distinguished) triangles (X, Y, Z, u, v, w) , (X', Y', Z', u', v', w') and a commutative diagram

$$\begin{array}{ccccc} X & \xrightarrow{u} & Y & \xrightarrow{v} & Z & \xrightarrow{w} & T(X) \\ \downarrow f & & \downarrow g & & & & \\ X' & \xrightarrow{u'} & Y' & \xrightarrow{v'} & Z' & \xrightarrow{w'} & T(X') \end{array}$$

there exists $h : Z \rightarrow Z'$ which makes a commutative diagram

$$\begin{array}{ccccc} X & \xrightarrow{u} & Y & \xrightarrow{v} & Z & \xrightarrow{w} & T(X) \\ \downarrow f & & \downarrow g & & \downarrow h & & \downarrow T(f) \\ X' & \xrightarrow{u'} & Y' & \xrightarrow{v'} & Z' & \xrightarrow{w'} & T(X') \end{array}$$

(TR4) (Octahedral axiom) For any two consecutive morphisms $u : X \rightarrow Y$ and $v : Y \rightarrow Z$, if we embed u , vu and v in (distinguished) triangles (X, Y, Z', u, i, i') , (X, Z, Y', vu, k, k') and (Y, Z, X', v, j, j') , respectively, then there exist morphisms $f : Z' \rightarrow Y'$, $g : Y' \rightarrow X'$ such that the following diagram commute

$$\begin{array}{ccccccc} X & \xrightarrow{u} & Y & \xrightarrow{i} & Z' & \xrightarrow{i'} & T(X) \\ \parallel & & \downarrow v & & \downarrow f & & \parallel \\ X & \xrightarrow{vu} & Z & \xrightarrow{k} & Y' & \xrightarrow{k'} & T(X) \\ & & \downarrow j & & \downarrow g & & \downarrow T(u) \\ & & X' & \xrightarrow{j'} & X' & \xrightarrow{j'} & TY \\ & & \downarrow j' & & \downarrow T(i)j' & & \\ & & T(Y) & \xrightarrow{T(i)} & T(Z') & & \end{array}$$

and the third column is a triangle.

Sometimes, we write $X[i]$ for $T^i(X)$.

Definition 1.2 (∂ -functor) Let $\mathcal{C}, \mathcal{C}'$ be triangulated categories. An additive functor $F : \mathcal{C} \rightarrow \mathcal{C}'$ is called a ∂ -functor (sometimes exact functor) provided that there is a functorial isomorphism $\alpha : FT_{\mathcal{C}} \xrightarrow{\sim} T_{\mathcal{C}'}F$ such that

$$F(X) \xrightarrow{F(u)} F(Y) \xrightarrow{F(v)} F(Z) \xrightarrow{\alpha_X F(w)} T_{\mathcal{C}'}(F(X))$$

is a triangle in \mathcal{C}' whenever $X \xrightarrow{u} Y \xrightarrow{v} Z \xrightarrow{w} T_{\mathcal{C}}(X)$ is a triangle in \mathcal{C} . Moreover, if a ∂ -functor F is an equivalence, then F is called a triangulated equivalence. In this case, we denote by $\mathcal{C} \stackrel{\Delta}{\cong} \mathcal{C}'$.

For $(F, \alpha), (G, \beta) : \mathcal{C} \rightarrow \mathcal{C}'$ ∂ -functors, a functorial morphism $\phi : F \rightarrow G$ is called a ∂ -functorial morphism if

$$(T_{\mathcal{C}'}\phi) \circ \alpha = \beta \circ \phi T_{\mathcal{C}}$$

We denote by $\partial(\mathcal{C}, \mathcal{C}')$ the collection of all ∂ -functors from \mathcal{C} to \mathcal{C}' , and denote by $\partial \text{Mor}(F, G)$ the collection of ∂ -functorial morphisms from F to G .

Proposition 1.3 Let $F : \mathcal{C} \rightarrow \mathcal{C}'$ be a ∂ -functor between triangulated categories. If $G : \mathcal{C}' \rightarrow \mathcal{C}$ is a right (or left) adjoint of F , then G is also a ∂ -functor.

Definition 1.4 A contravariant (resp., covariant) additive functor $H : \mathcal{C} \rightarrow \mathcal{A}$ from a triangulated category \mathcal{C} to an abelian category \mathcal{A} is called a homological functor (resp., a cohomological functor), if for any triangle (X, Y, Z, u, v, w) in \mathcal{C} the sequence

$$\begin{aligned} & H(T(X)) \rightarrow H(Z) \rightarrow H(Y) \rightarrow H(X) \\ (\text{resp., } & H(X) \rightarrow H(Y) \rightarrow H(Z) \rightarrow H(T(X))) \end{aligned}$$

is exact. Taking $H(T^i(X)) = H^i(X)$, we have the long exact sequence:

$$\begin{aligned} & \cdots \rightarrow H^{i+1}(X) \rightarrow H^i(Z) \rightarrow H^i(Y) \rightarrow H^i(X) \rightarrow \cdots \\ (\text{resp., } & \cdots \rightarrow H^i(X) \rightarrow H^i(Y) \rightarrow H^i(Z) \rightarrow H^{i+1}(X) \rightarrow \cdots) \end{aligned}$$

Proposition 1.5 The following hold.

1. If (X, Y, Z, u, v, w) is a triangle, then $vu = 0$, $wv = 0$ and $T(u)w = 0$.
2. For any $X \in \mathcal{C}$, $\text{Hom}_{\mathcal{C}}(-, X) : \mathcal{C} \rightarrow \mathfrak{Ab}$ (resp., $\text{Hom}_{\mathcal{C}}(X, -) : \mathcal{C} \rightarrow \mathfrak{Ab}$) is a homological functor (resp., a cohomological functor).
3. For any homomorphism of triangles

$$\begin{array}{ccccccc} X & \xrightarrow{u} & Y & \xrightarrow{v} & Z & \xrightarrow{w} & T(X) \\ \downarrow f & & \downarrow g & & \downarrow h & & \downarrow T(f) \\ X' & \xrightarrow{u'} & Y' & \xrightarrow{v'} & Z' & \xrightarrow{w'} & T(X') \end{array}$$

if two of f , g and h are isomorphisms, then the rest is also an isomorphism.

Definition 1.6 (Compact Object) Let \mathcal{C} be a triangulated category. An object $C \in \mathcal{C}$ is called a compact object in \mathcal{C} if the canonical morphism

$$\coprod_{i \in I} \text{Hom}_{\mathcal{C}}(C, X_i) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(C, \coprod_{i \in I} X_i)$$

is an isomorphism for any set $\{X_i\}_{i \in I}$ of objects (if $\coprod_{i \in I} X_i$ exists in \mathcal{C}).

For a triangulated category \mathcal{C} , a set \mathcal{S} of compact objects is called a generating set if $\text{Hom}_{\mathcal{C}}(\mathcal{S}, X) = 0 \Rightarrow X = 0$, and if $T(\mathcal{S}) = \mathcal{S}$. A triangulated category \mathcal{C} is compactly generated if \mathcal{C} contains arbitrary coproducts, and if it has a generating set.

Definition 1.7 (Homotopy Limit) Let \mathcal{C} be a triangulated category which contains arbitrary coproducts (resp., products). For a sequence $\{X_i \rightarrow X_{i+1}\}_{i \in \mathbb{N}}$ (resp., $\{X_{i+1} \rightarrow X_i\}_{i \in \mathbb{N}}$) of morphisms in \mathcal{C} , the homotopy colimit (resp., homotopy limit) of the sequence is the third (resp., second) term of the triangle

$$\coprod_i X_i \xrightarrow{1\text{-shift}} \coprod_i X_i \rightarrow \underline{\text{hocolim}} X_i \rightarrow T \left(\coprod_i X_i \right)$$

(resp., $T^{-1} \left(\prod_i X_i \right) \rightarrow \underline{\text{holim}} X_i \rightarrow \prod_i X_i \xrightarrow{1\text{-shift}} \prod_i X_i$)

where the above shift morphism is the coproduct (resp., product) of $X_i \xrightarrow{f_i} X_{i+1}$ (resp., $X_{i+1} \xrightarrow{f_i} X_i$) ($i \in \mathbb{N}$).

Proposition 1.8 Let \mathcal{C} be a triangulated category which contains arbitrary coproducts, $\{X_i \rightarrow X_{i+1}\}_{i \in \mathbb{N}}$ a sequence of morphisms in \mathcal{C} . For a compact object C in \mathcal{C} , we have

$$\text{Hom}(C, \underline{\text{hocolim}} X_i) \cong \varinjlim \text{Hom}(C, X_i)$$

Proof. We have an exact sequence

$$0 \rightarrow \coprod_i \text{Hom}(C, X_i) \rightarrow \coprod_i \text{Hom}(C, X_i) \rightarrow \text{Hom}(C, \underline{\text{hocolim}} X_i) \rightarrow 0 \quad \square$$

Theorem 1.9 (Brown Representability Theorem [Ne]) Let \mathcal{C} be a compactly generated triangulated category. If a homological functor $H : \mathcal{C} \rightarrow \mathfrak{Ab}$ sends coproducts to products, then it is representable, that is, there is an object $X \in \mathcal{C}$ such that $H \cong \text{Hom}_{\mathcal{C}}(-, X)$.

Sketch of Proof. Let \mathcal{S} be a generating set of \mathcal{C} . There exist a coproduct X_1 of objects of \mathcal{S} and a morphism $h_{X_1} \rightarrow H$ such that $\text{Hom}_{\mathcal{C}}(C, X_1) \rightarrow H(C)$ is surjective for any $C \in \mathcal{S}$. For a functor $K_1 = \text{Ker}(h_{X_1} \rightarrow H)$ there exists a coproduct Z_2 of objects in \mathcal{S} and a morphism $h_{Z_2} \rightarrow K_1$ such that $\text{Hom}_{\mathcal{C}}(C, Z_2) \rightarrow K_1(C)$ is surjective for any $C \in \mathcal{S}$. Then we have a triangle $Z_2 \rightarrow X_1 \rightarrow X_2 \rightarrow Z_2[1]$. Since H is a homological functor, we have a commutative diagram

$$\begin{array}{ccccc} H(X_2) & \longrightarrow & H(X_1) & \longrightarrow & H(Z_2) \\ \downarrow \wr & & \downarrow \wr & & \downarrow \wr \\ \text{Mor}(h_{X_2}, H) & \longrightarrow & \text{Mor}(h_{X_1}, H) & \longrightarrow & \text{Mor}(h_{Z_2}, H) \end{array}$$

Then there is a morphism $X_1 \rightarrow X_2$ satisfying a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_1 & \longrightarrow & \text{Hom}_{\mathcal{C}}(-, X_1) & \longrightarrow & H \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & K_2 & \longrightarrow & \text{Hom}_{\mathcal{C}}(-, X_2) & \longrightarrow & H \end{array}$$

and we have a morphism of exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_1(C) & \longrightarrow & \text{Hom}_{\mathcal{C}}(C, X_1) & \longrightarrow & H(C) \longrightarrow 0 \\ & & \downarrow 0 & & \downarrow & & \parallel \\ 0 & \longrightarrow & K_2(C) & \longrightarrow & \text{Hom}_{\mathcal{C}}(C, X_2) & \longrightarrow & H(C) \longrightarrow 0 \end{array}$$

for any $C \in \mathcal{S}$. By inductive step, we have a triangle

$$\coprod_i X_i \xrightarrow{1\text{-shift}} \coprod_i X_i \rightarrow \text{hocolim } X_i \rightarrow T\left(\coprod_i X_i\right)$$

and we have an exact sequence

$$\begin{array}{ccccc} H(\text{hocolim } X_i) & \longrightarrow & \prod_i H(X_i) & \longrightarrow & \prod_i H(X_i) \\ \downarrow \wr & & \downarrow \wr & & \downarrow \wr \\ \text{Mor}(\text{h}_{\text{hocolim } X_i}, H) & \longrightarrow & \prod_i \text{Mor}(\text{h}_{X_i}, H) & \longrightarrow & \prod_i \text{Mor}(\text{h}_{X_i}, H) \end{array}$$

Therefore there is a morphism $\text{Hom}_{\mathcal{C}}(-, \text{hocolim } X_i) \rightarrow H$ such that

$$\text{Hom}_{\mathcal{C}}(C, \text{hocolim } X_i) \cong H(C)$$

for any $C \in \mathcal{S}$. Hence we have $\text{Hom}_{\mathcal{C}}(-, \text{hocolim } X_i) \cong H$. \square

Corollary 1.10 (Adjoint Functor Theorem [Ne]) *Let \mathcal{C} be a compactly generated triangulated category. If a ∂ -functor $F : \mathcal{C} \rightarrow \mathcal{D}$ commutes with arbitrary coproducts, then there exists a ∂ -functor $G : \mathcal{D} \rightarrow \mathcal{C}$ which is a right adjoint of F .*

Proof. For any $Y \in \mathcal{D}$, the functor

$$\text{Hom}_{\mathcal{D}}(F(-), Y) : \mathcal{C} \rightarrow \mathfrak{Ab}$$

is a homological functor. By Brown representability theorem there is an object $GY \in \mathcal{C}$ such that

$$\text{Hom}_{\mathcal{D}}(F(-), Y) \cong \text{Hom}_{\mathcal{C}}(-, GY) \quad \square$$

Definition 1.11 (Multiplicative System) *Let S be a multiplicative system in a triangulated category \mathcal{C} which satisfies the following conditions:*

- (FR0) *For a morphism s in \mathcal{C} , if there exist f, g such that $sf, gs \in S$, then $s \in S$.*
- (FR1) *(1) $1_X \in S$ for every $X \in \mathcal{C}$.*
(2) For $s, t \in S$, if st is defined, then $st \in S$.

(FR2) Every diagram in \mathcal{C}

$$\begin{array}{ccc} X & \xrightarrow{s} & Y \\ f \downarrow & & \\ X' & & \end{array}$$

with $s \in S$, can be completed to a commutative square

$$\begin{array}{ccc} X & \xrightarrow{s} & Y \\ f \downarrow & & \downarrow g \\ X' & \xrightarrow{t} & Y' \end{array}$$

with $s, t \in S$. Ditto for the statement with all arrows reversed.

(FR3) For $f, g \in \text{Hom}_{\mathcal{C}}(X, Y)$ the following are equivalent.


- (1) There exists $s \in S$ such that $sf = sg$.
- (2) There exists $t \in S$ such that $ft = gt$.

(FR4) For a morphism u in \mathcal{C} , $u \in S$ if and only if $T(u) \in S$.

(FR5) For triangles (X, Y, Z, u, v, w) , (X', Y', Z', u', v', w') and morphisms $f : X \rightarrow X'$, $g : Y \rightarrow Y'$ in S with $gu = u'f$, there exists $h : Z \rightarrow Z'$ in S such that (f, g, h) is a homomorphism of triangles.

Definition 1.12 (Quotient Category) We define the quotient category $S^{-1}\mathcal{C}$ of \mathcal{C} , as follows:


1. $\text{Ob}(S^{-1}\mathcal{C}) = \text{Ob}(\mathcal{C})$.
2. For $X, Y \in \text{Ob}(\mathcal{C})$, let $V(X, Y) = \{(s, Y', f) \mid s : Y \rightarrow Y' \in S, f : X \rightarrow Y'\}$. In $V(X, Y)$, we define $(s, Y', f) \sim (s', Y'', f')$ if there is (s'', Y''', f') such that all triangles are commutative in the following diagram:



Then we define a morphism from X to Y by an equivalence class $s^{-1}f$ of (s, Y', f) .

3. For $s^{-1}f : X \rightarrow Y, t^{-1}g : Y \rightarrow Z$, by (FR2) there are $s' : Z' \rightarrow Z'' \in S$ and

$g' : Y' \rightarrow Z''$ such that $s'og = g'os$. Then we define $(t^{-1}g) \circ (s^{-1}f) = (s'ot)^{-1}g'of$.



Moreover, we define the quotient functor $Q : \mathcal{C} \rightarrow \mathcal{S}^{-1}\mathcal{C}$ by

(Q1) $Q(X) = X$ for $X \in \mathcal{C}$.

(Q2) $Q(f) = 1_Y^{-1}f$ for a morphism $f : X \rightarrow Y$ in \mathcal{C} .

Remark 1.13 Can we define (2) in the above?

Definition 1.14 (Épaisse Subcategory) Let \mathcal{C} be a triangulated category. A full subcategory \mathcal{U} of \mathcal{C} is called a full triangulated subcategory if $X \rightarrow Y$ is a morphism in \mathcal{U} , then there is a triangle $X \rightarrow Y \rightarrow Z \rightarrow TX$ with $Z \in \mathcal{U}$.

A full triangulated subcategory \mathcal{U} is called an épaisse subcategory if it is closed under direct summands. In this case, let $S(\mathcal{U})$ be the collection of morphisms s such that $X \xrightarrow{s} Y \rightarrow Z \rightarrow X[1]$ is a triangle with $Z \in \mathcal{U}$. Then $S(\mathcal{U})$ is a multiplicative system satisfying (FR0) - (FR5). We write $\mathcal{C}/\mathcal{U} = S(\mathcal{U})^{-1}\mathcal{C}$.

In the case that \mathcal{C} contains arbitrary coproducts, a full triangulated subcategory \mathcal{U} is called a localizing subcategory if it is closed under coproducts.

Remark 1.15 The above definition of an épaisse subcategory \mathcal{U} is the same as the original definition [Ve], that is, a full triangulated category satisfying that if $X \xrightarrow{u} Y$ factors through some object in \mathcal{U} and if there is a triangle $X \xrightarrow{u} Y \rightarrow Z \rightarrow T(X)$ with $Z \in \mathcal{U}$, then $X, Y \in \mathcal{U}$.

Proposition 1.16 ([BN]) Let \mathcal{C} be a triangulated category which contains arbitrary coproducts. Then any localizing subcategory is an épaisse subcategory.

Sketch of Proof. Let \mathcal{U} be a localizing subcategory, and $X \in \mathcal{U}$ with $X = Y \oplus Z$ in \mathcal{C} . We take a morphism $e : X \rightarrow Y \hookrightarrow X$, and consider the sequence of morphisms

$$(*) \quad X \xrightarrow{e} X \xrightarrow{e} X \xrightarrow{e} \dots$$

Then it is easy to see that $Y \cong \text{hocolim}(*) \in \mathcal{U}$. □

Proposition 1.17 Let \mathcal{C} be a triangulated category. For a multiplicative system S satisfying the conditions (FR0) - (FR5), let $\mathcal{U}(S)$ be the full triangulated subcategory consisting of objects Z which is in a triangle $X \xrightarrow{s} Y \rightarrow Z \rightarrow X[1]$ with $s \in S$. Then the following hold.

1. $S(\mathcal{U})$ and $\mathcal{U}(S)$ induce an 1 - 1 correspondence between the collection of multiplicative systems S satisfying the conditions (FR0) - (FR5) and the collection of épaisse subcategories \mathcal{U} .
2. For an épaisse subcategory \mathcal{U} , \mathcal{C}/\mathcal{U} is a triangulated category whose (distinguished) triangles are defined to be isomorphic to (distinguished) triangles of \mathcal{C} .
3. Assume \mathcal{C} contains arbitrary coproducts. For a localizing subcategory \mathcal{U} , \mathcal{C}/\mathcal{U} also contains arbitrary coproducts.

Definition 1.18 (stable t -structure) For full subcategories \mathcal{U} and \mathcal{V} of a triangulated category \mathcal{C} , $(\mathcal{U}, \mathcal{V})$ is called a stable t -structure in \mathcal{C} provided that

1. \mathcal{U} and \mathcal{V} are stable for translations.
2. $\text{Hom}_{\mathcal{C}}(\mathcal{U}, \mathcal{V}) = 0$.
3. For every $X \in \mathcal{C}$, there exists a triangle $U \rightarrow X \rightarrow V \rightarrow TU$ with $U \in \mathcal{U}$ and $V \in \mathcal{V}$.

Proposition 1.19 ([BBD], c.f. [Mi]) Let \mathcal{C} be a triangulated category, $(\mathcal{U}, \mathcal{V})$ a stable t -structure in \mathcal{C} , and $i_* : \mathcal{U} \rightarrow \mathcal{C}, j_* : \mathcal{V} \rightarrow \mathcal{C}$ the canonical embeddings. Then the following hold.

1. \mathcal{U} and \mathcal{V} is épaisse subcategories of \mathcal{C} .
2. i_* (resp., j_*) has a right adjoint $i^!$ (resp., a left adjoint j^*).
3. The adjunction arrows induce a triangle

$$i_*i^!X \xrightarrow{\alpha_X} X \xrightarrow{\beta_X} j_*j^*X \rightarrow i_*i^!X[1]$$

for any $X \in \mathcal{C}$.

4. \mathcal{C}/\mathcal{U} (resp., \mathcal{C}/\mathcal{V}) exists, and it is triangulated equivalent to \mathcal{V} (resp., \mathcal{U}).

$$\begin{array}{ccccc}
 & & \mathcal{C}/\mathcal{V} & & \\
 & & \uparrow & \swarrow & \\
 & & \mathcal{U} & \xrightarrow{i_*} & \mathcal{C} & \xrightarrow{j_*} & \mathcal{V} \\
 & & \xleftarrow{i^!} & \mathcal{C} & \xleftarrow{j_*} & \mathcal{V} & \downarrow i \\
 & & & & & & \mathcal{C}/\mathcal{U}
 \end{array}$$

Corollary 1.20 Let \mathcal{C} be a compactly generated triangulated category, and \mathcal{U} a localizing subcategory of \mathcal{C} . Then \mathcal{C}/\mathcal{U} can be defined if and only if there is a full triangulated subcategory \mathcal{V} such that $(\mathcal{U}, \mathcal{V})$ a stable t -structure in \mathcal{C} .

Proof. If \mathcal{C}/\mathcal{U} can be defined, then the quotient functor $Q : \mathcal{C} \rightarrow \mathcal{C}/\mathcal{U}$ commutes with coproducts. By Adjoint Functor Theorem, Q has a right adjoint $F : \mathcal{C}/\mathcal{U} \rightarrow \mathcal{C}$. By Proposition 1.19, it is easy to see that $(\mathcal{U}, \text{Im } F)$ is a stable t -structure in \mathcal{C} . \square

2 Derived Categories

Throughout this section, \mathcal{A} is an abelian category and \mathcal{B} is an additive subcategory of \mathcal{A} which is closed under isomorphisms.

Definition 2.1 (Complex) A (cochain) complex is a collection $X^\cdot = (X^n, d_X^n : X^n \rightarrow X^{n+1})_{n \in \mathbb{Z}}$ of objects and morphisms of \mathcal{B} such that $d_X^{n+1} d_X^n = 0$. A complex $X^\cdot = (X^n, d_X^n : X^n \rightarrow X^{n+1})_{n \in \mathbb{Z}}$ is called bounded below (resp., bounded above, bounded) if $X^n = 0$ for $n \ll 0$ (resp., $n \gg 0$, $n \ll 0$ and $n \gg 0$).

A complex $X^\cdot = (X^n, d_X^n)$ is called a stalk complex if there exists an integer n_0 such that $X^i = 0$ if $i \neq n_0$. We identify objects of \mathcal{B} with a stalk complexes of degree 0.

A morphism $f : X^\cdot \rightarrow Y^\cdot$ of complexes is a collection of morphisms $f^n : X^n \rightarrow Y^n$ which makes a commutative diagram

$$\begin{array}{ccccccc} \dots & \longrightarrow & X^n & \xrightarrow{d_X^n} & X^{n+1} & \longrightarrow & \dots \\ & & \downarrow f^n & & \downarrow f^{n+1} & & \\ \dots & \longrightarrow & Y^n & \xrightarrow{d_Y^n} & Y^{n+1} & \longrightarrow & \dots \end{array}$$

We denote by $\mathcal{C}(\mathcal{B})$ (resp., $\mathcal{C}^+(\mathcal{B})$, $\mathcal{C}^-(\mathcal{B})$, $\mathcal{C}^b(\mathcal{B})$) the category of complexes (resp., bounded below complexes, bounded above complexes, bounded complexes) of \mathcal{B} . An autofunctor $T : \mathcal{C}(\mathcal{B}) \rightarrow \mathcal{C}(\mathcal{B})$ is called translation if $(TX^\cdot)^n = X^{n+1}$ and $(Td_X)^n = -d_X^{n+1}$ for any complex $X^\cdot = (X^n, d_X^n)$.

In $\mathcal{C}(\mathcal{A})$, a morphism $u : X^\cdot \rightarrow Y^\cdot$ is called a quasi-isomorphism if $H_n(u)$ is an isomorphism for any n .

In this section, “*” means “nothing”, “+”, “-” or “b”.

Definition 2.2 For $u \in \text{Hom}_{\mathcal{C}(\mathcal{B})}(X^\cdot, Y^\cdot)$, the mapping cone of u is a complex $M^\cdot(u)$ with

$$\begin{aligned} M^n(u) &= X^{n+1} \oplus Y^n, \\ d_{M^\cdot(u)}^n &= \begin{bmatrix} -d_X^{n+1} & 0 \\ u^{n+1} & d_Y^n \end{bmatrix} : X^{n+1} \oplus Y^n \rightarrow X^{n+2} \oplus Y^{n+1}. \end{aligned}$$

Definition 2.3 (Homotopy Relation) Two morphisms $f, g \in \text{Hom}_{\mathcal{C}(\mathcal{B})}(X^\cdot, Y^\cdot)$ are said to be homotopic (denote by $f \underset{h}{\simeq} g$) if there is a collection of morphisms $h = (h^n)$, $h^n : X^n \rightarrow Y^{n+1}$ such that $f^n - g^n = d_Y^{n-1} h^n + h^{n+1} d_X^n$ for all $n \in \mathbb{Z}$.

Definition 2.4 (Homotopy Category) The homotopy category $K^*(\mathcal{B})$ of \mathcal{B} is defined by

1. $\text{Ob}(K^*(\mathcal{B})) = \text{Ob}(\mathcal{C}^*(\mathcal{B}))$,
2. $\text{Hom}_{K^*(\mathcal{B})}(X^\cdot, Y^\cdot) = \text{Hom}_{\mathcal{C}^*(\mathcal{B})}(X^\cdot, Y^\cdot) / \underset{h}{\simeq}$ for $X^\cdot, Y^\cdot \in \text{Ob}(K^*(\mathcal{B}))$.

Proposition 2.5 A category $K^*(\mathcal{B})$ is a triangulated category whose (distinguished) triangles are defined to be isomorphic to

$$X^\cdot \xrightarrow{u} Y^\cdot \rightarrow M(u) \rightarrow T(X^\cdot)$$

for any $u : X^\cdot \rightarrow Y^\cdot$ in $K^*(\mathcal{B})$.

Definition 2.6 (Derived Category) The derived category $D^*(\mathcal{A})$ of an abelian category \mathcal{A} is $K^*(\mathcal{A})/K^{*\phi}(\mathcal{A})$, where $K^{*\phi}(\mathcal{A})$ is the full subcategory of $K^*(\mathcal{A})$ consisting of null complexes, that is, complexes whose all cohomologies are 0.

Proposition 2.7 The following hold.

1. $D^*(\mathcal{A})$ is a triangulated category, and the canonical functor $Q : K^*(\mathcal{A}) \rightarrow D^*(\mathcal{A})$ is a ∂ -functor.
2. The i -th cohomology of complexes is a cohomological functor in the sense of Definition 1.4.

Proposition 2.8 If $0 \rightarrow X^\cdot \xrightarrow{u} Y^\cdot \xrightarrow{v} Z^\cdot \rightarrow 0$ is a exact sequence in $C(\mathcal{A})$, then it can be embedded in a triangle in $D(\mathcal{A})$

$$Q(X^\cdot) \xrightarrow{Q(u)} Q(Y^\cdot) \xrightarrow{Q(v)} Q(Z^\cdot) \rightarrow TQ(X^\cdot).$$

Definition 2.9 (K-injective Complex) A complex X^\cdot of $K(\mathcal{B})$ is called *K-injective* (resp., *K-projective*) if

$$\text{Hom}_{K(\mathcal{B})}(N^\cdot, X^\cdot) = 0 \quad (\text{resp.}, \text{Hom}_{K(\mathcal{B})}(X^\cdot, N^\cdot) = 0)$$

for any null complex N^\cdot .

Example 2.10 Let A be a ring, $\text{Mod } A$ the category of right A -modules, and $\text{Inj } A$ (resp., $\text{Proj } A$) the category of injective (resp., projective) right A -modules. Then any complex $I \in K^+(\text{Inj } A)$ (resp., $P \in K^-(\text{Proj } A)$) is a *K-injective* (resp., *K-projective*) complex in $K(\text{Mod } A)$.

Example 2.11 Let k be a field, $A = k[x]/(x^2)$, and

$$X^\cdot : \cdots \xrightarrow{x} A \xrightarrow{x} A \xrightarrow{x} \cdots$$

Then X^\cdot is a null complex of finitely generated projective-injective A -modules. But it is neither *K-projective* nor *K-injective*, because $\text{Hom}_{K(\text{Mod } A)}(X^\cdot, X^\cdot) \neq 0$.

Theorem 2.12 ([Sp], [Ne], [LAM], [Fr]) Let $K^{inj}(\text{Mod } A)$ (resp., $K^{proj}(\text{Mod } A)$) be the category of *K-injective* (resp., *K-projective*) complexes, then the following hold.

1. $(K^{proj}(\text{Mod } A), K^\phi(\text{Mod } A))$ is a stable t -structure in $K(\text{Mod } A)$, and hence $D(\text{Mod } A)$ exists and is triangulated equivalent to $K^{proj}(\text{Mod } A)$.

2. $(\mathbf{K}^\phi(\text{Mod } A), \mathbf{K}^{inj}(\text{Mod } A))$ is a stable t -structure in $\mathbf{K}(\text{Mod } A)$, and hence $\mathbf{D}(\text{Mod } A)$ is triangulated equivalent to $\mathbf{K}^{inj}(\text{Mod } A)$.
3. For a Grothendieck category \mathcal{A} , $(\mathbf{K}^\phi(\mathcal{A}), \mathbf{K}^{inj}(\mathcal{A}))$ is a stable t -structure in $\mathbf{K}(\mathcal{A})$, and hence $\mathbf{D}(\mathcal{A})$ exists and is triangulated equivalent to $\mathbf{K}^{inj}(\mathcal{A})$.

Proof. For a complex $X^\cdot = (X^i, d^i)$, we define the following truncations:

$$\begin{aligned}\sigma_{\leq n} X^\cdot &: \cdots \rightarrow X^{n-2} \rightarrow X^{n-1} \rightarrow \text{Ker } d^n \rightarrow 0 \rightarrow \cdots \\ \sigma_{\geq n} X^\cdot &: \cdots \rightarrow 0 \rightarrow \text{Cok } d^{n-1} \rightarrow X^{n+1} \rightarrow X^{n+2} \rightarrow \cdots\end{aligned}$$

(1) For any n , there is a complex $P_n^\cdot \in \mathbf{K}^-(\text{Proj } A)$ which has a quasi-isomorphism $P_n^\cdot \rightarrow \sigma_{\leq n} X^\cdot$. Then we have the following quasi-isomorphisms (qis)

$$X^\cdot \cong \varinjlim \sigma_{\leq n} X^\cdot \xleftarrow{\text{qis}} \text{hocolim } \sigma_{\leq n} X^\cdot \xleftarrow{\text{qis}} \text{hocolim } P_n^\cdot$$

Since $\text{Hom}_{\mathcal{C}}(\coprod_n P_n^\cdot, -) \cong \prod_n \text{Hom}_{\mathcal{C}}(P_n^\cdot, -)$, $\coprod_n P_n^\cdot$ is \mathbf{K} -projective. Here $\text{h}^M = \text{Hom}_{\mathcal{C}}(M, -)$ for any object M . It is easy to see that $\text{hocolim } P_n^\cdot$ is \mathbf{K} -projective by the following exact sequence

$$\text{h}\coprod_n P_n^\cdot \rightarrow \text{h}\coprod_n P_n^\cdot \rightarrow \text{h} \xrightarrow{\text{hocolim } P_n^\cdot} \text{h}^T(\coprod_n P_n^\cdot) \rightarrow \text{h}^T(\coprod_n P_n^\cdot)$$

(2) For any n , there is a complex $I_n^\cdot \in \mathbf{K}^+(\text{Inj } A)$ which has a quasi-isomorphism $\sigma_{\geq -n} X^\cdot \rightarrow I_n^\cdot$. Then we have the following quasi-isomorphisms (qis)

$$X^\cdot \cong \varprojlim \sigma_{\geq -n} X^\cdot \xrightarrow{\text{qis}} \text{holim } \sigma_{\geq -n} X^\cdot \xrightarrow{\text{qis}} \text{holim } I_n^\cdot$$

by the same reason of (1), we have the statement.

(3) Because there is a ring A such that \mathcal{A} is a localization of $\text{Mod } A$ (Gabriel-Popescu Theorem). See [LAM] or [Fr] for details. \square

Remark 2.13 If P^\cdot is a \mathbf{K} -projective complex (e.g. a bounded above complex of projective A -modules), then we have

$$\text{Hom}_{\mathbf{K}(\text{Mod } A)}(P^\cdot, X^\cdot) \cong \text{Hom}_{\mathbf{D}(\text{Mod } A)}(P^\cdot, X^\cdot)$$

for any complex X^\cdot . Similarly, for a \mathbf{K} -injective complex I^\cdot (e.g. bounded below complex of injective A -modules), then we have

$$\text{Hom}_{\mathbf{K}(\text{Mod } A)}(X^\cdot, I^\cdot) \cong \text{Hom}_{\mathbf{D}(\text{Mod } A)}(X^\cdot, I^\cdot)$$

for any complex X^\cdot . In particular, given A -modules M, N , we have

$$\text{Ext}_A^i(M, N) \cong \text{Hom}_{\mathbf{D}(\text{Mod } A)}(M, N[i])$$

Definition 2.14 (Double Complex, Total complex) A double complex $C^{\cdot,\cdot}$ is a bi-graded object $(C^{p,q})_{p,q \in \mathbb{Z}}$ of \mathcal{A} together with $d_I^{p,q} : C^{p,q} \rightarrow C^{p+1,q}$ and $d_{II}^{p,q} : C^{p,q} \rightarrow C^{p,q+1}$ such that

$$C^{\cdot,q} = (C^{p,q}, d_I^{p,q} : C^{p,q} \rightarrow C^{p+1,q}), \quad C^{p,\cdot} = (C^{p,q}, d_{II}^{p,q} : C^{p,q} \rightarrow C^{p,q+1})$$

are complexes satisfying $d_I^{p,q+1} d_{II}^{p,q} - d_{II}^{p+1,q} d_I^{p,q} = 0$. For a double complex $C^{\cdot,\cdot}$, we define the total complexes

$$\begin{aligned} \text{Tot}^{\text{II}} C^{\cdot,\cdot} &= (X^n, d^n), \text{ where } X^n = \coprod_{p+q=n} C^{p,q}, d^n = \coprod_{p+q=n} (d_I^{p,q} + (-1)^p d_{II}^{p,q}) \\ \text{Tot}^{\text{I}} C^{\cdot,\cdot} &= (Y^n, d^n), \text{ where } Y^n = \prod_{p+q=n} C^{p,q}, d^n = \prod_{p+q=n} (d_I^{p,q} + (-1)^p d_{II}^{p,q}). \end{aligned}$$

Definition 2.15 (Cartan-Eilenberg Resolution) For a complex $X^{\cdot} \in \text{D}(\text{Mod } A)$, let

$$\dots \rightarrow P^{-1\cdot} \rightarrow P^{0\cdot} \rightarrow X^{\cdot} \rightarrow 0 \quad (\text{resp.}, \quad 0 \rightarrow X^{\cdot} \rightarrow I^{0\cdot} \rightarrow I^{1\cdot} \rightarrow \dots)$$

be an exact sequence with P^n (resp., I^n) being a complex of projective (resp., injective) A -modules. We call $\dots \rightarrow P^{-1\cdot} \rightarrow P^{0\cdot}$ (resp., $I^{0\cdot} \rightarrow I^{1\cdot} \rightarrow \dots$) a Cartan-Eilenberg projective (resp., injective) resolution of X^{\cdot} if the induced complexes $\dots \rightarrow B^n(P^{-1\cdot}) \rightarrow B^n(P^{0\cdot})$ and $\dots \rightarrow H^n(P^{-1\cdot}) \rightarrow H^n(P^{0\cdot})$ (resp., $B^n(I^{0\cdot}) \rightarrow B^n(I^{1\cdot}) \rightarrow \dots$ and $H^n(I^{0\cdot}) \rightarrow H^n(I^{1\cdot}) \rightarrow \dots$) are also projective (resp., injective) resolutions of $B^n(X^{\cdot}), H^n(X^{\cdot})$, respectively.

Proposition 2.16 Under the setting of Definition 2.15, the following hold.

1. $\text{Tot}^{\text{II}} P^{\cdot,\cdot}$ is \mathbf{K} -projective, and the induced morphism of complexes $\text{Tot}^{\text{II}} P^{\cdot,\cdot} \rightarrow X^{\cdot}$ is a quasi-isomorphism.
2. $\text{Tot}^{\text{I}} I^{\cdot,\cdot}$ is \mathbf{K} -injective, and the induced morphism of complexes $X^{\cdot} \rightarrow \text{Tot}^{\text{I}} I^{\cdot,\cdot}$ is a quasi-isomorphism.

Sketch of Proof. We consider the following truncations

$$\sigma_{\leq n}^{\text{II}} P^{\cdot,\cdot} : \dots \rightarrow \sigma_{\leq n} P^{-1\cdot} \rightarrow \sigma_{\leq n} P^{0\cdot}, \quad \sigma_{\geq n}^{\text{I}} I^{\cdot,\cdot} : \sigma_{\geq n} I^{0\cdot} \rightarrow \sigma_{\geq n} I^{1\cdot} \rightarrow \dots$$

Then it is easy to see $\text{Tot}^{\text{II}} \sigma_{\leq n}^{\text{II}} P^{\cdot,\cdot}$ (resp., $\text{Tot}^{\text{I}} \sigma_{\geq n}^{\text{I}} I^{\cdot,\cdot}$) is \mathbf{K} -projective (resp., \mathbf{K} -injective), and that the induced morphism of complexes $\text{Tot}^{\text{II}} \sigma_{\leq n}^{\text{II}} P^{\cdot,\cdot} \rightarrow \sigma_{\leq n} X^{\cdot}$ (resp., $\sigma_{\geq n} X^{\cdot} \rightarrow \text{Tot}^{\text{I}} \sigma_{\geq n}^{\text{I}} I^{\cdot,\cdot}$) is a quasi-isomorphism. Therefore we have the following quasi-isomorphisms (qis)

$$\begin{aligned} X^{\cdot} &\xleftarrow{\text{qis}} \text{hocolim}_{\leftarrow} \sigma_{\leq n} X^{\cdot} \xleftarrow{\text{qis}} \text{hocolim}_{\leftarrow} \text{Tot}^{\text{II}} \sigma_{\leq n}^{\text{II}} P^{\cdot,\cdot} \xrightarrow{\text{qis}} \text{Tot}^{\text{II}} P^{\cdot,\cdot} \\ (\text{resp.}, X^{\cdot} &\xrightarrow{\text{qis}} \text{holim}_{\leftarrow} \sigma_{\geq -n} X^{\cdot} \xrightarrow{\text{qis}} \text{holim}_{\leftarrow} \text{Tot}^{\text{I}} \sigma_{\geq -n}^{\text{I}} I^{\cdot,\cdot} \xleftarrow{\text{qis}} \text{Tot}^{\text{I}} I^{\cdot,\cdot}) \end{aligned}$$

and $\text{Tot}^{\text{II}} P^{\cdot,\cdot}$ (resp., $\text{Tot}^{\text{I}} I^{\cdot,\cdot}$) is \mathbf{K} -projective (resp., \mathbf{K} -injective). □

Definition 2.17 (Right Derived Functor) For a ∂ -functor $F : K^*(\mathcal{A}) \rightarrow K(\mathcal{A}')$, the right derived functor of F is a ∂ -functor

$$\mathbf{R}^*F : D^*(\mathcal{A}) \rightarrow D(\mathcal{A}')$$

together with a functorial morphism of ∂ -functors

$$\xi \in \partial \text{Mor}(Q_{\mathcal{A}'} \circ F, \mathbf{R}^*F \circ Q_{\mathcal{A}}^*)$$

with the following property:

For $G \in \partial(D^*(\mathcal{A}), D(\mathcal{A}'))$ and $\zeta \in \partial \text{Mor}(Q_{\mathcal{A}'} \circ F, G \circ Q_{\mathcal{A}}^*)$, there exists a unique morphism $\eta \in \partial \text{Mor}(\mathbf{R}^*F, G)$ such that

$$\zeta = (\eta Q_{\mathcal{A}}^*)\xi.$$

In other words, we can simply write the above using functor categories. For triangulated categories $\mathcal{C}, \mathcal{C}'$, the ∂ -functor category $\partial(\mathcal{C}, \mathcal{C}')$ is the category (?) consisting of ∂ -functors from \mathcal{C} to \mathcal{C}' as objects and ∂ -functorial morphisms as morphisms. Then we have

$$\begin{array}{ccc} \partial \text{Mor}(Q_{\mathcal{A}'} \circ F, -Q_{\mathcal{A}}^*) \cong \partial \text{Mor}(\mathbf{R}^*F, -) & K^*(\mathcal{A}) & \xrightarrow{F} K(\mathcal{A}') \\ & Q_{\mathcal{A}} \downarrow & \downarrow Q_{\mathcal{A}'} \\ & D^*(\mathcal{A}) & \xrightarrow[\mathbf{R}^*F]{G} D(\mathcal{A}') \end{array}$$

as functors from $\partial(D^*(\mathcal{A}), D(\mathcal{A}'))$ to \mathfrak{Set} .

Proposition 2.18 Let $\mathcal{A}, \mathcal{A}'$ be abelian categories, $F : K(\mathcal{A}) \rightarrow K(\mathcal{A}')$ a ∂ -functor. If \mathcal{A} is a Grothendieck category, then we have the right derived functor $\mathbf{R}F : D(\mathcal{A}) \rightarrow D(\mathcal{A}')$ such that $F(X^\cdot) \cong \mathbf{R}F(X^\cdot)$ for any K -injective complex X^\cdot .

Remark 2.19 In the setting of Definition 2.17, the left derived functor $\mathbf{L}^*F : D^*(\mathcal{A}) \rightarrow D(\mathcal{A}')$ can be also defined by reversing arrows of ∂ -functorial morphisms. Let $\mathbf{R}^n F(X^\cdot) = H^n(\mathbf{R}F(X^\cdot))$, $\mathbf{L}^n F(X^\cdot) = H^n(\mathbf{L}F(X^\cdot))$, then $\mathbf{R}^n F$ (resp., $\mathbf{L}^n F$) coincides with the ordinary definition of the n -th right (resp., left) derived functor. According to Proposition 2.16, if F commutes with products (resp., coproducts), then the n -th hypercohomology $\mathbf{R}^n F$ (resp., hyperhomology $\mathbf{L}^n F$) coincides with $\mathbf{R}^n F$ (resp., $\mathbf{L}^n F$) (c.f. [CE], [Mc], [We]).

Definition 2.20 ($\text{Hom}_A^\cdot, \otimes_A$) Let X^\cdot, Y^\cdot be complexes in $C(\text{Mod } A)$, Z^\cdot a complex in $C(\text{Mod } A^{op})$. We define the complex $\text{Hom}_A^\cdot(X^\cdot, Y^\cdot)$ in $C(\mathfrak{Ab})$ by

$$\text{Hom}_A^\cdot(X^\cdot, Y^\cdot) = \prod_{j-i=n} \text{Hom}_A(X^i, Y^j), \quad d_{\text{Hom}^\cdot(X,Y)}^\cdot(f) = d_X \circ f - (-1)^n f \circ d_Y$$

for $f \in \text{Hom}_A^\cdot(X^\cdot, Y^\cdot)$. And we define the complex $X^\cdot \otimes_A Z^\cdot$ in $C(\mathfrak{Ab})$ by

$$X^\cdot \otimes_A Z^\cdot = \prod_{i+j=n} X^i \otimes_A Z^j, \quad d_{X \otimes Y}^\cdot = d_X \otimes 1 + (-1)^n 1 \otimes d_Z$$

Proposition 2.21 *Let A be a ring. Then we have a right derived functor*

$$\mathbf{R}\mathrm{Hom}_A^{\cdot} : \mathrm{D}(\mathrm{Mod} A)^{op} \times \mathrm{D}(\mathrm{Mod} A) \rightarrow \mathrm{D}(\mathfrak{Ab})$$

and a left derived functor

$$\dot{\otimes}_A^L : \mathrm{D}(\mathrm{Mod} A) \times \mathrm{D}(\mathrm{Mod} A^{op}) \rightarrow \mathrm{D}(\mathfrak{Ab})$$

Proposition 2.22 *Let A be a ring. For complexes X^{\cdot}, Y^{\cdot} , we have isomorphisms*

$$\begin{aligned} H^n(\mathrm{Hom}_A^{\cdot}(X^{\cdot}, Y^{\cdot})) &\cong \mathrm{Hom}_{\mathbf{K}(\mathrm{Mod} A)}(X^{\cdot}, Y^{\cdot}[n]) \\ H^n(\mathbf{R}\mathrm{Hom}_A^{\cdot}(X^{\cdot}, Y^{\cdot})) &\cong \mathrm{Hom}_{\mathrm{D}(\mathrm{Mod} A)}(X^{\cdot}, Y^{\cdot}[n]) \end{aligned}$$

Definition 2.23 (Perfect Complex) *Let A be a ring. A complex $X^{\cdot} \in \mathrm{D}(\mathrm{Mod} A)$ is called a perfect complex if X^{\cdot} is quasi-isomorphic to a bounded complex of finitely generated projective A -modules.*

Let X be a scheme, $\mathrm{D}(X)$ the derived category of sheaves of \mathcal{O}_X -modules. We denote by $\mathrm{D}_{qc}(X)$ the full subcategory of $\mathrm{D}(X)$ consisting of complexes whose cohomologies are quasi-coherent sheaves. A complex $X^{\cdot} \in \mathrm{D}_{qc}(X)$ is called a perfect complex if X^{\cdot} is locally quasi-isomorphic to a bounded complex of vector bundles (See [TT]).

We denote by $\mathrm{D}_{pf}(\mathcal{A})$ the full triangulated subcategory of $\mathrm{D}(\mathcal{A})$ consisting of perfect complexes.

Proposition 2.24 ([Rd1], [Ne]) *For a ring A , the following hold.*

1. *A complex $X^{\cdot} \in \mathrm{D}(\mathrm{Mod} A)$ is perfect if and only if it is a compact object in $\mathrm{D}(\mathrm{Mod} A)$.*
2. *$\mathrm{D}(\mathrm{Mod} A)$ is compactly generated.*

Theorem 2.25 ([BV]) *Let X be a quasi-compact quasi-separated scheme, then the following hold.*

1. *A complex $X^{\cdot} \in \mathrm{D}_{qc}(X)$ is perfect if and only if it is a compact object in $\mathrm{D}_{qc}(X)$.*
2. *$\mathrm{D}_{qc}(X)$ is compactly generated.*

Theorem 2.26 ([BN]) *Let X be a quasi-compact separated scheme, then the canonical functor $\mathrm{D}(\mathrm{Qcoh} X) \rightarrow \mathrm{D}_{qc}(X)$ is a triangulated equivalence, where $\mathrm{Qcoh} X$ is the category of quasi-coherent sheaves of \mathcal{O}_X -modules.*

Corollary 2.27 ([BV]) *Let X be smooth over a field, then we have*

$$\mathrm{D}^b(\mathrm{coh} X) \stackrel{\Delta}{\cong} \mathrm{D}_{pf}(X).$$

where $\mathrm{coh} X$ is the category of coherent sheaves of \mathcal{O}_X -modules.

For a ring A , we denote by $\text{proj } A$ the category of finitely generated projective A -modules.

Theorem 2.28 ([Rd1], [Rd2]) *Let A, B be algebras over a field k . The following are equivalent.*

1. $D(\text{Mod } A) \stackrel{\Delta}{\cong} D(\text{Mod } B)$.
2. $K^b(\text{proj } A) \stackrel{\Delta}{\cong} K^b(\text{proj } B)$.
3. *There is a perfect complex $T \in D(\text{Mod } A)$ such that*
 - (a) $B \cong \text{End}_{D(\text{Mod } A)}(T)$,
 - (b) $\text{Hom}_{D(\text{Mod } A)}(T, T[i]) = 0$ for $i \neq 0$,
 - (c) $\{T[i] \mid i \in \mathbb{Z}\}$ is a generating set in $D(\text{Mod } A)$.
4. *There is a complex V of B - A -bimodules such that*

$$\mathbf{R}\text{Hom}_A(V, -) : D(\text{Mod } A) \rightarrow D(\text{Mod } B)$$

is an equivalence.

In this case, T is called a tilting complex for A , V is called a two-sided tilting complex, and $\mathbf{R}\text{Hom}_A(V, -)$ is called a standard equivalence.

Theorem 2.29 ([BO]) *Let X be a smooth irreducible projective variety with ample canonical or anticanonical sheaf. If X' is a smooth algebraic variety such that $D^b(\text{coh } X) \stackrel{\Delta}{\cong} D^b(\text{coh } X')$, then X' is isomorphic to X .*

Theorem 2.30 ([Be]) *Let $\mathbf{P} = \mathbf{P}_k^n$ be the n -dimensional projective space over a field k , and let $\mathcal{T}_1 = \bigoplus_{i=0}^n \mathcal{O}(i)$, $\mathcal{T}_2 = \bigoplus_{i=0}^n \Omega(-i)$, and $B_1 = \text{End}_{\mathbf{P}}(\mathcal{T}_1)$, $B_2 = \text{End}_{\mathbf{P}}(\mathcal{T}_2)$. Then B_i is a finite dimensional k -algebra of finite global dimension, and*

$$D^b(\text{coh } \mathbf{P}) \stackrel{\Delta}{\cong} D^b(\text{mod } B_i)$$

where $\text{mod } B_i$ is the category of finitely generated B_i -modules ($i = 1, 2$).

Definition 2.31 *Let A be an algebra over a field k . The derived Picard group of A (relative to k) is*

$$\text{DPic}_k(A) := \frac{\{\text{two-sided tilting complexes } T \in D^b(\text{Mod } A^e)\}}{\text{isomorphism}}$$

with identity element A , product $(T_1, T_2) \mapsto T_1 \otimes_A^L T_2$ and inverse $T \mapsto T^\vee := \mathbf{R}\text{Hom}_A(T, A)$. Given any k -linear triangulated category \mathcal{C} we let

$$\text{Out}_k^\Delta(\mathcal{C}) := \frac{\{k\text{-linear triangulated self-equivalences of } \mathcal{C}\}}{\text{isomorphism}}.$$


Theorem 2.32 ([MY]) *Let k be an algebraically closed field, and A a finite dimensional hereditary k -algebra. Then we have*

$$\mathrm{DPic}_k(A) = \mathrm{Out}_k^\Delta(\mathrm{D}^b(\mathrm{Mod} A)) = \mathrm{Out}_k^\Delta(\mathrm{D}^b(\mathrm{mod} A))$$

M. Kontsevich and A. Rosenberg introduced the notion of non-commutative projective spaces \mathbf{NP}^n [KR], and showed that

$$\begin{aligned} \mathrm{D}^b(\mathrm{Qcoh} \mathbf{NP}^n) &\cong \mathrm{D}^b(\mathrm{Mod} kQ_n) \\ \mathrm{D}^b(\mathrm{coh} \mathbf{NP}^n) &\cong \mathrm{D}^b(\mathrm{mod} kQ_n) \end{aligned}$$

where Q_n is the quiver



Corollary 2.33 ([MY]) *For non-commutative projective spaces \mathbf{NP}^n , we have*

$$\begin{aligned} \mathrm{Out}_k^\Delta(\mathrm{D}^b(\mathrm{Qcoh} \mathbf{NP}^n)) &\cong \mathrm{Out}_k^\Delta(\mathrm{D}^b(\mathrm{coh} \mathbf{NP}^n)) \\ &\cong \mathbb{Z} \times (\mathbb{Z} \ltimes \mathrm{PGL}_{n+1}(k)) \end{aligned}$$

Theorem 2.34 ([BO]) *Let X be a smooth irreducible projective variety with ample canonical or anticanonical sheaf. Then $\mathrm{Out}_k^\Delta(\mathrm{D}^b(\mathrm{coh} X))$ is generated by the automorphisms of variety, the twists by invertible sheaves and the translations, and hence $\mathrm{Out}_k^\Delta(\mathrm{D}^b(\mathrm{coh} X)) \cong (\mathrm{Aut}_k X \ltimes \mathrm{Pic} X) \times \mathbb{Z}$.*

References

- [Be] A.A. Beilinson, Coherent sheaves on \mathbf{P}^n and problems of linear algebra, *Func. Anal. Appl.* **12** (1978), 214–216.
- [BBD] A. A. Beilinson, J. Bernstein and P. Deligne, *Faisceaux Pervers*, *Astérisque* **100** (1982).
- [BN] M. Bökstedt and A. Neeman, Homotopy Limits in Triangulated Categories, *Compositio Math.* **86** (1993), 209–234.
- [BO] A. Bondal and D. Orlov, Reconstruction of a variety from the derived category and groups of autoequivalences, *Compositio Math.* **125** (2001), no. 3, 327–344.
- [BV] A. Bondal, M. Van den Bergh, Generators and representability of functors in commutative and noncommutative geometry, *math.AG/0204218*.
- [CE] H. Cartan, S. Eilenberg, “Homological Algebra,” Princeton Univ. Press, 1956.
- [Fr] J. Franke, On the Brown representability theorem for triangulated categories. *Topology* **40** (2001), no. 4, 667–680.

- [KR] M. Kontsevich and A. Rosenberg, Noncommutative smooth spaces, preprint; eprint math.AG/9812158.
- [LAM] Leovigildo Alonso Tarrío, Ana Jeremías López, María José Souto Salorio, Localization in categories of complexes and unbounded resolutions, *Canad. J. Math.* **52** (2000), no. 2, 225–247.
- [Mc] S. Mac Lane, “Homology,” Springer-Verlag, Berlin, 1963.
- [Mi] J. Miyachi, Localization of Triangulated Categories and Derived Categories, *J. Algebra* **141** (1991), 463–483.
- [MY] J. Miyachi, A. Yekutieli, Derived Picard groups of finite-dimensional hereditary algebras. *Compositio Math.* **129** (2001), no. 3, 341–368.
- [Ne] A. Neeman, The Grothendieck duality theorem via Bousfield’s techniques and Brown representability, *J. American Math. Soc.* **9** (1996), 205–236.
- [RD] R. Hartshorne, “Residues and Duality,” *Lecture Notes in Math.* **20**, Springer-Verlag, Berlin, 1966.
- [Rd1] J. Rickard, Morita Theory for Derived Categories, *J. London Math. Soc.* **39** (1989), 436–456.
- [Rd2] J. Rickard, Derived Equivalences as Derived Functors, *J. London Math. Soc.* **43** (1991), 37–48.
- [RZ] R. Rouquier and A. Zimmermann, Picard Groups for Derived Module Categories, *Proc. London Math. Soc.* (3) **87** (2003), no. 1, 197–225.
- [TT] R. W. Thomason, T. Trobaugh, Higher algebraic K -theory of schemes and of derived categories, *The Grothendieck Festschrift, Vol. III*, 247–435, *Progr. Math.*, 88, Birkhäuser Boston, Boston, MA, 1990.
- [We] C. A. Weibel, “An Introduction to Homological Algebra,” *Cambridge studies in advanced mathematics.* **38**, Cambridge Univ. Press, 1995.
- [Sp] N. Spaltenstein, Resolutions of Unbounded Complexes, *Composition Math.* **65** (1988), 121–154.
- [Ve] J. Verdier, “Catéories Déivées, état 0”, pp. 262-311, *Lecture Notes in Math.* **569**, Springer-Verlag, Berlin, 1977.