TITLE:

Fermentation Ability of Gut Microbiota of Wild Japanese Macaques in the Highland and Lowland Yakushima: In Vitro Fermentation Assay and Genetic Analyses

AUTHOR(S):

Hanya, Goro; Tackmann, Janko; Sawada, Akiko; Lee, Wanyi; Pokharel, Sanjeeta Sharma; de Castro Maciel, Valdevino Gisele; Toge, Akito; ... Hayakawa, Takashi; Shimizu, Kentaro K.; Ushida, Kazunari

CITATION:

Hanya, Goro ...[et al]. Fermentation Ability of Gut Microbiota of Wild Japanese Macaques in the Highland and Lowland Yakushima: In Vitro Fermentation Assay and Genetic Analyses. *Microbial Ecology* 2020, 80: 459-474

ISSUE DATE:

2020-8

URL:

<http://hdl.handle.net/2433/253562>

RIGHT:

This is a post-peer-review, pre-copyedit version of an article published in 'Microbial Ecology'. The final authenticated version is available online at: <https://doi.org/10.1007/s00248-020-01515-8>; The full-text file will be made open to the public on 24 April 2021 in accordance with publisher's 'Terms and Conditions for Self-Archiving'. ; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

Fermentation by gut microbe of Japanese macaques
Hanya et al.

1

1 Fermentation ability of gut microbiota of wild Japanese macaques in the
2 highland and lowland Yakushima: *in vitro* fermentation assay and genetic
3 analyses
4
5 Goro Hanya¹, Janko Tackmann², Akiko Sawada^{1,3}, Lee Wanyi¹, Sanjeeta
6 Sharma Pokharel⁴, Valdevino Gisele de Castro Maciel⁵, Akito Toge¹, Kota
7 Kuroki¹, Ryoma Otsuka⁶, Ryoma Mabuchi⁷, Jie Liu⁸, Masaomi Hatakeyama^{9,10},
8 Eri Yamasaki⁹, Christian von Mering², Rie Shimizu-Inatsugi⁹, Takashi
9 Hayakawa^{1,11,12}, Kentaro K. Shimizu^{9,13}, Kazunari Ushida³

10

- 11 ¹ Primate Research Institute, Kyoto University, Inuyama, Japan
12 ² Institute of Molecular Life Sciences, University of Zurich, Zurich, Switzerland
13 ³ Chubu University Academy of Emerging Sciences, Kasugai, Japan
14 ⁴ Centre for Ecological Sciences, Indian Institute of Science, India
15 ⁵ Amazon National Institute of Research, Manaus, Brazil
16 ⁶ Graduate School of Asian and African Area Studies, Kyoto University, Kyoto,
17 Japan
18 ⁷ Graduate School of Science, Kyoto University, Kyoto, Japan
19 ⁸ Wildlife Research Center, Kyoto University, Kyoto, Japan
20 ⁹ Department of Evolutionary Biology and Environmental Studies, University of
21 Zurich, Zurich, Switzerland
22 ¹⁰ Functional Genomics Center Zurich, Zurich, Switzerland
23 ¹¹ Faculty of Environmental Earth Science, Hokkaido University, Sapporo, Japan
24 ¹² Japan Monkey Centre, Inuyama, Japan
25 ¹³ Kihara Institute for Biological Research, Yokohama City University, Yokohama,

Fermentation by gut microbe of Japanese macaques
Hanya et al.

2

26 Japan

27

28 Correspondence to: G. Hanya: E-mail: hanya.goro.5z@kyoto-u.ac.jp, ORCID29 iD: <https://orcid.org/0000-0002-8612-659X>

30

31 Acknowledgements

32 We would like to thank our friends and colleagues in Yakushima, the fellow
33 students and staff who joined the Yakushima Field Science Course in the spring
34 of 2016, and the administrative staff who made the course possible behind the
35 scenes. Drs. Misako Yamazaki, Ayako Izuno, Catherine Aquino, Lucas Mohn,
36 Sirisha Aluri, Sayaka Tsuchida, Masayuki Suzuki, and members of the
37 Department of Evolutionary Biology and Environmental Studies, University of
38 Zurich, and Functional Genomics Center Zurich helped us with genetic and
39 SCFA analyses. The Sarugoya Committee and Wildlife Research Center of
40 Kyoto University offered us excellent facilities for fieldwork. Permission to
41 conduct this study was given by the Yakushima Forest Ecosystem Conservation
42 Center and Kagoshima Prefecture. The study was financed by the Leading
43 Graduate Program of Primatology and Wildlife Science of Kyoto University,
44 MEXT Grant-in-Aid for Promotion of Joint International Research (Fostering
45 Joint International Research) (15KK0256 and 19KK0186) and for Scientific
46 Research B (25291100 and 18H02508) to GH, and by the University Research
47 Priority Program of Evolution in Action from University of Zurich.

48

49 The nucleotide sequence data reported are available in the DDBJ database
50 under the accession numbers DRA009238 and DRA009275.

Fermentation by gut microbe of Japanese macaques
Hanya et al.

3

51

52 Author Contributions

53 Goro Hanya, Akiko Sawada, Christian von Mering, Rie Shimizu-Inatsugi,

54 Kentaro K. Shimizu, and Kazunari Ushida designed the research. Goro Hanya,

55 Akiko Sawada, Sanjeeta Sharma Pokharel, Valdevino Gisele de Castro Maciel,

56 Akito Toge, Kota Kuroki, Ryoma Otsuka, Ryoma Mabuchi, Jie Liu, Takashi

57 Hayakawa and Kazunari Ushida collected samples and performed the *in vitro*

58 fermentation assay. Goro Hanya, Akiko Sawada, Lee Wanyi, Eri Yamasaki, Rie

59 Shimizu-Inatsugi, and Takashi Hayakawa conducted the genetic analyses,

60 Janko Tackmann and Masaomi Hatakeyama analyzed the sequence data. Goro

61 Hanya wrote the manuscript.

62

Fermentation by gut microbe of Japanese macaques

Hanya et al.

4

63 Abstract

64 Wild Japanese macaques (*Macaca fuscata* Blyth) living in the highland and
65 lowland areas of Yakushima are known to have different diets, with highland
66 individuals consuming more leaves. We aim to clarify whether and how these
67 differences in diet are also reflected by gut microbial composition and
68 fermentation ability. Therefore, we conduct an *in vitro* fermentation assay using
69 fresh feces from macaques as inoculum and dry leaf powder of *Eurya japonica*
70 Thunb. as a substrate. Fermentation activity was higher for feces collected in the
71 highland, as evidenced by higher gas and butyric acid production and lower pH.
72 Genetic analysis indicated separation of highland and lowland in terms of both
73 community structure and function of the gut microbiota. Comparison of feces and
74 suspension after fermentation indicated that the community structure changed
75 during fermentation, and the change was larger for lowland samples. Analysis of
76 the 16S rRNA V3-V4 barcoding region of the gut microbiota showed that
77 community structure was clearly clustered between the two areas. Furthermore,
78 metagenomic analysis indicated separation by gene and pathway abundance
79 patterns. Two pathways (glycogen biosynthesis I and D-galacturonate
80 degradation I) were enriched in lowland samples, possibly related to the
81 fruit-eating lifestyle in the lowland. Overall, we demonstrated that the more
82 leaf-eating highland Japanese macaques harbor gut microbiota with higher leaf
83 fermentation ability compared to the more fruit-eating lowland ones. Broad,
84 non-specific taxonomic and functional gut microbiome differences suggest that
85 this pattern may be driven by a complex interplay between many taxa and
86 pathways rather than single functional traits.

Fermentation by gut microbe of Japanese macaques

Hanya et al.

5

87 Keywords: diet, *in vitro* fermentation assay, meta-16S analysis, digestion,
88 generalists

89

90 Introduction

91 There are three steps in feeding—searching, processing and digestion—and
92 animals adapt in various ways to maximize nutritional intake and/or minimize
93 intake of toxic compounds at each step. Many animal traits, including locomotion
94 [1], morphology [2], vision [3], gustation [4], and digestion enzymes [5] are, at
95 least partly, a result of this adaptation process, and some of these traits are
96 genetically fixed. While specialization to a particular food type opens up normally
97 inaccessible resources [6], such specialization may, on the other hand, limit
98 overall resource range [7]. Since environmental fluctuations can occur at much
99 shorter time scales compared to adaptive evolution, a generalist strategy may
100 require sufficient flexibility to respond to rapidly changing food conditions [8].
101 Flexibility can, for instance, be achieved by behavioral adaptation, including
102 changes in activity budget [9], ranging [10], and feeding techniques [11], which
103 have been intensively studied among large-brained, behaviorally flexible
104 generalist animals such as primates [12]. These studies mainly focused on the
105 question of how such behavioral adjustments can increase net food intake.

106 Recently, as another class of feeding-related flexible adaptations, much
107 attention has been given to the role of the gut microbiome. A typical human
108 individual harbors 10-100 trillion symbiotic microorganisms, the majority of which
109 are gut microbes [13]. Human gut microorganisms are estimated to possess 3.3
110 million non-redundant genes [14], compared to only around 22,000 genes found
111 in the human genome [15]. Host and microbial symbiont genomes often have

Fermentation by gut microbe of Japanese macaques

Hanya et al.

6

complementary roles in digestion [16]. Cellulose, xylan, and many other polysaccharides are fermented by gut microbes to produce short-chain fatty acids (SCFA), an important energy source for vertebrates [17]. In fact, energy gain through SCFAs is estimated to reach as high as 10% in humans [18] and 31% in a non-human primate species [19]. Furthermore, the genomic diversity of gut microbes among different hosts is much larger than that for the hosts themselves [13], which allows gut microbiota to respond more quickly to environmental fluctuations than their hosts. Another function of the gut microbe is detoxification, i.e. degradation of a plant's toxic and anti-nutritional compounds [20, 21].

Enabled by the development of next-generation sequencing techniques, a rapidly increasing number of studies on the flexibility of the gut microbiome have been published [22]. Many of these studies sequence hypervariable regions of the microbial 16S rRNA gene to assess the taxonomic composition of any given gut community. Community composition can change flexibly with regard to habitat [23-25], age and sex of the host [26, 27], social contact with other animals [28], and season [29-31]. Observational studies of wild populations [32] as well as feeding experiments on captive animals [33] show that the gut microbiome composition adjusts flexibly in response to changes in diet. However, data on community composition is often difficult to link directly to function because many functional genes, including those involved in digestion, are shared across diverse microbial groups [34]. As a result, even though the feeding behavior of non-human primates has been studied in detail, there are few studies, as described below, that clarified the digestive effectiveness of the gut microbiota of wild animals in a direct, straightforward experimental setting.

Fermentation by gut microbe of Japanese macaques
Hanya et al.

7

137 One way to reveal the specific gut microbiome adaptations is to study
138 the presence and/or expression patterns of functional genes through
139 metagenomic and metatranscriptomic analyses [16, 35]. Combined with
140 metabolomic analysis [36], this approach can enable researchers to track the
141 metabolic pathways involved in digesting specific foods into absorbable nutrients,
142 thus providing a deeper understanding of the role of gut microbes in these
143 processes at the molecular level. However, each food contains various nutrients
144 and toxins, and numerous microbial species with large numbers of genes are
145 involved in their digestion. Data obtained by these omics approaches are thus
146 immense and complex; consequently, extracting system-level insights on the
147 interplay between hosts and their microbial symbionts can be a considerable
148 challenge [34].

149 Another way to answer an ecologically important, simple question on
150 the effectiveness of the gut microbe in food digestion is *in vitro* fermentation
151 assay [37]. In such assays, live gut microbes contained in fresh feces of animals
152 are used as inoculum with a specific food as a substrate, which is subsequently
153 fermented under *in vitro* conditions (e.g. anaerobic and 37°C). If one type of
154 microbial community is more efficient at fermenting the substrate than others,
155 the overall fermentation activity, such as gas or SCFA production, is expected to
156 be larger. While this method is an established practice in the area of animal
157 husbandry, it is rarely used in wildlife studies. Indeed, previous work on
158 non-human primates has so far been mostly limited to zoo animals [38-41], with
159 the exception of a single study revealing the fermentation ability of plant gum by
160 the gut microbiota of wild chimpanzees [42]. These studies on captive animals
161 found that the relationship between diet and fermentation ability was complex

Fermentation by gut microbe of Japanese macaques
Hanya et al.

8

and non-specific. However, gut microbiome composition can be significantly modified under captivity [43, 44], and recent studies also highlighted the importance of study *in natura*, or in natural environments [45]. Thus, applying *in vitro* fermentation to wild animals *in natura* provides a unique opportunity to study microbiome-related fermentation responses under natural conditions.

The subject of this study, the Japanese macaque (*Macaca fuscata* Blyth), is an ideal species for studying the digestive role of the gut microbiome in a fluctuating environment, since they modify their diet in response to regional and seasonal variations in food availability [46, 47]. In our study site, Yakushima, there is an elevational gradient in fruit production [48]: while the main foods in the fruit-poor highland are fiber-rich foods, such as mature leaves [49], diet in the fruit-rich lowland features more fruits and seeds [50]. These two areas are only about 7 km apart and genetic variation between macaque populations is generally low [51]. Therefore, non-genetic, flexible adjustments are required to survive under the differing food conditions provided by these two contrasting habitats. Since macaques are habituated to human presence in many places in Yakushima [52], the collection of fresh feces is feasible. To further minimize disturbance of the fecal microbiota, *in vitro* fermentation assays can be performed directly on-site at a research station of Kyoto University.

In this study, we examined the hypothesis that the gut microbiome of a generalist host is shaped by the host's flexible feeding behavior and contributes to digestion in a food-type-specific manner. In particular, we tested the prediction that the gut microbiome of the more leaf-eating highland macaques is more capable of fermenting leaves than that of the more fruit-eating lowland macaques. We evaluated the fermentation ability by quantifying gas and SCFA

Fermentation by gut microbe of Japanese macaques

Hanya et al.

9

production in an *in vitro* fermentation assay. Furthermore, we conducted genetic analyses of the fecal samples used in the fermentation assay to better understand which microbial taxa and genes might be responsible for the observed differences in fermentation ability. To this end, we first conducted a meta-16S rRNA gene analysis to determine gut microbiome community profiles in both the highland and lowland, which allowed us to examine whether community composition significantly differs between regions and to identify differentially abundant taxa for each region. We also compared between feces and corresponding suspension samples after fermentation to reveal the changes in the community structure during the assay. Subsequently, we conducted a complementary whole-genome shotgun analysis of the fecal samples to additionally examine functional differences (i.e. in terms of gene and pathway composition). Apart from investigating broad functional differences, we also specifically compared the abundance of 37 genes known to be involved in fermentation. These genes participate in most upstream and downstream stages of the synthesis of SCFAs from polysaccharides.

Methods

Study sites

Yakushima is an island in southern Japan (30 °N, 131 °E) with an area of 503 km², with the highest peak being 1936 m in elevation. In the lowland, warm temperate evergreen broad-leaved trees, such as *Castanopsis cuspidata*, *Quercus salicina*, and *Distylium racemosum*, mix with subtropical plants. In the highland, warm-temperate evergreen broad-leaved trees, such as *Q. acuta*, *Q. salicina*, and *D. racemosum*, are interspersed with conifers (including

Fermentation by gut microbe of Japanese macaques

Hanya et al.

10

Cryptomeria japonica, *Abies firma*, and *Tsuga sieboldii*). Further details of the study site are described in Hanya et al. [48].

Study subjects

The subjects of this study, Japanese macaques, have a home range of ca 0.5-1 km² [53] and their daily path length is 2.2-2.6 km [54]. The collection sites in the highland and lowland were more than 3 km apart from each other, so the two sites were used by different groups. In May, when we collected samples for this study, the main foods were reported to be sapfruits of *Myrica rubra* in the lowland and mature leaves of *Symplocos myrtacea* and *Eurya japonica* in the highland [49, 50], which was consistent with our *ad libitum* observation at the time of collection. Neutral detergent fiber content is higher for leaves of *S. myrtaceae* (34.1%) and *E. japonica* (50.8%) than the pulp of *M. rubra* (22.4%) [55].

Sample collection

On May 22-24, 2016, we collected fresh feces of wild Japanese macaques in the eastern and western highland (730-1330 m a.s.l.; N=12), as well as the western lowland (150-190 m a.s.l.; N=15), of Yakushima (Fig. 1). Defecation was directly observed for the majority of samples, and in all other cases, feces were judged to be fresh (<1 h defecation) because a monkey group was still in the proximity and the surface of the feces showed moistness. During sampling, we first collected a small amount of fecal materials with a cotton swab and kept it in a 2 mL tube filled with RNAlater [56]. To avoid environmental contamination, these genetic samples were collected only from the inner part of the feces, which did not touch the soil. One sample was collected for genetic analysis but could not

Fermentation by gut microbe of Japanese macaques

Hanya et al.

11

be included as an inoculum of the subsequent fermentation assay. We then put all remaining fecal material in a sealed plastic bag, which was semi-vacuumized using a kitchen vacuumizer. The plastic bags were labelled and kept in a jar filled with blue ice. We recorded the sampling locations with a GPS device (GPSMAP 60CSx ®Garmin) and brought the samples to the field station of the Wildlife Research Center of Kyoto University in Nagata village within four hours of sample collection.

In vitro fermentation assay

In the laboratory, one part of the fecal samples was mixed with four parts of McDougall's buffer (ingredients (g/L): NaHCO₃ (9.8), Na₂HPO₄·2H₂O (2.44), KCl (0.57), NaCl (0.47), MgSO₄·7H₂O (0.12), CaCl₂·2H₂O (0.16)) which was pre-heated at 39°C and saturated with CO₂ gas. We confirmed the pH of the buffer before use as 6.8. We squeezed the mixture through two layers of sterile gauzes to remove large particles in the feces. We subsampled 5 mL of the filtrate, then centrifuged it at 2,600 x g for 5 min at 4°C. The supernatant was stored at -30°C for subsequent measurement of SCFA at time “zero” of the incubation. However, we mistakenly neglected to collect two samples from highland and those seven from lowland, which were therefore excluded from the SCFA measurement. As a substrate, leaves of *Eurya japonica* Thunb. were previously collected in Inuyama city, central Japan. We chose this species because the macaques in both highland and lowland eat this [49]. The leaves were dried at 40°C and then ground by a mill (Wonder Blender WB-1 ®Osaka Chemical) before use. A portion of the filtrate (100 ml) and one gram dry weight of powdered *E. japonica* leaves were poured into a serum bottle. Then, the bottles

Fermentation by gut microbe of Japanese macaques

Hanya et al.

12

were closed with butyl rubber stoppers and aluminum seals after replacing the head-space gas with 100% CO₂. Bottles were placed in a water bath at 37°C with continuous shaking for 24 h. Gas production, an indicator of fermentation activity, was measured at 6 h, 12 h, 18 h and 24 h of incubation by displacing a plunger of a 10-ml glass syringe in a manner similar to the method established by Blümmel and Ørskov [57]. After 24 h, the pH of inoculum was measured and additional samples were collected for DNA analysis of the suspension after fermentation (SAF, hereafter). Then, it was centrifuged at 2,600 x g for 5 min at room temperature, and the supernatant was collected for SCFA analysis.

SCFA analysis

Analysis of SCFA was conducted by TechnoPro R&D, Inc., Nagoya, Japan. The samples were centrifuged at 10,000 rpm for 5 min at 4°C, and the supernatant was filtered with 0.45 µm PVDF filter (Millex-HV filter, Merk Millipore). The filtrates were diluted with the same volume of ethanol. Carboxylic acids were pre-labelled with 2-nitrophenylhydrazine using the Short- and Long-Chain Fatty Acid Analysis Kit (YMC, Co. Ltd, Kyoto, Japan). The SCFA derivatives were subsequently extracted by n-hexane and diethyl ether and finally evaporated to a dry state. The residue was dissolved in methanol and filtered through a 0.45-µm membrane filter. A portion (10 µl) of filtrate was injected into an HPLC system with YMC-Pack FA column (250 × 6.0 mm; YMC Co., Ltd.). We applied the isocratic elution mode with the acetonitrile/methanol/water mobile phase (30:16:54, v/v) at 50°C. Six SCFAs (acetic, propionic, butyric, isobutyric, valeric and isovaleric acids) were measured for absorbance at 400 nm. The sum of individual SCFA concentrations was regarded as total SCFA.

Fermentation by gut microbe of Japanese macaques
Hanya et al.
13

287

288 Genetic analysis

289 Approximately one week after collecting the samples, we centrifuged the sample
290 tubes at 14000 rpm for 10 min and discarded the supernatant (RNA later
291 solution). This was carried at Primate Research Institute, Kyoto University,
292 Japan. The pelleted samples were then frozen at -80°C until extraction, except
293 at the time of export to Switzerland on September 29 (31 hours, temperature <= 4°C
294 at all times). Further analysis was conducted at the Department of
295 Evolutionary Biology and Environmental Studies, University of Zurich, and the
296 Functional Genomics Center Zurich. We added four zirconia beads (3 mm in
297 diameter), 0.1 mg zirconia/silica beads (0.1 mm in diameter), and 500 µL of
298 Inhibit Ex buffer of QIAamp Fast DNA Stool Mini Kit® QIAGEN into the tubes
299 and crushed the sample substrate with a bead crusher at 1800 rpm for 5 min.
300 DNA was extracted by QIAamp Fast DNA Stool Mini Kit following the
301 manufacturer's protocol. We quantified the DNA concentration with Qubit dsDNA
302 HS Assay Kit and a Qubit fluorometer® Thermo Fisher Scientific.

303 We conducted meta-16S analysis for both feces and SAF samples. We
304 amplified the V3-V4 region of the 16S rRNA gene with the following primer set:
305 S-D-Bact-0341-b-S-17 (forward) CCT ACG GGN GGC WGC AG and
306 S-D-Bact-0785-a-A-21 (reverse) GAC TAC HVG GGT ATC TAA TCC [58]. We
307 chose this region because it has already been used for a previous study of
308 Japanese macaques and it can represent the composition of gut microbe
309 community as well as other regions, such as V1-V2 [56]. We purified the PCR
310 amplicons using Agencourt AMPure XP beads® Beckman Coulter, Inc. and then
311 performed a second PCR (using the Illumina Nextera XT Index Kit) to attach

Fermentation by gut microbe of Japanese macaques
Hanya et al.
14

312 specific dual indices and sequencing adapters to each amplicon. DNA
313 concentration of the 2nd PCR products were quantified and pooled so that the
314 molarities of all samples were equalized. We evaluated the fragment size
315 distribution with TapeStation ®Agilent Technologies, Inc. After Phi X spike-in, the
316 library was sequenced on the Illumina Miseq platform (2 x 300 bp). Further
317 details of the protocol are described in Hayakawa et al. [56].

318 We conducted the shotgun metagenomics analysis for only feces
319 samples. We fragmented the DNA with Covaris E220 ®M&S Instruments, Inc.,
320 following the manufacturer's recommended setting to make the average DNA
321 size 200 bp (http://covarisinc.com/wp-content/uploads/pn_010308.pdf). After
322 confirming the fragment size by TapeStation, we prepared a library using Ultra™
323 II DNA Library Prep for Illumina ®New England Biolabs Inc. and sequenced it
324 with the Illumina Hiseq 4000 platform (2 x 150 bp).

325

326 Data analysis

327 The difference in gas and SCFA production between the highland and lowland
328 were tested by t-test using R 3.2.2.

329 For the 16S rRNA gene analysis, demultiplexing yielded 7,424,940
330 paired-end reads for downstream analysis (median/minimum per sample:
331 244,875/78,467). Reads were denoised with dada2 (version 1.12.1, [59]) using
332 read truncation lengths of '280, 250', maximum expected errors of 8, pooling
333 option 'pseudo,' maximum allowed mismatches when merging pairs of 3, and
334 default values for all other options. Denoising resulted in 25,518 Amplicon
335 Sequencing Variants (ASVs), which were subsequently aligned using
336 INFERNAL (version 1.1.2, [60]) with the microbial secondary structure model

Fermentation by gut microbe of Japanese macaques
Hanya et al.
15

SSU-ALIGN. Based on this alignment, ASVs were further clustered into 2,541 *de novo* OTUs (97% sequence identity) using HPC-CLUST (version 1.2.1, [61]) with the average linkage option. Taxonomic classification was performed by mapping all ASVs with MAPseq (version 1.2.1, [62]) against its default reference database (MAPref 2.2b, see [63] for details on how it was generated). ASV-specific classifications were then summarized into consensus classifications per OTU by retaining only classifications with >90% assigned representatives (MAPseq confidence cutoff ≥ 0.5).

For phylogenetic tree construction, representative ASVs with the highest abundance were chosen for each OTU. Representatives were aligned using INFERNAL (see above), and the resulting alignment was subsequently used to construct a phylogenetic tree with FastTree (version 2.1.9, [64]) using the GTR substitution model and otherwise default options. Alpha- and Beta-diversity indices were computed using the python package scikit-bio (version 0.5.1, <http://scikit-bio.org/>), and standard statistical tests (Student's t-test for independent samples, Mann-Whitney U test) were performed with the python package scipy (version 0.19.1; <https://www.scipy.org/>). Differential abundance analysis on all OTUs and taxa found in at least five samples was conducted using edgeR (version 3.24.3, [65]) with robust dispersion (function `estimate GLMRobustDisp`) and normalization factor option TMM.

For the shotgun metagenomics analysis, de-multiplexing yielded 395,051,334 paired-end reads for downstream analysis. Read pairs were joined and filtered using mothur (version 1.38.1, [66]) using the “make.contigs” and “screen.seqs” methods (the latter with options `maxambig 10`, `maxhomop 10`, `maxlength 480`), which resulted in 373,262,340 filtered reads (5.5% removed).

Fermentation by gut microbe of Japanese macaques
Hanya et al.
16

362 To remove putative eukaryotic sequences, all filtered reads were mapped
363 against KEGG (release 59, [67]) with the DIAMOND tool (version 0.8.38, [68],
364 options “sensitive” and “-b 20”) and all reads with a eukaryotic best hit were
365 discarded (16,091,001, or 4.3%).

366 Based on all remaining reads, gene and pathway abundances were
367 generated with HUMAnN2 (version 0.11.1, [69]) using the recommended
368 UniRef90 database [70] for gene definitions and the MetaCyc database [71] for
369 pathway definitions (both databases downloaded in February 2018), as well as
370 relative abundance normalization (option “relab”). Level 4 EC identifiers were
371 assigned based on HUMAnN2 utility mapping files. Standard statistical tests
372 were performed using the python package scipy (see above).

373

374 Compliance with Ethical Standards

375 During the fieldwork, we adhered to the “Guideline for field research of
376 non-human primates” of the Primate Research Institute, Kyoto University.
377 Furthermore, our procedure complied with ARRIVE guidelines for the use of
378 animals in research (<http://www.nc3rs.org.uk/ARRIVE>), as well as the legal
379 requirements of Japan. No prior consent from the Japanese government is
380 required to export biological samples from Japan in the context of Convention on
381 Biological Diversity (<http://www.env.go.jp/en/nature/biodiv/abs/index.html>).

382

383 Results

384 *In vitro* fermentation assay

385 We found the overall fermentation activity to be higher for fecal samples from the
386 highland compared to the lowland, as evident from both significantly higher gas

Fermentation by gut microbe of Japanese macaques Hanya et al. 17

production (6 h, 12 h, 18 h, and 24 h; Fig. 2a) and significantly lower pH (Fig. 2b). There was no significant difference in total SCFA production between the highland and lowland (Fig. 2c). Among the four SCFAs (acetic acid, propionic acid, butyric+isobutyric acids, and valeric+iso valeric acids), only the production of butyric acid (including isobutyric acid) was significantly higher for highland samples compared to lowland (Fig. 3). Prior to fermentation (time point 0 h), we found no significant difference in total SCFA concentration or for any of the four SCFA classes.

Comparisons of community structure of the fecal microbiota We obtained 7,424,940 reads (median/minimum per sample: 244,875/78,467) which were successfully demultiplexed, filtered and *de novo* clustered into 2541 operational taxonomic units (OTUs) using a 97% sequence identity cutoff (see Methods). Out of these OTUs, 1311 (51.6%) were classifiable at the phylum level or deeper. A summary of the phylogenetic assignment of the OTUs in each sample is shown in Supplemental Material 3.

The differences in relative abundance of major taxa (>1% at the phylum, class, order, family and genus levels) were all insignificant, except for significant enrichment of Anaeroplasmataceae in the highland (Supplemental Material 3, 4). Differential abundance analysis revealed 130 OTUs that were significantly enriched in highland (54) or lowland (76) samples. Enriched bacterial OTUs in the highlands were 12 OTUs of phylum Firmicutes, class Clostridia, order Clostridiales, one OUT of *Bacteroides uniformis* in the phylum Bacteroidetes, family Bacteroidaceae and the remaining 39 OTUs could not be mapped to any taxa. Enriched OTUs in the lowland included 16 OTUs of Bacteroidetes (all were

Fermentation by gut microbe of Japanese macaques Hanya et al. 18

412 class Bacteroidia, order Bacteroidales), eight OTUs of Firmicutes (order
413 Clostridia and Negativicutes), two OTU of Proteobacteria, 23 OTUs of
414 Proteobacteria (all were *Treponema berlinense*) and 27 unmapped OTUs.

415 Average alpha diversity did not differ between the lowland and highland
416 as differences in observed OTUs: Chao1, Shannon and Faith's Phylogenetic
417 Diversity indices were all insignificant (Supplemental Material 1).

418 Community structure clearly differed between the highland and lowland,
419 based on pairwise Bray-Curtis dissimilarity (PERMANOVA, $P = 0.001$). In
420 addition to clustering separately from lowland samples, highland samples
421 furthermore showed increased variability and sub-structure (Fig. 4). This
422 sub-structure could not be explained by longitudinal differences, since eastern
423 and western samples did not cluster separately.

424
425 Comparisons of community structure between the feces and suspension after
426 fermentation (SAF)
427 Community structure of the SAF was significantly different from the
428 corresponding fecal samples (Fig. 4, PERMANOVA, highland: pseudo- $F=1.86$,
429 $P=0.022$; lowland: pseudo- $F=3.06$, $P=0.001$). Judging from the F and P values,
430 the difference in community structure between SAF and feces in the lowland was
431 larger than that in the highland.

432 The number of taxa showing significant changes in abundance
433 between SAF and feces was larger in the lowland (three phyla, four classes, six
434 orders, eight families, two genera and 104 OTUs) than in the highland (one
435 phylum, no class, one order, two families, one genus and 47 OTUs) (Table 1).
436 Among the major taxa ($>1\%$ of relative abundance in feces or SAF), significantly

Fermentation by gut microbe of Japanese macaques Hanya et al. 19

437 more enrichment in SAF than in feces was detected for Proteobacteria (highland
438 and lowland) and Spirochaetes at the phylum level and Streptococcaceae
439 (highland and lowland), Succinivibrionaceae (lowland), and Pasteurellaceae
440 (lowland) at the family level and *Streptococcus* (highland and lowland) at the
441 genus level.

442 None of the alpha diversity indices differed significantly between feces
443 and SAF in either the highland or the lowland (Supplemental Material 1).

444

445 Metagenomic analysis of the fecal microbiota

446 After having assessed the community composition, we next focused on
447 systematic and specific functional (gene and pathway) differences between
448 highland and lowland microbiota. To this end, we conducted whole-genome
449 shotgun sequencing on all fecal samples, which yielded 373,262,340 reads, or
450 14,356,243 reads/sample, after de-multiplexing and filtering. PERMANOVA
451 analysis (based on pairwise Bray-Curtis dissimilarity) showed a statistically
452 significant separation between lowland and highland samples in terms of both
453 gene abundance ($P = 0.001$; Fig. 5a) and pathway abundance ($P = 0.034$; Fig.
454 5b). In contrast to abundance, pathway coverage showed no significant
455 lowland-highland separation ($P = 0.11$; Fig. 5c).

456 Among 37 selected genes involved in the metabolism of
457 polysaccharides and the synthesis of SCFAs from pyruvate, the abundance was
458 not significantly different for any of the genes.

459 When mapping the presence and absence of the selected genes on
460 polysaccharide metabolism and SCFA synthesis pathways, most genes were
461 uniformly present or uniformly absent in both the lowland and highland samples

Fermentation by gut microbe of Japanese macaques
Hanya et al.
20

(Supplemental Material 2). Regarding polysaccharide metabolism, most gut microbiota possessed genes to catalyze cellulose, xylan, mannan, and pectin degradation, all of which are lacking in the genome of rhesus macaques (the most closely related species of Japanese macaques with a published genome). Pathways for the synthesis of acetate, butyrate, and propionate from pyruvate were all inter-connected for most fecal metagenomes, both in the lowland and highland, and also within the macaque reference genome.

Out of 263 tested MetaCyc [71] pathways (present in > 10 samples), two showed a significant difference in abundance between highland and lowland samples: GLYCOGENSYNTH-PWY (glycogen biosynthesis I, from ADP-D-Glucose); Mann-Whitney U test, $P = 0.041$, Benjamini-Hochberg adjusted) and GALACTUROCAT-PWY (D-galacturonate degradation I; Mann-Whitney U test, $P = 0.044$, Benjamini-Hochberg adjusted), both of which were more abundant in lowland.

Discussion

Difference in fermentation ability between highland and lowland

The *in vitro* fermentation assay clearly indicated that the fermentation ability of leaves was higher for highland gut microbiota compared to lowland communities. In the highland, where the macaque diet is leaf-based, the corresponding communities produced significantly more gas and induced a significantly larger decrease in pH during fermentation. While they also showed a trend toward increased SCFA production, this difference was not significant. We note, however, that this may be caused by power issues due to decreased sample size compared to the gas and pH measurements (nine samples were not

Fermentation by gut microbe of Japanese macaques
Hanya et al.
21

included, see Methods). SCFAs are utilized as an energy source by the host, and they constitutes a considerable portion in both human and non-human primates' energy intake [18, 19]. Taken together, our results indicate that the higher leaf-fermentation potential of highland gut communities may facilitate digestion of the leaf-based diet of highland macaques, compared to the lowland animals. While more research is necessary, this finding highlights the potential importance of the gut microbiome for generalist hosts (such as the macaques studied here) to flexibly respond to and benefit from changing food conditions. Interference experiments on humans show that dietary changes can modify the gut microbiome within only a few days (2-5) [72], so the gut microbiota of macaques in the highland and lowland, living separately for years (in the case of philopatric females) or months (in the case of immigrant males), likely have sufficient transition time to respond to different dietary regimes.

The assay also suggested that the fermentation system was different between the two areas. When investigating individual classes of SCFAs, we found butyric acid production to be significantly increased in highland compared to lowland samples. Many bacteria are capable of acetic acid production, but propionic or butyric acid producers are generally less common [34]. Furthermore, biosynthesis pathways for butyric acid are highly diverse and can often vary even within a single bacterial family [34]. Some butyrate producers can produce butyrate with the aid of other bacteria. For example, lactate is produced by other species, and the resultant lactate is converted to butyrate by lactate users, such as *Megasphaera* [73]. Increased butyric acid production has been reported in many different contexts: for example, butyric acid in feces has been reported to be higher when energy intake decreased in howler monkeys [31], while butyric

Fermentation by gut microbe of Japanese macaques

Hanya et al.

22

acid production increased with higher fruit consumption in gorillas [74]. For human hosts, the situation is generally mixed, since some interference experiments in which additional fiber was added to the normal diet led to increased fecal butyrate concentration, while other studies did not report such increases [75]. Our finding of increased butyric acid production in highland microbiota (exposed to leaf-based diets) contributes to this growing body of research and, while the overall picture is still complex, provides additional evidence for a link between high-fiber diets and the production of this SCFA.

Meta-16S analysis: Gut microbiota community structure

Gut community structure was clearly clustered between the highland and lowland populations in Yakushima. A previous study using the meta-16S analysis of wild and captive Japanese macaques included samples collected in the highland and lowland Yakushima (N=18), which also indicated clear differentiation of gut microbe community structure between the two areas [44] (Supplemental Materials 5). We did not directly compare those results with the current study because the methods of that previous study were not the same as the method used in the current work, nevertheless that study provided further confirmation of the validity of our findings. While we observed higher OTU profile variation and considerable sub-structure within highland samples compared to lowland ones, this sub-clustering did not correspond to longitudinal differences in sampling locations (i.e. east vs. west). This is interesting because western highland macaques live in the immediate geographical vicinity of the lowland population (in contrast to the eastern highland population), but this co-locality is not reflected by a higher similarity in the gut microbe community. It thus seems

Fermentation by gut microbe of Japanese macaques
Hanya et al.
23

537 that habitat similarity, rather than physical contact between hosts (typically via
538 male immigration and emigration) [76], is responsible for shaping the gut
539 communities. Unmeasured factors related to group identity may provide an
540 alternative explanation for the sub-structure within highland samples, but more
541 data would be needed for definite conclusions.

542 Comparisons between feces and suspension after fermentation (SAF)
543 indicated that the community structure of the microbiota was significantly altered
544 during fermentation than it used to be in the gut. This alteration seems to be
545 larger for lowland than highland, evidenced by the F and P statistics and the
546 larger number of taxa showing changes in abundance. This suggests that the
547 difference between the experimental and original conditions in the gut was larger
548 for lowland than for highland microbiota, which is consistent with the fruit-based
549 diet in the lowland.

550 The comparisons of differential abundance between the feces and SAF
551 may give us a clue to the bacterial taxa that are responsible for the fermentation
552 activity. For example, Succinivibrionaceae (phylum Proteobacteria, class
553 Gammaproteobacteria, order Aeromonadales) increased during fermentation.
554 This family is reported to increase in the feces of wild Tibetan macaques during
555 winter, when they eat more leaves [29]. Interestingly, however, in the rumen of
556 cows, this family increased when the animals are supplied with low-fiber diet [77].
557 Another example is *Streptococcus* (phylum Firmicutes, class Bacilli, order
558 Lactobacillales, family Streptococcaceae). This genus includes the species with
559 tanning-binding ability [20, 21], which may be useful for digesting leaves. Feces
560 of feral pigs with high feed efficiency, and thus with a higher ability to degrade
561 cellulose, contain more *Streptococcus* [78]. These studies are consistent with

Fermentation by gut microbe of Japanese macaques
Hanya et al.
24

our results that this genus increased during the assay using mature leaves as a substrate. However, in wild howler monkeys, this genus increased when they ate low-fiber foods (fruits) [31]. The *Succinivibrionaceae* and *Streptococcus* examples show that the same taxa may exhibit a different function in fermentation in different hosts. We note, however, that even one species of microorganisms has a wide variety of genes and thus metabolic potential [79] and, therefore, we can expect the deviations at the higher taxonomic levels. Furthermore, it has been shown for different environments that stable community function can be retained even under high species turnover [80, 81]. For the same trait—especially if it features many species and pathways, such as in fiber digestion—taxonomic composition may therefore differ widely. Our results further underscore this notion and caution against generalized assumptions on community function that are based solely on taxonomy.

In contrast to a recent study on fecal microbiota of the same host species [56], a large fraction of OTUs (48.4%) could not be taxonomically classified even at the phylum level in our analysis. This difference is due to the more precise mapping approach we used here (MAPseq; [62]), which assigns low confidence to a read if multiple taxa have similar alignment scores and can thus not be confidently distinguished. In various benchmarks, this approach was shown to yield better classifications than the commonly used less conservative alternatives [62, 82]. The high fraction of largely unclassified OTUs may be an indication that much is still to be learned about the macaque gut microbiome.

Metagenomic analysis: Difference with respect to the genes and pathway composition

Fermentation by gut microbe of Japanese macaques
Hanya et al.
25

Metagenomic analysis indicated that highland and lowland populations also differed on the level of genes and pathways. This shows that the function of the gut microbe varies between the two areas, in which macaques experience different food and thermal conditions [46, 83].

For example, we found two pathways that differed in abundance between highland and lowland samples, which may be related to the lifestyle of the host animals. For example, the glycogen biosynthesis I pathway was more abundant in the lowland samples. Glycogen is synthesized to store excess glucose temporarily in the liver [84]. The main food in the sampled season in the lowland of Yakushima is the sapfruit of *Myrica rubra* [85], which is considered one of the most sugar-rich fruits available in Yakushima [86]. Therefore, lowland macaques may experience situations of having excess glucose more often than leaf-eating highland macaques. Genes in this pathway could thus be beneficial for storing excess glucose and using it as an energy source in the subsequent food-scarce season. In contrast, in the highland, saving the function of the pathway may be advantageous when the macaques do not eat fruits. The D-galacturonate degradation I pathway was also enriched in the lowland; D-galacturonate is an oxidized form of galactose, and it is included in many polysaccharides in plants, including pectin [87]. This pathway may also be related to the fruit-dominated diet in the lowland, but this remains unknown without detailed knowledge of the metabolomics profile of their foods.

We did not find a genetic difference which can directly account for the observed differences in fermentation ability between highland and lowland. Among the 37 genes encoding enzymes that catalyze the majority of upstream and downstream reactions in the degradation of polysaccharides to SCFAs, we

Fermentation by gut microbe of Japanese macaques

Hanya et al.

26

found no difference in enrichment patterns for any of the genes. It was evident that the gut microbiota in both regions possess genes necessary for the synthesis of SCFAs from polysaccharides but are lacking in the host genome (assuming it is the same as that of rhesus macaques). Although not as important as in the highland, leaves are also one of the main foods in the lowland [50], and all of these genes may be necessary for survival in both areas.

Implications of the flexibility in digestive ability

In this study, we demonstrated that the gut microbiome of macaques is shaped according to habitat and diet, in terms of both community structure (taxa, OTUs) and the function (genes, pathways). Furthermore, we showed that these differences affect the ability to digest leaves and thus may play an important adaptive role in this generalist host. Since similar alterations in the gut microbial community composition in response to dietary changes are a well-known fact in many hosts [22], we predict similar microbiome-mediated increases in digestive ability for commonly consumed foods in many other generalist animals.

We can point out a number of limitations in the interpretations of the current study. The inoculum of this assay is feces, not gut contents. Even though fecal samples are usually used as a representative of 'gut' (colon) microbiota, these microbiota are subject to changes immediately after defecation due to biotic and abiotic factors [88]. We found no difference in the fecal SCFA, or 0 h concentrations between the highland and lowland. It is possible that this was simply due to the smaller sample size than gas production reduced by mistake, but it is also possible that the difference was a real one. Fecal SCFA concentration is determined not only by the microbiota but also by the digesta,

Fermentation by gut microbe of Japanese macaques

Hanya et al.

27

which are derived from many different kinds of foods in the case of wild animals. Some bacteria in feces in the lowland may have worked to produce as much SCFAs *in vivo* as in the highland in the presence of natural foods, but not when supplied with only leaves of *E. japonica*. Therefore, different substrates should be tested in the future to confirm our conclusion.

The above limitations of the current *in vitro* fermentation assay suggest that further improvement of this method is possible as a way to reveal the fermentation ability of the gut microbe. In this study, we examined only one type of substrate, but it is possible to divide the fecal samples into multiple portions and use these as an inoculum for different substrate types. Similarly, we investigated leaf-fermentation ability only for one plant species eaten frequently in the highland but only rarely in the lowland; consequently, it would also be interesting to test the fermentation of substrates commonly eaten in the lowland, since these may be processed more efficiently by microbial communities from lowland hosts. In our assay, the fermentation time was set to 24 h, which approximates the typical retention time in the hindgut of Japanese macaques (total gut retention time can be up to 35 h when feeding on high-fiber foods; [89]). However, retention time can vary based on different factors, including the body size of the animal, and this should be taken into consideration in future studies. Furthermore, our time-series data on gas production suggest that the fermentation process is time-dependent, so it may be possible to mimic the *in vivo* digestion process much better via a more dynamic and flexible experimental setup, including substrate preparation and the duration of fermentation. In future studies, it will also be necessary to evaluate the relative contribution of the fermentation by the gut microbiome compared to the enzymatic digestion by the

Fermentation by gut microbe of Japanese macaques

Hanya et al.

28

host. This is necessary to better understand how gut microbial shifts can affect the digestion and survival of the host. Another interesting avenue of study would be to complement the metagenomics data used here with approaches based on metabolomics [36, 74] and metatranscriptomics [16, 90] approaches, which could provide deeper insights into the molecular basis and mechanisms affecting the fermentation potential. It would also be interesting to isolate bacteria species from the feces and to explore its function, for example, by sequencing the whole genome. We have already isolated and sequenced the genome of *Sarcina ventriculi* from the feces of Japanese macaques in the highland [91]. The genome includes potentially important genes for the digestion of leaves, such as cyanate metabolism. Interestingly, this bacteria cannot be isolated from feces collected in the lowland. Using combinations of these multi-level approaches would clarify the mechanism that produces different fermentation abilities among the host animals living in different environments.

In conclusion, we demonstrated that gut microbe community structure differed considerably between the two investigated habitats, which may in part be explained by habitat-specific diets. In support of this, via an *in vitro* fermentation assay, we showed that gut microbiota of the more leaf-eating highland Japanese macaques have a higher fermentation ability for leaves than those of the more fruit-eating lowland animals. Taken together, our results indicate that gut microbiota may help generalist hosts to improve their digestive ability in response to the variations on food availability.

Conflict of Interest: The authors declare that they have no conflict of interest.

Fermentation by gut microbe of Japanese macaques Hanya et al. 29

687 Ethical approval: During the fieldwork, we adhered to the “Guideline for field
688 research of non-human primates” of the Primate Research Institute, Kyoto
689 University. According to the guideline, our study is purely non-invasive and does
690 not need approval from an ethical committee.

691

692 References

- 693 1. Vogel ER, Haag L, Mitra-Setia T, van Schaik CP, Dominy NJ (2009)
694 Foraging and ranging behavior during a fallback episode: *Hylobates*
695 *albibarbis* and *Pongo pygmaeus wurmbii* compared. Am J Phys Anthropol
696 140: 716-726. doi: 10.1002/ajpa.21119
- 697 2. Ledevin R, Koyabu D (2019) Patterns and Constraints of Craniofacial
698 Variation in Colobine Monkeys: Disentangling the Effects of Phylogeny,
699 Allometry and Diet. Evol Biol 46: 14-34. doi:
700 10.1007/s11692-019-09469-7
- 701 3. Kries K, Barros MAS, Duytschaever G, Orkin JD, Janiak MC, Pessoa
702 DMA, Melin AD (2018) Colour vision variation in leaf-nosed bats
703 (Phyllostomidae): Links to cave roosting and dietary specialization. Mol
704 Ecol 27: 3627-3640. doi: 10.1111/mec.14818
- 705 4. Hayakawa T, Suzuki-Hashido N, Matsui A, Go Y (2014) Frequent
706 Expansions of the Bitter Taste Receptor Gene Repertoire during Evolution
707 of Mammals in the Euarchontoglires Clade. Mol Biol Evol 31: 2018-2031.
708 doi: 10.1093/molbev/msu144
- 709 5. Pajic P, Pavlidis P, Dean K, Neznanova L, Romano RA, Garneau D,
710 Daugherty E, Globig A, Ruhl S, Gokcumen O (2019) Independent
711 amylase gene copy number bursts correlate with dietary preferences in

Fermentation by gut microbe of Japanese macaques Hanya et al. 30

- 712 mammals. eLife 8: 22. doi: 10.7554/eLife.44628
- 713 6. Toju H, Sota T (2006) Phylogeography and the geographic cline in the
714 armament of a seed-predatory weevil: effects of historical events vs.
715 natural selection from the host plant. Mol Ecol 15: 4161-4173.
- 716 7. Grant PR, Grant BR (2002) Unpredictable evolution in a 30-year study of
717 Darwin's finches. Science 296: 707-711. doi: 10.1126/science.1070315
- 718 8. Hanya G, Bernard H (2013) Functional response to fruiting seasonality by
719 a primate seed predator, red leaf monkey (*Presbytis rubicunda*). Trop
720 Ecol 54: 383-395.
- 721 9. Hanya G (2004) Seasonal variations in the activity budget of Japanese
722 macaques in the coniferous forest of Yakushima: Effects of food and
723 temperature. Am J Primatol 63: 165-177.
- 724 10. Hanya G, Bernard H (2016) Seasonally consistent small home range and
725 long ranging distance in *Presbytis rubicunda* in Danum Valley, Borneo. Int
726 J Primatol 37: 390-404.
- 727 11. Schuppli C, Forss SIF, Meulman EJM, Zweifel N, Lee KC, Rukmana E,
728 Vogel ER, van Noordwijk MA, van Schaik CP (2016) Development of
729 foraging skills in two orangutan populations: needing to learn or needing
730 to grow? Front Zool 13: 17. doi: 10.1186/s12983-016-0178-5
- 731 12. Hemingway C, Bynum N (2005) The influence of seasonality on primate
732 diet and ranging. In: Brockman, DK, van Schaik, CP (eds.) Seasonality in
733 Primates: Studies of Living and Extinct Human and Non-Human Primates.
734 Cambridge University Press, Cambridge, pp. 57-104
- 735 13. Ursell LK, Metcalf JL, Parfrey LW, Knight R (2012) Defining the human
736 microbiome. Nutr Rev 70: S38-S44. doi:

Fermentation by gut microbe of Japanese macaques
Hanya et al.
31

- 737 10.1111/j.1753-4887.2012.00493.x
- 738 14. Qin JJ, Li RQ, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen
739 T, Pons N, Levenez F, Yamada T, Mende DR, Li JH, Xu JM, Li SC, Li DF,
740 Cao JJ, Wang B, Liang HQ, Zheng HS, Xie YL, Tap J, Lepage P, Bertalan
741 M, Batto JM, Hansen T, Le Paslier D, Linneberg A, Nielsen HB, Pelletier E,
742 Renault P, Sicheritz-Ponten T, Turner K, Zhu HM, Yu C, Li ST, Jian M,
743 Zhou Y, Li YR, Zhang XQ, Li SG, Qin N, Yang HM, Wang J, Brunak S,
744 Dore J, Guarner F, Kristiansen K, Pedersen O, Parkhill J, Weissenbach J,
745 Bork P, Ehrlich SD, Wang J (2010) A human gut microbial gene catalogue
746 established by metagenomic sequencing. *Nature* 464: 59-U70. doi:
747 10.1038/nature08821
- 748 15. Collins FS, Lander ES, Rogers J, Waterston RH, Int Human Genome
749 Sequencing C (2004) Finishing the euchromatic sequence of the human
750 genome. *Nature* 431: 931-945. doi: 10.1038/nature03001
- 751 16. Scully ED, Geib SM, Carlson JE, Tien M, McKenna D, Hoover K (2014)
752 Functional genomics and microbiome profiling of the Asian longhorned
753 beetle (*Anoplophora glabripennis*) reveal insights into the digestive
754 physiology and nutritional ecology of wood feeding beetles. *BMC*
755 *Genomics* 15: 21. doi: 10.1186/1471-2164-15-1096
- 756 17. Cummings JH, Rombeau JL, Sakata T (1995) Physiological and clinical
757 aspects of short chain fatty acids. Cambridge University Press,
758 Cambridge
- 759 18. McNeil NI (1984) The contribution of the large-intestine to energy supplies
760 in man. *Am J Clin Nutr* 39: 338-342.
- 761 19. Milton K, McBee RH (1983) Rates of fermentative digestion in the howler

Fermentation by gut microbe of Japanese macaques Hanya et al. 32

- 762 monkey, *Alouatta palliata* (Primates, Ceboidea). Comp Biochem Physiol
763 A-Physiol 74: 29-31. doi: 10.1016/0300-9629(83)90706-5
- 764 20. Osawa RO (1990) Formation of a clear zone on tannin-treated brain heart
765 infusion agar by a *Streptococcus* sp isolated from feces of koalas. Appl
766 Environ Microbiol 56: 829-831.
- 767 21. Tsuchida S, Murata K, Ohkuma M, Ushida K (2017) Isolation of
768 *Streptococcus gallolyticus* with very high degradability of condensed
769 tannins from feces of the wild Japanese rock ptarmigans on Mt. Tateyama.
770 J Gen Appl Microbiol 63: 195-198. doi: 10.2323/jgam.2016.09.003
- 771 22. Clayton JB, Gomez A, Amato K, Knights D, Travis DA, Blekhman R,
772 Knight R, Leigh S, Stumpf R, Wolf T, Glander KE, Cabana F, Johnson TJ
773 (2018) The gut microbiome of nonhuman primates: Lessons in ecology
774 and evolution. Am J Primatol 80: 27. doi: 10.1002/ajp.22867
- 775 23. Amato KR, Martinez-Mota R, Righini N, Raguet-Schofield M, Corcione FP,
776 Marini E, Humphrey G, Gogul G, Gaffney J, Lovelace E, Williams L, Luong
777 A, Dominguez-Bello MG, Stumpf RM, White B, Nelson KE, Knight R,
778 Leigh SR (2016) Phylogenetic and ecological factors impact the gut
779 microbiota of two Neotropical primate species. Oecologia 180: 717-733.
780 doi: 10.1007/s00442-015-3507-z
- 781 24. Zhao JS, Yao YF, Li DY, Xu HM, Wu JY, Wen AX, Xie M, Ni QY, Zhang
782 MW, Peng GN, Xu HL (2018) Characterization of the gut microbiota in six
783 geographical populations of Chinese rhesus macaques (*Macaca mulatta*),
784 implying an adaptation to high-altitude environment. Microbial Ecology
785 76: 565-577. doi: 10.1007/s00248-018-1146-8
- 786 25. Hayakawa T, Nathan S, Stark DJ, Saldivar DAR, Sipangkui R, Goossens

Fermentation by gut microbe of Japanese macaques Hanya et al. 33

- 787 B, Tuuga A, Clauss M, Sawada A, Fukuda S, Imai H, Matsuda I (2018)
788 First report of foregut microbial community in proboscis monkeys: are
789 diverse forests a reservoir for diverse microbiomes? Environ Microbiol
790 Rep 10: 655-662. doi: 10.1111/1758-2229.12677
- 791 26. Amato KR, Leigh SR, Kent A, Mackie RI, Yeoman CJ, Stumpf RM, Wilson
792 BA, Nelson KE, White BA, Garber PA (2014) The Role of Gut Microbes in
793 Satisfying the Nutritional Demands of Adult and Juvenile Wild, Black
794 Howler Monkeys (*Alouatta pigra*). Am J Phys Anthropol 155: 652-664. doi:
795 10.1002/ajpa.22621
- 796 27. Springer A, Fichtel C, Al-Ghalith GA, Koch F, Amato KR, Clayton JB,
797 Knights D, Kappeler PM (2017) Patterns of seasonality and group
798 membership characterize the gut microbiota in a longitudinal study of wild
799 Verreaux's sifakas (*Propithecus verreauxi*). Ecol Evol 7: 5732-5745. doi:
800 10.1002/ece3.3148
- 801 28. Raulo A, Ruokolainen L, Lane A, Amato K, Knight R, Leigh S, Stumpf R,
802 White B, Nelson KE, Baden AL, Tecot SR (2018) Social behaviour and
803 gut microbiota in red-bellied lemurs (*Eulemur rubriventer*): In search of
804 the role of immunity in the evolution of sociality. J Anim Ecol 87: 388-399.
805 doi: 10.1111/1365-2656.12781
- 806 29. Sun BH, Wang X, Bernstein S, Huffman MA, Xia DP, Gu ZY, Chen R,
807 Sheeran LK, Wagner RS, Li JH (2016) Marked variation between winter
808 and spring gut microbiota in free-ranging Tibetan Macaques (*Macaca*
809 *thibetana*). Sci Rep 6: 8. doi: 10.1038/srep26035
- 810 30. Gomez A, Rothman JM, Petrzalkova K, Yeoman CJ, Vlckova K, Umana
811 JD, Carr M, Modry D, Todd A, Torralba M, Nelson KE, Stumpf RM, Wilson

Fermentation by gut microbe of Japanese macaques Hanya et al. 34

- 812 BA, Blekhman R, White BA, Leigh SR (2016) Temporal variation selects
813 for diet-microbe co-metabolic traits in the gut of *Gorilla* spp. *Isme J* 10:
814 514-526. doi: 10.1038/ismej.2015.146
- 815 31. Amato KR, Leigh SR, Kent A, Mackie RI, Yeoman CJ, Stumpf RM, Wilson
816 BA, Nelson KE, White BA, Garber PA (2015) The Gut Microbiota Appears
817 to Compensate for Seasonal Diet Variation in the Wild Black Howler
818 Monkey (*Alouatta pigra*). *Microbial Ecology* 69: 434-443. doi:
819 10.1007/s00248-014-0554-7
- 820 32. Mallott EK, Amato KR, Garber PA, Malhi RS (2018) Influence of fruit and
821 invertebrate consumption on the gut microbiota of wild white-faced
822 capuchins (*Cebus capucinus*). *Am J Phys Anthropol* 165: 576-588. doi:
823 10.1002/ajpa.23395
- 824 33. Hao YT, Wu SG, Xiong F, Tran NT, Jakovlic I, Zou H, Li WX, Wang GT
825 (2017) Succession and Fermentation Products of Grass Carp
826 (*Ctenopharyngodon idellus*) Hindgut Microbiota in Response to an
827 Extreme Dietary Shift. *Front Microbiol* 8: 12. doi:
828 10.3389/fmicb.2017.01585
- 829 34. Rowland I, Gibson G, Heinken A, Scott K, Swann J, Thiele I, Tuohy K
830 (2018) Gut microbiota functions: metabolism of nutrients and other food
831 components. *Eur J Nutr* 57: 1-24. doi: 10.1007/s00394-017-1445-8
- 832 35. Sanders JG, Beichman AC, Roman J, Scott JJ, Emerson D, McCarthy JJ,
833 Girguis PR (2015) Baleen whales host a unique gut microbiome with
834 similarities to both carnivores and herbivores. *Nature Communications* 6:
835 8. doi: 10.1038/ncomms9285
- 836 36. Amato KR, Ulanov A, Ju KS, Garber PA (2017) Metabolomic data suggest

Fermentation by gut microbe of Japanese macaques Hanya et al. 35

- 837 regulation of black howler monkey (*Alouatta pigra*) diet composition at the
838 molecular level. Am J Primatol 79: 10. doi: 10.1002/ajp.22616
- 839 37. Tilley JMA, Terry RA (1963) A two-stage technique for the in vitro
840 digestion of forage crops. Jour Brit Grassland Soc 18: 104-111.
- 841 38. Schmidt DA, Kerley MS, Dempsey JL, Porton IJ, Porter JH, Griffin ME,
842 Eilersieck MR, Sadler WC (2005) Fiber digestibility by the orangutan
843 (*Pongo abelii*): In vitro and in vivo. J Zoo Wildl Med 36: 571-580. doi:
844 10.1638/04-103.1
- 845 39. Kišidayová S, Váradyová Z, Pristaš P, Piknová M, Nigutová K,
846 Petrzelková KJ, Profousová I, Schovancová K, Kamler J, Modry D (2009)
847 Effects of high- and low-fiber diets on fecal fermentation and fecal
848 microbial populations of captive chimpanzees. Am J Primatol 71: 548-557.
849 doi: 10.1002/ajp.20687
- 850 40. Lambert JE, Fellner V (2012) *In vitro* fermentation of dietary
851 carbohydrates consumed by African apes and monkeys: preliminary
852 results for interpreting microbial and digestive strategy. Int J Primatol 33:
853 263-281. doi: 10.1007/s10764-011-9559-y
- 854 41. Campbell JL, Williams CV, Eisemann JH (2002) Fecal inoculum can be
855 used to determine the rate and extent of in vitro fermentation of dietary
856 fiber sources across three lemur species that differ in dietary profile:
857 *Varecia variegata*, *Eulemur fulvus* and *Hapalemur griseus*. J Nutr 132:
858 3073-3080.
- 859 42. Ushida K, Fujita S, Ohashi G (2006) Nutritional significance of the
860 selective ingestion of *Albizia zygia* gum exudate by wild chimpanzees in
861 Bossou, Guinea. Am J Primatol 68: 143-151. doi: 10.1002/ajp.20212

Fermentation by gut microbe of Japanese macaques Hanya et al. 36

- 862 43. Clayton JB, Vangay P, Huang H, Ward T, Hillmann BM, Al-Ghalith GA,
863 Travis DA, Long HT, Tuan BV, Minh VV, Cabana F, Nadler T, Toddes B,
864 Murphy T, Glander KE, Johnson TJ, Knights D (2016) Captivity
865 humanizes the primate microbiome. Proc Natl Acad Sci U S A 113:
866 10376-10381. doi: 10.1073/pnas.1521835113
- 867 44. Lee W, Hayakawa T, Kiyono M, Yamabata N, Hanya G (2019) Gut
868 microbiota composition of Japanese macaques associates with extent of
869 human encroachment. Am J Primatol 81: e23072.
- 870 45. Shimizu KK, Kudoh H, Kobayashi MJ (2011) Plant sexual reproduction
871 during climate change: gene function in natura studied by ecological and
872 evolutionary systems biology. Ann Bot 108: 777-787. doi:
873 10.1093/aob/mcr180
- 874 46. Hanya G, Noma N, Agetsuma N (2003) Altitudinal and seasonal variations
875 in the diet of Japanese macaques in Yakushima. Primates 44: 51-59.
- 876 47. Agetsuma N, Nakagawa N (1998) Effects of habitat differences on
877 feeding behaviors of Japanese monkeys: Comparison between
878 Yakushima and Kinkazan. Primates 39: 275-289.
- 879 48. Hanya G, Yoshihiro S, Zamma K, Matsubara H, Ohtake M, Kubo R, Noma
880 N, Agetsuma N, Takahata Y (2004) Environmental determinants of the
881 altitudinal variations in relative group densities of Japanese macaques on
882 Yakushima. Ecol Res 19: 485-493.
- 883 49. Hanya G (2004) Diet of a Japanese macaque troop in the coniferous
884 forest of Yakushima. Int J Primatol 25: 55-71.
- 885 50. Hill DA (1997) Seasonal variation in the feeding behavior and diet of
886 Japanese macaques (*Macaca fuscata yakui*) in lowland forest of

Fermentation by gut microbe of Japanese macaques Hanya et al. 37

- 887 Yakushima. Am J Primatol 43: 305-322.
- 888 51. Hayaishi S, Kawamoto Y (2006) Low genetic diversity and biased
889 distribution of mitochondrial DNA haplotypes in the Japanese macaque
890 (*Macaca fuscata yakui*) on Yakushima Island. Primates 47: 158-164. doi:
891 10.1007/s10329-005-0169-1
- 892 52. Yamagiwa J (2008) History and present scope of field studies on *Macaca*
893 *fuscata yakui* at Yakushima Island, Japan. Int J Primatol 29: 49-64.
- 894 53. Hanya G, Kiyono M, Yamada A, Suzuki K, Furukawa M, Yoshida Y,
895 Chijiwa A (2006) Not only annual food abundance but also fallback food
896 quality determines the Japanese macaque density: evidence from
897 seasonal variations in home range size. Primates 47: 275-278.
- 898 54. Kurihara Y, Hanya G (2015) Comparison of feeding behavior between two
899 different-sized groups of Japanese macaques (*Macaca fuscata yakui*).
900 Am J Primatol 77: 986-1000. doi: 10.1002/ajp.22429
- 901 55. Kurihara Y, Kinoshita K, Shiroishi I, Hanya G (in press) Seasonal variation
902 in energy balance of wild Japanese macaques (*Macaca fuscata yakui*) in
903 a warm-temperate forest: a preliminary assessment in the coastal forest
904 of Yakushima. Primates.
- 905 56. Hayakawa T, Sawada A, Tanabe AS, Fukuda S, Kishida T, Kurihara Y,
906 Matsushima K, Liu J, Akomo-Okoue EF, Gravena W, Kashima M, Suzuki
907 M, Kadowaki K, Suzumura T, Inoue E, Sugiura H, Hanya G, Agata K
908 (2018) Improving the standards for gut microbiome analysis of fecal
909 samples: insights from the field biology of Japanese macaques on
910 Yakushima Island. Primates 59: 423-436. doi:
911 10.1007/s10329-018-0671-x

Fermentation by gut microbe of Japanese macaques Hanya et al. 38

- 912 57. Blümmel M, Ørskov ER (1993) Comparison of invitro gas-production and
913 nylon bag degradability of roughages in predicting feed-intake in cattle.
914 Anim Feed Sci Technol 40: 109-119. doi: 10.1016/0377-8401(93)90150-i
- 915 58. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M,
916 Glockner FO (2013) Evaluation of general 16S ribosomal RNA gene PCR
917 primers for classical and next-generation sequencing-based diversity
918 studies. Nucleic Acids Res 41: 11. doi: 10.1093/nar/gks808
- 919 59. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes
920 SP (2016) DADA2: High-resolution sample inference from Illumina
921 amplicon data. Nat Methods 13: 581-+. doi: 10.1038/nmeth.3869
- 922 60. Nawrocki EP, Eddy SR (2013) Infernal 1.1: 100-fold faster RNA homology
923 searches. Bioinformatics 29: 2933-2935. doi:
924 10.1093/bioinformatics/btt509
- 925 61. Rodrigues JFM, von Mering C (2014) HPC-CLUST: distributed
926 hierarchical clustering for large sets of nucleotide sequences.
927 Bioinformatics 30: 287-288. doi: 10.1093/bioinformatics/btt657
- 928 62. Rodrigues JFM, Schmidt TSB, Tackmann J, von Mering C (2017)
929 MAPseq: highly efficient k-mer search with confidence estimates, for
930 rRNA sequence analysis. Bioinformatics 33: 3808-3810. doi:
931 10.1093/bioinformatics/btx517
- 932 63. Tackmann J, Frederico J, Rodrigues M, von Mering C (2019) Rapid
933 Inference of Direct Interactions in Large-Scale Ecological Networks from
934 Heterogeneous Microbial Sequencing Data. Cell Syst 9: 286-+. doi:
935 10.1016/j.cels.2019.08.002
- 936 64. Price MN, Dehal PS, Arkin AP (2010) FastTree 2-Approximately

Fermentation by gut microbe of Japanese macaques
Hanya et al.
39

- 937 Maximum-Likelihood Trees for Large Alignments. PLoS One 5: 10. doi:
938 10.1371/journal.pone.0009490
- 939 65. Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor
940 package for differential expression analysis of digital gene expression
941 data. Bioinformatics 26: 139-140. doi: 10.1093/bioinformatics/btp616
- 942 66. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB,
943 Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B,
944 Thallinger GG, Van Horn DJ, Weber CF (2009) Introducing mothur:
945 Open-Source, Platform-Independent, Community-Supported Software for
946 Describing and Comparing Microbial Communities. Appl Environ
947 Microbiol 75: 7537-7541. doi: 10.1128/aem.01541-09
- 948 67. Kanehisa M, Goto S (2000) KEGG: Kyoto Encyclopedia of Genes and
949 Genomes. Nucleic Acids Res 28: 27-30. doi: 10.1093/nar/28.1.27
- 950 68. Buchfink B, Xie C, Huson DH (2015) Fast and sensitive protein alignment
951 using DIAMOND. Nat Methods 12: 59-60. doi: 10.1038/nmeth.3176
- 952 69. Franzosa EA, Mclver LJ, Rahnava G, Thompson LR, Schirmer M,
953 Weingart G, Lipson KS, Knight R, Caporaso JG, Segata N, Huttenhower C
954 (2018) Species-level functional profiling of metagenomes and
955 metatranscriptomes. Nat Methods 15: 962-+. doi:
956 10.1038/s41592-018-0176-y
- 957 70. Suzek BE, Wang YQ, Huang HZ, McGarvey PB, Wu CH, UniProt C
958 (2015) UniRef clusters: a comprehensive and scalable alternative for
959 improving sequence similarity searches. Bioinformatics 31: 926-932. doi:
960 10.1093/bioinformatics/btu739
- 961 71. Caspi R, Altman T, Billington R, Dreher K, Foerster H, Fulcher CA,

Fermentation by gut microbe of Japanese macaques Hanya et al. 40

- 962 Holland TA, Keseler IM, Kothari A, Kubo A, Krummenacker M,
963 Latendresse M, Mueller LA, Ong Q, Paley S, Subhraveti P, Weaver DS,
964 Weerasinghe D, Zhang PF, Karp PD (2014) The MetaCyc database of
965 metabolic pathways and enzymes and the BioCyc collection of
966 Pathway/Genome Databases. Nucleic Acids Res 42: D459-D471. doi:
967 10.1093/nar/gkt1103
- 968 72. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe
969 BE, Ling AV, Devlin AS, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ,
970 Turnbaugh PJ (2014) Diet rapidly and reproducibly alters the human gut
971 microbiome. Nature 505: 559-+. doi: 10.1038/nature12820
- 972 73. Hashizume K, Tsukahara T, Yamada K, Koyama H, Ushida K (2003)
973 *Megasphaera elsdenii* JCM1772(T) normalizes hyperlactate production in
974 the large intestine of fructooligosaccharide-fed rats by stimulating
975 butyrate production. J Nutr 133: 3187-3190.
- 976 74. Gomez A, Petrzekova K, Yeoman CJ, Vlckova K, Mrazek J, Koppova I,
977 Carbonero F, Ulanov A, Modry D, Todd A, Torralba M, Nelson KE,
978 Gaskins HR, Wilson B, Stumpf RM, White BA, Leigh SR (2015) Gut
979 microbiome composition and metabolomic profiles of wild western
980 lowland gorillas (*Gorilla gorilla gorilla*) reflect host ecology. Mol Ecol 24:
981 2551-2565. doi: 10.1111/mec.13181
- 982 75. Shortt C, Hasselwander O, Meynier A, Nauta A, Fernandez EN, Putz P,
983 Rowland I, Swann J, Turk J, Vermeiren J, Antoine JM (2018) Systematic
984 review of the effects of the intestinal microbiota on selected nutrients and
985 non-nutrients. Eur J Nutr 57: 25-49. doi: 10.1007/s00394-017-1546-4
- 986 76. Suzuki S, Hill DA, Sprague DS (1998) Intertroop transfer and dominance

Fermentation by gut microbe of Japanese macaques
Hanya et al.
41

- 987 rank structure of nonnatal male Japanese macaques in Yakushima,
988 Japan. *Int J Primatol* 19: 703-722.
- 989 77. Deusch S, Camarinha-Silva A, Conrad J, Beifuss U, Rodehutsord M,
990 Seifert J (2017) A Structural and Functional Elucidation of the Rumen
991 Microbiome Influenced by Various Diets and Microenvironments. *Front*
992 *Microbiol* 8: 21. doi: 10.3389/fmicb.2017.01605
- 993 78. Quan JP, Cai GY, Yang M, Zang ZH, Ding RR, Wang XW, Zhuang ZW,
994 Zhou SP, Li SY, Yang HQ, Li ZC, Zheng EG, Huang W, Yang J, Wu ZF
995 (2019) Exploring the Fecal Microbial Composition and Metagenomic
996 Functional Capacities Associated With Feed Efficiency in Commercial
997 DLY Pigs. *Front Microbiol* 10: 12. doi: 10.3389/fmicb.2019.00052
- 998 79. Porter NT, Martens EC (2017) The Critical Roles of Polysaccharides in
999 Gut Microbial Ecology and Physiology. In: Gottesman, S (ed.) *Annual*
1000 *Review of Microbiology*, Vol 71. Annual Reviews, Palo Alto, pp. 349-369
- 1001 80. Consortium THMP (2012) Structure, function and diversity of the healthy
1002 human microbiome. *Nature* 486: 207-214. doi: 10.1038/nature11234
- 1003 81. Louca S, Jacques SMS, Pires APF, Leal JS, Srivastava DS, Parfrey LW,
1004 Farjalla VF, Doebeli M (2017) High taxonomic variability despite stable
1005 functional structure across microbial communities. *Nat Ecol Evol* 1: 12.
1006 doi: 10.1038/s41559-016-0015
- 1007 82. Almeida A, Mitchell AL, Tarkowska A, Finn RD (2018) Benchmarking
1008 taxonomic assignments based on 16S rRNA gene profiling of the
1009 microbiota from commonly sampled environments. *Gigascience* 7: 10.
1010 doi: 10.1093/gigascience/giy054
- 1011 83. Hanya G, Kiyono M, Hayaishi S (2007) Behavioral thermoregulation of

Fermentation by gut microbe of Japanese macaques Hanya et al. 42

- 1012 wild Japanese macaques: Comparisons between two subpopulations. Am
- 1013 J Primatol 69: 802-815. doi: 10.1002/ajp.20397|ISSN 0275-2565
- 1014 84. Shulman RG, Rothman DL (2001) ¹³C NMR of intermediary metabolism:
- 1015 Implications for systemic physiology. Annu Rev Physiol 63: 15-48. doi:
- 1016 10.1146/annurev.physiol.63.1.15
- 1017 85. Agetsuma N, Noma N (1995) Rapid shifting of foraging pattern by
- 1018 Yakushima macaques (*Macaca fuscata yakui*) in response to heavy
- 1019 fruiting of *Myrica rubra*. Int J Primatol 16: 247-260.
- 1020 86. Noma N, Yumoto T (1997) Fruiting phenology of animal-dispersed plants
- 1021 in response to winter migration of frugivores in a warm temperate forest
- 1022 on Yakushima Island, Japan. Ecol Res 12: 119-129.
- 1023 87. Kuivanen J, Biz A, Richard P (2019) Microbial hexuronate catabolism in
- 1024 biotechnology. AMB Express 9: 11. doi: 10.1186/s13568-019-0737-1
- 1025 88. Hale VL, Tan CL, Niu KF, Yang YQ, Cui DY, Zhao HX, Knight R, Amato
- 1026 KR (2016) Effects of field conditions on fecal microbiota. J Microbiol
- 1027 Methods 130: 180-188. doi: 10.1016/j.mimet.2016.09.017
- 1028 89. Sawada A, Sakaguchi E, Hanya G (2011) Digesta Passage Time,
- 1029 Digestibility, and Total Gut Fill in Captive Japanese Macaques (*Macaca*
- 1030 *fuscata*): Effects Food Type and Food Intake Level. Int J Primatol 32:
- 1031 390-405. doi: 10.1007/s10764-010-9476-5
- 1032 90. Lu HP, Liu PY, Wang YB, Hsieh JF, Ho HC, Huang SW, Lin CY, Hsieh CH,
- 1033 Yu HT (2018) Functional Characteristics of the Flying Squirrel's Cecal
- 1034 Microbiota under a Leaf-Based Diet, Based on Multiple Meta-Omic
- 1035 Profiling. Front Microbiol 8: 13. doi: 10.3389/fmicb.2017.02622
- 1036 91. Ushida K, Tsuchida S, Ogura Y, Hayashi T, Sawada A, Hanya G (2016)

Fermentation by gut microbe of Japanese macaques
Hanya et al.
43

- 1037 Draft Genome Sequences of *Sarcina ventriculi* Strains Isolated from Wild
- 1038 Japanese Macaques in Yakushima Island. Microbiol Resour Ann 4: 2. doi:
- 1039 10.1128/genomeA.01694-15
- 1040
- 1041
- 1042

Fermentation by gut microbe of Japanese macaques
Hanya et al.
44

1043 Legends for figures

1044 Fig. 1. Map of the study site and location of sample collection. Open squares are
1045 highland samples and filled squares are lowland samples. Note that several
1046 locations are too close to be distinguished on the map. Contours are drawn
1047 every 300 m.

1048 Fig. 2. Difference in fermentation activity between the highland and lowland. (a)
1049 Gas production, (b) pH values after fermentation, and (c) total short-chained
1050 fatty acid (SCFA) production.

1051 Fig. 3. Profile of short-chained fatty acids produced during fermentation.

1052 Fig. 4. Beta-diversity of the community composition of the feces and suspension
1053 after fermentation (SAF) samples in the highland and lowland based on
1054 Bray-Curtis similarity index.

1055 Fig. 5. Beta-diversity of (a) abundance patterns of genes, (b) abundance
1056 patterns of pathway, and (c) coverage patterns of pathway between the
1057 highland and lowland.

1058

Fermentation by gut microbe of Japanese macaques
Hanya et al.
45

1059

1060 Fig. 1

1061

Fermentation by gut microbe of Japanese macaques Hanya et al. 46

Fig. 2

Fermentation by gut microbe of Japanese macaques
Hanya et al.
47

1065

1066 Fig. 3

1067

1068

Fermentation by gut microbe of Japanese macaques Hanya et al. 48

1069

1070 Fig. 4

1071

Fermentation by gut microbe of Japanese macaques Hanya et al. 49

1072

1073

1074 Fig. 5

Table 1. Significantly enriched taxa in feces or suspension after fermentation (SAF) samples

(a) Highland

Taxonomic level	Taxa	Enriched in	P (after FDR)	Relative abundance Feces	Relative abundance SAF
Phylum	Proteobacteria	SAF	0.0059	0.0686	0.2958
Order	Lactobacillales	SAF	0.0106	0.001	0.0324
Family	Streptococcaceae	SAF	0.0105	0.0009	0.0323
	Comamonadaceae	SAF	0.0112	2E-07	0.0041
Genus	<i>Streptococcus</i>	SAF	<0.0001	0.0009	0.0317

(b) Lowland

Taxonomic level	Taxa	Enriched in	P (after FDR)	Relative abundance Feces	Relative abundance SAF
Phylum	Proteobacteria	SAF	0.0004	0.0412	0.1643
	Tenericutes	Feces	0.0159	0.0183	0.0052
	Spirochaetes	SAF	0.0159	0.0625	0.1453
Class	Bacilli	SAF	<0.0001	0	0.0086
	Gammaproteobacteria	SAF	<0.0001	0.0231	0.1552
	Spirochaetia	SAF	<0.0001	0.0625	0.1453
	Negativicutes	SAF	0.0319	0.0104	0.0254
Order	Pasteurellales	SAF	<0.0001	3E-05	0.0336
	Aeromonadales	SAF	<0.0001	0.0168	0.1008
	Lactobacillales	SAF	<0.0001	0	0.0086
	Spirochaetales	SAF	0.0022	0.0624	0.1453
	Acholeplasmatales	Feces	0.006	0.0067	0.0031
	Selenomonadales	SAF	0.0396	0.0025	0.0118
Family	Pasteurellaceae	SAF	<0.0001	3E-05	0.0336
	Streptococcaceae	SAF	<0.0001	0	0.0086
	Oscillospiraceae	SAF	<0.0001	0	0.0029
	Ruminococcaceae	Feces	0.0008	0.0191	0.0073
	Succinivibrionaceae	SAF	0.0015	0.0136	0.0969
	Lachnospiraceae	Feces	0.007	0.0469	0.0167
	Clostridiaceae	Feces	0.0205	0.0016	4E-05
	Anaeroplasmataceae	Feces	0.0308	0.01	0.0017
Genus	<i>Streptococcus</i>	SAF	<0.0001	0	0.0086
	<i>Holdemanella</i>	SAF	0.004	1E-05	0.0011