

1 **Greater Lumbar Extension During Dolphin Kick and Psoas Major Tightness in**  
2 **Swimmers with Low Back Pain**

3

4 **Authors:**

5 Gakuto, Kitamura PT, B.S<sup>1</sup>, Hiroshige, Tateuchi PT, Ph.D<sup>1</sup>, Noriaki, Ichihashi  
6 PT, Ph.D<sup>1</sup>

7 1. Department of Physical Therapy, Human Health Science, Graduate School of  
8 Medicine, Kyoto University,

9 **Correspondence**

10 Name: Gakuto, Kitamura, PT, B.S

11 Address: Department of Human Health Science, Graduate School of Medicine,  
12 Kyoto University, 53, kawara-cho, shogoin, sakyo-ku, Kyoto 606-8507, Japan

13 Telephone: +81-075-751-3964

14 FAX: +81-075-751-3964

15 Email: kitamura.gakuto.82m@st.kyoto-u.ac.jp

16

17

18

19

20 **Abstract**

21 Context: In competitive swimming, many swimmers experience lower back pain (LBP).

22 Lumbar hyperextension may cause LBP and tight hip flexor muscle may cause lumbar  
23 extension during swimming.

24 Objective: The purpose of this study was to clarify the features of the elastic moduli of the  
25 muscles and the lumbar extension when swimmers with low back pain (LBP) perform a  
26 dolphin kick (DK).

27 Design: Cross sectional study.

28 Setting: Single center.

29 Participants: Eleven male college swimmers were enrolled as the LBP group (who have LBP  
30 when swimming and during a lumbar extension), and 21 male college swimmers were recruited  
31 as control group (no LBP).

32 Interventions: The elastic moduli of the psoas major, iliacus, teres major, latissimus dorsi,  
33 pectoralis major, and pectoralis minor were measured through ultrasonic shear wave  
34 elastography. The lumbar and hip extension angles during a DK were measured using a video  
35 camera. The passive hip extension and shoulder flexion range of motion (ROM) were measured  
36 using a goniometer.

37 Main Outcome Measures: Muscle elastic moduli and lumbar extension angles during DK.

38 Results: The characteristics, muscle elastic moduli, DK motion, and ROM were compared  
39 between the two groups. LBP group demonstrated significantly higher elastic modulus of the  
40 psoas major and lower modulus of pectoralis minor compared to the control group. Also, LBP  
41 group showed greater lumbar extension during a DK and less hip extension ROM than the  
42 control group.

43 Conclusion: The higher elastic modulus of the psoas major and greater lumbar extension during  
44 a DK may be related to the LBP in swimmers.

45

46 Keywords: prevention, swimming, ultrasound, posture, injury management

47 In various sports, low back pain (LBP) is frequently seen in athletes including volleyball players,  
48 rhythmic gymnasts, and competitive swimming.<sup>1</sup> This often leads to reductions in training time and/or  
49 competitive opportunities. It has been reported that about half of all competitive swimmers experience LBP.<sup>2</sup>  
50 Therefore, the prevention of LBP in competitive swimmers is important. The physical characteristics of  
51 competitive swimmers is different from those of ordinary people. To prevent LBP in swimmers, it is  
52 necessary to know their physical characteristics. Competitive swimmers have a high risk of spinal  
53 deformities such as spine asymmetry, thoracic kyphosis, or lumbar lordosis.<sup>3</sup> Intervertebral disc degeneration  
54 is frequently found in the lower area of the lumbar spine (mainly at the L5/S1 level, and occasionally at the  
55 L4/L5 level) and this tendency is not associated with the swimming style.<sup>4</sup>

56  
57 Several studies have been conducted on the swimming kinematics of swimmers without LBP. It has  
58 been reported that the hip joint extends to about 10 degrees during a front crawl<sup>5</sup> and may be larger during a  
59 dolphin kick (DK) owing to its performance characteristics. In a previews study, it was reported that the  
60 tightness of the hip flexor muscle can reduce hip extension that create a lumbar hyperextension and pelvic  
61 anterior tilt in various movement in water.<sup>6</sup> Pelvic anterior tilting can make the pelvis at a lower position  
62 than normal in water.<sup>6</sup> A study examined the swimmers experiencing LBP and reported that repetitive lumbar  
63 hyperextensions during swimming may cause LBP.<sup>7</sup> The tight hip flexor muscle may extend lumbar spine  
64 during DK and cause LBP in swimmers. Furthermore, excessive training may also cause LBP in swimmers.  
65 However, the physical and kinematic characteristics of swimmers with LBP have not been clarified.

66  
67 Competitive swimmers have more general joint laxity than normal, but the range of motion (ROM) of  
68 their internal and external shoulder rotations are smaller.<sup>8</sup> Furthermore, it has been reported that the ROM  
69 of the internal shoulder rotations decrease as competitive swimmers continue their careers from their youth  
70 level to the college level.<sup>9</sup> Therefore, although joint flexibility is required in competitive swimming,  
71 swimming training imposes a burden on the muscles around the shoulder joint, which can tighten. The basic  
72 posture in swimming used to achieve the least amount of water resistance, called a “streamline,” is a posture  
73 with the trunk and hip at a 0 degree extension, and a shoulder flexion of greater than 180 degrees, with both  
74 hands placed above the head. As described above, swimming training can burden the muscles and tighten  
75 shoulder muscles. If a swimmer experiences tightness in the shoulder muscles, a trunk extension will occur

76 as a compensation of the shoulder flexion during a streamline posture, which is frequently seen in practice.  
77 As described above, repetitive lumbar extensions can cause LBP.<sup>7</sup> Therefore, tightness of shoulder muscles  
78 may cause wrong streamline posture that is related to LBP in swimmers.

79

80 Currently, shear wave elastography is known to be able to assess muscle stiffness noninvasively.  
81 However, no studies have compared the elastic moduli of the hip flexor and shoulder muscle or the lumbar  
82 extension angle during a DK between swimmers with and without LBP. Therefore, the aim of the present  
83 study was to compare the elastic moduli of the hip flexor and shoulder muscles as well as the ROM of the  
84 passive shoulder flexion and hip extension between swimmers with and without LBP, and to clarify the  
85 features of their muscle stiffness, hip and shoulder ROM, and lumbar motion during a DK. There were three  
86 hypotheses. First, the elastic modulus of hip flexor muscles is higher in swimmers with LBP than those  
87 without LBP. Second, the elastic modulus of shoulder muscles is higher in swimmers with LBP than those  
88 without LBP. Third, the lumbar extension angle during DK is greater in swimmers with LBP than those  
89 without LBP.

90

91

## 92 **METHODS**

93

### 94 **Design**

95

96 A cross sectional study was used.

97

98

### 99 **Participants**

100

101 Thirty-nine male college swimmers were initially enrolled in this study. The inclusion criteria of the  
102 LBP group were subjects with more than 30 mm of LBP in VAS while both swimming and during a lumbar  
103 extension.<sup>10</sup> The inclusion criteria of the control group were subjects currently with no LBP (0 mm on the  
104 VAS). As a result, 11 subjects were enrolled in the LBP group ( $21.1 \pm 1.5$  years), and 21 subjects were

105 enrolled in the control group ( $20.6 \pm 1.5$  years). Three subjects were excluded because they experienced LBP  
106 when swimming but not during a lumbar extension, and four were excluded because their LBP was less than  
107 30 mm in VAS. The subjects were provided a full explanation, based on the Helsinki declaration, regarding  
108 the aim, methods, and risks of the study, and their agreement was received. This study was conducted with  
109 the approval of the ethics committee of institution of authors.

110

## 111 **Procedures**

112

113 The subjects answered a questionnaire after an explanation of the study and an agreement to participate.  
114 In the questionnaire, the presence and intensity of their LBP, their swimming style, and the number of their  
115 training years were determined. The intensity of their LBP was assessed as being between zero and 100  
116 based on the visual analogue scale (VAS). The elastic moduli of the psoas major, iliacus, teres major,  
117 latissimus dorsi, pectoralis major, and pectoralis minor muscles were measured using ultrasonic shear wave  
118 elastography. Their lumbar and hip extension angles during a DK were measured using an underwater  
119 camera. In addition, their passive hip extension and shoulder flexion ROM were measured using a  
120 goniometer.

121

122 In this study, the elastic moduli of the iliacus, psoas major were analyzed since they may restrict hip  
123 extension and the teres major, latissimus dorsi, pectoralis major clavicular, pectoralis major sternocostal, and  
124 pectoralis minor muscles were measured because they may restrict shoulder flexion. The elastic moduli of  
125 these muscles were measured using ultrasonic shear wave elastography (Aixplorer, Supersonic imagine, Aix-  
126 en-Provence, France) with a linear array probe (SL10-2, Super Sonic Imagine, Aix-en-Provence, France).  
127 Using this device, the muscle stiffness can be measured quantitatively based on elastic moduli, and the elastic  
128 moduli of the muscles as measured through ultrasonic shear wave elastography were found in a study on  
129 cadavers to be strongly related to the degree of muscle expansion.<sup>11</sup> In a living body, passive joint movement  
130 or passive tension of a muscle are strongly related to its elastic moduli.<sup>12</sup> The elastic moduli (G) reflects the  
131 elasticity or stiffness of a tissue, which is calculated based on the propagation velocity (V) of the shear wave  
132 in the tissue, as described in the following expressions.<sup>13</sup>

133

134  $G=\rho V^2$

135

136 In this expression,  $\rho$  is the density of the tissue, which for muscle has been reported to be 1000 kg/m<sup>3</sup>.<sup>14</sup>

137

138 Measurements of the elastic moduli of the psoas major and iliacus muscles were conducted in a relaxed  
139 supine position with the hip joint extended at 0 degrees, and the pelvis of the subject was fastened using a  
140 band to avoid anterior tilting. The measurement location of the psoas major and iliacus are shown in Figure  
141 1. This location was determined to be 3 to 5 cm below the anterior superior iliac spine (ASIS) and distal of  
142 the inguinal ligament, based on a previous study.<sup>15</sup> The psoas major and iliacus were then identified as lateral  
143 to the femoral artery using ultrasonic images.

144 The shoulder position and locations of the elastic moduli measurements of the shoulder muscles are  
145 shown in Figures 2A–2C. The teres major and latissimus dorsi were measured during a relaxed sitting  
146 position and with the shoulder flexed at 90 degrees, rotated at 0 degrees, and the elbow flexed at 90 degrees.  
147 The measurement location of the teres major and latissimus dorsi was defined as 10 cm below the acromion  
148 (Figure 2A). The pectoralis major and minor were measured during a relaxed sitting position and with the  
149 shoulder abducted at 90 degrees, rotated at 0 degrees, and the elbow flexed at 90 degrees. The pectoralis  
150 major clavicular and sternocostal parts were measured at the midpoint between the greater tubercle and the  
151 acromioclavicular joint and at midpoint between the greater tubercle and the fourth sternocostal joint,  
152 respectively. The pectoralis minor was measured at the midpoint between the coracoid process and the fourth  
153 sternocostal joint in the same way as in a previous study.<sup>16</sup>

154 Before measuring the elastic moduli, the state of each muscle was confirmed through ultrasonic  
155 imaging. For all muscle measurements, the region of interest (ROI), in which area the elastic moduli can be  
156 measured, was set at the center of the muscle. The elastic moduli of all muscles were measured by placing  
157 an ultrasonic probe parallel to the muscle fiber. The elastic moduli of all muscles considered were measured  
158 three times, and the mean value was used for a statistical analysis.

159

160 To evaluate the reliability of the elastic moduli measurement using ultrasonic elastography with a liner  
161 probe, the elastic moduli of all muscles for ten of the asymptomatic subjects were measured twice to assess  
162 the intra-class correlation coefficients (ICC 1.1). Consequently, the ICC 1.1 value of each muscle was as

163 follows; psoas major = 0.888, iliacus = 0.924, teres major = 0.622, latissimus dorsi = 0.756, pectoralis major  
164 clavicular = 0.755, pectoralis major = 0.897, and pectoralis minor = 0.642. According to a report by Landis  
165 et al., if the ICC value is higher than 0.81, the reliability of the measurement may be nearly perfect.<sup>17</sup>  
166 Therefore, the number (k) of measurements to meet the requirement of the lowest ICC value of the teres  
167 major, i.e., greater than 0.8, was calculated using the Spearman-Brown formula (a).

168

$$169 \quad k = \frac{\rho_1(1 - \rho_2)}{\rho_2(1 - \rho_1)} \quad (a)$$

170

171 In this formula,  $\rho_1$  is the target ICC value, and  $\rho_2$  is the resulting ICC value. As a result, all measurements  
172 of the elastic moduli were conducted three times because k was approximately 2.57.

173

174 An underwater video camera (O10-SC13001-W-1, Lerving Technology, Kwai Chung N.T., Hong  
175 Kong; 1920×1080 resolution, at 60 fps) and a Kinovea video player (version 0.8.15, available for download  
176 at <http://www.kinovea.org>) were used to measure the lumbar angle during a DK. The reliability of the joint  
177 angle measurement in the sagittal plane as assessed from a digital image was obtained in a previous study.<sup>18</sup>  
178 The subjects performed an underwater DK for 15 m with full effort, and the video was taken at a point 7.5  
179 m away along the sagittal plane. The subjects were instructed to perform the DK at around 0.5 m from the  
180 surface of the water. The camera was set at 0.5 m below the surface of the water, 7.5 m from the starting side  
181 wall of the pool, and 4 m from the subject's swimming lane.<sup>19</sup> Markers were put on the subject's spinous  
182 process at T10, L3, S2, iliac crest (IC), greater trochanter (GT), and lateral epicondyle of the femur (LEF)  
183 to measure the angle of the lumbar and hip joints. The lumbar angle ( $\theta_1$ ) was defined as an angle made by  
184 lines A and B. Line A was produced by the markers at T10 and L3, whereas line B was produced by the  
185 markers at L3 and S2 (Figure 3).<sup>20</sup> The hip angle ( $\theta_2$ ) was defined as the angle made by two lines, one  
186 produced by the IC and GT, and the other produced by the GT and LEF.<sup>19</sup> The lumbar angle was measured  
187 when the angle of the hip extension was largest when performing a DK. The average values from three  
188 appropriate trials were used in the statistical analysis and false trials in which the subjects did not pass in  
189 front of the video camera were excluded. The time for the 15 m DK was measured with a stopwatch.

190

191 The passive shoulder flexion and hip extension ROM were assessed using a goniometer. The shoulder  
192 flexion ROM was assessed in a sitting position, and the hip extension ROM was assessed in a prone position.  
193 All measurement of joint angle during DK and passive ROM were analyzed by the same researcher.

194

## 195 **Statistical Analyses**

196

197 A statistical analysis was conducted using SPSS (version 22.0, SPSS Japan, Inc., Tokyo, Japan). The  
198 normality of all data obtained was confirmed through a Shapiro-Wilk test. The age, height, weight, number  
199 of training years, elastic modulus of each muscle, lumbar extension, and hip extension angle applied when  
200 performing a DK, as well as the passive ROM of the shoulder flexion and hip extension between the LBP  
201 and asymptomatic groups were compared using a Mann-Whitney U-test. Spearman's rank correlation  
202 coefficient test was conducted to investigate the relationship between swimming performance (best record  
203 of 50 m crawl or 15 m DK time) and degree of lumbar extension or muscle elastic modulus. The differences  
204 were shown to be statistically significant at an alpha level of 0.05.

205

## 206 **RESULTS**

207

208 All measurements were completed for each subject. The characteristics of the subjects are shown in  
209 Table 1. There were no differences in age, height, weight or number of training years between both groups.

210 The results of the elastic moduli are shown in Table 2. The LBP group showed a higher value in the  
211 elastic modulus of the psoas major compared with the control group. For the pectoralis minor, the LBP group  
212 showed a lower value compared with the control group. There were no differences in the elastic moduli of  
213 the other muscles between both groups.

214 The results of the lumbar and hip angle when performing a DK, along with the passive shoulder flexion  
215 and hip extension ROM, are shown in Table 3. The LBP group showed a larger lumbar extension angle when  
216 performing a DK than the control group.

217 The passive hip extension ROM of the LBP group was smaller than that of the control group. There  
218 were no significant differences in the other joint angles between both groups.

219 The result of Spearman's rank correlation coefficient test is shown in Table 4. No significant



220 correlation between swimming performance and lumbar extension angle or muscle elastic modulus was  
221 found (Table 4).

222

## 223 **DISCUSSION**

224

225 The LBP group showed a higher elastic modulus of the psoas major and lower elastic modulus of the  
226 pectoralis minor than control group. Additionally, LBP group indicated a larger lumbar extension angle when  
227 performing a DK and smaller passive hip extension angle than the control group.

228

229 This is the first study comparing the elastic moduli of the muscles and the joint angle when performing  
230 a DK between swimmers with and without LBP.

231

232 The first hypothesis of this study was that the LBP group would show a higher elastic modulus of the  
233 hip flexor muscle than the control group. This hypothesis was partially supported because the results of this  
234 study demonstrated that, for the psoas major, the LBP group showed a higher elastic modulus than the control  
235 group, although no significant difference was shown in the elastic modulus of the iliacus. The psoas major  
236 orients from the lumbar spine, and attaches to the lesser trochanter. This muscle is a hip flexor and has an  
237 extension moment arm at levels L1 through L3 of the lumbar spine at standing and lumbar extended  
238 positions.<sup>21</sup> Furthermore, because the spine is extended to maintain a streamline position, which is a basic  
239 position for swimming, the extension stress from the psoas major increases at a higher level of the lumbar  
240 spine. In addition, because the psoas major has a flexion moment arm at a low level of the lumbar spine, and  
241 produces a lumbar moment in a different direction between the high and low levels of the lumbar spine, it is  
242 thought that the L3 vertebra moves forward and the L1 vertebra moves backward, such that the entire lumbar  
243 spine becomes extended.<sup>21, 22</sup> Therefore, LBP may likely occur if the elastic modulus of the psoas major is  
244 high because the lumbar spine will be more extended. Moreover, the psoas major causes a stronger shear  
245 force at the low level area of the lumbar spine than at a high level.<sup>21</sup> It is known from studies on rats and  
246 rabbits that lumbar intervertebral disc degeneration occurs if the vertebral disc is stressed from a continual  
247 shear force.<sup>23, 24</sup> In humans, an overload of the intervertebral disc may be related to disc degeneration.<sup>25</sup> A  
248 previous study reported that intervertebral disc degeneration, such as a decrease in the intervertebral disc

249 thickness, is related to the frequency, duration, and intensity of the LBP.<sup>26</sup> With the subjects of our current  
250 study, an increase in the shear force at the low level of the lumbar spine may have occurred owing to the  
251 high elastic modulus of the psoas major, which could be the cause of their intervertebral disc degeneration  
252 and LBP. Although the iliacus has a hip flexion moment arm, as with psoas major, the iliacus is oriented  
253 from the pelvis and not from the lumbar spine. Therefore, if the elastic modulus of the iliacus is high, a  
254 pelvic anterior tilt and subsequent lumbar extension stress will occur; however, the stress to the lumbar spine  
255 might be less compared to that of the psoas major muscle. For this reason, it is rational for the iliacus to not  
256 be related to LBP in the present study, unlike with the psoas major.

257

258         The second hypothesis of this study was that the LBP group will show higher elastic moduli in the  
259 shoulder muscles than the control group; however, a lower elastic modulus was found in the pectoralis minor  
260 muscle for the LBP group than for the control group, and there were no significant differences in the elastic  
261 moduli of the other shoulder muscles. Therefore, our hypothesis was not supported. The pectoralis minor is  
262 oriented from the second to the fifth rib, and attached to the coracoid process, and the scapula is anterior  
263 tilted, rotated downward, protracted, and depressed when this muscle is activated.<sup>27</sup> Although, the normal  
264 scapula movement during a shoulder flexion has an upward rotation, a posterior tilt, and an external  
265 rotation,<sup>28</sup> the short length of the pectoralis minor normalized by height can cause a scapula anterior tilt and  
266 an internal rotation compared with the longer length of the pectoralis minor normalized by height.<sup>29</sup> To the  
267 best of our knowledge, there have been no studies investigating the relationship between the stiffness of the  
268 pectoralis minor and scapula kinematics; however, if the pectoralis minor is tight, the scapula kinematics  
269 may be similar to the kinematics under a short pectoralis minor. Conversely, the scapular kinematics under  
270 less stiffness of the pectoralis minor may be similar to that of a long pectoralis minor. In this study, a low  
271 elastic modulus of the pectoralis minor in the LBP group may help the scapula tilt at the posterior and rotate  
272 externally during a shoulder flexion, and the LBP subjects may be able to compensate for a glenohumeral  
273 flexion by increasing the movement of their scapula. Therefore, to prevent LBP from occurring owing to a  
274 lumbar extension, the LBP group may compensate for the lumbar extension during a shoulder flexion while  
275 in the streamline position through an increased scapula movement. We initially hypothesized that the LBP  
276 group would show higher elastic moduli of the shoulder muscles than the control group because the LBP  
277 subjects would compensate for a shoulder flexion occurring from the high elastic moduli of their shoulder

278 muscles by increasing their trunk extension, thereby causing LBP. However, the LBP group showed a lower  
279 elastic modulus of the pectoralis minor than the control group, and accordingly, it is thought that the low  
280 elastic modulus of the pectoralis minor in the LBP group is not the cause of, but rather a compensation for,  
281 LBP. In addition to the pectoralis minor, the elastic moduli of the latissimus dorsi, teres major, and pectoralis  
282 major, as the shoulder muscles considered in this study, were measured. These muscles have a shoulder  
283 extension moment in a shoulder flexed position,<sup>30</sup> and are thought to restrict a shoulder flexion, induce a  
284 lumbar extension, and result in LBP if their elastic moduli are high. However, there was no significant  
285 difference in the elastic moduli of the latissimus dorsi, teres major, and pectoralis major between the LBP  
286 and control groups. Therefore, it is thought that these muscles did not affect the LBP in the swimmers  
287 considered in this study.

288

289         The third hypothesis was that the LBP group would show a greater lumbar extension when performing  
290 a DK than the control group. The LBP group showed a significantly higher lumbar extension angle during a  
291 DK than the control group, which is congruent with this hypothesis. Because repetitive lumbar  
292 hyperextensions while swimming can induce LBP,<sup>7</sup> the large angle of the lumbar extension during a DK can  
293 cause LBP. In this study, the passive hip extension ROM in the LBP group was significantly lower than that  
294 in the control group, and the subjects in the LBP group may have compensated for a hip extension when  
295 performing a DK through a lumbar extension. However, because the angle of the hip extension during a DK  
296 was about 9 degrees in both groups, it is thought that the hip was not extended to the limit of a passive hip  
297 extension ROM when performing a DK. Therefore, the significant difference in the lumbar extension angle  
298 during a DK may be influenced by another factor. The results of this study indicate that the elastic modulus  
299 of the psoas major was higher in the LBP group than in the control. Because the high elastic modulus of the  
300 psoas major can cause a lumbar lordosis, and the hip will extend the psoas major when performing a DK,  
301 thereby increasing the elastic modulus of psoas major, the high lumbar extension angle during a DK in the  
302 LBP group may be caused by the high elastic modulus of the psoas major. An approach to decrease the elastic  
303 modulus of the psoas major, or exercising the hip extension without a trunk extension may be useful in  
304 treating or preventing LBP.

305

306         There was no correlation between swimming performance and degree of lumbar extension during DK,

307 as demonstrated by the Spearman's rank correlation coefficient test. Therefore, increased lumbar extension  
308 during DK does not necessarily improve the swimming performance. However, it may increase the lumbar  
309 spine load. Thus, higher lumbar extension degree during DK is not an appropriate technique to improve  
310 swimming performance or reduce lumbar spine load.

311

312 There may be several factors causing LBP in swimmers; extended training period being one of them.  
313 However, no significant relationship was found between LBP and number of training years using Mann-  
314 Whitney U-test. These results may be partially explained by the participants' team. In this study, the  
315 participants were the members of a swimming club. Thus, their practice frequency and training year were  
316 similar. So, the effects of extended training on swimmers with LBP are unclear.

317

318 There are certain limitations to the present study. First, as a cross-sectional study, it is impossible to  
319 indicate the cause and effect relationship between LBP and the elastic moduli of the muscles or the lumbar  
320 angle during a DK. Second, the inclusion criteria for the LBP in this study differ from those of many other  
321 previous studies because many other studies include LBP patients whose LBP has continued for longer than  
322 3 months, and thus the results of this study are incommensurable with those of other studies. Third, the  
323 stiffness of the psoas major is thought to cause a lumbar extension when swimming; however, swimming  
324 styles other than a DK were not assessed. In competitive swimming, swimmers spend a significant amount  
325 of their training time using a crawl stroke regardless of their particular swimming style. Therefore, the  
326 lumbar motion during a crawl stroke may be related to LBP. To clarify the cause and effect relationship  
327 between the elastic moduli of the muscles and LBP, or the same criteria of LBP as applied in other studies,  
328 further longitudinal researches are needed.

329

## 330 **CONCLUSION**

331

332 This study indicated that swimmers suffering from LBP during lumbar extension have a higher  
333 elastic modulus of their psoas major, lower elastic modulus of their pectoralis minor, and a greater lumbar  
334 extension during a DK than swimmers without LBP. This study also showed that lumbar extension during  
335 DK and muscle elastic modulus are not related to swimming performance. Therefore, these issues in

336 swimmers with LBP must be resolved to prevent LBP.


337

338 **Reference**

- 339 1. Vidal-conti J. The relationship between low back pain and sport practice in young people. *International*  
340 *Journal of Orthopedics and Rehabilitation*. 2014;1:67-73.
- 341 2. Masiero S, Carraro E, Celia A, Sarto D, Ermani M. Prevalence of nonspecific low back pain in  
342 schoolchildren aged between 13 and 15 years. *Acta Paediatr*. 2008;97(2):212-216.
- 343 3. Zaina F, Donzelli S, Lusini M, Minnella S, Negrini S. Swimming and spinal deformities: A cross-  
344 sectional study. *J Pediatr*. 2015;166(1):163-167.
- 345 4. Kaneoka K, Shimizu K, Hangai M, et al. Lumbar intervertebral disk degeneration in elite competitive  
346 swimmers: a case control study. *Am J Sports Med*. 2007;35(8):1341-1345.
- 347 5. SANDERS RH. Kinematics, coordination, variability, and biological noise in the prone flutter kick at  
348 different levels of a “learn-to-swim” programme. *J Sports Sci*. 2007;25(2):213-227.
- 349 6. Kenal KA, Knapp LD. Rehabilitation of injuries in competitive swimmers. *Sports Medicine*.  
350 1996;22(5):337-347.
- 351 7. Wanivenhaus F, Fox AJS, Chaudhury S, Rodeo SA. Epidemiology of injuries and prevention strategies  
352 in competitive swimmers. *Sports health*. 2012;4(3):246-251.
- 353 8. Jansson A, Saartok T, Werner S, Renström P. Evaluation of general joint laxity, shoulder laxity and  
354 mobility in competitive swimmers during growth and in normal controls. *Scand J Med Sci Sports*.  
355 2005;15(3):169-176.
- 356 9. Riemann BL, Witt J DG. Glenohumeral joint rotation range of motion in competitive swimmers. *J*  
357 *Sports Sci*. 2015;29:1191-1199.
- 358 10. Teodorczyk-Injeyan JA, McGregor M, Triano JJ, Injeyan HS. Elevated Production of Nociceptive CC-  
359 chemokines and sE-selectin in Patients with Low Back Pain and the Effects of Spinal Manipulation.  
360 *Clin J Pain*. 2017;0:1.
- 361 11. Eby SF, Song P, Chen S, Chen Q, Greenleaf JF, An KN. Validation of shear wave elastography in  
362 skeletal muscle. *J Biomech*. 2013;46(14):2381-2387.
- 363 12. Maïsetti O, Hug F, Bouillard K, Nordez A. Characterization of passive elastic properties of the human  
364 medial gastrocnemius muscle belly using supersonic shear imaging. *J Biomech*. 2012;45(6):978-984.

- 365 13. Hug F, Tucker K, Gennisson JL, Tanter M, Nordez A. Elastography for muscle biomechanics: Toward  
366 the estimation of individual muscle force. *Exerc Sport Sci Rev.* 2015;43(3):125-133.
- 367 14. Gennisson JL, Cornu C, Catheline S, Fink M, Portero P. Human muscle hardness assessment during  
368 incremental isometric contraction using transient elastography. *J Biomech.* 2005;38(7):1543-1550.
- 369 15. Jiroumaru T, Kurihara T, Isaka T. Establishment of a recording method for surface electromyography in  
370 the iliopsoas muscle. *J Electromyogr Kinesiol.* 2014;24(4):445-451.
- 371 16. Umehara J, Nakamura M, Fujita K, et al. Shoulder horizontal abduction stretching effectively increases  
372 shear elastic modulus of pectoralis minor muscle. *J Shoulder Elbow Surg.* 2017;26(7):1159-1165.
- 373 17. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics.*  
374 1977;33(1):159-174.
- 375 18. Moral-Muñoz JA, Esteban-Moreno B, Arroyo-Morales M, Cobo MJ H-VE. Agreement between face-  
376 to-face and free software video analysis for assessing hamstring flexibility in adolescents. *J Strength*  
377 *Cond Res.* 2015;29(9):2661-2665.
- 378 19. Atkison RR, Dickey JP, Dragunas A, Nolte V. Importance of sagittal kick symmetry for underwater  
379 dolphin kick performance. *Hum Mov Sci.* 2014;33(1):298-311.
- 380 20. Claus AP, Hides JA, Moseley GL, Hodges PW. Is “ideal” sitting posture real?: Measurement of spinal  
381 curves in four sitting postures. *Man Ther.* 2009;14(4):404-408.
- 382 21. Bogduk N, Pearcy M, Hadfield G. Anatomy and biomechanics of psoas major. *Clin Biomech (Bristol,*  
383 *Avon).* 1992;7(2):109-119.
- 384 22. Penning L. Psoas muscle and lumbar spine stability: A concept uniting existing controversies. Critical  
385 review and hypothesis. *Eur Spine J.* 2000;9(6):577-585.
- 386 23. Kim J, Yang SJ, Kim H, Kim Y, Park JB, Dubose C LT. Effect of shear force on intervertebral disc  
387 ( IVD ) degeneration : An In vivo rat study. *Ann Biomed Eng.* 2012;40(9):1996-2004.
- 388 24. Xia DD, Lin SL, Wang XY, Wang YL, Xu HM, Zhou F TJ. Effects of shear force on intervertebral  
389 disc : an in vivo rabbit study. *Eur Spine J.* 2015;24:1711-1719.
- 390 25. Stokes IAF, Iatridis JC. Mechanical conditions that accelerate intervertebral disc degeneration: Overload  
391 versus immobilization. *Spine.* 2004;29(23):2724-2732.
- 392 26. Videman T, Battié MC, Gibbons LE, Maravilla K, Manninen H, Kaprio J. Associations between back  
393 pain history and lumbar MRI findings. *Spine.* 2003;28(6):582-588.

- 394 27. Schenkman M, Rugo de Cartaya V. Kinesiology of the shoulder complex. *J Orthop Sports Phys Ther.*  
395 1987;8(9):438-450.
- 396 28. McClure PW, Michener LA, Sennett BJ, Karduna AR. Direct 3-dimensional measurement of scapular  
397 kinematics during dynamic movements in vivo. *J Shoulder Elbow Surg.* 2001;10(3):269-277.
- 398 29. Borstad JD, Ludewig PM. The effect of long versus short pectoralis minor resting length on scapular  
399 kinematics in healthy individuals. *J Orthop Sports Phys Ther.* 2005;35(4):227-238.
- 400 30. Kuechle DK, Newman SR, Itoi E, Morrey BF. Kuechle--Shoulder muscle moment arms during  
401 horizontal flexion and elevation. *J Shoulder Elbow Surg.* 1997;6(5):429-439.
- 402



403

404 **Figure 1.** Measurement sites of shear elastic modulus for psoas major and iliacus muscles

405

406 The measurement locations of the psoas major (PM) and iliacus (IL) are shown. First, the femoral artery  
407 was identified through ultrasonic imaging as 3 to 5 cm below the anterior superior iliac spine (ASIS) and  
408 distal of the inguinal ligament (InL). These muscles were then identified lateral to the femoral artery. At  
409 this location, the probe was set parallel to the muscle fiber, and the elastic moduli were measured.






410

411 **Figure 2A.** Measurement position for the shear elastic modulus of the teres major and latissimus dorsi

412

413 The measurement position for the teres major and latissimus dorsi is shown in Figure 2A. The subject sat

414 with a shoulder flexed at 90 degrees, rotated at 0 degrees, and the elbow flexed at 90 degrees.




415

416 **Figure 2B.** Measurement position for the shear elastic modulus of the pectoralis major and minor shoulder  
417 muscles

418

419 The position used during the measurement of the pectoralis major and minor is shown in Figure 2B. The  
420 position was the shoulder abducted at 90 degrees, rotated at 0 degrees, and with the elbow flexed at 90  
421 degrees.

422




423

424 **Figure 2C.** Measurement sites of shear elastic modulus for pectoralis major clavicular part, sternocostal  
425 part, and pectoralis minor muscles

426

427 The measurement locations of the pectoralis major clavicular (PMaC), sternocostal (PMaS), and pectoralis  
428 minor (PMi) are shown in Figure 2C. The pectoralis major clavicular part was measured at the midpoint  
429 between the greater tubercle (GT) and acromioclavicular joint (ACJ), the pectoralis minor (PMi) was  
430 measured at the midpoint between the coracoid process (CP) and fourth sternocostal joint (4th SCJ), and  
431 the sternocostal (PMaS) was measured at the midpoint between the greater tubercle and the fourth  
432 sternocostal joint.



433

434 **Figure 3.** Measurement method of lumbar and hip joint angle during a DK

435

436 The measurement method of the lumbar and hip joint angle is shown in Figure 3. The lumbar angle ( $\theta_1$ )

437 was defined as the angle created by two lines, one produced by the markers at T10 and L3, and the other

438 produced by the markers at L3 and S2. The hip angle ( $\theta_2$ ) was defined as the angle made by two lines, one

439 produced by the iliac crest (IC) and greater trochanter (GT), and the other produced by the GT and lateral

440 epicondyle of the femur (LEF).

441

442 **Table 1. Characteristics of the participants**

443

	<b>LBP group (n = 11)</b>	<b>Control group (n = 21)</b>	<b><i>P</i> value</b>
age, y	21.1 (1.5)	20.6 (1.5)	.558
height, cm	176.4 (5.7)	173.2 (5.9)	.208
body mass, kg	68.3 (8.0)	65.9 (7.2)	.238
training years, y	12.6 (3.5)	12.1 (3.4)	.785

444 Abbreviations: LBP, low back pain.

445

446

447 **Table 2. Shear elastic modulus of each muscle**

448

<b>Shear elastic modulus, kPa</b>				
<b>Muscles</b>	<b>LBP group (n = 11)</b>	<b>Control group (n = 21)</b>	<b>Effect size (<i>r</i>)</b>	<b><i>P</i> value</b>
psoas major	17.9 (4.3)	13.1 (2.6)	0.516	.003
iliacus	14.2 (4.7)	11.6 (2.5)	0.228	.208
teres major	9.4 (3.6)	8.5 (1.7)	0.140	.434
latissimus dorsi	8.8 (2.6)	8.1 (1.6)	0.088	.639
pectoralis major	10.3 (3.8)	10.4 (3.4)	0.025	.907
clavicular part				
pectoralis major	8.5 (3.7)	8.3 (2.8)	0.025	.907
sterncostal part				
pectoralis minor	5.4 (2.1)	7.6 (3.5)	0.389	.027

449 Abbreviations: LBP, low back pain.

450

451 **Table 3. Angle of each joint**

452

<b>Joint angle, deg</b>				
	<b>LBP group (n = 11)</b>	<b>Control group (n = 21)</b>	<b>Effect size (<i>r</i>)</b>	<b><i>P</i> value</b>
Lumbar extension	22.1 (4.0)	15.3 (2.6)	0.701	<.001
during DK				
Hip extension	9.5 (4.3)	8.7 (5.3)	0.070	.690
during DK				
Passive	199.5 (11.4)	202.9 (8.8)	0.107	.546
shoulder flexion				
Passive	23.6 (7.4)	29.3 (4.9)	0.364	.040
hip extension				

453 Abbreviations: DK, dolphin kick; LBP, low back pain.

454 **Table 4. Correlation Between Swimming Performance and Lumbar Extension or Muscle Elastic**  
 455 **Modulus**

Variables	Correlation coefficient		Correlation coefficient	
	Best records of 50m crawl	<i>P</i> value	15m DK time	<i>P</i> value
Lumbar extension angle	.088	.630	.101	.583
Psoas major	-.033	.859	.037	.839
Iliacus	-.009	.962	.158	.387
Teres major	.175	.337	.031	.865
Latissimus dorsi	.219	.229	.229	.207
Pectoralis major clavicular part	-.084	.647	-.060	.745
Pectoralis major sterncostal part	.041	.826	-.012	.949
Pectoralis minor	.129	.483	.063	.734

456 Abbreviations: DK, dolphin kick.