TITLE:

Successful treatment of Hodgkin lymphoma-like EBV-associated post-transplant lymphoproliferative disorder following allogeneic hematopoietic stem cell transplantation with nivolumab

AUTHOR(S):

Wada, Fumiya; Kondo, Tadakazu; Nakamura, Momoko; Chonabayashi, Kazuhisa; Nishikori, Momoko; Hishizawa, Masakatsu; Yamashita, Kouhei; Takaori-Kondo, Akifumi

CITATION:

Wada, Fumiya ...[et al]. Successful treatment of Hodgkin lymphoma-like EBV-associated post-transplant lymphoproliferative disorder following allogeneic hematopoietic stem cell transplantation with nivolumab. *Annals of Hematology* 2020, 99: 887-889

ISSUE DATE:

2020-04

URL:

<http://hdl.handle.net/2433/254675>

RIGHT:

This is a post-peer-review, pre-copyedit version of an article published in *Annals of Hematology*. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s00277-020-03960-4>; The full-text file will be made open to the public on 12 February 2021 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

**Successful treatment of Hodgkin Lymphoma-like EBV-associated post-transplant
lymphoproliferative disorder following allogeneic hematopoietic stem cell
transplantation with nivolumab**

Fumiya Wada¹, Tadakazu Kondo^{1*}, Momoko Nakamura¹, Kazuhisa Chonabayashi¹,
Momoko Nishikori¹, Masakatsu Hishizawa¹, Kouhei Yamashita¹, and Akifumi
Takaori-Kondo¹

¹Department of Hematology and Oncology, Graduate School of Medicine, Kyoto
University, Kyoto, Japan

* To whom correspondence should be addressed: Tadakazu Kondo, Department of

Hematology and Oncology, Graduate School of Medicine, Kyoto University

54 Shogoin Kawaharacho Sakyo-ku, Kyoto 606-8507, Japan

Tel: +81-75-751-4946; FAX: +81-75-751-4963; E-mail: tadakazu@kuhp.kyoto-u.ac.jp

Running title: Treatment of EBV-PTLD with nivolumab

18 **Keywords:** Epstein-Barr virus, PTLN, allogeneic hematopoietic stem cell

19 transplantation, nivolumab, immune checkpoint inhibitors

20

21 **Type of manuscript:** Letter to the Editor

22

23 Figure: 1

24 Reference count: 8

25

26

27

28

29

30

31

32

33

34

35

36 Dear Editor,

37 Epstein-Barr virus (EBV)-associated post-transplantation lymphoproliferative
38 disorders (PTLD) represent a heterogeneous group of EBV-driven abnormal lymphoid
39 proliferations occurring after allogeneic hematopoietic stem cell transplantation [1-3].

40 Although several therapeutic strategies, such as reduction of immunosuppressive agents,
41 rituximab monotherapy, multiagent chemotherapy, and infusion of EBV-specific
42 cytotoxic T-lymphocytes (CTLs) have somewhat improved the outcome of patients with
43 PTLD, there is a concern that PTLD is still significantly associated with high mortality.
44 The use of immune checkpoint inhibitors (iCPIs) has emerged as a promising strategy
45 because of its potential effectiveness in enhancing anti-tumor immunity.

46 Recently several case reports have been published describing the use of iCPIs to treat
47 patients with aggressive hematological malignancies such as Hodgkin lymphoma [4-6].

48 However, there is a paucity of data on the safety and efficacy regarding the use of iCPIs
49 before or after allo-HSCT. Although iCPIs can induce an anti-tumor immune response,
50 there is also a risk of graft-versus-host disease (GVHD). Ijaz A et al. have reported that
51 14% of the patients who receive iCPIs after allo-HSCT develop acute GVHD and 9%
52 develop chronic GVHD [7]. Therefore, development of GVHD should be considered
53 while applying these agents in the pre- or post-allo-HSCT settings. Despite the

abundant data on the use of iCPIs for the treatment of cHL, only one pediatric case report is available on the use of iCPIs for PTLD [8]. Here we report the first case of successful iCPI use for the treatment of an adult patient with cHL-like PTLD following allo-HSCT.

A 58-year-old woman was diagnosed with myelodysplastic syndrome and refractory cytopenia with multilineage dysplasia (MDS-RCMD) with complex karyotype in June 2013. She received bone marrow transplantation from an HLA8/8 allele-matched unrelated donor in March 2014. In March 2018, ^{18}F fluorodeoxyglucose-positron emission tomography (^{18}F -FDG-PET) revealed left supraclavicular and abdominal paraaortic lymphadenopathies (Figure 1a and b). Biopsy of the supraclavicular lymph node was performed for diagnosis, and the histology showed EBV-associated cHL-like PTLD with the presence of CD15, CD30, and EBER-positive tumor cells.

We started administering brentuximab vedotin (BV) every three weeks. After six courses of BV, ^{18}F -FDG-PET/CT showed complete remission. After four months, follow-up CT study showed paraaortic lymphadenopathy of 20 mm in diameter. The CT-guided biopsy of the lymph node demonstrated the relapse of cHL-like PTLD. ^{18}F -FDG-PET/CT revealed no other lymph node involvement (Ann Arbor stage I). Following the regulatory approval, biweekly administration of nivolumab was started.

The starting dose was set to 240 mg/every other week. After four courses, liver toxicity (grade 3) developed, thus we postponed the next course. One month later, the liver function was improved. Therefore, we restarted administering nivolumab at the same dose. After six courses, cutaneous pruritus developed as an immune-related adverse event (irAE) of nivolumab, and thus we started administering 15mg prednisolone. Final evaluation with ^{18}F -FDG-PET after ten courses of nivolumab administration showed complete remission of PTLD (Figure 1c, d).

Although nivolumab is considered as a rational treatment option in cHL-like PTLD, to the best of our knowledge, this is the first case report of its use in an adult patient of EBV-associated PTLD after allo-HSCT. Ijaz A et al have demonstrated the high risk of acute and chronic GVHD among patients who received iCPIs after allo-HSCT [7]. In fact, a treatment for skin rash and liver damage after using iCPIs is needed, although it is difficult to distinguish between GVHD and irAE.

In conclusion, this case report shows the effectiveness of iCPIs on EBV-associated cHL-like PTLD after allo-SCT. However, the high risk of GVHD and irAE should be carefully considered while treating with iCPIs. To evaluate the efficacy and safety of iCPIs for PTLD, further analysis with a larger number of patients must be conducted.

90 **Acknowledgments**

91 We are grateful to the medical, nursing, and laboratory staff of the participating
92 departments for their contributions to this study.

93

94 **Compliance with ethical standards**

95 **Ethical approval:** All procedures performed in studies involving human participants
96 were in accordance with the ethical standards of the institutional and/or national
97 research committee and with the 1964 Helsinki declaration and its later amendments or
98 comparable ethical standards.

99 **Funding:** none

100 **Conflict of interest:** The authors declare no conflicts of interest.

101 **Informed consent:** Informed consent was obtained from the patient included in the
102 study.

103

104 **References**

- 105 1. Rashe L, Kapp M, Einsele H, Mielke S (2014) EBV-induced post transplant
106 lymphoproliferative disorders: a persisting challenge in allogeneic hematopoietic
107 SCT. Bone Marrow Transplant. 49: 163-167

- 108 2. Loren AW, Porter DL, Stadtmauer EA, Tsai DE (2003) Post-transplant
109 lymphoproliferative disorder:a review. Bone Marrow Transplant. 31: 145-155
- 110 3. Dierickx D, Habermann TM (2018) Post-transplantation lymphoproliferative
111 disorders in adults. N Engl J Med. 378: 549-562
- 112 4. Soiffer RJ (2019) Checkpoint inhibition to prevent or treat relapse in allogeneic
113 hematopoietic cell transplantation. Bone Marrow Transplant. 54:798-802
- 114 5. Merryman RW, Kim HT, Zinzani PL, Cario-Stella C, Ansell SM, Perales MA, et al
115 (2017) Safety and efficacy of allogeneic hematopoietic stem cell transplant after
116 PD-1 blockade in relapsed/refractory lymphoma. Blood. 129: 1380-1388.
- 117 6. Herbaux C, Gauthier J, Brice P, et al (2017) Efficacy and tolerability of nivolumab
118 after allogeneic transplantation for relapsed Hodgkin lymphoma. Blood. 129:
119 2471-2478.
- 120 7. Ijaz A, Khan AY, Malik SU, Faridi W, Fraz MA, Usman M, et al (2019) Significant
121 risk of graft-versus-host disease with exposure to checkpoint inhibitors before and
122 after allogeneic transplantation. Bio Blood Marrow Transplant. 25: 94-99
- 123 8. Kassa C, Remenyi P, Sinko J, Kallay K, kertes G, Kricvan G (2018) Successful
124 nivolumab therapy in an allogeneic stem cell transplant child with post-transplant
125 lymphoproliferative disorder. Pediatr Transplant. 22: e13302

126

127 **Figure. 1**

128 Response to nivolumab in a patient with Hodgkin lymphoma-like PTLN after alloSCT.

129 ^{18}F -FDG-PET obtained before (a and b) and after 10 courses of nivolumab

130 administration (c and d). The maximum intensity projections (a and c) and axial

131 reconstructions of the CT/PET fusions (b and d) are shown. The pre-treatment images (a

132 and b) show a metabolically active lymph node in the left inner hilum of the kidney.

133 The post-treatment scans (c and d) reveal the complete resolution of the metabolic

134 activity in the lesion.

135

136

137

138

A Self-archived copy in

京都大学学術情報リポジトリ

KURENAI

Kyoto University Research Information Repository