TITLE:

Kinetic analysis of inhibition of α -glucosidase by leaf powder from *Morus australis* and its component iminosugars

AUTHOR(S):

Qiao, Ying; Nakayama, Juri; Ikeuchi, Takeaki; Ito, Masaaki; Kimura, Toshiyuki; Kojima, Kenji; Takita, Teisuke; Yasukawa, Kiyoshi

CITATION:

Qiao, Ying ...[et al]. Kinetic analysis of inhibition of α -glucosidase by leaf powder from *Morus australis* and its component iminosugars. *Bioscience, Biotechnology, and Biochemistry* 2020, 84(10): 2149-2156

ISSUE DATE:

2020-10-2

URL:

<http://hdl.handle.net/2433/255582>

RIGHT:

This is an Accepted Manuscript of an article published by Taylor & Francis Group in 'Bioscience, Biotechnology, and Biochemistry' on 2020, available online at: <http://www.tandfonline.com/10.1080/09168451.2020.1783991>; The full-text file will be made open to the public on 13 Jul 2021 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。

Bioscience, Biotechnology, and Biochemistry

Running Title: Inhibition of α -glucosidase by *Morus australis* extracts

**Kinetic analysis of inhibition of α -glucosidase by leaf powder from *Morus australis*
and its component iminosugars**

Ying Qiao¹, Juri Nakayama¹, Takeaki Ikeuchi¹, Masaaki Ito², Toshiyuki Kimura³,
Kenji Kojima¹, Teisuke Takita¹, and Kiyoshi Yasukawa^{1,*}

¹*Division of Food Science and Biotechnology, Graduate School of Agriculture, Kyoto
University, Sakyo-ku, Kyoto 606-8502, Japan;* ²*Department of Bioresources Engineering,
National Institute of Technology, Okinawa College, 905, Henoko, Nago 905-2192,
Japan;* ³*Food Research Institute (NFRI), National Agriculture and Food Research
Organization (NARO), Tsukuba, Ibaraki, Japan*

*Corresponding author. Email: yasukawa@kais.kyoto-u.ac.jp

Abbreviations: 1-DNJ, 1-deoxynojirimycin; GAL-DNJ, 2-O- α -D-galactopyranosyl-DNJ

20 Mulberry leaves contain iminosugars, such as 1-deoxynojirimycin (1-DNJ), fagomine,
21 and 2-*O*- α -D-galactopyranosyl deoxynojirimycin (GAL-DNJ) that inhibit α -glucosidase.
22 In this study, we quantified iminosugars in *Morus australis* leaves and made the kinetic
23 analysis in the hydrolysis of maltose by α -glucosidase. By LC-MS/MS, the concentrations
24 of 1-DNJ, fagomine, and GAL-DNJ in the powdered leaves were 4.0, 0.46, and 2.5 mg/g,
25 respectively, and those in the roasted ones were 1.0, 0.24, and 0.73 mg/g, respectively,
26 suggesting that the roasting process degraded iminosugars. Steady-state kinetic analysis
27 revealed that the powdered and roasted leaves exhibited competitive inhibition. At pH 6.0
28 at 37°C, the IC₅₀ values of the extracts from the boiled powdered or roasted leaves were
29 0.36 and 1.1 mg/mL, respectively. At the same condition, the IC₅₀ values of 1-DNJ,
30 fagomine, and GAL-DNJ were 0.70 μ g/mL, 0.18 mg/mL, and 2.9 mg/mL, respectively.
31 These results suggested that in *M. australis*, 1-DNJ is a major inhibitor of α -glucosidase.
32
33 **Key words:** α -glucosidase; fagomine; iminosugar; 1-DNJ; *Morus australis*.

34 Introduction

35 α -Glucosidase [EC 3.2.1.20] catalyzes the hydrolysis reaction of α -1,4 glycosidic
36 bonds at the non-reducing end of the substrate. It is involved in the digestion of sugars.
37 One of the strategies to treat type-2 diabetes is inhibition of α -glucosidase. Acarbose,
38 miglitol, and voglibose are clinically used for this purpose.¹⁾ 1-Deoxynojirimycin (1-
39 DNJ) strongly inhibits α -glucosidase activity and is anticipated for clinical use.²⁾ In
40 addition, various α -glucosidase inhibitors have been identified in plants. They include
41 alkaloids such as vasicine,³⁾ flavonoids such as quercetin,⁴⁾ polyphenols such as *p*-
42 hydroxycinnamic acid and protocatechuic acid,⁵⁾ and iminosugars such as 1-DNJ,^{2,6-9)}
43 fagomine,⁷⁾ and 2-*O*- α -D-galactopyranosyl deoxynojirimycin (GAL-DNJ).⁷⁾

44 *Morus* (commonly known as mulberry) is distributed in temperate regions in the
45 world. It is cultivated as a crop for silkworm feed, fruit, and timber. *Morus* contains
46 various α -glucosidase inhibitors, among which 1-DNJ is the most characterized.⁷⁻¹⁰⁾
47 *Morus australis* is distributed in Ryukyu islands and is widely cultivated in Urasoe-shi,
48 Okinawa, Japan. It was reported that the 1-DNJ concentrations in *M. australis* leaves were
49 4 to 9-fold higher than those in *Morus alba* leaves distributed in Japanese mainland.^{10,11)}
50 Powdered and roasted *M. australis* leaves were developed as a tea with a function to
51 inhibit α -glucosidase activity. The ingestion of the tea suppressed sucrose-induced
52 elevation of blood glucose and insulin levels, indicating that the tea inhibited postprandial
53 elevation of blood glucose.¹¹⁾ In the inhibition of the hydrolysis of sucrose by sucrase
54 from rat intestine, boiled water extract of powdered *M. australis* leaves exhibited the IC₅₀
55 value of 12 ± 1.3 μ g/mL.¹¹⁾ These results suggest that a high intake of *M. australis* tea
56 might be effective for the prevention of type-2 diabetes. Recently, various physiological
57 activities of 1-DNJ have been reported, such as suppression of the elevation of

postprandial blood glucose,¹³⁾ postprandial glycemic control in subjects with impaired glucose metabolism,¹⁴⁾ stimulation of adiponectin and GLUT4 expressions adipocytes,¹⁵⁾ hypoglycemic effect,¹⁶⁾ prevention of diet-induced obesity through increases in adiponectin,¹⁷⁾ and suppression of lipid accumulation through activation of the β -oxidation system.¹⁸⁾ To pursue the application of *M. australis* leaves, enzyme chemical analysis of the inhibitory effects of *M. australis* leaves and its inhibitory components toward α -glucosidase is required. In this study, we selected three iminosugars 1-DNJ, fagomine, and GAL-DNJ as a research target. We quantified their concentrations in *M. australis* leaves by LC-MS/MS and made the kinetic analysis of their inhibitory effects toward the hydrolysis of maltose by α -glucosidase.

Materials and methods

Materials. Rat intestinal acetone powder was purchased from Sasaki Chemical (Kyoto, Japan) and used as the preparation of α -glucosidase. Maltose, 1-DNJ, and Glucose CII test were purchased from Fujifilm Wako Pure Chemical (Osaka, Japan). Fagomine and GAL-DNJ were purified from mulberry leaves as reported previously.^{9,12)} Powdered and roasted *M. australis* leaves were purchased from Urasoe-shi silver human resources center (Urasoe, Japan).

Preparation of extracts of *M. australis* leaves. Boiled water (50 mL) was added to powdered or roasted *M. australis* leaves (500 mg) and stirred for 5 min. The solution was suction-filtrated, and the filtrate was collected and used as the leaf extract.

LC-MS/MS analysis. LC-MS/MS analysis was performed as previously described.¹²⁾ Briefly, 50 mg of powdered or roasted *M. australis* leaves were dissolved in 5.0 mL of 50% ethanol. After sonication for 5 min, the mixture was centrifuged at 10,000

x g for 5 min at 4°C. The supernatant was separated and filtered (0.2 µm pore size; Sartorius Stedim Biotech, Goettingen, Germany). The filtrate was diluted with 0.1% formic acid, 50% acetonitrile to be 1% v/v for the quantification of 1-DNJ and GAL-DNJ and to be 10% for the quantification of fagomine. Standard 1-DNJ, fagomine, and GAL-DNJ were dissolved in 0.1% formic acid, 50% acetonitrile to be 200–1,000 ng/mL.

Conditions of LC-MS/MS are as follows: apparatus, LC-20A and LCMS-8045 (Shimadzu, Kyoto, Japan); column, TSK gel Amide-80 (particle size 5 µm; 100 mm × 2.0 mm i.d., Tosoh, Tokyo, Japan); column oven temperature, 40°C; mobile phase, 0.1% formic acid in acetonitrile (A) and 0.1% formic acid in water (B); mobile phase flow rate, 0.2 mL/min; injection volume, 5 µL; ion source, electrospray ionization (positive mode); drying gas, nitrogen (180°C, 7 L/min); nebulizing gas, nitrogen (1.6 bar); capillary voltage, -4,500 V; hexapole RF, 100 Vpp; quadrupole ion energy, 5 eV; collision gas, nitrogen (1.6 bar); collision energy, 10 eV; collision RF, 100 Vpp; and mass range, *m/z* 50–1500. Elution gradients are as follows: 0–2.0 min, 20%–60% B; 2.0–5.5 min, 60% B; 5.5–5.6 min, 60%–20% B; and 5.6–8.0 min, 20% B. 1-DNJ, fagomine, and GAL-DNJ were detected individually in the postcolumn by MS/MS with multiple reaction monitoring (MRM) for transition of the parent ions to the product ions. The concentrations of 1-DNJ, fagomine, and GAL-DNJ in powdered and roasted *M. australis* leaf extracts were calculated from calibration curves using standard 1-DNJ, fagomine, and GAL-DNJ.

Preparation of crude α-glucosidase solution. Rat intestinal acetone powder (0.5 g) was suspended with 5 mL of water followed by sonication. The solution was centrifuged at 10,000 x g for 15 min at 4°C, and the supernatant was collected. Then, the solution was diluted with water to be 35 mg/mL in protein concentration and used as the α-glucosidase

solution.

Measurement of α -glucosidase activity. 1-DNJ, fagomine, and GAL-DNJ were dissolved in water. Protocol for the reaction at pH 6.0 at 37°C is as follows. Pre-incubation (90 μ l) was initiated by mixing 80 μ l of the maltose solution in 0.1 M maleate buffer (pH 6.0) and 10 μ l of the extracts of powdered (0–10 mg/mL) or roasted (0–20 mg/mL) *M. australis* leaves or 10 μ l of the 1-DNJ (0–1 mg/mL), fagomine (0–40 mg/mL), and GAL-DNJ (0–100 mg/mL) solution in water. After the pre-incubation at 37°C for 10 min, the reaction was initiated by adding 10 μ l of α -glucosidase solution and continued at 37°C. Blank solution was prepared by adding 10 μ l of 0.1 M maleate buffer (pH 6.0). After 1 or 2 min, the solution was boiled for 5 min to stop reaction. To 13.3 μ l of the solution, 200 μ l of the coloring solution in Glucose CII test was added. The absorbance at 505 nm was measured with an EnSight multimodal plate reader (PerkinElmer, Waltham, MA). From the reaction rate, the Michaelis constant in the absence of inhibitor (K_m), the Michaelis constant in the presence of inhibitor (K_{mapp}), and maximum velocity (V_{max}) were calculated based on using Hanes-Woolf plot with Microsoft Excel.

The inhibitor constant (K_i) of reversible competitive inhibitor was calculated by the following equation:

$$\frac{K_{mapp}}{K_m} = 1 + \frac{[I]_o}{K_i} \quad (1)$$

where $[I]_o$ is the initial inhibitor concentration.

Thermodynamic analysis. The enthalpy change, ΔH° , of the dissociation of the

complex of α -glucosidase and inhibitor was determined from a van't Hoff plot according to Eq. 2, as described previously.¹⁹⁾ The Gibbs free energy change of dissociation, ΔG° and the entropy change of dissociation, ΔS° , at certain temperature were determined according to Eqs. 3 and 4, respectively.

$$\ln(K_i) = A - (\Delta H^\circ/R)(1/T) \quad (2)$$

$$\Delta G^\circ = -RT \ln K_i \quad (3)$$

$$\Delta S^\circ = (\Delta H^\circ - \Delta G^\circ) / T \quad (4)$$

where A , R , and T are the constant term, the gas constant ($= 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), and absolute temperature in degrees Kelvin.

Results and discussion

Quantification of iminosugars in M. australis leaves by LC-MS/MS

For quantification of iminosugars in *M. australis* leaves, we applied a standard 1-DNJ, fagomine, and GAL-DNJ to LC-MS/MS to make a calibration curve. The total ion scanning profile showed an intense molecular ions at m/z 164.20 $[M+H]^+$ for 1-DNJ, 148.20 $[M+H]^+$ for fagomine, and 326.20 $[M+H]^+$ for GAL-DNJ. The product ion scanning for these ions showed intense molecular ions at m/z 69.20 $[M+H-95]^+$, 80.20 $[M+H-84]^+$, and 146.20 $[M+H-H_2O]^+$ for 1-DNJ (Fig. S1a), at m/z 86.15 $[M+H-62]^+$, 112.15 $[M+H-2H_2O]^+$, and 130.15 $[M+H-H_2O]^+$ for fagomine (Fig. S1b), and at m/z 61.10 $[M+H-265]^+$, 146.15 $[M+H-180]^+$, and 164.20 $[M+H-162]^+$ for GAL-DNJ (Fig. S1c). The MRM chromatogram of the total of these three ions showed a peak at 5.4 min for 1-DNJ (Fig. S2a), at 4.0 min for fagomine (Fig. S2b), and at 4.3 min for GAL-DNJ (Fig. S2c).

The peak intensity increased linearly with increasing concentration of 1-DNJ (0–1,000 ng/mL) (Fig. S3a), fagomine (0–1,000 ng/mL) (Fig. S3b) or GAL-DNJ (0–1,000 ng/mL) (Fig. S3c).

We applied the extracts of powdered or roasted *M. australis* leaves to determine the concentrations of 1-DNJ, fagomine, and GAL-DNJ in the extracts of powdered (Fig. 1a–c) and roasted (Fig. 1d–f) leaves. The MRM chromatogram showed one peak at 5.4 min for the analysis of 1-DNJ (Fig. 1a and d), two peaks at 3.4 and 4.0 min, respectively, for the analysis of fagomine (Fig. 1b and e), and one peak at 4.3 min for the analysis of GAL-DNJ (Fig. 1c and f). In the analysis of fagomine, the peak at 4.0 min was used (Fig. 1b and e), according to the MRM chromatogram of standard fagomine (Fig. S2b). The structure of the substance corresponding to the peak at 3.4 min is unknown. Based on the calibration curves (Fig. S3), the concentrations of 1-DNJ, fagomine, and GAL-DNJ in the powdered leaf extracts were calculated to be 40, 4.6, and 25 $\mu\text{g/mL}$, respectively, and those in the roasted leaf extracts were calculated to be 10, 2.4, and 7.3 $\mu\text{g/mL}$, respectively. Thus, the concentrations of these three iminosugars in the leaves were calculated to be 4.0, 0.46, and 2.5 mg/g, respectively, for the powdered ones and 1.0, 0.24, and 0.73 mg/g, respectively, for the roasted ones. These results indicated that the iminosugar concentrations in the roasted leaves were 25–52% of those in the powdered leaves, suggesting that the roasting process considerably degraded iminosugars. These results also indicated that the 1-DNJ concentration in the leaves was 1.6–8.7 and 1.4–4.2 fold higher than the fagomine and GAL-DNJ concentrations, respectively.

The 1-DNJ concentrations of mulberry leaves in *M. alba* and *M. bombycis* were reported to be 1.0–1.4 mg/g, and those of mulberry leaf products on the market were 1.3–4.8 mg/g.¹⁰⁾ Thus, it seems that the 1-DNJ concentration of *M. australis* leaf is

relatively high. However, it should be noted that 1-DNJ concentration in mulberry leaf varies depending on seasons and collection site.¹⁹⁾

*Kinetic analysis of inhibition of α -glucosidase by the extracts of *M. australis* leaves*

The extract we prepared from rat intestinal acetone powder and used as the α -glucosidase solution was thought to contain maltase-glucoamylase and sucrase-isomaltase.²⁰⁾ In this study, we characterized the inhibitory effects of the extracts of *M. australis* leaves and imminosugars toward the maltase-mediated maltose-hydrolyzing activity and regarded the effects toward α -glucosidase activity.

We examined the reaction rates of α -glucosidase in the hydrolysis of 50 mM maltose in the presence of various concentrations of the extracts of powdered (0–1.0 mg/mL) or roasted (0–2.0 mg/mL) *M. australis* leaves at pH 6.0 at 37°C (Fig. 2a). The reaction rate decreased with increasing the concentration of each extract. The IC₅₀ values, which were determined using intersection at 50% activity in line graphs, of the extract of powdered leaves and that of roasted leaves were 0.36 and 1.1 mg/mL, respectively, indicating that the inhibitory effect of the powdered leaves was 3-fold higher than that of the roasted leaves.

We measured the reaction rates of α -glucosidase in the hydrolysis of various concentrations (0–75 mM) of maltose in the presence of 0.6 mg/mL of powdered or roasted leaf extract at pH 6.0 at 37°C. All plots showed saturated profiles (Fig. 2b). Hanes-Woolf plot showed parallel lines intersecting at the Y-axis (Fig. 2c), indicating that the manner of inhibition was competitive.

Kinetic analysis of inhibition of α -glucosidase by 1-DNJ, fagomine, and GAL-DNJ

Figure 3a and b show the reaction rates of α -glucosidase in the hydrolysis of 50 mM maltose in the presence of various concentrations of 1-DNJ (0–10 μ g/mL), fagomine (0–4 mg/mL), and GAL-DNJ (0–8 mg/mL) at pH 6.0 at 37°C. The reaction rates decreased with increasing the concentration of each iminosugar. The activity was almost completely inhibited by 1-DNJ and fagomine, while 20% activity remained by GAL-DNJ, suggesting that the inhibition by GAL-DNJ is partial. The IC_{50} values were 0.70 μ g/mL for 1-DNJ, 0.18 mg/mL for fagomine, and 2.9 mg/mL for GAL-DNJ, indicating that the inhibitory effect of 1-DNJ was 250- and 4,000- fold higher than that of fagomine or GAL-DNJ, respectively.

Figure 3c shows the reaction rates of α -glucosidase in the hydrolysis of various concentrations (0–75 mM) of maltose in the presence of 0.25 μ g/mL of 1-DNJ, 0.1 mg/mL of fagomine, or 1.5 mg/mL of GAL-DNJ at pH 6.0 at 37°C. Like the case with the reaction in the presence of the leaf extracts, all plots showed saturated profiles (Fig. 2b). Hanes-Woolf plot showed parallel lines intersecting at the Y-axis (Fig. 3d), indicating competitive inhibition. The K_m value was calculated to be 1.9 mM in the absence of iminosugar, and the K_{mapp} values were calculated to be the 9.0 mM in the presence of 0.25 μ g/mL 1-DNJ, 16 mM in the presence of 0.1 mg/mL fagomine, and 30 mM in the presence of 1.5 mg/mL GAL-DNJ.

The K_i values of 1-DNJ (molecular weight: 163), fagomine (147), and GAL-DNJ (330) calculated from the K_m and K_{mapp} values using Eq. 1 were 0.068 μ g/mL ($= 4.1 \times 10^{-7}$ M), 14 μ g/mL ($= 9.2 \times 10^{-5}$ M), and 102 μ g/mL ($= 3.1 \times 10^{-4}$ M), respectively. The K_i values of 1-DNJ and fagomine thus obtained were similar to those previously reported (1.12×10^{-6} M for 1-DNJ and 3.24×10^{-4} M for fagomine).²¹⁾

Insight into the contribution of 1-DNJ in *M. australis* leaves to the inhibition of α -glucosidase

As described above, the concentrations of IC₅₀ value of the powdered leaf extract, roasted leaf extract, and 1-DNJ were 360, 1,100, and 0.70 $\mu\text{g/mL}$, respectively, and the concentrations of 1-DNJ in the powdered and roasted leaves were 4.0 and 1.0 mg/g, respectively. Nakanishi et al. showed that dependences on season of 1-DNJ amount and the α -glucosidase inhibitory activity of mulberry leaves exhibited similar profiles.²²⁾ In this study, the inhibitory effects of fagomine and GAL-DNJ to α -glucosidase were thought to be marginal compared with 1-DNJ. Assuming that 1-DNJ is a sole component that inhibits α -glucosidase in mulberry leaf, it was calculated that 360 $\mu\text{g/mL}$ powdered leaf extract contained 1.4 $\mu\text{g/mL}$ 1-DNJ, and 1,100 $\mu\text{g/mL}$ roasted leaf extract contained 1.1 $\mu\text{g/mL}$ 1-DNJ. These concentrations correspond to 200% and 150%, respectively, of the IC₅₀ value of 1-DNJ (0.70 $\mu\text{g/mL}$), suggesting that in *M. australis*, the inhibitory effects of 1-DNJ were suppressed in the powdered and roasted leaf extracts. We speculate that GAL-DNJ, which exhibited partial inhibition, might suppress the inhibitory effect of 1-DNJ.

Effects of temperature and pH on the inhibition of α -glucosidase by the extracts of *M. australis* leaves and 1-DNJ

To explore the role of 1-DNJ on the inhibition of α -glucosidase by the extracts of *M. australis* powdered leaves, we examined the effects of temperature (27–47°C) and pH (6.0–8.0) on the inhibition. Figure S4 shows the effects of reaction temperature and pH on the α -glucosidase activity. The highest activity was observed at 62°C (Fig. S4a), and the relatively high activity was observed at pH 5–8 (Fig. S4b), suggesting that α -

glucosidase did not lose activity at pH 6.0–8.0 or at 27–47°C. Figure S5a–d show the reaction rates of α -glucosidase in the hydrolysis of various concentrations (0–75 mM) of maltose in the presence of 0.15 mg/mL of the powdered leaf extracts or 0.25 μ g/mL 1-DNJ at pH 6.0–8.0, at 27–47°C. All plots showed saturated profiles.

The K_i values were calculated from Eq. 1. Figure 4a and b show the dependence of K_i values at pH 6.0 on reaction temperature and the dependence of K_i values at 37°C on pH, respectively. The profiles of the extracts of powdered leaves and 1-DNJ were very similar, suggesting that 1-DNJ is a major inhibitory component in the powdered leaves.

The K_i values of the dissociation depended on temperature (Fig. 4a). Figure 4c shows van't Hoff plot of K_i values. Enthalpy changes (ΔH°) at pH 6 were calculated from the slope to be -87.6 ± 6.4 kJ mol⁻¹ for the extracts of powdered leaves and -72.8 ± 12.9 kJ mol⁻¹ for 1-DNJ. The negative ΔH° values of the dissociation indicate that the dissociation was exothermic. For 1-DNJ, Gibbs free energy change of dissociation (ΔG°) at pH 6.0 at 37°C were calculated be 38.1 kJ mol⁻¹ for 1-DNJ, and entropy change of dissociation (ΔS°) at pH 6.0 at 37°C were calculated to be -351.7 J mol⁻¹ K⁻¹ from Eq. 3. Since ΔH° is negative and $-T\Delta S^\circ$ is positive, the dissociation is enthalpy-driven. The K_i values of the dissociation also depended on pH (Fig. 4b). This might be due to that changes in ionization state of the amino acid residues of α -glucosidase involved in the binding.

In conclusion, *M. australis* leaves contain 1-DNJ a lot and strongly inhibit α -glucosidase activity. The K_i values of the extract of *M. australis* powdered leaves and 1-DNJ are temperature- and pH-dependent. By combining LC-MS/MS, the kinetic analysis of leaf extracts and its components might be valuable to evaluate the inhibitory activity toward α -glucosidase of various plant products.

271

272 **Author contribution**

273 M.I., T.K. and K.Y. designed research; Y.Q., J.N., and K.Y. performed research; Y.Q.,
274 J.N., T.I., M.I., T.K., K.K., T.T, and K.Y. analyzed data; Y.Q., M.I., T.K., and K.Y. wrote
275 the manuscript.

276

277 **Disclosure statement**

278 No potential conflict of interest was reported by the authors.

279

280 **Funding**

281 This study was supported in part by the Tojuro Iijima Foundation for Food Science
282 and Technology for Y.Q. and K.Y. and Grants-in-Aid for Scientific Research (nos.
283 18K19839 for K.Y. and 18KK0285 for K.K., T.T., and K.Y.) from the Japan Society for
284 the Promotion of Science.

285

286 **Acknowledgments**

287 We appreciate Dr. J. Wada of Kyoto Integrated Science & Technology BIO-
288 Analysis Center for instructing LC-MS/MS.

289

290 **References**

291

- 292 [1] Assefa ST, Yang EY, Chae SY, et al. Alpha glucosidase inhibitory activities of
293 plants with focus on common vegetables. Plants (Basel) 2019;18:E2.
- 294 [2] Nakagawa K. Studies targeting α -glucosidase inhibition, antiangiogenic effects, and

- lipid modification regulation: background, evaluation, and challenges in the development of food ingredients for therapeutic purposes. *Biosci. Biotechnol. Biochem.* 2013;77:900–908.
- [3] Gao H, Huang YN, Gao B, et al. Inhibitory effect on α -glucosidase by *Adhatoda vasica* Nees. *Food Chem.* 2008;108:965–972.
- [4] Tadera K, Minami Y, Takamatsu K, et al. Inhibition of α -glucosidase and α -amylase by flavonoids. *J. Nutr. Sci. Vitaminol. (Tokyo)* 2006;52:149–153.
- [5] Nile SH, Park SW. Antioxidant, α -glucosidase and xanthine oxidase inhibitory activity of bioactive compounds from maize (*Zea mays* L.). *Chem. Biol. Drug Des.* 2014;83:119–125.
- [6] Asano N. Glycosidase inhibitors: update and perspectives on practical use. *Glycobiology* 2003;13:93R–104R.
- [7] Asano N, Yamashita T, Yasuda K, et al. Polyhydroxylated alkaloids isolated from mulberry trees (*Morus alba* L.) and silkworms (*Bombyx mori* L.). *J. Agric. Food Chem.* 2001;49:4208–4213.
- [8] Asano N, Tomioka E, Kizu H, et al. Sugars with nitrogen in the ring isolated from the leaves of *Morus bombycis*. *Carbohydr. Res.* 1994;253:235–245.
- [9] Asano N, Oseki K, Tomioka E, et al. *N*-containing sugars from *Morus alba* and their glycosidase inhibitory activities. *Carbohydr. Res.* 1994;259:243–255.
- [10] Kimura T, Nakagawa K, Saito Y, et al. Determination of 1-deoxynojirimycin in mulberry leaves using hydrophilic interaction chromatography with evaporative light scattering detection. *J. Agric. Food Chem.* 2004;52:1415–1418.
- [11] Kume D, Fukamizu A, Kuraya E, et al. Inhibitory effect of leaf powder from *Morus australis* on postprandial blood glucose elevation. *Food Sci. Technol. Res.*

- 319 2019;66:52–56.
- 320 [12] Nakagawa K, Ogawa K, Higuchi O, et al. Determination of iminosugars in mulberry
321 leaves and silkworms using hydrophilic interaction chromatography-tandem mass
322 spectrometry. *Anal. Biochem.* 2010;15:217–222.
- 323 [13] Kimura T, Nakagawa K, Kubota H, et al. Food-grade mulberry powder enriched
324 with 1-deoxynojirimycin suppresses the elevation of postprandial blood glucose in
325 humans. *J. Agric. Food Chem.* 2007;55:5869–5874.
- 326 [14] Asai A, Nakagawa K, Higuchi O, et al. Effect of mulberry leaf extract with enriched
327 1-deoxynojirimycin content on postprandial glycemic control in subjects with
328 impaired glucose metabolism. *J. Diabetes Investig.* 2011;2:318–323.
- 329 [15] Seung-Min J, Do HJ, Shin MJ, et al. 1-Deoxynojirimycin isolated from a *Bacillus*
330 *subtilis* stimulates adiponectin and GLUT4 expressions in 3T3-L1 adipocytes. *J.*
331 *Microbiol. Biotechnol.* 2013;23:637–643.
- 332 [16] Vichasilp C, Nakagawa K, Sookwong P, et al. A novel gelatin crosslinking method
333 retards release of mulberry 1-deoxynojirimycin providing a prolonged
334 hypoglycaemic effect. *Food Chem.* 2012;134:1823–1830.
- 335 [17] Tsuduki T, Kikuchi I, Kimura T, et al. Intake of mulberry 1-deoxynojirimycin
336 prevents diet-induced obesity through increases in adiponectin in mice. *Food Chem.*
337 2013;139:16–23.
- 338 [18] Tsuduki T, Nakamura Y, Honma T, et al. Intake of 1-deoxynojirimycin suppresses
339 lipid accumulation through activation of the β -oxidation system in rat liver. *J. Agric.*
340 *Food Chem.* 2009;57: 11024–11029.
- 341 [19] Horn JR, Russell D, Lewis EA, et al. Van't Hoff and calorimetric enthalpies from
342 isothermal titration calorimetry: are there significant discrepancies? *Biochemistry*

343 2001;40:1774–1778.

344 [20] Byung-Hoo Lee BH, Hamaker BR. Maltase has most versatile α -hydrolytic activity
345 among the mucosal α -glucosidases of the small intestine. J. Pediatr. Gastroenterol.
346 Nutr. 2018;66:S7–S10.

347 [21] Liu Z, Yang Y, Dong W, et al. Investigation on the enzymatic profile of mulberry
348 alkaloids by enzymatic study and molecular docking. Molecules 2019;24:E1776.

349 [22] Nakanishi H, Onose S, Kitahara E, et al. Effect of environmental conditions on the
350 α -glucosidase inhibitory activity of mulberry leaves. Biosci. Biotechnol. Biochem.
351 2011;75:2293–2296.

352

353

Figure Legends

Fig. 1. MRM chromatogram of the extracts of *M. australis* leaves.

Notes: The extracts of powdered (a–c) or roasted (d–f) leaves were applied. MRM chromatogram of the total of ions at m/z 69.20 $[M+H-95]^+$, 80.20 $[M+H-84]^+$, and 146.20 $[M+H-H_2O]^+$ for 1-DNJ (a, d), at 86.15 $[M+H-62]^+$, 112.15 $[M+H-2H_2O]^+$, and 130.15 $[M+H-H_2O]^+$ for fagomine (b, e), and at 61.10 $[M+H-265]^+$, 146.15 $[M+H-180]^+$, and 164.20 $[M+H-162]^+$ for GAL-DNJ (c, f) are shown.

Fig. 2. Effects of the extracts of *M. australis* leaves on the hydrolysis of maltose by α -glucosidase.

Notes. (a) Dependence on the concentration of leaf extracts on the reaction rate. The reaction was carried out with 50 mM maltose and varying concentrations of powdered (open triangle) or roasted (open square) leaf extracts at pH 6.0 at 37°C. (b, c) Dependence on the substrate concentration of the reaction rate. The reaction was carried out with varying concentrations of maltose in the absence of iminosugar (open circle) or in the presence of 0.6 mg/mL powdered leaf extracts (open triangle) or roasted leaf extracts (open square) at pH 6.0 at 37°C. (b) Reaction rate (v_o) vs. maltose concentration ($[maltose]$). Solid line represents the best fit to the experimental data using the Michaelis-Menten equation using Microsoft Excel. (c) Hanes-Woolf plot ($[maltose] / v_o$ vs. $[maltose]$). The K_m and V_{max} values were calculated to be 1.9 mM and 0.48 $\mu M/min$, respectively, in the absence of leaf extracts, and the K_{mapp} and V_{max} values were calculated to be 30 mM and 0.37 $\mu M/min$, respectively, in the presence of 0.6 mg/mL of powdered leaf extract, and 18 mM and 0.45 $\mu M/min$, respectively, in the presence of 0.6

mg/mL of roasted leaf extract. Error bars indicate SD values. The average of triplicate determination is shown.

Fig. 3. Effects of 1-DNJ, fagomine, and GAL-DNJ on the hydrolysis of maltose by α -glucosidase.

Notes. (a, b) Dependence on the concentration of 1-DNJ, fagomine, and GAL-DNJ on the reaction rate. The reaction was carried out with 50 mM maltose and varying concentrations of 1-DNJ (open triangle) (a), fagomine (open square) (b), or GAL-DNJ (open diamond) (b) at pH 6.0 at 37°C. (c, d) Dependence on the substrate concentration of the reaction rate. The reaction was carried out with varying concentrations of maltose in the absence of iminosugar (open circle) or in the presence of 0.25 μ g/mL of 1-DNJ (closed triangle), 0.1 mg/mL of fagomine (closed square), or 1.5 mg/mL of GAL-DNJ (closed diamond) at pH 6.0 at 37°C. (d) v_o vs. [maltose]. Solid line represents the best fit to the experimental data using the Michaelis-Menten equation using Microsoft Excel. (d) Hanes-Woolf plot ([maltose] / v_o vs. [maltose]). The K_m and V_{max} values were calculated to be 1.9 mM and 0.48 μ M/min, respectively, in the absence of iminosugar, and the K_{mapp} and V_{max} values were calculated to be 9.0 mM and 0.44 μ M/min, respectively, in the presence of 0.25 μ g/mL of 1-DNJ, 16 mM and 0.44 μ M/min, respectively, in the presence of 0.1 mg/mL of fagomine, and 30 mM and 0.46 μ M/min, respectively, in the presence of 1.5 mg/mL of GAL-DNJ. Error bars indicate SD values. The average of triplicate determination is shown.

Fig. 4. Effects of reaction temperature and pH on the inhibitory effects of the extracts of *M. australis* powdered leaves and 1-DNJ toward the hydrolysis of maltose by α -

402 glucosidase.

403 Notes: (a) Dependence of K_i values at pH 6.0 on reaction temperature. The K_i
404 values at 27°C, 37°C, and 47°C are 0.21 ± 0.01 , 0.040 ± 0.002 , and 0.024 ± 0.003 mg/mL
405 for the extracts of powdered leaves and 0.22 ± 0.02 , 0.067 ± 0.001 , and 0.035 ± 0.008
406 $\mu\text{g/mL}$ for 1-DNJ. (b) Dependence of K_i values at 37°C on pH. The K_i values at pH 6, 7,
407 and 8 are 0.04 ± 0.01 , 0.049 ± 0.001 , and 0.18 ± 0.01 mg/mL for the extracts of powdered
408 leaves and 0.067 ± 0.001 , 0.098 ± 0.003 , and 0.30 ± 0.05 $\mu\text{g/mL}$ for 1-DNJ. (c) van't Hoff
409 plot of K_i values. K_i values were plotted against the reciprocal of the absolute temperature.
410 Enthalpy changes (ΔH°) were calculated from the slope to be -87.6 ± 6.4 kJ mol⁻¹ for the
411 extracts of powdered leaves and -72.8 ± 12.9 kJ mol⁻¹ for 1-DNJ. Error bars indicate SD
412 values. The average of triplicate determination is shown.

Fig. 1

Fig. 1

a

b

c

Fig. 2

Fig. 3

a

b

c

Fig. 4

a

b

c

Fig. S1. Product ion spectra of 1-DNJ, fagomine, and GAL-DNJ.

Notes: Standard 1-DNJ (5 μL in 1.0 $\mu\text{g}/\mu\text{L}$) (a), fagomine (5 μL in 1.0 $\mu\text{g}/\mu\text{L}$) (b), GAL-DNJ (5 μL in 1.0 $\mu\text{g}/\mu\text{L}$) (c) were applied. Product ion spectra of the ions at m/z 164.20 $[\text{M}+\text{H}]^+$ for 1-DNJ (a), at m/z 148.20 $[\text{M}+\text{H}]^+$ for fagomine (b), and at m/z 326.20 $[\text{M}+\text{H}]^+$ for GAL-DNJ (c) are shown.

Fig. S2. MRM chromatogram of 1-DNJ, fagomine, and GAL-DNJ

Notes: Standard 1-DNJ (5 μ L in 1.0 μ g/ μ L) (a), fagomine (5 μ L in 1.0 μ g/ μ L) (b), GAL-DNJ (5 μ L in 1.0 μ g/ μ L) (c) were applied. MRM chromatogram of the total of ions at m/z 69.20 $[M+H-95]^+$, 80.20 $[M+H-84]^+$, and 146.20 $[M+H-H_2O]^+$ for 1-DNJ (a), at 86.15 $[M+H-62]^+$, 112.15 $[M+H-2H_2O]^+$, and 130.15 $[M+H-H_2O]^+$ for fagomine (b), and at 61.10 $[M+H-265]^+$, 146.15 $[M+H-180]^+$, and 164.20 $[M+H-162]^+$ for GAL-DNJ (c) are shown.

a

b

c

Fig. S3. Calibration curves of 1-DNJ, fagomine, and GAL-DNJ.

Notes: Calibration curves of 1-DNJ (0–1000 ng/mL) (a), fagomine (0–1000 ng/mL) (b), and GAL-DNJ (0–1000 ng/mL) (c) are shown.

Fig. S4. Effects of reaction temperature and pH on the α -glucosidase activity.

Notes: (a) Temperature dependence. The activity at 62°C was defined as 100%. The reaction was carried out with 3.5 mg/mL extract from rat acetone powder, 50 mM maltose at pH 6 at 27–92°C. (b) pH dependence. The activity at pH 6 was defined as 100%. The reaction was carried out with 3.5 mg/mL extract, 50 mM maltose at pH 4–6 at 37°C.

Fig. S5. Effects of reaction temperature and pH on the inhibitory effects of the extracts of *M. australis* powdered leaves and 1-DNJ toward the hydrolysis of maltose by α -glucosidase.

Notes: (a–d) Dependence on the substrate concentration of the reaction rate. The reaction was carried out with varying concentrations of maltose in the absence of the extracts of *M. australis* leaves or 1-DNJ (open circle) or in the presence of 0.15 mg/mL powdered leaf extracts (open square) or 0.25 $\mu\text{g}/\text{mL}$ 1-DNJ (open triangle) at pH 6.0 at 27°C (a), at pH 6.0 at 47°C (b), at pH 7.0 at 37°C (c), and at pH 8.0 at 37°C (d). Solid line represents the best fit to the experimental data using the Michaelis-Menten equation with the nonlinear least-squares methods. Error bars indicate SD values.