TITLE:

Ni-Catalyzed Carboxylation of C(sp²)-S Bonds with CO: Evidence for the Multifaceted Role of Zn

AUTHOR(S):

Yanagi, Tomoyuki; Somerville, J. Rosie; Nogi, Keisuke; Martin, Ruben; Yorimitsu, Hideki

CITATION:

Yanagi, Tomoyuki ...[et al.], Ni-Catalyzed Carboxylation of C(sp²)-S Bonds with CO: Evidence for the Multifaceted Role of Zn. ACS Catalysis 2020, 10(3): 2117-2123

ISSUE DATE:

2020-02-07

URL:

<http://hdl.handle.net/2433/255855>

RIGHT:

This document is the Accepted Manuscript version of a Published Work that appeared in final form in ACS Catalysis, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <https://doi.org/10.1021/acscatal.9b05141>; The full-text file will be made open to the public on 15 January 2021 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。

Ni-Catalyzed Carboxylation of C(sp²)-S bonds with CO₂: Evidence for the Multifaceted Role of Zn

Tomoyuki Yanagi,^{† ⊥} Rosie J. Somerville,^{‡ § ⊥} Keisuke Nogi,[†] Ruben Martin*^{‡ ||} and Hideki Yori-
mitsu*[†]

[†] Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan

[‡] Institute of Chemical Research of Catalonia (ICIQ), The Barcelona Institute of Science and Technology, Av. Països Catalans 16, 43007 Tarragona, Spain

[§] Universitat Rovira i Virgili, Departament de Química Analítica i Química Orgànica, c/Marcel·lí Domingo, 1, 43007 Tarragona, Spain.

^{||} ICREA, Passeig Lluís Companys, 23, 08010 Barcelona, Spain

KEYWORDS: Carboxylation, Nickel, C-S cleavage, Carbon dioxide, Zinc.

ABSTRACT: Nickel-catalyzed reductive carboxylation reactions of aryl electrophiles typically require the use of metallic reducing agents. At present, the prevailing perception is that these serve as both a source of electrons and as sources of Lewis acids that may aid CO₂ insertion into the Ni-C bond. Herein, we provide evidence for the in situ formation of organometallic species from the metallic reductant, a step that has either been ruled out or has been unexplored in catalytic carboxylation reactions with metal powder reductants. Specifically, we demonstrate that Zn(0) acts as a reductant and that Zn(II) generates arylzinc species that might play a role in the C(sp²)-S carboxylation of arylsulfonium salts. Overall, the reductive Ni-catalyzed C(sp²)-S carboxylation reaction proceeds under mild conditions in a non-amide solvent, displays a wide substrate scope, and can be applied to a formal *para* C-H carboxylation of arenes.

Nickel-catalyzed reductive carboxylation reactions of organic (pseudo)halides with CO₂ have recently gained considerable momentum as routes to carboxylic acids, privileged molecules in a myriad of biologically relevant scaffolds (Scheme 1).^{1,2} Despite the advances realized, the intricacies of these reactions still remain speculative, probably due to the difficulty in accessing well-defined catalytically relevant species and the non-trivial roles exerted by the ligands and/or additives.³ Although Zn(0) and Mn(0) are typically employed in these endeavors as a source of electrons in order to access Ni(0)L_n and Ni(I)L_n species within the catalytic cycle,² Hazari has recently shown that Mn(II) salts derived from these reductions might also play a beneficial role in the carboxylation of aryl chlorides.^{3d,4} Overall, these observations underscore our limited mechanistic knowledge of carboxylation processes and suggest that a better understanding of the roles of the metallic reductants may in turn lay the foundation for improving these reactions.

Scheme 1. Metallic reductants in carboxylation reactions.

Scheme 2. Carboxylation of arylsulfonium salts.

Driven by the ready availability and synthetic versatility of organosulfur compounds, we wondered whether it would be possible to develop a Ni-catalyzed reductive carboxylation of arylsulfonium salts via C(sp²)-S cleavage (Scheme 2).⁵⁻⁸ The choice of this electrophile is not arbitrary.⁹ First, arylsulfonium salts are bench-stable compounds that are easily accessed on a large scale from simple aryl sulfides upon treatment with an appropriate electrophilic partner. Second, the neutral R₂S leaving group of the sulfonium moiety should not compete for binding at the nickel center.^{10,11} Furthermore, the electron-deficient nature of the arylsulfonium salt should facilitate oxidative addition at Ni(0)L_n.¹² If successful, we recognized that such a reaction would not only provide an unrealized opportunity to enable a catalytic C-S functionalization/CO₂ fixation, but would also offer the possibility to study mechanistic aspects of the reaction, such as the role of the metal reductant in the targeted carboxylation event. As part of our ongoing interest in catalytic carboxylations^{13,14} and C-S bond functionalization,^{7d,11b,f,g} we report herein the successful realization of a mild catalytic carboxylation of arylsulfonium salts. Importantly, we provide evidence for the intermediacy of a Ni(I) species and the involvement of organozinc species, thus revealing a multifaceted role played by

Zn that may have implications in related reductive cross-coupling endeavors.

Our investigations began by evaluating the catalytic carboxylation of **2a**, which is readily obtained on a multigram scale from the corresponding aryl sulfide **1a** by treatment with MeOTf. After some experimentation, we found that **2a** underwent carboxylation with CO₂ (1 atm) in the presence of a Ni(II) precatalyst bearing **L1** and Zn powder to give **3a** in 84% isolated yield at room temperature (Table 1, entry 1). Of particular note was the use of DMSO as the solvent (entries 8 and 9): our protocol is therefore a rare example of a Ni-catalyzed reductive carboxylation that avoids toxic amide solvents such as DMF, DMA, or NMP.^{3d,15} As anticipated, the nature of the ligand was critical to success, with phenanthroline ligands possessing substituents adjacent to the nitrogen atoms providing the best results (entries 2-6). Strikingly, Zn ($E^\circ = -0.76$ V vs SCE)¹⁶ was uniquely suited for the carboxylation of **2a** and no acid was obtained with stronger reductant Mn ($E^\circ = -1.32$ V vs SCE, entry 11).¹⁷ Likewise, the use of homogeneous reductants such as tetrakis(dimethylamino)ethylene (TDAE; $E^\circ = -0.62$ V vs SCE)¹⁶, DMAP-OED ($E^\circ = -1.24$ V vs SCE)¹⁸, or CoCp₂ ($E^\circ = -0.87$ V vs SCE)^{19,20} did not deliver **3a** (entries 12-14). Importantly, direct reduction of **2g** by Zn only gave a 3% yield of aryl sulfide **1g**, while reaction between DMAP-OED and **2f** gave a 60% yield of **1f**.²¹ The use of Mn did not result in the formation of **1a**. Together, these experiments suggest that Zn does not solely act as a reductant during the catalytic cycle and that other scenarios might come into play. Control experiments revealed that both Ni and Zn were critical for success (entry 15). In particular, we observed that *p*-tolylSMe did not undergo

carboxylation, ruling out the requirement for aryl sulfide intermediates during the carboxylation of arylsulfonium salts.

Table 1. Optimization of the Reaction Conditions.

entry	deviation from standard conditions	3 (%) ^b
1	none	87 (84) ^c
2	L2 instead of L1	48
3	L3 instead of L1	33
4	L4 instead of L1	56
5	L5 instead of L1	47
6	NiCl ₂ (dppf) instead of NiBr ₂ L1 / L1	3
7	Ni(COD) ₂ / L1 instead of NiBr ₂ L1	86
8	DMA (DMF) instead of DMSO	13 (28)
9	NMP as solvent	15
10	<i>p</i> -C ₆ H ₄ SMe (1a) instead of 2a	0
11	Mn instead of Zn	0
12	TDAE (3 equiv) instead of Zn	0 ^d
13	DMAP-OED (3 equiv) instead of Zn	0 ^{d,e}
14	CoCp ₂ (3 equiv) instead of Zn	0
15	no Zn or no Ni/ L1	0

^a **2a** (0.30 mmol), NiBr₂**L1** (2.5 mol %), **L1** (3.5 mol %), Zn powder (0.90 mmol), CO₂ (1 atm), DMSO (1.5 mL) at rt for 16 h followed by acidic workup. ^b NMR yields using 1,1,2,2-tetrabromoethane as internal standard. ^c Isolated yield. ^d A similar result was found with the addition of 1 equiv Zn(OTf)₂. ^e THF solvent.

Scheme 3. Scope of the Catalytic Carboxylation of Arylsulfonium Salts.

^a Isolated yield from arylsulfonium salts. Yields in one-pot carboxylation of aryl sulfides are shown in parentheses. ^b Yield was determined by NMR using 1,1,2,2-tetrabromoethane as an internal standard. ^c 5.0 mol % NiBr₂L1 and 7.0 mol % L1 was used. ^d 5% of **2b** was also formed. ^e Obtained as 4-acetylbenzoic acid after acidic workup.

With optimized conditions in hand, we investigated the generality of the reaction (Scheme 3). Overall, electron-neutral or electron-rich arylsulfonium salts showed good-to-excellent reactivity regardless of the counterion.^{21,22} Likewise, groups that are also reactive in Ni-catalyzed reactions – esters (**2k**, **2m**), aryl fluorides (**2e**), phosphonates (**2l**), ethers (**2f**, **2g**, **2h** and **2q**) and alkyl chlorides (**2q**) – were all well tolerated. We note that reactions of electron-deficient substrates (**2d**, **2j** and **2k**) were carried out with higher catalyst loadings due to competitive reductive demethylation by Zn.²¹ As expected, a substrate possessing an aryl methyl sulfide group was selectively carboxylated at the sulfonium moiety (**2o**). Surprisingly, the carboxylation reaction could be conducted in the presence of an organotin group (**2p**), providing ample room for further derivatization via orthogonal Stille-Migita-Kosugi cross-coupling reactions. Unfortunately, pyridyl sulfides could not be transformed to sulfonium salts with MeOTf due to preferential N-methylation. As shown for **2n**, the use of an *ortho*-substituted arylsulfonium salts did not hinder the reaction.²² Similar results were also obtained regardless of the alkyl substituents on the sulfonium moiety (**2t-2v**). Note, however, that non-negligible amounts of alkyl carboxylic acids were sometimes obtained via C(sp³)-S bond cleavage. Interestingly, the reaction can also be extended to diarylsulfonium salts (**2w**). For unsymmetrically substituted analogues (**2x**), carboxylation occurred preferentially at the most electron-deficient C-S bond. As demonstrated by the successful carboxylation of β -styrylsulfonium **2y**, C(sp²)-S carboxylation is not limited to arylsulfonium salts. Unfortunately, the reaction could not be extended to trialkylsulfonium salts (**2z**). Particularly noteworthy was the observation that the catalytic carboxylation of a variety of arylsulfonium salts can be carried out in a one-pot procedure from the parent aryl sulfide after simply removing volatiles upon completion of the reaction with MeOTf. The versatility and synthetic applicability of our protocol is further illustrated in Scheme 4. As shown, **6** can easily be obtained from **4** by telescoping the formation of the arylsulfonium salt **5** without isolation. Indeed, the successful preparation of **6** indicates that *sulfonium salts might be used as linchpins in formal para-C(sp²)-H carboxylation reactions*.²³ This is particularly important, as it might lead to new knowledge in synthetic design for incorporating CO₂ at remote C(sp²)-H bonds.

Scheme 4. Formal *para*-C(sp²)-H carboxylation of arenes.

Encouraged by the preparative aspects of our catalytic carboxylation reaction, we turned our attention to studying the mechanistic intricacies of the reaction. Interestingly, exposing **2g** to the optimized conditions *in the absence of CO₂* followed by a D₂O quench led to **7** in 55% yield and with >90% D incorporation (Scheme 5, *top*). This suggested a build-up of basic aryl organometallic species in solution.²⁴ Considering that *the Ni catalyst is required for the carboxylation to occur, the*

formation of 3 suggests that a Ni-to-Zn transmetalation might occur prior to CO₂ insertion.^{25,26} This notion was investigated by attempting the carboxylation of phenylzinc triflate with CO₂ in order to determine whether arylzinc species were competent reaction intermediates (Scheme 5, *bottom*). Low conversions to **3b** were observed in the absence of Ni(II) or Zn(0), and a high yield was only observed in the presence of *both* the Ni catalyst and Zn(0). Although Zn(OTf)₂ may promote carboxylation of PhZnOTf as a Lewis acid,²⁷ its addition did not result in an increased yield of **3b**. These results provide evidence that the carboxylation event is assisted by the Ni catalyst regardless of whether or not arylzinc species are formed.^{15a,27}

Scheme 5. Intermediacy of Arylzinc Species.

Aiming at shedding light on the origin of C(sp²)-S cleavage, we next turned our attention to investigating the formation of Ni(0)L_n from Ni(II)L_n prior to oxidative addition. In order to mimic the optimized catalytic carboxylation conditions, we monitored the reduction of NiBr₂L1 with Zn(0) using ¹H NMR spectroscopy. The formation of DMSO-ligated Ni(II) species was observed upon dissolving NiBr₂L1 in DMSO-*d*₆ and the formation of a DMSO adduct was confirmed by X-ray crystallography.²⁸ In the presence of Zn(0), NiBr₂L1 was transformed within 10 minutes into a deep blue paramagnetic Ni(I) complex in 74% yield (Scheme 6). X-ray crystallography identified this complex as [Ni(I)(L1)₂]Br, with the halide anion being outer-sphere.^{29,30} After longer reaction times, we observed that the yield [Ni(I)(L1)₂]Br had reduced to only 16%, suggesting the formation of further reduced Ni(0)L1 species. Unfortunately, these Ni(0) species could not be quantified, probably due to the very poor solubility of Ni(L1)₂ in DMSO.³¹ However, evidence for the presence of Ni(0) was obtained by addition of Ph₂S₂ or 1,5-cyclooctadiene to the reaction mixture, leading to the observation of oxidized products and COD-bearing complexes, respectively.³² These experiments provided indirect evidence for the formation of Ni(0) species from the overall reaction between NiBr₂L1 and Zn. This contrasts with a recent report by Diao where the formation of Ni(0) from a related Ni(bipyridine)/Zn system was ruled out.³³

Scheme 6. Formation of a Ni(I) Complex.

Prompted by the results shown in Scheme 6, we investigated the catalytic relevance of $[\text{Ni}(\text{I})(\text{L}1)_2]^+$ complexes in the carboxylation of arylsulfonium salts. Indeed, analyzing the catalytic carboxylation of **2f** by both NMR and EPR spectroscopy allowed us to detect the presence of $[\text{Ni}(\text{I})(\text{L}1)_2]\text{OTf}$ during catalysis. An aliquot removed after 1 h of reaction showed that of the initial catalyst loading (5 mol %), 60% of the Ni existed as $[\text{Ni}(\text{I})(\text{L}1)_2]\text{OTf}$. After 3.5 h this was 44%.³⁰ This finding is particularly relevant as it suggests that $[\text{Ni}(\text{L}1)_2]\text{OTf}$ may be a catalyst resting state. With evidence for both Ni(I) and Ni(0) intermediates in hand, stoichiometric experiments were carried out with $\text{Ni}(\text{L}1)_2$ in order to study oxidative addition of the $\text{C}(\text{sp}^2)\text{--}\text{S}$ bond (Scheme 7). First, reaction of **2f** with $\text{Ni}(\text{L}1)_2$ and $\text{Zn}(\text{OTf})_2$ – produced during every turnover of the catalyst – resulted in 50% conversion of **2f** and a 44% yield of Me_2S , with $[\text{Ni}(\text{L}1)_2]\text{OTf}$ (70%) being the only Ni species detected by ^1H NMR spectroscopy.³⁴ Interestingly, the presence of $\text{Zn}(0)$ led to full conversion of **2f** and an 80% yield of Me_2S , indicating that Zn could generate Ni(0) species from $[\text{Ni}(\text{L}1)_2]\text{OTf}$.

Overall, these experiments provide further evidence that the targeted $\text{C}(\text{sp}^2)\text{--}\text{S}$ bond cleavage occurs at Ni(0). Although the isolation of an oxidative addition complex of an arylsulfonium salt to $\text{Ni}(\text{L}1)_2$ proved to be elusive, we were able to track the fate of the aryl group after $\text{C}(\text{sp}^2)\text{--}\text{S}$ cleavage. Specifically, reactions between **2f** and $\text{Ni}(\text{L}1)_2$ carried out in the presence of $\text{Zn}(\text{OTf})_2$ revealed new ^1H NMR signals that were assigned to arylzinc species based on the identical ^1H NMR data obtained upon reaction of ZnAr_2 with $\text{Zn}(\text{OTf})_2/\text{L}1$ (Scheme 7).³⁵ Putting everything into perspective, we believe these experiments provide a strong argument for the formation of arylzinc species after $\text{C}(\text{sp}^2)\text{--}\text{S}$ cleavage by $\text{Ni}(0)\text{L}_n$.²⁶

Scheme 7. Stoichiometric Experiments with $\text{Ni}(\text{L}1)_2$.

■ Stoichiometric reactions ($\text{Ar} = p\text{-MeOC}_6\text{H}_4$)

2f + $\text{Ni}(\text{L}1)_2$	Zn(OTf) ₂ (2 equiv) Zn (x equiv) DMSO-d ₆		2f + Me ₂ S +		In situ arylzinc species "ArZn ⁺ " ← Zn(OTf) ₂ , L1 → Ar ₂ Zn
	x = 0	x = 3	50%	44%	
	traces	85%	40%	90%	

In light of our results, we propose the mechanistic scenario depicted in Scheme 8, which includes *Ni-to-Zn* transmetalation steps. At this stage, it is unclear whether the product of $\text{C}(\text{sp}^2)\text{--}\text{SMe}_2^+$ oxidative addition (**I**) undergoes transmetalation with $\text{Zn}(\text{II})$ to give Ni(II) complex **II** or whether **I** is first reduced to Ar-Ni(I) species **III** prior to transmetalation en route to **IV** and ArZnX . The latter may be more feasible when comparing the nucleophilicities of the nickel centers of **I** and **III**.^{3a,35} Whether **IV** indeed corresponds to $[\text{Ni}(\text{L}1)_2]\text{OTf}$ – identified in the catalytic carboxylation reaction – requires further investigations.^{36,37} After CO_2 insertion to Ar-Ni(I) species **III**, the resulting carboxylate complex **V** is either directly reduced to $\text{Ni}(0)\text{L}_n$ or undergoes anion exchange with ZnX_2 to form **IV** prior to reduction.^{3d,25} Given that carboxylation of PhZnOTf required the presence of both Ni and Zn for yields to surpass ca. 10%, we

suggest that ArZnX species may serve as a reservoir of the aryl group rather than as species at which significant amounts of carboxylation occurs. This observation is particularly relevant in the context of Ni-catalyzed cross-electrophile coupling reactions mediated by Zn, because these results contribute to the perception that the role of the metallic reductant in these endeavors might not be as straightforward as one might initially anticipate and that further investigations might be required to unravel the intricacies of these processes.³⁸

Scheme 8. Proposed Mechanistic Rationale.

In conclusion, we have developed a mild Ni-catalyzed reductive carboxylation of arylsulfonium salts that is mediated by Zn and offers reactivity complementary to that observed with aryl (pseudo)halides. Our studies have provided evidence for the formation of organozinc intermediates during the reaction, suggesting that Ni-to-Zn transmetalation is plausible and that a greater variety of mechanistic pathways may need to be studied depending on the aryl electrophile.

AUTHOR INFORMATION

Corresponding Author

*rmartinromo@iciq.es

*yori@kuchem.kyoto-u.ac.jp

Note

[†]T.Y. and R.J.S. contributed equally

Supporting Information

Experimental procedures, crystallographic data and spectral data (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

ACKNOWLEDGMENT

H.Y. thanks JSPS KAKENHI Grant Numbers JP16H04109, JP18H04254, JP18H04409, and JP19H00895. K.N. thanks JSPS KAKENHI Grand Number JP18K14212. T.Y. acknowledges a JSPS Predoctoral Fellowship and JSPS Overseas Challenge Program for Young Researchers. This work was partly supported by JST CREST Grant Number JPMJCR19R4, Japan. R.M. and R.J.S. thank ICIQ, FEDER/MICIU – AEI/PGC2018- 096839-B-100, and funding from “la Caixa” Foundation (ID 100010434) under the agreement LCF/BQ/SO15/52260010 for financial support. We also acknowledge the X-ray diffraction Research Support Area at ICIQ and Dr. Georgiana Stoica for her help with EPR studies. R.J.S. sincerely thanks “la Caixa” for a predoctoral fellowship.

REFERENCES

- (1) (a) Carbon Dioxide as Chemical Feedstock; Aresta, M., Ed.; Wiley-VCH: Weinheim, 2010. (b) Sakakura, T.; Choi, J.-C.; Yasuda, H. Transformation of Carbon Dioxide. *Chem. Rev.* **2007**, *107*, 2365–2387. (c) Artz, J.; Müller, T. E.; Thenert, K.; Kleinekorte, J.; Meys, R.; Sternberg, A.; Bardow, A.; Leitner, W. Sustainable Conversion of Carbon Dioxide: An Integrated Review of Catalysis and Life Cycle Assessment. *Chem. Rev.* **2018**, *118*, 434–504.
- (2) Selected reviews: (a) Cokoja, M.; Bruckmeier, D.-C. C.; Rieger, B.; Herrmann, W. A.; Kühn, F. E. Transformation of Carbon Dioxide with Homogeneous Transition-Metal Catalysts: A Molecular Solution to a Global Challenge? *Angew. Chem. Int. Ed.* **2011**, *50*, 8510–8537. (b) Huang, K.; Sun, C.-L.; Shi, Z.-J. Transition-metal-catalyzed C–C bond formation through the fixation of carbon dioxide. *Chem. Soc. Rev.* **2011**, *40*, 2435–2452. (c) Tortajada, A.; Juliá-Hernández, F.; Börjesson, M.; Moragas, T.; Martin, R. Transition Metal-Catalyzed Carboxylation Reactions with Carbon Dioxide. *Angew. Chem. Int. Ed.* **2018**, *57*, 15948–15982. (d) Burkart, M. D.; Hazari, N.; Tway, C. L.; Zeitler, E. L. Opportunities and Challenges for Catalysis in Carbon Dioxide Utilization. *ACS Catal.* **2019**, *9*, 7937–7956.
- (3) (a) Sayyed, F. B.; Tsuji, Y.; Sakaki, S. The Crucial Role of a Ni(I) Intermediate in Ni-Catalyzed Carboxylation of Aryl Chloride with CO₂: A Theoretical Study. *Chem. Commun.* **2013**, *49*, 10715. (b) Sayyed, F. B.; Sakaki, S. The Crucial Roles of MgCl₂ as a Non-Innocent Additive in the Ni-Catalyzed Carboxylation of Benzyl Halide with CO₂. *Chem. Commun.* **2014**, *50*, 13026–13029. (c) Obst, M.; Pavlovic, L.; Hopmann, K. H. Carbon-Carbon Bonds with CO₂: Insights from Computational Studies. *J. Organomet. Chem.* **2018**, *864*, 115–127. (d) Charboneau, D. J.; Brudvig, G. W.; Hazari, N.; Lant, H. M. C.; Saydjari, A. K. Development of an Improved System for the Carboxylation of Aryl Halides through Mechanistic Studies. *ACS Catal.* **2019**, *9*, 3228–3241. (e) Mohadjer Beromi, M.; Brudvig, G. W.; Hazari, N.; Lant, H. M. C.; Mercado, B. Q. Synthesis and Reactivity of Paramagnetic Nickel Polypyridyl Complexes Relevant to C(sp²)–C(sp³) Coupling Reactions. *Angew. Chem. Int. Ed.* **2019**, *58*, 6094–6098.
- (4) See ref. 3 and, for a specific focus on metallic reducing agents, see: Richmond, E.; Moran, J. Recent Advances in Nickel Catalysis Enabled by Stoichiometric Metallic Reducing Agents. *Synthesis* **2018**, *50*, 499–513.
- (5) (a) Ruano, J. L. G.; de la Plata, B. C. In *Organosulfur Chemistry I*; Page, P. C. B., Ed.; Springer: Heidelberg, 1999; p 1. (b) Furukara, N.; Sato, S. In *Organosulfur Chemistry II*; Page, P. C. B., Ed.; Springer: Heidelberg, 1999; p 89. (c) Rayner, C. M.; Advances in Sulfur Chemistry; JAI Press: Greenwich, 2000, Vol.2. (d) Sulfur Compounds: Advances in Research and Application; Acton, A. Q., Ed.; Scholarly Eds.: Atlanta, GA, 2012. (e) Feng, M.; Tang, B.; Liang, S. H.; Jiang, X. Sulfur Containing Scaffolds in Drugs: Synthesis and Application in Medicinal Chemistry. *Curr. Top. Med. Chem.* **2016**, *16*, 1200–1216.
- (6) Sun, S.; Yu, J.-T.; Cheng, J. Copper(I)-Catalyzed Desulfinate Carboxylation of Sodium Sulfinate using Carbon Dioxide. *Adv. Synth. Catal.* **2015**, *357*, 2022–2026.
- (7) Related carbonylation with CO gas via aromatic C–S bond cleavage: (a) Shim, S. C.; Alper, H. Desulfurization and carbonylation of mercaptans. *J. Org. Chem.* **1985**, *50*, 147–149. (b) Kamigata, N.; Satoh, A.; Yoshida, M.; Kameyama, M. The Reaction of Arylazo Aryl Sulfones with Carbon Monoxide and Nucleophiles Catalyzed by the Palladium(0) Complex. *Bull. Chem. Soc. Jpn.* **1989**, *62*, 605–607. (c) Li, X.; Liang, D.; Huang, W.; Zhou, H.; Li, Z.; Wang, B.; Ma, Y.; Wang, H. Visible light-induced carbonylation of indoles with arylsulfonyl chlorides and CO. *Tetrahedron* **2016**, *72*, 8442–8448. (d) Minami, H.; Nogi, K.; Yorimitsu, H. Palladium-Catalyzed Alkoxy-carboxylation of Arylsulfoniums. *Org. Lett.* **2019**, *21*, 2518–2522.
- (8) Reviews: (a) Modha, S. G.; Mehta, V. P.; Van der Eycken, E. V. Transition metal-catalyzed C–C bond formation via C–S bond cleavage. *Chem. Soc. Rev.* **2013**, *42*, 5042–5055. (b) Pan, F.; Shi, Z. J. Recent Advance in Transition-Metal-Catalyzed C–S Activation: From Thioester to (Hetero)aryl Thioether. *ACS Catal.* **2014**, *4*, 280–288. (c) Gao, K.; Otsuka, S.; Baralle, A.; Nogi, K.; Yorimitsu, H.; Otsuka, A. Cross-coupling of Aryl Sulfides Powered by N-Heterocyclic Carbene Ligands. *J. Synth. Org. Chem., Jpn.* **2016**, *74*, 1119–1127. (d) Otsuka, S.; Nogi, K.; Yorimitsu, H. C–S Bond Activation. *Top. Curr. Chem.* **2018**, *376*, 13.
- (9) Pioneering report: Srogl, J.; Allred, G. D.; Liebeskind, L. S. Sulfonium Salts. Participants par Excellence in Metal-Catalyzed Carbon–Carbon Bond-Forming Reactions. *J. Am. Chem. Soc.* **1997**, *119*, 12376–12377.
- (10) Reviews on cross-coupling reactions of arylsulfonium salts: (a) Tian, Z.-Y.; Hu, Y.-T.; Teng, H.-B.; Zhang, C.-P. Application of arylsulfonium salts as arylation reagents. *Tetrahedron Lett.* **2018**, *59*, 299–309. (b) Kaiser, D.; Klose, I.; Oost, R.; Neuhaus, J.; Maulide, N. Bond-Forming and -Breaking Reactions at Sulfur(IV): Sulfoxides, Sulfonium Salts, Sulfur Ylides, and Sulfinate Salts. *Chem. Rev.* **2019**, *119*, 8701–8780.
- (11) Selected recent examples: (a) Cowper, P.; Jin, Y.; Turton, M. D.; Kociok-Köhn, G.; Lewis, S. E. Azulenensulfonium Salts: Accessible, Stable, and Versatile Reagents for Cross-Coupling. *Angew. Chem. Int. Ed.* **2016**, *55*, 2564–2568. (b) Kawashima, H.; Yanagi, T.; Wu, C.-C.; Nogi, K.; Yorimitsu, H. Regioselective C–H Sulfanylation of Aryl Sulfoxides by Means of Pummerer-Type Activation. *Org. Lett.* **2017**, *19*, 4552–4555. (c) Tian, Z.-Y.; Wang, S.-M.; Jia, S.-J.; Song, H.-X.; Zhang, C.-P. Sonogashira Reaction Using Arylsulfonium Salts as Cross-Coupling Partners. *Org. Lett.* **2017**, *19*, 5454–5457. (d) Hock, K. J.; Hommelsheim, R.; Mertens, L.; Ho, J.; Nguyem, T. V.; Koenig, R. M. Corey-Chaykovsky Reactions of Nitro Styrenes Enable *cis*-Configured Trifluoromethyl Cyclopropanes. *J. Org. Chem.* **2017**, *82*, 8220–8227. (e) Aukland, M. H.; Talbot, F. J. T.; Fernández-Salas, J. A.; Ball, M.; Pulis, A. P.; Procter, D. J. An Interrupted Pummerer/Nickel-Catalyzed Cross-Coupling Sequence. *Angew. Chem. Int. Ed.* **2018**, *57*, 9785–9789. (f) Uno, D.; Minami, H.; Otsuka, S.; Nogi, K.; Yorimitsu, H. Palladium-Catalyzed Mizoroki-Heck-Type Alkenylation of Mono-aryldialkylsulfoniums. *Chem. Asian J.* **2018**, *13*, 2397–2400. (g) Minami, H.; Otsuka, S.; Nogi, K.; Yorimitsu, H. Palladium-Catalyzed Borylation of Arylsulfoniums with Diborons. *ACS Catal.* **2018**, *8*, 579–583. (h) Tian, Z.-Y.; Zhang, C.-P. Ullmann-type N-arylation of anilines with alkyl(aryl)sulfonium salts. *Chem. Commun.* **2019**, *55*, 11936–11939.
- (12) We reported the carboxylation of benzylamines via inert C–N bond activation by conversion to the corresponding ammonium salts: Moragas, T.; Gaydou, M.; Martin, R. Nickel-Catalyzed Carboxylation of Benzylic C–N bond with CO₂. *Angew. Chem. Int. Ed.* **2016**, *55*, 5053–5057.
- (13) Selected examples: (a) Correa, A.; Martin, R. Palladium-Catalyzed Direct Carboxylation of Aryl Bromides with Carbon Dioxide. *J. Am. Chem. Soc.* **2009**, *131*, 15974–15975. (b) Börjesson, M.; Moragas, T.; Martin, R. Ni-Catalyzed Carboxylation of Unactivated Alkyl Chlorides with CO₂. *J. Am. Soc. Chem.* **2016**, *138*, 7504–7507. (c) Juliá-Hernández, F.; Moragas, T.; Cornella, J.; Martin, R. Remote Carboxylation of Halogenated Aliphatic Hydrocarbons with Carbon Dioxide. *Nature* **2017**, *545*, 84–88.
- (14) Selected examples from other groups: (a) Fujihara, T.; Nogi, K.; Xu, T.; Terao, J.; Tsuji, Y. Nickel-Catalyzed Carboxylation of Aryl and Vinyl Chlorides Employing Carbon Dioxide. *J. Am. Chem. Soc.* **2012**, *134*, 9106–9109. (b) Meng, Q.-Y.; Wang, S.; König, B. Carboxylation of Aromatic and Aliphatic Bromides and Triflates with CO₂ by Dual Visible - Light - Nickel Catalysis. *Angew. Chem. Int. Ed.* **2017**, *56*, 13426–13430. (c) Gui, Y.-Y.; Hu, N.; Chen, X.-W.; Liao, L.-L.; Ju, T.; Ye, J.-H.; Zang, Z.; Li, J.; Yu, D.-G. Highly Regio- and Enantioselective Copper-Catalyzed Reductive Hydroxymethylation of Styrenes and 1,3-Dienes with CO₂. *J. Am. Chem. Soc.* **2017**, *139*, 17011–17014. (d) Hou, J.; Ee, A.; Cao, H.; Ong, H.-W.; Xu, J.-H.; Wu, J. Visible-Light-Mediated Metal - Free Difunctionalization of Alkenes with CO₂ and Silanes or C(sp³)-H Alkanes. *Angew. Chem. Int. Ed.* **2018**, *57*, 17220–17224. (e) Ma, C.; Zhao, C.-Q.; Xu, X.-T.; Li, Z.-M.; Wang, X.-Y.; Zhang, K.; Mei, T. S. Nickel-Catalyzed Carboxylation of Aryl and Heteroaryl Fluorosulfates Using Carbon Dioxide. *Org. Lett.* **2019**, *21*, 2464–2467.
- (15) (a) Williams, C. M.; Johnson, J. B.; Rovis, T. Nickel-Catalyzed Reductive Carboxylation of Styrenes Using CO₂. *J. Am. Chem. Soc.* **2008**, *130*, 14936–14937. (b) See ref. 23a–b for Ni-catalyzed carboxylation reactions of organozinc species in THF and DME. (c) For a Cu-

- catalyzed carboxylation reaction carried out in DMSO, see: Tran-Vu, H.; Daugulis, O. Copper-Catalyzed Carboxylation of Aryl Iodides with Carbon Dioxide. *ACS Catal.* **2013**, *3*, 2417–2420. (d) For an example of a Pd-catalyzed carboxylation conducted in diglyme: Sasano, K.; Takaya, J.; Iwasawa, N. Palladium(II)-Catalyzed Direct Carboxylation of Alkenyl C–H Bonds with CO₂. *J. Am. Chem. Soc.* **2013**, *135*, 10954–10957. (e) For an example of a redox-neutral Rh-catalyzed carboxylation in dioxane: Ukai, K.; Aoki, M.; Takaya, J.; Iwasawa, N. Rhodium(I)-Catalyzed Carboxylation of Aryl- and Alkenylboronic Esters with CO₂. *J. Am. Chem. Soc.* **2006**, *128*, 8706–8707.
- (16) (a) Wiberg, N. Tetraaminoäthylene Als Starke Elektronendonoren. *Angew. Chem.* **1968**, *80*, 809–856. (b) Broggi, J.; Terme, T.; Vanelle, P. Organic Electron Donors as Powerful Single-Electron Reducing Agents in Organic Synthesis. *Angew. Chem. Int. Ed.* **2014**, *53*, 384–413.
- (17) Birbilis, N.; Buchheit, R. G. Electrochemical Characteristics of Intermetallic Phases in Aluminum Alloys: An Experimental Survey and Discussion. *J. Electrochem. Soc.* **2005**, *152*, 140–151.
- (18) (a) Murphy, J. A.; Gamier, J.; Park, S. R.; Schoenebeck, F.; Zhou, S. Z.; Turner, A. T. Super-Electron Donors: Bis-Pyridinylidene Formation by Base Treatment of Pyridinium Salts. *Org. Lett.* **2008**, *10*, 1227–1230. (b) Garnier, J.; Kennedy, A. R.; Berlouis, L. E. A.; Turner, A. T.; Murphy, J. A. Structure and Reactivity in Neutral Organic Electron Donors Derived from 4-Dimethylaminopyridine. *Beilstein J. Org. Chem.* **2010**, *6*, 4–11.
- (19) Organometallics and Catalysis: An Introduction; Bochmann, M., Ed.; Oxford University Press: Oxford, 2015; p 189.
- (20) Use of 3.0 equiv CoCp₂ with 1.0 equiv Zn(OTf)₂ in the carboxylation of sulfonium salt **2g** gave **3g** in 18% yield. See the Supporting Information for further information.
- (21) For single electron reduction of dialkylarylsulfonium salts, see: Beak, P.; Sullivan, T. A. One-electron chemical reductions of phenylalkylsulfonium salts. *J. Am. Chem. Soc.* **1982**, *104*, 4450.
- (22) For unsuccessful and low-yielding substrates, see the Supporting Information Scheme S2.
- (23) Selected examples of catalytic C–H carboxylation: (a) Mizuno, H.; Takaya, J.; Iwasawa, N. Rhodium(I)-Catalyzed Direct Carboxylation of Arenes with CO₂ via Chelation-Assisted C–H Bond Activation. *J. Am. Chem. Soc.* **2011**, *133*, 1251–1253. (b) Boogaerts, I. I. F.; Nolan, S. P. Carboxylation of C–H Bonds Using N-Heterocyclic Carbene Gold(I) Complexes. *J. Am. Chem. Soc.* **2010**, *132*, 8858–8859. (c) Luo, J.; Larrosa, I. C–H Bond Carboxylation with Carbon Dioxide. *ChemSusChem* **2017**, *10*, 3317–3332.
- (24) For deuterium or acidic quenching experiments in the context of reductive carboxylations involving organozinc reagents, see ref. 15 and: (a) Nogi, K.; Fujihara, T.; Terao, J.; Tsuji, Y. Carboxyzincation Employing Carbon Dioxide and Zinc Powder: Cobalt-Catalyzed Multicomponent Coupling Reactions with Alkynes. *J. Am. Chem. Soc.* **2016**, *138*, 5547–5550. For activation of C–SMe bonds by Zn: (b) Begouin, J.-M.; Rivard, M.; Gosmini, C. Cobalt-Catalyzed C–SMe Bond Activation of Heteroaromatic Thioethers. *Chem. Commun.* **2010**, *46*, 5972–5974.
- (25) (a) Conan, A.; Sibille, S.; Périchon, J. Metal Exchange between an Electrogenenerated Organonickel Species and Zinc Halide: Application to an Electrochemical, Nickel-Catalyzed Reformatsky Reaction. *J. Org. Chem.* **1991**, *56*, 2018–2024. (b) Sibille, S.; Ratovelomanana, V.; Périchon, J. Electrochemical Conversion of Functionalised Aryl Chlorides and Bromides to Arylzinc Species. *J. Chem. Soc., Chem. Commun.* **1992**, 283–284.
- (26) For Ni-catalyzed carbozincation, see: (a) Stüdemann, T.; Knochel, P. New Nickel-Catalyzed Carbozincation of Alkynes: A Short Synthesis of (Z)-Tamoxifen. *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 93–95. (b) Stüdemann, T.; Ibrahim-Ouali, M.; Knochel, P. A Nickel-Catalyzed Carbozincation of Aryl-Substituted Alkynes. *Tetrahedron* **1998**, *54*, 1299–1316. For reviews of carbozincation, see: Murakami, K.; Yorimitsu, Y. Recent Advances in Transition-Metal-Catalyzed Intermolecular Carbomagnesiation and Carbozincation. *Beilstein J. Org. Chem.* **2013**, *9*, 278–302.
- (27) (a) Yeung, C. S.; Dong, V. M. Beyond Aresta's Complex: Ni- and Pd-Catalyzed Organozinc Coupling with CO₂. *J. Am. Chem. Soc.* **2008**, *130*, 7826–7827. (b) Ochiai, H.; Jang, M.; Hirano, K.; Yorimitsu, H.; Oshima, K. Nickel-Catalyzed Carboxylation of Organozinc Reagents with CO₂. *Org. Lett.* **2008**, *10*, 2681–2683. (c) Kobayashi, K.; Kondo, Y. Transition-Metal-Free Carboxylation of Organozinc Reagents Using CO₂ in DMF Solvent. *Org. Lett.* **2009**, *11*, 2035–2037. (d) Metzger, A.; Bernhardt, S.; Manolikakes, G.; Knochel, P. MgCl₂-Accelerated Addition of Functionalized Organozinc Reagents to Aldehydes, Ketones, and Carbon Dioxide. *Angew. Chem. Int. Ed.* **2010**, *49*, 4665–4668.
- (28) For similar reactivity in an NiCl₂(**L3**) complex, see: Brewer, B.; Brooks, N. R.; Sykes, A. G. Characterization of an Asymmetric M(chelate)₂(μ-Cl)₂ MCl₂ Dimer and Isolation of Corresponding DMSO Adducts: X-Ray Crystal Structures of Co(phen)₂(μ-Cl)₂CoCl₂·C₄H₈O and M(chelate)Cl₂(DMSO)_x (Co, X = 1; Ni, X = 2) Complexes. *J. Chem. Crystallogr.* **2003**, *33* (9), 663–668.
- (29) Bond distances between the cationic Ni(I) centre and **L1** and spectroscopic features (NMR, EPR) were comparable to those of analogous complexes previously reported in the literature: ref. 3(e) and: Powers, D. C.; Anderson, B. L.; Nocera, D. G. Two-Electron HCl to H₂ Photocycle Promoted by Ni(II) Polypyridyl Halide Complexes. *J. Am. Chem. Soc.* **2013**, *135*, 18876–18883.
- (30) The Ni(I) complexes [Ni(I)(**L1**)₂]Br and [Ni(I)(**L1**)₂]OTf were synthesized by oxidation of Ni(**L1**)₂ with the corresponding AgX salt as reported in ref. 29.
- (31) Oxidation states in NiL₂ complexes have been studied in: Wang, M.; England, J.; Weyhermüller, T.; Wieghardt, K. Electronic Structures of “Low-Valent” Neutral Complexes [NiL₂]⁰ (S = 0; L = bpy, phen, tpy) – An Experimental and DFT Computational Study. *Eur. J. Inorg. Chem.* **2015**, 1511–1523.
- (32) See Supporting Information for details.
- (33) Lin, Q.; Diao, T. Mechanism of Ni-Catalyzed Reductive 1,2-Dicarbofunctionalization of Alkenes. *J. Am. Chem. Soc.* **2019**, *141*, 17937–17948.
- (34) In the absence of Zn and Zn(II), this reaction gave a very low conversion of **2f** (**9f**) to **1f** (**6f**), Me₂S (**5f**), biaryl (**2f**), and [Ni(**L1**)₂]OTf (**8f**).
- (35) In the electrochemical zincation of aryl halides under cobalt catalysis, one-electron reduction of ArCo(III) to ArCo(II) prior to Co-to-Zn transmetalation was proposed. See: Seka, S.; Buriez, O.; Nédélec, J.-Y.; Périchon, J. Mechanism of the Electrochemical Conversion of Aryl Halides to Arylzinc Compounds by Cobalt Catalysis in DMF/Pyridine. *Chem. Eur. J.* **2002**, *8*, 2534–2538.
- (36) The majority of Ni(I) halide complexes bearing phenanthroline ligands have been found to have an outer sphere counteranion and a [NiL₂]X formula. See references 3(e) and 29. Diao recently reported a Ni(I) complex LNiBr where only one ligand is coordinated to Ni and the bromide ligand is bound directly to Ni (ref. 33).
- (37) Recent work by Diao has shown that insertion into a Ni(I)-alkyl bond is more rapid than insertion into the analogous Ni(II)-alkyl bond: Dicciani, J. B.; Hu, C. T.; Diao, T. Insertion of CO₂ Mediated by a (Xantphos)Ni(II)-Alkyl Species. *Angew. Chem. Int. Ed.* **2019**, *58*, 13865–13868.
- (38) For example, see the following reference for a discussion of Zn insertion during a cross-electrophile coupling between aryl bromides and aryl chlorides with alkyl bromides: Everson, D. A.; Jones, B. A.; Weix, D. J. Replacing Conventional Carbon Nucleophiles with Electrophiles: Nickel-Catalyzed Reductive Alkylation of Aryl Bromides and Chlorides. *J. Am. Chem. Soc.* **2012**, *134*, 6146–6159.