

Title	Demonstrations of geometric algebra for mathematicians (Innovative Teaching of Mathematics with Geometric Algebra)
Author(s)	Ishi, Hideyuki
Citation	数理解析研究所講究録 (2004), 1378: 1-6
Issue Date	2004-05
URL	http://hdl.handle.net/2433/25642
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Demonstrations of geometric algebra for mathematicians

伊師 英之 (Hideyuki ISHI)
横浜市立大学大学院 総合理学研究科
Department of mathematical sciences,
Yokohama City University

§ Introduction. The notion of the Clifford algebra is widely known among mathematicians. But it is hardly known (particularly in Japan) that recently some people study this algebra with great enthusiasm, and a large number of applications to physics and engineering have been developed. The members of the research groups call their own object *geometric algebra* rather than Clifford algebra because of its “geometric nature”, returning the original terminology by Clifford himself. One of their key concepts is to consider the Grassmann algebra inside the Clifford algebra. In this article, we present briefly what we can do with this idea in elementary linear algebra. Our main reference is [2, Section 1].

§1. Preliminary. Let V be an n -dimensional Euclidean space, and $C(V)$ the Clifford algebra of V . Namely, $C(V)$ is the associative algebra generated by V together with the real number field \mathbb{R} with a relation $x^2 = \|x\|^2$ ($x \in V$). Then $(xy + yx)/2$ equals the scalar product $(x|y)$ for vectors $x, y \in V$. Thus, if x and y are orthogonal, we have $xy = -yx$.

For $v_1, v_2, \dots, v_r \in V$, we define their *outer product* by

$$(1) \quad v_1 \wedge v_2 \wedge \dots \wedge v_r := \frac{1}{r!} \sum_{\sigma \in \mathfrak{S}_r} (\text{sgn } \sigma) v_{\sigma(1)} v_{\sigma(2)} \dots v_{\sigma(r)} \in C(V)$$

and call such an element of $C(V)$ a *simple r -vector* or an *r -blade*. A linear combination of simple r -vectors is called an *r -vector*, and the set of r -vectors is denoted by $C^r(V)$. Like the ordinary exterior product of the Grassmann algebra, the outer product is anti-commutative. If the vectors v_1, \dots, v_r are mutually orthogonal, the outer product equals the Clifford product: $v_1 \wedge v_2 \wedge \dots \wedge v_r = v_1 v_2 \dots v_r$ because $(\text{sgn } \sigma) v_{\sigma(1)} v_{\sigma(2)} \dots v_{\sigma(r)} = v_1 v_2 \dots v_r$ for each $\sigma \in \mathfrak{S}_r$ in this case.

Let $\{e_1, \dots, e_n\}$ be an orthonormal basis of V . For a set $I = \{i_1, i_2, \dots, i_r\}$ ($1 \leq i_1 < i_2 < \dots < i_r \leq n$), we write e_I for the product $e_{i_1} e_{i_2} \dots e_{i_n} \in C(V)$, and for

$I = \emptyset$ we put $e_I := 1 \in C(V)$. Then $\{e_I\}_{I \subset \{1, \dots, n\}}$ is a basis of $C(V)$ as a linear space. Since e_I is equal to the outer product $e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_r}$, we have $C^r(V) = \sum_{\#I=r}^{\oplus} \mathbb{R}e_I$, so that we get a direct sum decomposition of $C(V)$ as a linear space:

$$(2) \quad C(V) = \mathbb{R} \oplus V \oplus C^2(V) \oplus \dots \oplus C^n(V).$$

Let $\dagger : C(V) \rightarrow C(V)$ be the canonical anti-automorphism (“reverse”), that is, the linear map on $C(V)$ with the properties

$$1^\dagger = 1, \quad v^\dagger = v \quad (v \in V), \quad (AB)^\dagger = B^\dagger A^\dagger \quad (A, B \in C(V)).$$

For elements A, B of $C(V)$, we denote by $(A|B)$ the scalar part of $AB^\dagger \in C(V)$ with respect to the decomposition (2). If $A = \sum_I a_I e_I$ and $B = \sum_I b_I e_I$, we have $(A|B) = \sum_I a_I b_I$, so that $(\cdot|\cdot)$ defines a positive definite scalar product on $C(V)$ extending the original scalar product on V . For $A \in C(V)$, we write $\|A\|$ for the Euclidean norm $(A|A)^{1/2}$ of A .

§2. Inverse of a simple r vector. In what follows, we assume that vectors $v_1, v_2, \dots, v_r \in V$ are linearly independent, which is equivalent to $v_1 \wedge v_2 \wedge \dots \wedge v_r \neq 0$.

Proposition 1. (i) *The norm $\|v_1 \wedge v_2 \wedge \dots \wedge v_r\|$ gives the volume of the parallelotope spanned by v_1, v_2, \dots, v_r .*

(ii) *The element $A := v_1 \wedge v_2 \wedge \dots \wedge v_r$ is invertible as an element of $C(V)$. Indeed, one has*

$$A^{-1} = \frac{1}{\|A\|^2} A^\dagger = \frac{1}{\|v_1 \wedge v_2 \wedge \dots \wedge v_r\|^2} v_r \wedge \dots \wedge v_2 \wedge v_1.$$

For the proof of Proposition 1, we state the following lemma, which is equivalent to the validity of the Gram-Schmidt orthogonalization (see Proposition 3 (ii)).

Lemma 2. *For linearly independent vectors v_1, v_2, \dots, v_r , we can take a unique family of coefficients $c_{ij} \in \mathbb{R}$ ($i < j$) for which the vectors*

$$(3) \quad u_1 := v_1, \quad u_j := v_j + \sum_{i=1}^{j-1} c_{ij} v_i \quad (j = 2, \dots, r)$$

are mutually orthogonal.

Now let us prove Proposition 1. From (3) and the orthogonality of u_1, \dots, u_r , we see that

$$(4) \quad A = v_1 \wedge v_2 \wedge \dots \wedge v_r = u_1 \wedge u_2 \wedge \dots \wedge u_r = u_1 u_2 \dots u_r.$$

Then we have

$$AA^\dagger = (u_1 u_2 \cdots u_r)(u_r \cdots u_2 u_1) = \|u_1\|^2 \|u_2\|^2 \cdots \|u_r\|^2.$$

Since AA^\dagger belongs to \mathbb{R} , we have $AA^\dagger = \|A\|^2$. Thus $\|A\| = \|u_1\| \|u_2\| \cdots \|u_r\|$, which implies the assertion (i). On the other hand, we observe

$$A^\dagger A = (u_r \cdots u_2 u_1)(u_1 u_2 \cdots u_r) = \|u_1\|^2 \|u_2\|^2 \cdots \|u_r\|^2 = \|A\|^2.$$

Therefore $A(\frac{1}{\|A\|^2} A^\dagger) = (\frac{1}{\|A\|^2} A^\dagger)A = 1$, which means the assertion (ii). \square

As we shall see below, it is a great advantage over the ordinary Grassmann algebra that we can take the inverse of the simple r -vector $v_1 \wedge \cdots \wedge v_r$ in the Clifford algebra. The argument in the proof of Proposition 1 tells us that the formula $X^{-1} = X^\dagger / \|X\|^2$ holds for any Clifford product $X = a_1 a_2 \cdots a_r \in C(V)$ of non-zero vectors $a_1, \dots, a_r \in V$. However this is not true for a general non-zero element of $C(V)$. For example, since $(e_1 e_2 + e_3 e_4)(e_1 e_2 - e_3 e_4) = 0$, neither $e_1 e_2 + e_3 e_4$ nor $e_1 e_2 - e_3 e_4$ is invertible.

§3. Gram-Schmidt orthogonalization. The vectors u_1, \dots, u_r in Lemma 2 are given by a simple formula.

Proposition 3. (i) $u_k = (v_1 \wedge \cdots \wedge v_{k-1})^{-1}(v_1 \wedge \cdots \wedge v_k)$ ($k = 2, \dots, r$).

(ii) If one puts $\tilde{e}_k := u_k / \|u_k\|$ ($k = 1, \dots, r$), then $\{\tilde{e}_1, \dots, \tilde{e}_r\}$ is nothing but the orthonormal family obtained by the Gram-Schmidt orthogonalization from $\{v_1, \dots, v_r\}$.

Proof. Similarly to (4), we have for $k = 2, \dots, r$,

$$\begin{aligned} v_1 \wedge \cdots \wedge v_{k-1} \wedge v_k &= u_1 \wedge \cdots \wedge u_{k-1} \wedge u_k = (u_1 \cdots u_{k-1})u_k \\ &= (v_1 \wedge \cdots \wedge v_{k-1})u_k, \end{aligned}$$

whence (i) follows. The assertion (ii) follows from (3). \square

§4. Reflection and projection. We denote by $V(v_1, \dots, v_r)$ the subspace of V spanned by the vectors v_1, \dots, v_r , and by $V(v_1, \dots, v_r)^\perp$ its orthogonal complement in V . We note that any vector $x \in V$ is decomposed uniquely as $x = x_\parallel + x_\perp$ with $x_\parallel \in V(v_1, \dots, v_r)$ and $x_\perp \in V(v_1, \dots, v_r)^\perp$. As in Proposition 1, we put $A := v_1 \wedge v_2 \wedge \cdots \wedge v_r$.

Proposition 4. *One has $(-1)^r Ax A^{-1} = -x_{\parallel} + x_{\perp}$. In other words, the linear map $R_A : V \ni x \mapsto (-1)^r Ax A^{-1} \in V$ gives the reflection with respect to the space $V(v_1, \dots, v_r)^{\perp}$.*

For the case $r = 1$, Proposition 4 means a well-known description of the reflection with respect to a hyperplane.

Proof. We recall again the vectors u_1, \dots, u_r in Lemma 2. If $x = x_{\perp} \in V(v_1, \dots, v_r)^{\perp}$, then we have $xu_k = -u_k x$ ($k = 1, \dots, r$), so that

$$\begin{aligned} xA &= xu_1u_2 \cdots u_r = -u_1xu_2 \cdots u_r = (-1)^2 u_1u_2xu_3 \cdots u_r = \dots \\ &= (-1)^r u_1u_2 \cdots u_r x = (-1)^r Ax. \end{aligned}$$

Thus we obtain $R_A(x) = (-1)^r Ax A^{-1} = x$ in the case $x = x_{\perp}$. On the other hand, if $x = x_{\parallel} \in V(v_1, \dots, v_r)$ and $x \neq 0$, we can take vectors $y_1, \dots, y_{r-1} \in V(v_1, \dots, v_r)$ for which $\{x, y_1, \dots, y_{r-1}\}$ forms an orthogonal basis of $V(v_1, \dots, v_r)$. Then we have

$$v_1 \wedge v_2 \wedge \cdots \wedge v_r = c_0 x \wedge y_1 \wedge \dots \wedge y_{r-1} = c_0 xy_1 \cdots y_{r-1}$$

for some scalar $c_0 \in \mathbb{R}$. Since $xy_k = -y_k x$ ($k = 1, \dots, r-1$), we observe

$$\begin{aligned} xA &= x(c_0 xy_1y_2 \cdots y_{r-1}) = -c_0 xy_1xy_2 \cdots y_{r-1} = (-1)^2 c_0 xy_1y_2xy_3 \cdots y_{r-1} = \dots \\ &= (-1)^{r-1} c_0 xy_1y_2 \cdots y_{r-1} x = (-1)^{r-1} Ax. \end{aligned}$$

Thus we obtain $R_A(x) = (-1)^r Ax A^{-1} = -x$ in the case $x = x_{\parallel} \neq 0$. But this is valid also for $x = 0$. Therefore, for general $x = x_{\parallel} + x_{\perp} \in V$, we have $R_A(x) = R_A(x_{\parallel}) + R_A(x_{\perp}) = -x_{\parallel} + x_{\perp}$. \square

Since the map $C(V) \ni X \mapsto AXA^{-1} \in C(V)$ is an algebra automorphism on $C(V)$, we have for vectors $x_1, x_2, \dots, x_s \in V$,

$$Ax_1A^{-1} \wedge Ax_2A^{-1} \wedge \cdots \wedge Ax_sA^{-1} = A(x_1 \wedge x_2 \wedge \cdots \wedge x_s)A^{-1}.$$

Thus, the reflection of the s -vector $x_1 \wedge \cdots \wedge x_s$ is given by the following formula:

Lemma 5. $R_A(x_1) \wedge R_A(x_2) \wedge \cdots \wedge R_A(x_s) = (-1)^{rs} A(x_1 \wedge x_2 \wedge \cdots \wedge x_s)A^{-1}$.

From Proposition 4, we obtain the following results.

Proposition 6. *One has $x_{\parallel} = (x - (-1)^r Ax A^{-1})/2$ and $x_{\perp} = (x + (-1)^r Ax A^{-1})/2$.*

Proposition 7. (i) $V(v_1, \dots, v_r) = \{x \in V; xA = (-1)^{r-1}Ax\}$.

(ii) $V(v_1, \dots, v_r)^{\perp} = \{x \in V; xA = (-1)^r Ax\}$.

§5. Inner and outer product of a vector with an r -vector. For a vector $x \in V$ and an r -vector B , we define

$$x \rfloor B := (xB - (-1)^r Bx)/2, \quad x \wedge B := (xB + (-1)^r Bx)/2.$$

As in the previous sections, we assume that the vectors v_1, \dots, v_r are linearly independent, and that $x = x_{\parallel} + x_{\perp}$ ($x_{\parallel} \in V(v_1, \dots, v_r)$, $x_{\perp} \in V(v_1, \dots, v_r)^{\perp}$), $A = v_1 \wedge \dots \wedge v_r = u_1 \cdots u_r$.

Proposition 8. (i) *One has*

$$(5) \quad x \rfloor (v_1 \wedge \dots \wedge v_r) = \sum_{i=1}^r (-1)^{i-1} (x|v_i) (v_1 \wedge \dots \wedge \check{v}_i \wedge \dots \wedge v_r) \in C^{r-1}(V),$$

$$(6) \quad x \wedge (v_1 \wedge \dots \wedge v_r) = x \wedge v_1 \wedge \dots \wedge v_r \in C^{r+1}(V),$$

where \check{v}_i in (5) means that the i -th vector is omitted from the outer product.

(ii) *One has* $x_{\parallel} = (x \rfloor A)A^{-1}$ and $x_{\perp} = (x \wedge A)A^{-1}$.

(iii) *The subspaces are described as* $V(v_1, \dots, v_r) = \{x \in V; x \wedge A = 0\}$ and $V(v_1, \dots, v_r)^{\perp} = \{x \in V; x \rfloor A = 0\}$.

Proof. We observe that

$$\begin{aligned} xv_1 v_2 \cdots v_r &= (xv_1 + v_1 x)v_2 \cdots v_r \\ &\quad - v_1(xv_2 + v_2 x)v_3 \cdots v_r + (-1)^2 v_1 v_2 (xv_3 + v_3 x)v_4 \cdots v_r + \dots \\ &\quad + (-1)^{r-1} v_1 \cdots v_{r-1} (xv_r + v_r x) + (-1)^r v_1 v_2 \cdots v_r x, \end{aligned}$$

which is rewritten as

$$\begin{aligned} &(xv_1 v_2 \cdots v_r - (-1)^r v_1 v_2 \cdots v_r x)/2 \\ &= (x|v_1)v_2 \cdots v_r - (x|v_2)v_1 v_3 \cdots v_r + (-1)^2 (x|v_3)v_1 v_2 v_4 \cdots v_r + \dots \\ &\quad + (-1)^{r-1} (x|v_r)v_1 \cdots v_{r-1}, \end{aligned}$$

because $(xv_i + v_i x)/2 = (x|v_i)$ ($i = 1, \dots, r$). Replacing the indices $1, 2, \dots, r$ by $\sigma(1), \sigma(2), \dots, \sigma(r)$ for $\sigma \in \mathfrak{S}_r$, and summing up each term as in (1), we obtain (5).

We have $x_{\perp} A = (xA + (-1)^r Ax)/2 = x \wedge A$ by Proposition 6. On the other hand,

$$x_{\perp} A = x_{\perp} u_1 u_2 \cdots u_r = x_{\perp} \wedge u_1 \wedge u_2 \wedge \cdots \wedge u_r = x \wedge v_1 \wedge v_2 \wedge \cdots \wedge v_r,$$

where the last equality follows from (3) and $x_{\parallel} \in V(v_1, \dots, v_r)$. Thus we get (6).

The assertions (ii) and (iii) follow from Propositions 6 and 7 respectively. \square

In view of (5) and (6), we call $x \rfloor B$ (resp. $x \wedge B$) the *inner* (resp. *outer*) *product* of x with B . By definition we have

$$(7) \quad xB = x \rfloor B + x \wedge B.$$

Note that the Clifford product is invertible if B is a non-zero simple r -vector, while the inner nor the outer product is not in general.

§6. Dual basis. A basis $\{w_1, \dots, w_r\}$ of the subspace $V(v_1, \dots, v_r)$ is said to be *dual* to $\{v_1, \dots, v_r\}$ if $(v_i | w_j) = \delta_{ij}$ ($1 \leq i, j \leq r$), where δ_{ij} is Kronecker's delta.

Proposition 9. *The dual basis $\{w_1, \dots, w_r\}$ is given by the following formula:*

$$w_k = (-1)^{k-1} (v_1 \wedge \dots \check{v}_k \dots \wedge v_r) (v_1 \wedge \dots \wedge v_r)^{-1}.$$

Proof. Since $w_k \in V(v_1, \dots, v_r)$, Proposition 8 (iii) together with (7) tells us that $w_k A = w_k \rfloor A$. On the other hand, we have by (5)

$$\begin{aligned} w_k \rfloor A &= \sum_{i=1}^r (-1)^{i-1} (w_k | v_i) (v_1 \wedge \dots \check{v}_i \dots \wedge v_r) \\ &= (-1)^{k-1} v_1 \wedge \dots \check{v}_k \dots \wedge v_r \end{aligned}$$

because $(w_k | v_i) = \delta_{ki}$. Therefore we have $w_k (v_1 \wedge \dots \wedge v_r) = (-1)^{k-1} v_1 \wedge \dots \check{v}_k \dots \wedge v_r$, whence the formula follows. \square

§7. Concluding comments. We have seen that various calculations and formulas are described quite simply by using the Clifford algebra if we just introduce a few of notions such as the outer product of vectors. But they are merely a small part of concepts which enrich the Clifford algebra (see [2]). For example, the notion of *Clifford analytic function* (or *monogenic function*) extends naturally the theory of complex analysis ([1], [2]). On the other hand, if one utilizes the Clifford algebra of signature $(3, 1)$ (*the spacetime algebra*), the formulas in special relativity and electromagnetism like the Maxwell equation are rewritten impressively ([1], [4]).

REFERENCES

- [1] S. Gull, A. Lasenby, and C. Doran, *Imaginary Numbers are not Real — the Geometric Algebra of Spacetime*, Found. Phys. **23** (1993), 1175–1201.
- [2] D. Hestenes and G. Sobczyk, "Clifford algebra to geometric calculus," D. Reidel, 1984.
- [3] D. Hestenes, "New foundations for classical physics," D. Reidel, 1986.
- [4] P. Lounesto, "Clifford algebras and spinors", London Mathematical Society Lecture Note Series **286**, Cambridge University Press, 2001.