

Description of a New Freshwater Goby from Japan¹⁾

By

Nobuhiko MIZUNO²⁾

Zoological Institute, College of Science, University of Kyoto

(Received September 24, 1960)

***Tukugobius flumineus* sp. nov.**

Head 3.60 in total length without caudal; snout 2.73 in head; eye 5.1; interorbital 8.4; caudal peduncle 1.07; depth of the same 2.27. D. VII-I, 7; A. I, 7; P. 17; V. I, 4; C. 16; thirty-five scales in a lateral series, fourteen scales in a transverse series between the origin of second dorsal and anal.

Body elongate, cylindrical anteriorly; caudal peduncle low and compressed; head long and depressed, with bulging cheeks; eyes on top of head; mouth subterminal and slightly oblique, with thick lips; lower jaw slightly included, the posterior angle of maxillary not extending to eye; jaws with moderately broad bands of small conical teeth; the outer row of these teeth in upper jaw enlarged, including some small caninoid teeth which are not seen in some specimens; the same in lower jaw consisting of similar but much smaller teeth; behind outer row in each jaw, about two rows of minute teeth; tongue with round tip; (glossohyal fan-shaped; gills 4, gill rakers on first gill-arch 2+5, short)³⁾.

Two dorsals well separated; first dorsal with strongly curved margin, inserted before middle of pectoral, in some males third and fourth spines long and filamentous in the spawning season; second dorsal higher than first dorsal, with subequal rays and gently curved margin; anal inserted below third or fourth ray of second dorsal, with gently curved margin, last ray ending little before that of second dorsal which does not reach caudal when depressed; pectoral without free silk-like rays on the upper margin, with broadly rounded posterior margin, extending below the tip of fourth spine of first dorsal when depressed; ventrals joining each other, to form a circular adhesive disk, in some specimens broader than long, extending a half way to anus; caudal broadly rounded.

1) The expenses of the present work have been defrayed in part by the Grant in Aid for Fundamental Scientific Research of the Ministry of Education.

2) Present address: Biological Laboratory, Osaka Gakuji University.

3) These characters were examined with the other specimens.

Body covered with firm ctenoid scales which become smaller towards belly and nape; entire head and nuchal region naked; pectoral bases and the entire region about ventrals naked, including breast and a median stripe behind ventrals which sometimes extend to anus; no lateral line; anal papilla distinct.

Colour in formalin yellowish brown, lighter below, with five longitudinal bands of very faint darkish colour along the middle of each side which unite to be a longitudinal streak or line in some specimens, without distinct marking on head and paired fins; vertical fins faint dusky; brownish spots on rays of second dorsal and caudal arranged in several longitudinal rows.

Type-specimens: Holotype: ♀ (48.8 mm in total length), paratypes: 3 ♂♂, 2 ♀♀ (30-VIII-1959, collected by N. MIZUNO).

All the type-specimens are preserved in my collection.

Type-locality: Sôshagawa river at Oshita, in Ochigun District, Ehime Prefecture, Shikoku, Japan.

Habitat: Mountain streams in the southwestern region of Japan, including the Riukiu Islands, as reported on the previous pages.

Remarks: This fish is very variable as seen in Table 1, and has been frequently included in *Rhinogobius similis* GILL, from which it differs in such characters as shown in Table 2. These details were reported on the previous pages.

Table 1. Measurements of *Tukugobius flumineus* collected in several stations.

Station	Head	Snout	Eye	Inter-orbital	Caudal peduncle	D.	A.	P.	C.	Scales	Body length
R. Sôshagawa	3.60	2.73	5.1	8.4	1.07	VII-I,9	I, 7	17	16	35	48.8
	3.70	3.12	4.8	9.1	1.09	VI -I,9	I, 7	16	15	34	46.2
R. Yuragawa	3.26	3.70	4.9	8.8	1.21	VI -I,8	I, 7	17	16	34	59.0
	3.38	2.87	4.6	6.7	1.16	VI -I,8	I, 7	17	16	34	52.4
	3.43	2.98	5.3	4.6	1.15	VI -I,8	I, 8	17	16	35	58.2
	3.55	4.52	4.5	7.5	1.15	VI -I,9	I, 8	17	15	36	52.6
	3.69	4.80	4.8	9.2	1.04	VI -I,9	I, 8	17	16	35	50.9
R. Inukaigawa	3.17	2.95	6.2	7.3	1.24	VII-I,8	I, 8	17	16	34	48.6
	3.33	2.95	5.5	6.3	1.10	VII-I,8	I, 7	16	16	34	50.8
	3.40	2.95	4.8	6.3	1.09	VI -I,8	I, 7	16	15	34	50.8
	3.63	3.51	4.8	6.2	1.05	VI -I,7	I, 7	17	15	36	50.5
R. Kamogawa	3.04	2.64	6.5	6.6	1.15	VI -I,8	I, 7	17	15	33	57.4
	3.32	2.61	5.9	6.7	1.16	VI -I,8	I, 7	16	17	33	57.6
	3.24	2.96	5.5	5.1	1.16	VI -I,8	I, 7	16	15	35	58.3
	3.64	3.48	4.9	6.6	0.97	VI -I,8	I, 7	17	15	35	48.8




Fig. 1. *Tukugobius flumineus* sp. nov., ♀, of the Sōshagawa river.

Table 2. Distinctive features between *Tukugobius flumineus* and *Rhinogobius similis*.

	<i>Tukugobius flumineus</i>	<i>Rhinogobius similis</i>
Number of first dorsal rays	VI or VII	VI
Number of pectoral rays	15~18	18~22
Number of vertebrae	28 (rarely 27)	26
Number of gill rakers	2+5	2+8~9
Size of eggs (mm)	1.4~2.1	0.4~0.8
Number of eggs in a female	Ca. 70~130	Ca. 1000~7000
Size of newly hatched larvae (mm)	Ca. 7.5	Ca. 4.0
Mode of life	fluvial life	Amphidromous life or Lacustrine life