



TITLE:

# Rigid Fibers of Spinning Tops

AUTHOR(S):

KAWASAKI, Morimichi; ORITA, Ryuma

---

CITATION:

KAWASAKI, Morimichi ...[et al]. Rigid Fibers of Spinning Tops. 2020: 1-17: RIMS-1922.

ISSUE DATE:

2020-08

URL:

<http://hdl.handle.net/2433/261820>

RIGHT:

RIMS-1922

**Rigid Fibers of Spinning Tops**

By

Morimichi KAWASAKI and Ryuma ORITA

August 2020



京都大学 数理解析研究所

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

# RIGID FIBERS OF SPINNING TOPS

MORIMICHI KAWASAKI AND RYUMA ORITA

ABSTRACT. (Non-)displaceability of fibers of integrable systems has been an important problem in symplectic geometry. In this paper, for a large class of classical Liouville integrable systems containing the Lagrangian top, the Kovalevskaya top and the C. Neumann problem, we find a non-displaceable fiber for each of them. Moreover, we show that the non-displaceable fiber which we detect is the unique fiber which is non-displaceable from the zero-section. As a special case of this result, we also show that a singular level set of a convex Hamiltonian is non-displaceable from the zero-section. To prove these results, we use the notion of superheaviness introduced by Entov and Polterovich.

## CONTENTS

1. Introduction	1
1.1. Backgrounds	1
1.2. Main results	3
2. Applications	5
2.1. Relationship with Mañé's critical values	5
2.2. Classical integrable systems	6
2.3. On the existence of infinitely many non-displaceable fibers	9
3. Preliminaries	9
3.1. Conventions and notation	9
3.2. Partial symplectic quasi-states	10
4. Proofs of the main results	11
4.1. Proof of Theorem 1.10	11
4.2. Proof of Theorem 1.11	13
4.3. Proof of Proposition 1.13	14
5. Proof of Proposition 2.14	14
Acknowledgments	15
References	15

## 1. INTRODUCTION

**1.1. Backgrounds.** Let  $(M, \omega)$  be a symplectic manifold. A subset  $X \subset M$  is called *displaceable* from a subset  $Y \subset M$  if there exists a Hamiltonian  $H: [0, 1] \times M \rightarrow \mathbb{R}$  with compact support such that  $\varphi_H(X) \cap \bar{Y} = \emptyset$ , where  $\varphi_H$  is the *Hamiltonian diffeomorphism* generated by  $H$  (see Section 3.1 for the definition) and  $\bar{Y}$  is

---

*Date:* October 7, 2019.

*2010 Mathematics Subject Classification.* Primary 57R17, 53D12; Secondary 53D20, 53D40, 53D45.

*Key words and phrases.* Symplectic manifolds, groups of Hamiltonian diffeomorphisms, moment maps, symplectic quasi-states, heavy subsets.

This work has been supported by JSPS KAKENHI Grant Numbers JP18J00765, JP18J00335.

the topological closure of  $Y$ . Otherwise,  $X$  is called *non-displaceable* from  $Y$ . For simplicity, we call  $X$  (non-)displaceable if  $X$  is (non-)displaceable from  $X$  itself.

The problem of (non-)displaceability of a subset of a symplectic manifold (from another subset or from itself) has attracted much attention in symplectic geometry. Non-displaceability results often pinpoint symplectic rigidity, namely the difference between symplectic topology and differential topology, and lead to interesting results in symplectic topology and Hamiltonian dynamics, see for example [PPS]. In this paper, all symplectic manifolds are cotangent bundles  $T^*N$  over closed smooth manifolds  $N$ , equipped with the standard symplectic form. These are the phase spaces of classical mechanics. The first result on non-displaceability in cotangent bundles was non-displaceability of the zero-section [Gr, LS, Ho, Fl]. The traditional tools (Morse theory for generating functions,  $J$ -holomorphic curves, and Floer homology) work only when the set in question is a submanifold. However, many dynamically relevant subsets of cotangent bundles are not submanifolds. Examples are energy levels of autonomous Hamiltonians at which the qualitative behavior of the dynamics changes, like Mañé’s critical values, and certain subsets therein. In [EP06], Entov and Polterovich used Floer homology to construct a function theoretical method that is designed to detect the non-displaceability of arbitrary closed subsets (we refer to [En, PR] as good surveys). This theory was adapted by Monzner, Vichery, and Zapolsky [MVZ] to cotangent bundles. In this paper we use their theory to prove the non-displaceability of fibers of classical integrable systems or the energy level corresponding to Mañé’s critical value.

We also note that there are some extrinsic applications of non-displaceability. Polterovich [Po14] proved the existence of a invariant measure of some Hamiltonian flow using non-displaceability of some subset in certain situations. He [Po98] also constructed a Hamiltonian diffeomorphism with arbitrary large Hofer’s norm using non-displaceability of (the equator)  $\times 0_{S^1}$  in  $S^2 \times T^*S^1$ , where  $0_{S^1}$  is the zero-section of  $T^*S^1$ . In [Ka17], the first author posed some generalization of Bavard’s duality theorem. Combing it with Polterovich’s above result, he pointed out that the existence of stably non-displaceable fibers might be related to the existence of partial quasi-morphisms on the group of Hamiltonian diffeomorphisms.

Let  $k$  be a positive integer. We call a smooth map  $\Phi = (\Phi_1, \dots, \Phi_k): M \rightarrow \mathbb{R}^k$  a *moment map* if  $\{\Phi_i, \Phi_j\} = 0$  for all  $1 \leq i, j \leq k$ , where  $\{\cdot, \cdot\}$  denotes the Poisson bracket on  $(M, \omega)$ . A moment map  $\Phi = (\Phi_1, \dots, \Phi_k): M \rightarrow \mathbb{R}^k$  is called a *Liouville integrable system* if  $k = \dim M/2$  and  $(d\Phi_1)_x, \dots, (d\Phi_k)_x$  are linearly independent almost everywhere.

Many researchers have studied (non-)displaceable fibers of Liouville integrable system associated with toric structures. For example, see [BEP, Ch, EP09, Mc, FOOO10, FOOO11, FOOO12, AM, ABM, KLS, AFOOO]. Recently some researchers study (non-)displaceable fibers of “moment maps” associated with various generalizations of toric structure like Gelfand–Cetlin systems, semi-toric structures and so on (see, e.g., [NNU, Wu, Vi, CKO, KO19b]).

In this paper, we deal with classical Liouville integrable systems on cotangent bundles. We study (non-)displaceable fibers of moment maps on the cotangent bundle of the two-sphere  $S^2$  or the three-dimensional rotation group  $\mathrm{SO}(3)$  which appear in classical mechanics, for example, the spherical pendulum, the Lagrange top and the Kovalevskaya top. As a previous research in a similar direction, we refer to Albers–Frauenfelder’s work [AF08]. They proved non-displaceability of the Polterovich torus in  $T^*S^2$  which can be regarded as a fiber of some Liouville integrable system.

As a general fact on (non-)displaceability of fibers of moment maps, Entov and Polterovich [EP06] proved the following theorem.

**Theorem 1.1** ([EP06, Theorem 2.1]). *Let  $(M, \omega)$  be a closed symplectic manifold and  $\Phi = (\Phi_1, \dots, \Phi_k): M \rightarrow \mathbb{R}^k$  a moment map. Then, there exists  $y_0 \in \Phi(M)$  such that  $\Phi^{-1}(y_0)$  is non-displaceable.*

To prove Theorem 1.1, Entov and Polterovich [EP06] introduced the concept of partial symplectic quasi-state (see Definition 3.1). In [EP09], they introduced the notion of heaviness of closed subsets in terms of partial symplectic quasi-states. Let  $C_c(M)$  denote the set of continuous functions on  $M$  with compact supports.

**Definition 1.2** ([EP09, Definition 1.3]). Let  $\zeta: C_c(M) \rightarrow \mathbb{R}$  be a partial symplectic quasi-state on  $(M, \omega)$ . A compact subset  $X$  of  $M$  is said to be  $\zeta$ -heavy (resp.  $\zeta$ -superheavy) if

$$\zeta(H) \geq \inf_X H \quad \left( \text{resp. } \zeta(H) \leq \sup_X H \right)$$

for any  $H \in C_c(M)$ .

Here we collect properties of (super)heavy subsets.

**Theorem 1.3** ([EP09, Theorem 1.4]). *Let  $\zeta: C_c(M) \rightarrow \mathbb{R}$  be a partial symplectic quasi-state on  $(M, \omega)$ .*

- (i) *Every  $\zeta$ -superheavy subset is  $\zeta$ -heavy.*
- (ii) *Every  $\zeta$ -heavy subset is non-displaceable.*
- (iii) *Every  $\zeta$ -heavy subset is non-displaceable from every  $\zeta$ -superheavy subset. In particular, every  $\zeta$ -heavy subset intersects every  $\zeta$ -superheavy subset.*

Relating to Theorem 1.1, Entov and Polterovich posed the following problem.

**Problem 1.4** ([EP09, Section 1.8.2], see also [En, Question 4.9]). *Let  $(M, \omega)$  be a closed symplectic manifold and  $\Phi = (\Phi_1, \dots, \Phi_k): M \rightarrow \mathbb{R}^k$  a moment map. Let  $\zeta: C(M) \rightarrow \mathbb{R}$  be a partial symplectic quasi-state on  $(M, \omega)$  made from the Oh-Schwarz spectral invariant (see [Sch, Oh05]). Then, does there exist  $y_0 \in \Phi(M)$  such that  $\Phi^{-1}(y_0)$  is  $\zeta$ -heavy?*

**1.2. Main results.** In this paper we prove that some classical integrable systems (e.g., Lagrange top and Kovalevskaya top) admit superheavy fibers. We consider the cotangent bundle  $(T^*N, \omega_0)$  of a closed smooth  $n$ -dimensional manifold  $N$  where  $\omega_0$  is the standard symplectic form on  $T^*N$ . Let  $(q, p)$  be canonical coordinates on  $T^*N$  where  $q \in N$  and  $p \in T_q^*N$ . Let  $\pi: T^*N \rightarrow N$  denote the natural projection.

**Definition 1.5.** A (time-independent) Hamiltonian  $H: T^*N \rightarrow \mathbb{R}$  satisfies *condition*  $(\star)$  if the following conditions hold.

- (i) For any  $c \in \mathbb{R}$  the sublevel set  $H^{-1}((-\infty, c]) \subset T^*N$  is compact.
- (ii) For any  $q \in N$ ,

$$H(q, 0) = \min_{p \in T_q^*N} H(q, p).$$

For a Hamiltonian  $H: T^*N \rightarrow \mathbb{R}$ , we set

$$(1) \quad m_H = \max_{q \in N} \min_{p \in T_q^*N} H(q, p) \quad (\text{if exists})$$

and then define

$$(2) \quad S_H = \{ (q, p) \in T^*N \mid H(q, p) = m_H \}.$$

If  $H$  satisfies condition  $(\star)$ , then we have

$$m_H = \max_{q \in N} H(q, 0) \quad \text{and} \quad S_H = \{ (q, 0) \in T^*N \mid H(q, 0) = m_H \}.$$

Typical examples of Hamiltonians satisfying condition  $(\star)$  are *convex Hamiltonians*

$$(3) \quad H(q, p) = \frac{1}{2} \|p\|_g^2 + U(q),$$

where  $\|\cdot\|_g$  is the dual norm of a Riemannian metric  $g$  on  $N$  and  $U: N \rightarrow \mathbb{R}$  is a smooth potential. In this case, the value  $m_H$  equals the *Mañé critical value*  $\max_N U$  and

$$S_H = \left\{ (q, 0) \in T^*N \mid U(q) = \max_N U \right\}.$$

In Section 2.2, we provide classical examples satisfying the assumption of Theorem 1.7.

To prove non-displaceability of a fiber of some integrable systems, we use the following partial symplectic quasi-state. In [Oh97, Oh99], Oh constructed a spectral invariant on  $(T^*N, \omega_0)$  in terms of the Lagrangian Floer theory of the zero-section  $0_N$  of  $T^*N$ . In [MVZ], Monzner, Vichery, and Zapolsky constructed a partial symplectic quasi-state on  $(T^*N, \omega_0)$ , denoted by  $\zeta_{\text{MVZ}}: C_c(T^*N) \rightarrow \mathbb{R}$ , as the asymptotization of Oh's Lagrangian spectral invariant. In this paper, the following property of  $\zeta_{\text{MVZ}}$  is crucial.

**Proposition 1.6** ([MVZ, Example 1.19]). *The zero-section  $0_N \subset T^*N$  is  $\zeta_{\text{MVZ}}$ -superheavy.*

Now we are in a position to state the main result of this paper.

**Theorem 1.7.** *Let  $N$  be a closed manifold and  $\Phi = (\Phi_1, \dots, \Phi_k): T^*N \rightarrow \mathbb{R}^k$  a moment map. Assume that  $\Phi_1$  satisfies condition  $(\star)$  and that the set  $\Phi(S_{\Phi_1})$  is a singleton, i.e.,  $\Phi(S_{\Phi_1}) = \{y_0\}$  for some  $y_0 \in \mathbb{R}^k$ . Then, the fiber  $\Phi^{-1}(y_0) \subset T^*N$  is  $\zeta_{\text{MVZ}}$ -superheavy.*

Theorem 1.7 gives a partial answer to Problem 1.4 for a large class of moment maps on cotangent bundles.

By Theorem 1.3 and Proposition 1.6, the fiber  $\Phi^{-1}(y_0)$  is non-displaceable from itself and from the zero-section  $0_N$ . Moreover, we can prove that every fiber of  $\Phi$ , other than  $\Phi^{-1}(y_0)$ , is displaceable from  $0_N$ . To refine Theorem 1.7, we introduce the notion of *X-stems*.

**Definition 1.8** ([Ka18]). Let  $(M, \omega)$  be a symplectic manifold and  $X$  a compact subset of  $M$ . A compact subset  $Y$  of  $M$  is called an *X-stem* if there exists a moment map  $\Phi = (\Phi_1, \dots, \Phi_k): M \rightarrow \mathbb{R}^k$  satisfying the following conditions:

- (i)  $Y = \Phi^{-1}(p)$  for some  $p \in \Phi(M)$ .
- (ii) Every fiber of  $\Phi$ , other than  $Y$ , is displaceable from itself or from  $X$ .

Entov and Polterovich [EP06] introduced the notion of *stems* (i.e., every fiber of  $\Phi$ , other than  $\Phi^{-1}(y_0)$ , is displaceable, where  $\Phi: T^*N \rightarrow \mathbb{R}^k$  is a moment map) and proved that stems are superheavy with respect to any partial symplectic quasi-state [EP09, Theorem 1.8]. We note that every stem is an *X-stem* for any compact subset  $X$ . Concerning *X-stems*, we have the following result.

**Theorem 1.9.** *Let  $(M, \omega)$  be a symplectic manifold,  $\zeta: C_c(M) \rightarrow \mathbb{R}$  a partial symplectic quasi-state on  $(M, \omega)$ , and  $X$  a  $\zeta$ -superheavy subset of  $M$ . Then every *X-stem* is  $\zeta$ -superheavy.*

We prove Theorem 1.9 in Section 3.2. Then we can refine Theorem 1.7 as follows.

**Theorem 1.10.** *Let  $N$  be a closed manifold and  $\Phi = (\Phi_1, \dots, \Phi_k): T^*N \rightarrow \mathbb{R}^k$  a moment map. Assume that  $\Phi_1$  satisfies condition  $(\star)$  and that the set  $\Phi(S_{\Phi_1})$  is a singleton, i.e.,  $\Phi(S_{\Phi_1}) = \{y_0\}$  for some  $y_0 \in \mathbb{R}^k$ . Then, every fiber of  $\Phi$ ,*

other than  $\Phi^{-1}(y_0)$ , is displaceable from the zero-section  $0_N$ . In particular, the fiber  $\Phi^{-1}(y_0)$  is a  $0_N$ -stem. Hence, by Theorem 1.9 and Proposition 1.6,  $\Phi^{-1}(y_0)$  is  $\zeta_{\text{MVZ}}$ -superheavy.

We prove Theorem 1.10 in Section 4.1. By Theorem 1.3, we see that  $\Phi^{-1}(y_0)$  is the unique fiber which is non-displaceable from  $0_N$ . On the other hand, it is a natural question to ask whether  $\Phi^{-1}(y_0)$  is a stem. In Conjecture 2.15, the authors expect that  $\Phi: T^*N \rightarrow \mathbb{R}^k$  has infinitely many non-displaceable fibers, in particular,  $\Phi^{-1}(y_0)$  is not a stem in a more general situation. For evidences supporting Conjecture 2.15, see Section 2.3.

Here we provide two other applications of our arguments.

**Theorem 1.11.** *Let  $H_1, \dots, H_k: T^*N \rightarrow \mathbb{R}$  be Hamiltonians satisfying condition  $(\star)$  and  $\{H_i, H_j\} = 0$  for all  $1 \leq i, j \leq k$ . Then,  $\bigcap_{i=1}^k S_{H_i} \neq \emptyset$ .*

For example, the functions  $H$  and  $G$  in Example 2.8 satisfy condition  $(\star)$  and one can confirm that  $S_H \cap S_G \neq \emptyset$ . As another example, the functions  $H$  and  $G$  in Example 2.12 also satisfy condition  $(\star)$  and we have  $S_H \cap S_G \neq \emptyset$ . We prove Theorem 1.11 in Section 4.2. The authors do not know another proof of this mysterious theorem without using the Floer theory.

**Proposition 1.12.** *Let  $\Phi = (\Phi_1, \dots, \Phi_k): T^*N \rightarrow \mathbb{R}^k$  be a moment map. Assume that  $\Phi_1$  satisfies condition  $(\star)$  and that the set  $\Phi(S_{\Phi_1})$  is a singleton, i.e.,  $\Phi(S_{\Phi_1}) = \{y_0\}$  for some  $y_0 \in \mathbb{R}^k$ . Then,  $\pi(\Phi^{-1}(y_0)) = N$ .*

When  $k = 1$ , the proof of Proposition 1.12 is straightforward by the definition of  $m_{\Phi_1}$ . Proposition 1.12 follows immediately from Theorem 1.7 and the following proposition.

**Proposition 1.13.** *If  $X$  is a  $\zeta_{\text{MVZ}}$ -superheavy subset of  $T^*N$ , then  $\pi(X) = N$ .*

We prove Proposition 1.13 in Section 4.3.

## 2. APPLICATIONS

In this section, we deal with some classical integrable systems satisfying the assumption of Theorem 1.7 and detect superheavy fibers of them.

**2.1. Relationship with Mañé's critical values.** We provide an application of our main theorem when a moment map is a function.

Let  $(N, g)$  be a closed Riemannian manifold. We equip the cotangent bundle  $T^*N$  with the standard symplectic form  $\omega_0$ . In the context of Mañé's critical values, Cieliebak, Frauenfelder, and Paternain [CFP] proved the following theorem.

**Theorem 2.1** ([CFP, Theorem 1.2]). *Let  $(N, g)$  be a closed Riemannian manifold and  $H: T^*N \rightarrow \mathbb{R}$  a convex Hamiltonian (see (3) for the definition). Then, the level set  $H^{-1}(m_H) \subset T^*N$  is non-displaceable.*

As a corollary of our main theorem (Theorem 1.7), we can prove that the level set  $H^{-1}(m_H)$  in Theorem 2.1 is non-displaceable from the zero-section  $0_N$  in a more general setting.

**Corollary 2.2.** *Let  $N$  be a closed manifold and  $H: T^*N \rightarrow \mathbb{R}$  a Hamiltonian satisfying condition  $(\star)$ . Then, the level set  $H^{-1}(m_H) \subset T^*N$  is non-displaceable from itself and from the zero-section  $0_N$ .*

*Remark 2.3.* Actually, Cieliebak, Frauenfelder, and Paternain [CFP] proved the non-displaceability of  $H^{-1}(c)$  for any  $c > m_H$  using the Rabinowitz Floer theory. Hence they obtained Theorem 2.1 as its corollary. On the other hand, as stated in Theorem 1.10,  $H^{-1}(c)$  is displaceable from  $0_N$  for any  $c \neq m_H$ .

**Example 2.4** (Pendulum). The pendulum is the Hamiltonian system with one degree of freedom on the cotangent bundle  $T^*S^1$  of the unit circle  $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ . We define a function  $H: T^*S^1 \rightarrow \mathbb{R}$  by

$$H(q, p) = \frac{1}{2}p^2 + (1 - \cos q).$$

Then,  $H$  satisfies condition  $(\star)$  and

$$m_H = \max_{q \in S^1} (1 - \cos q) = 2.$$

By Theorem 1.7, the level set  $H^{-1}(2) \subset T^*S^1$  is  $\zeta_{\text{MVZ}}$ -superheavy.  $H^{-1}(2)$  is homeomorphic to the figure eight. Note that the  $\zeta_{\text{MVZ}}$ -superheaviness of  $H^{-1}(2)$  also follows from [MVZ, Proposition 1.22].

**Example 2.5** (Double spherical pendulum). The double spherical pendulum [MS] consists of two coupled spherical pendula (see also [HLS]). Let

$$(4) \quad S^2 = \{q = (q_1, q_2, q_3) \in \mathbb{R}^3 \mid q_1^2 + q_2^2 + q_3^2 = 1\}$$

denote the unit two-sphere in  $\mathbb{R}^3$ . Let  $q = (q_1, q_2, q_3) \in S^2$  be the position vector of the first spherical pendulum relative to  $(0, 0, 0) \in \mathbb{R}^3$ . Let  $q' = (q'_1, q'_2, q'_3) \in S^2$  be the position vector of the second spherical pendulum relative to the first one. We define a function  $\underline{H}: T(S^2 \times S^2) \rightarrow \mathbb{R}$  by

$$\underline{H}(q, q', v, v') = \frac{1}{2}\|v - v'\|_{g_0}^2 + \frac{1}{2}\|v'\|_{g_0}^2 + 2q_3 + q'_3.$$

Let  $\Psi: T(S^2 \times S^2) \rightarrow T^*(S^2 \times S^2)$  denote the Legendre transformation of  $\underline{H}$ . We then define a function on  $T^*(S^2 \times S^2)$  by  $H = \underline{H} \circ \Psi^{-1}$ . Then  $H$  satisfies condition  $(\star)$  and

$$m_H = \max_{(q, q') \in S^2 \times S^2} (2q_3 + q'_3) = 3.$$

By Theorem 1.7, the level set  $H^{-1}(3) \subset T^*(S^2 \times S^2)$  is  $\zeta_{\text{MVZ}}$ -superheavy. It is homeomorphic to the unit cotangent bundle  $S_{g_0, 1}^*(S^2 \times S^2)$  with the fiber over the point  $(n, n) \in S^2 \times S^2$  being collapsed to the single point  $((n, n), (0, 0)) \in T^*(S^2 \times S^2)$ , where  $n = (0, 0, 1) \in S^2$  is the north pole.

## 2.2. Classical integrable systems.

**Example 2.6** (Spherical pendulum). The spherical pendulum [La] describes a motion of a particle moving on the unit two-sphere  $S^2 \subset \mathbb{R}^3$  under a gravitational force. Let  $g_0$  denote the standard Riemannian metric on  $S^2$ . We define functions  $\underline{H}, \underline{G}: TS^2 \rightarrow \mathbb{R}$  by

$$\underline{H}(q, v) = \frac{1}{2}\|v\|_{g_0}^2 + q_3 \quad \text{and} \quad \underline{G}(q, v) = q_1 v_2 - q_2 v_1$$

for  $(q, v) = (q_1, q_2, q_3, v_1, v_2, v_3) \in TS^2 \subset T\mathbb{R}^3 \cong \mathbb{R}^3 \times \mathbb{R}^3$ , respectively. Let  $\Psi: TS^2 \rightarrow T^*S^2$  denote the Legendre transformation of  $\underline{H}$ . We then define functions on  $T^*S^2$  by  $H = \underline{H} \circ \Psi^{-1}$  and  $G = \underline{G} \circ \Psi^{-1}$ . Then,  $\{H, G\} = 0$  and the function  $H$  satisfies condition  $(\star)$ . We set  $\Phi = (H, G): T^*S^2 \rightarrow \mathbb{R}^2$ . Since  $S_H = \{(0, 0, 1, 0, 0, 0)\}$ , we have  $\Phi(S_H) = \{(1, 0)\}$ . By Theorem 1.7, the fiber  $\Phi^{-1}(1, 0) \subset T^*S^2$  is  $\zeta_{\text{MVZ}}$ -superheavy. In particular,  $\Phi^{-1}(1, 0)$  is non-displaceable from itself and from  $0_{S^2}$ . We note that the value  $(1, 0)$  corresponds to the focus-focus singularity of this system and the fiber  $\Phi^{-1}(1, 0)$  is homeomorphic to the two-dimensional torus pinched at a single point (see [CB, Section IV.3.4]).

*Remark 2.7.* Brendel, Kim, and Schlenk [BKS] proved that the fiber  $\Phi^{-1}(c, 0)$  is non-displaceable for any  $c > 1$ . Thus, the non-displaceability of  $\Phi^{-1}(1, 0)$  immediately follows. On the other hand, as stated in Theorem 1.10,  $\Phi^{-1}(c, 0)$  is displaceable from  $0_{S^2}$  for any  $c > 1$ .



The authors do not know whether there exist a Hamiltonian  $H$  satisfying condition  $(\star)$  and a real number  $c$  with  $c > m_H$  such that  $H^{-1}(c)$  is displaceable.

**Example 2.8** (C. Neumann problem). Let  $a_1, a_2, a_3$  be positive numbers satisfying  $a_1 < a_2 < a_3$ . Let  $S^2 \subset \mathbb{R}^3$  denote the unit two-sphere as in (4). In [Neu], C. Neumann introduced a Hamiltonian system on  $T^*S^2$  which describes the motion of a particle on the unit two-sphere  $S^2$  under the influence of the linear force  $-(a_1q_1, a_2q_2, a_3q_3)$ . We define functions  $\underline{H}, \underline{G}: TS^2 \rightarrow \mathbb{R}$  by

$$\underline{H}(q, v) = \frac{1}{2}\|v\|_{g_0}^2 + \frac{1}{2}(a_1q_1^2 + a_2q_2^2 + a_3q_3^2)$$

and

$$\underline{G}(q, v) = \frac{1}{2}\sum_{i=1}^3 a_i v_i^2 + \frac{1}{2}\|v\|_{g_0}^2 \sum_{i=1}^3 a_i q_i^2 + \frac{1}{2}\sum_{i=1}^3 a_i^2 q_i^2$$

for  $(q, v) = (q_1, q_2, q_3, v_1, v_2, v_3) \in TS^2 \subset T\mathbb{R}^3 \cong \mathbb{R}^3 \times \mathbb{R}^3$ , respectively. Let  $\Psi: TS^2 \rightarrow T^*S^2$  denote the Legendre transformation of  $\underline{H}$ . We then define functions  $H, G: T^*S^2 \rightarrow \mathbb{R}$  by  $H = \underline{H} \circ \Psi^{-1}$  and  $G = \underline{G} \circ \Psi^{-1}$ . Then,  $\{H, G\} = 0$  and the function  $H$  satisfies condition  $(\star)$ . We set  $\Phi = (H, G): T^*S^2 \rightarrow \mathbb{R}^2$ . Since  $S_H = \{(0, 0, \pm 1, 0, 0, 0)\}$ , we have  $\Phi(S_H) = \{(a_3/2, a_3^2/2)\}$ . By Theorem 1.7, the fiber  $\Phi^{-1}(a_3/2, a_3^2/2) \subset T^*S^2$  is  $\zeta_{\text{MVZ}}$ -superheavy.

2.2.1. *Spinning tops.* We consider the motion of tops. Let  $q_1 \cdot q_2$  (resp.  $q_1 \times q_2$ ) denote the dot (resp. cross) product of  $q_1$  and  $q_2$  in  $\mathbb{R}^3$ . Let

$$\text{SO}(3) = \{(q_1, q_2, q_3) \in \text{M}_3(\mathbb{R}) \mid q_1, q_2, q_3 \in S^2, q_1 \cdot q_2 = 0, q_3 = q_1 \times q_2\}$$

denote the three-dimensional rotation group, where  $S^2$  is the unit two-sphere in  $\mathbb{R}^3$ . Let  $(e_1, e_2, e_3)$  denote the identity matrix. Given a point  $(q_1, q_2, q_3) \in \text{SO}(3)$ , we set  $n_i = q_i \cdot e_3$  for each  $i = 1, 2, 3$ .

Let  $(q, \omega) = (q_1, q_2, q_3, \omega_1, \omega_2, \omega_3)$  be the canonical coordinates on  $T\text{SO}(3)$  defined in terms of the angular velocity (see, for example, [Ar, Section 26]). Let  $0_{\text{SO}(3)}$  denote the zero-section of  $T^*\text{SO}(3)$ .

Let  $I_1, I_2, I_3$  be positive numbers and  $f: [-1, 1] \times [-1, 1] \times [-1, 1] \rightarrow \mathbb{R}$  a smooth function. We define functions  $\underline{H}, \underline{L}_z: T\text{SO}(3) \rightarrow \mathbb{R}$  by

$$(5) \quad \underline{H}(q, \omega) = \frac{1}{2}(I_1\omega_1^2 + I_2\omega_2^2 + I_3\omega_3^2) + f(n_1, n_2, n_3)$$

and

$$(6) \quad \underline{L}_z(q, \omega) = I_1n_1\omega_1 + I_2n_2\omega_2 + I_3n_3\omega_3,$$

respectively. Let  $\Psi: T\text{SO}(3) \rightarrow T^*\text{SO}(3)$  denote the Legendre transformation of  $\underline{H}$ . Note that  $\Psi: T\text{SO}(3) \rightarrow T^*\text{SO}(3)$  is the metric dual operation with respect to the Riemannian metric  $g$  on  $\text{SO}(3)$  defined by

$$g_q(\omega, \omega') = I_1\omega_1\omega'_1 + I_2\omega_2\omega'_2 + I_3\omega_3\omega'_3$$

for  $q \in \text{SO}(3)$  and  $\omega = (\omega_1, \omega_2, \omega_3)$ ,  $\omega' = (\omega'_1, \omega'_2, \omega'_3) \in T_q\text{SO}(3)$ .

We then define functions on  $T^*\text{SO}(3)$  by  $H = \underline{H} \circ \Psi^{-1}$  and  $L_z = \underline{L}_z \circ \Psi^{-1}$ . Then,  $\{H, L_z\} = 0$  and the function  $H$  satisfies condition  $(\star)$ . We note that

$$(7) \quad S_H = \left\{ (q, 0) \in T^*\text{SO}(3) \mid H(q, 0) = \max_{\text{SO}(3)} f \circ \nu \right\},$$

where  $\nu: \text{SO}(3) \rightarrow [-1, 1] \times [-1, 1] \times [-1, 1]$  is the map defined by  $\nu(q_1, q_2, q_3) = (n_1, n_2, n_3)$ . Hence  $L_z(S_H) \subset L_z(0_{\text{SO}(3)}) = \{0\}$ .

**Example 2.9.** We set  $\Phi = (H, L_z): T^*\text{SO}(3) \rightarrow \mathbb{R}^2$ . Then,

$$\Phi(S_H) = \left\{ \left( \max_{\text{SO}(3)} f \circ \nu, 0 \right) \right\}.$$

By Theorem 1.7, the fiber  $\Phi^{-1}(\max_{\text{SO}(3)} f \circ \nu, 0)$  is  $\zeta_{\text{MVZ}}$ -superheavy.

**Example 2.10** (Lagrange top). The *Lagrange top* [La, Ar] is a top such that  $I_1 = I_2$  and  $f(x, y, z) = cz$  for some real number  $c$ . We define another function  $\underline{G}: T\text{SO}(3) \rightarrow \mathbb{R}$  by

$$\underline{G}(q, \omega) = I_3 \omega_3,$$

and set  $G = \underline{G} \circ \Psi^{-1}$ . Then,  $\{H, G\} = 0$  and  $\{L_z, G\} = 0$ . We set  $\Phi = (H, L_z, G): T^*\text{SO}(3) \rightarrow \mathbb{R}^3$ . By (7),  $H(S_H) = \{|c|\}$  and  $G(S_H) = \{0\}$ . Therefore,  $\Phi(S_H) = \{(|c|, 0, 0)\}$ . By Theorem 1.7, the fiber  $\Phi^{-1}(|c|, 0, 0)$  is  $\zeta_{\text{MVZ}}$ -superheavy. If  $|c| \neq 0$ , the fiber  $\Phi^{-1}(|c|, 0, 0)$  is homeomorphic to a 3-torus with a normal crossing along an  $S^1$ . For more precise description of this fiber and its singularity, see [CB, Section V.6].

**Example 2.11** (Kovalevskaya top). The *Kovalevskaya top* [Ko] is a top such that  $I_1 = I_2 = 2I_3$  and  $f(x, y, z) = ax$  for some real number  $a$ . We define another function  $\underline{G}: T\text{SO}(3) \rightarrow \mathbb{R}$  by

$$\underline{G}(q, \omega) = \left( \omega_1^2 - \omega_2^2 - \frac{2a}{I_1} n_1 \right)^2 + \left( 2\omega_1 \omega_2 - \frac{2a}{I_1} n_2 \right)^2,$$

and set  $G = \underline{G} \circ \Psi^{-1}$ . Then,  $\{H, G\} = 0$  and  $\{L_z, G\} = 0$ . We set  $\Phi = (H, L_z, G): T^*\text{SO}(3) \rightarrow \mathbb{R}^3$ . By (7),  $H(S_H) = \{|a|\}$ . If  $a \neq 0$ , then

$$S_H = \{ (q_1, q_2, q_3, 0, 0, 0) \in T^*\text{SO}(3) \mid q_1 = \text{sgn}(a)e_3 \},$$

where  $\text{sgn}(a)$  is the signature of  $a$ , and hence  $G(S_H) = \{4a^2/I_1^2\}$ . If  $a = 0$ , then  $S_H = 0_{\text{SO}(3)}$ , and hence  $G(S_H) = \{0\}$ .

Therefore, given  $a \in \mathbb{R}$ , we have  $\Phi(S_H) = \{(|a|, 0, 4a^2/I_1^2)\}$ . By Theorem 1.7, the fiber  $\Phi^{-1}(|a|, 0, 4a^2/I_1^2)$  is  $\zeta_{\text{MVZ}}$ -superheavy.

**Example 2.12** (Clebsch top). The *Clebsch top* [Cl] is a top such that  $I_1 < I_2 < I_3$  and

$$f(x, y, z) = \frac{1}{2I_1 I_2 I_3} (I_1 x^2 + I_2 y^2 + I_3 z^2).$$

This system describes a motion of a rigid body, fixed in its center of gravity, in an ideal fluid. We define another function  $\underline{G}: T\text{SO}(3) \rightarrow \mathbb{R}$  by

$$\underline{G}(q, \omega) = \frac{1}{2} (I_1^2 \omega_1^2 + I_2^2 \omega_2^2 + I_3^2 \omega_3^2) - \frac{1}{2I_1 I_2 I_3} (I_2 I_3 n_1^2 + I_3 I_1 n_2^2 + I_1 I_2 n_3^2),$$

and set  $G = \underline{G} \circ \Psi^{-1}$ . Then,  $\{H, G\} = 0$  and  $\{L_z, G\} = 0$ . We set  $\Phi = (H, L_z, G): T^*\text{SO}(3) \rightarrow \mathbb{R}^3$ . Since  $I_1 < I_2 < I_3$ , by (7),

$$S_H = \{ (q_1, q_2, q_3, 0, 0, 0) \in T^*\text{SO}(3) \mid n_3 = \pm 1 \}.$$

Then,

$$\Phi(S_H) = \left\{ \left( \frac{1}{2I_1 I_2}, 0, -\frac{1}{2I_3} \right) \right\}.$$

By Theorem 1.7, the fiber  $\Phi^{-1}((2I_1 I_2)^{-1}, 0, -(2I_3)^{-1})$  is  $\zeta_{\text{MVZ}}$ -superheavy.

*Remark 2.13.* We can also apply our main theorem to other famous Liouville integrable systems such as the Euler top [Eu, Ar]. However, the corresponding  $\zeta_{\text{MVZ}}$ -superheavy fiber of the Euler top contains the zero-section which is already known to be  $\zeta_{\text{MVZ}}$ -superheavy. In this sense, our theorem gives only trivial results for such examples.

**2.3. On the existence of infinitely many non-displaceable fibers.** It is a natural question to ask whether a Liouville integrable system has infinitely many non-displaceable fibers. Along this line, we have the following result.

Let  $(N, g)$  be a closed Riemannian manifold. Given a positive number  $r$ , let

$$S_{g,r}^*N = \{(q, p) \in T^*N \mid \|p\|_g = r\} \quad \text{and} \quad B_{g,r}^*N = \{(q, p) \in T^*N \mid \|p\|_g < r\}$$

denote the sphere subbundle of radius  $r$  and the open ball subbundle of radius  $r$ , respectively.

**Proposition 2.14.** *Let  $(N, g)$  be a closed Riemannian manifold. Assume that for any positive number  $r$  there exist a positive number  $R$  with  $R > r$  and a partial symplectic quasi-state  $\zeta_R: C_c(T^*N) \rightarrow \mathbb{R}$  such that  $S_{g,R}^*N$  is  $\zeta_R$ -superheavy. Let  $H: T^*N \rightarrow \mathbb{R}$  be a Hamiltonian such that  $H^{-1}((-\infty, c])$  is compact for any  $c \in \mathbb{R}$ . Then, every moment map  $\Phi = (\Phi_1, \dots, \Phi_k): T^*N \rightarrow \mathbb{R}^k$  with  $\Phi_1 = H$  has infinitely many non-displaceable fibers.*

We prove Proposition 2.14 in Section 5.

The authors do not know examples of Riemannian manifolds satisfying the assumption of Proposition 2.14. However, the authors expect that any closed Riemannian manifold satisfies the assumption due to the following reason. Given a Riemannian metric  $g$  on  $N$  and a positive number  $R$ , it is known that the Rabinowitz Floer homology of  $S_{g,R}^*N$  is non-trivial [CFO]. Thus, one can construct a Rabinowitz spectral invariant (with respect to the fundamental class) from the Rabinowitz Floer homology through Albers–Fauenfelder’s construction [AF10]. We expect that the asymptotization  $\zeta$  of that spectral invariant is a partial symplectic quasi-state and  $S_{g,R}^*N$  is  $\zeta$ -superheavy since  $\zeta$  is constructed from the Rabinowitz Floer theory of  $S_{g,R}^*N$ .

By Proposition 2.14 and the above expectation, we pose the following conjecture.

**Conjecture 2.15.** *Let  $N$  be a closed manifold. Let  $H: T^*N \rightarrow \mathbb{R}$  be a Hamiltonian such that  $H^{-1}((-\infty, c])$  is compact for any  $c \in \mathbb{R}$ . Then, every moment map  $\Phi = (\Phi_1, \dots, \Phi_k): T^*N \rightarrow \mathbb{R}^k$  with  $\Phi_1 = H$  has infinitely many non-displaceable fibers.*

Actually, this conjecture is true when  $\Phi$  is the spherical pendulum (Remark 2.7) or a convex Hamiltonian (Remark 2.3).

### 3. PRELIMINARIES

In this section, we first set conventions and notation. Then we define partial symplectic quasi-states. Let  $(M, \omega)$  be a symplectic manifold.

**3.1. Conventions and notation.** Let  $H$  be a one-periodic in time Hamiltonian with compact support, i.e., a smooth function  $H: [0, 1] \times M \rightarrow \mathbb{R}$  with compact support. We set  $H_t = H(t, \cdot)$  for  $t \in [0, 1]$ . The *Hamiltonian vector field*  $X_{H_t} \in \mathfrak{X}(M)$  associated to  $H_t$  is defined by

$$\iota_{X_{H_t}}\omega = -dH_t.$$

The *Hamiltonian isotopy*  $\{\varphi_H^t\}_{t \in \mathbb{R}}$  associated to  $H$  is defined by

$$\begin{cases} \varphi_H^0 = \text{id}, \\ \frac{d}{dt}\varphi_H^t = X_{H_t} \circ \varphi_H^t \quad \text{for all } t \in \mathbb{R}, \end{cases}$$

and its time-one map  $\varphi_H = \varphi_H^1$  is referred to as the *Hamiltonian diffeomorphism with compact support* generated by  $H$ . Let  $\text{Ham}(M)$  denote the group of Hamiltonian diffeomorphisms of  $M$  with compact supports.

**3.2. Partial symplectic quasi-states.** Let  $C_c^\infty(M)$  denote the set of smooth functions on  $M$  with compact supports.

**Definition 3.1** ([EP06, FOOO19, PR, KO19b]). A *partial symplectic quasi-state* on  $(M, \omega)$  is a functional  $\zeta: C_c(M) \rightarrow \mathbb{R}$  satisfying the following conditions.

**Normalization:** There exists a non-empty compact subset  $K_\zeta$  of  $M$  such that  $\zeta(F) = 1$  for any function  $F \in C_c(M)$  with  $F|_{K_\zeta} \equiv 1$ .

**Stability:** For any  $H_1, H_2 \in C_c(M)$ , we have

$$\min_M(H_1 - H_2) \leq \zeta(H_1) - \zeta(H_2) \leq \max_M(H_1 - H_2).$$

In particular, **Monotonicity** holds:  $\zeta(H_1) \leq \zeta(H_2)$  if  $H_1 \leq H_2$ .

**Semi-homogeneity:**  $\zeta(sH) = s\zeta(H)$  for any  $H \in C_c(M)$  and any  $s > 0$ .

**Hamiltonian Invariance:**  $\zeta(H \circ \phi) = \zeta(H)$  for any  $H \in C_c(M)$  and any  $\phi \in \text{Ham}(M)$ .

**Vanishing:**  $\zeta(H) = 0$  for any  $H \in C_c(M)$  whose support is displaceable.

**Quasi-subadditivity:**  $\zeta(H_1 + H_2) \leq \zeta(H_1) + \zeta(H_2)$  for any  $H_1, H_2 \in C_c^\infty(M)$  satisfying  $\{H_1, H_2\} = 0$ .

*Remark 3.2.* There are different definitions of partial symplectic quasi-state. Our definition is based on [KO19b], but our definition is slightly different from that one. In [KO19b], they consider the different normalization condition  $\zeta(a) = a$  for every real number  $a$ . In this paper, since we consider open symplectic manifolds and functions with compact supports, we cannot define  $\zeta(a)$  unless  $a = 0$ . This is why we take a slightly different normalization condition. One can easily prove that our definition and the original one are equivalent when  $M$  is closed.

To prove Theorem 1.9, we require the following propositions.

**Proposition 3.3** ([EP09, Proposition 4.1]). *Let  $X$  be a compact subset of  $M$ .*

- (i)  *$X$  is  $\zeta$ -heavy if and only if  $\zeta(H) = 0$  for any  $H \in C_c(M)$  satisfying  $H \leq 0$  and  $H|_X \equiv 0$ .*
- (ii)  *$X$  is  $\zeta$ -superheavy if and only if  $\zeta(H) = 0$  for any  $H \in C_c(M)$  satisfying  $H \geq 0$  and  $H|_X \equiv 0$ .*

**Proposition 3.4** ([KO19a, Proposition 3.16]). *Let  $U$  be an open subset of  $M$  that is displaceable from a  $\zeta$ -superheavy subset. Then  $\zeta(F) = 0$  for any function  $F \in C_c(M)$  with support in  $U$ .*

The proof of Theorem 1.9 is similar to that of [KO19b, Theorem 2.5]. We provide the proof for readers' convenience. We also note that we can apply the same argument as [Ka18] to prove Theorem 1.9 when  $X$  is the zero-section.

*Proof of Theorem 1.9.* Let  $Y = \Phi^{-1}(p)$ ,  $p \in \Phi(M)$ , be an  $X$ -stem. Take any function  $H: \mathbb{R}^k \rightarrow \mathbb{R}$  with compact support which vanishes on an open neighborhood  $V$  of  $p$ . First we claim that  $\zeta(\Phi^*H) \leq 0$ .

Consider a finite open covering  $\mathcal{U} = \{U_1, \dots, U_d\}$  of  $\Phi(M) \cap \text{supp}(H)$  so that each  $\Phi^{-1}(U_i)$  is displaceable from itself or from  $X$ . Here we note that  $\text{supp}(H) \cap V = \emptyset$ . Take a partition of unity  $\{\rho_1, \dots, \rho_d\}$  subordinated to  $\mathcal{U}$ . Namely,  $\text{supp}(\rho_i) \subset U_i$  for any  $i$  and

$$\sum_{i=1}^d \rho_i|_{\Phi(M) \cap \text{supp}(H)} \equiv 1.$$

Since  $\text{supp}(\Phi^*(\rho_i H)) \subset \Phi^{-1}(U_i)$ , by the vanishing properties of  $\zeta$  (from Definition 3.1 and Proposition 3.4),

$$\zeta(\Phi^*(\rho_i H)) = 0$$

for any  $i$ . Since  $\{\Phi^*(\rho_i H), \Phi^*(\rho_j H)\} = 0$  for any  $i$  and  $j$ , by the quasi-subadditivity,

$$\zeta(\Phi^* H) = \zeta\left(\sum_{i=1}^d \Phi^*(\rho_i H)\right) \leq \sum_{i=1}^d \zeta(\Phi^*(\rho_i H)) = 0,$$

and this completes the proof of the claim.

Now given any function  $G \in C_c(M)$  satisfying  $G \geq 0$  and  $G|_Y \equiv 0$ , one can find a function  $H: \mathbb{R}^k \rightarrow \mathbb{R}$  with compact support and an open neighborhood  $V$  of  $p$  with  $H|_V \equiv 0$  such that  $G \leq \Phi^* H$ . By the monotonicity and the claim,

$$0 = \zeta(0) \leq \zeta(G) \leq \zeta(\Phi^* H) \leq 0.$$

Here the fact that  $\zeta(0) = 0$  follows from the vanishing property since  $\text{supp}(0) = \emptyset$  is displaceable.

Therefore,  $\zeta(G) = 0$ . By Proposition 3.3 (ii),  $Y$  is  $\zeta$ -superheavy.  $\square$

We obtain the following corollary of Theorem 1.9 which is an analogue of the main result in [KO19b].

**Corollary 3.5.** *Let  $(M, \omega)$  be a symplectic manifold. Let  $\zeta: C_c(M) \rightarrow \mathbb{R}$  be a partial symplectic quasi-state on  $(M, \omega)$ , and  $X$  a  $\zeta$ -superheavy subset of  $M$ . Let  $H: M \rightarrow \mathbb{R}$  be a Hamiltonian such that  $H^{-1}((-\infty, c])$  is compact for any  $c \in \mathbb{R}$ . Then, every moment map  $\Phi = (\Phi_1, \dots, \Phi_k): M \rightarrow \mathbb{R}^k$  with  $\Phi_1 = H$  has a fiber that is non-displaceable from itself and from  $X$ .*

*Proof.* Arguing by contradiction, assume that every fiber of  $\Phi$  is displaceable from itself or from  $X$ . By the assumption on  $H$ , every fiber of  $\Phi$  is compact. Then, every fiber is an  $X$ -stem. Since  $X$  is  $\zeta$ -superheavy, by Theorem 1.9, every fiber is  $\zeta$ -superheavy. Since all fibers are mutually disjoint, it contradicts Theorem 1.3 (i) and (iii).  $\square$

#### 4. PROOFS OF THE MAIN RESULTS

In this section, we prove the main results stated in Section 1.2. Let  $N$  be a closed manifold. Let  $\pi: T^*N \rightarrow N$  denote the natural projection. We equip  $T^*N$  with the standard symplectic form  $\omega_0$ .

**4.1. Proof of Theorem 1.10.** For the sake of applications in Sections 2.3 and 5, we generalize condition  $(\star)$  as follows.

**Definition 4.1.** Let  $\Sigma$  be a compact subset of  $T^*N$ . A (time-independent) Hamiltonian  $H: T^*N \rightarrow \mathbb{R}$  satisfies *condition  $(\star)_\Sigma$*  if the following conditions hold.

- (i) For any  $c \in \mathbb{R}$  the sublevel set  $H^{-1}((-\infty, c]) \subset T^*N$  is compact.
- (ii) For any  $q \in N$ ,

$$H|_{T_q^*N \cap \Sigma} \equiv \min_{p \in T_q^*N} H(q, p).$$

We note that condition  $(\star)_\Sigma$  is equivalent to condition  $(\star)$  when  $\Sigma = 0_N$ . If a Hamiltonian  $H: T^*N \rightarrow \mathbb{R}$  satisfies condition  $(\star)_\Sigma$ , then the value  $m_H$  exists and the set  $S_H \subset T^*N$  is defined (see (1) and (2) for the definitions).

In this section, we prove the following theorem which generalizes Theorem 1.10.

**Theorem 4.2.** *Let  $N$  be a closed manifold,  $\Sigma$  a compact subset of  $T^*N$ , and  $\Phi = (\Phi_1, \dots, \Phi_k): T^*N \rightarrow \mathbb{R}^k$  a moment map. Assume that  $\Phi_1$  satisfies condition  $(\star)_\Sigma$  and that the set  $\Phi(S_{\Phi_1})$  is a singleton, i.e.,  $\Phi(S_{\Phi_1}) = \{y_0\}$  for some  $y_0 \in \mathbb{R}^k$ . Then, every fiber of  $\Phi$ , other than  $\Phi^{-1}(y_0)$ , is displaceable from  $\Sigma$ . In particular, the fiber  $\Phi^{-1}(y_0)$  is a  $\Sigma$ -stem. Hence, by Theorem 1.9,  $\Phi^{-1}(y_0)$  is  $\zeta$ -superheavy for any partial symplectic quasi-state  $\zeta$  on  $(T^*N, \omega_0)$  such that  $\Sigma$  is  $\zeta$ -superheavy.*

Therefore, applying Theorem 4.2 for  $\Sigma = 0_N$  yields Theorem 1.10. To prove Theorem 4.2, we require the following lemma.

**Lemma 4.3.** *Let  $\Sigma$  be a compact subset of  $T^*N$  and  $H: T^*N \rightarrow \mathbb{R}$  a Hamiltonian satisfying condition  $(\star)_\Sigma$ . Then, for any  $c \in \mathbb{R}$  with  $c < m_H$ , the level set  $H^{-1}(c) \subset T^*N$  is displaceable from  $\Sigma$ .*

Before proving Lemma 4.3, we show the following well-known fact.

**Lemma 4.4.** *Let  $X$  be a compact subset of  $T^*N$  and  $f: N \rightarrow \mathbb{R}$  a smooth function on  $N$ . Then the set*

$$\Gamma_f(X) = \{ (q, p + df_q) \in T^*N \mid (q, p) \in X \}$$

*is Hamiltonian isotopic to  $X$ .*

*Proof.* Let  $f: N \rightarrow \mathbb{R}$  be a smooth function. Let  $U \subset T^*N$  be an open neighborhood of  $\bigcup_{t \in [0,1]} \Gamma_{tf}(X)$ . Choose a smooth function  $\rho: T^*N \rightarrow \mathbb{R}$  with compact support such that  $\rho|_U \equiv 1$ . Then, the (time-independent) Hamiltonian  $\rho \cdot (f \circ \pi): T^*N \rightarrow \mathbb{R}$  has a compact support and gives the desired Hamiltonian isotopy between  $X$  and  $\Gamma_f(X)$ . Indeed, for any  $(q, p) \in X$  and any  $t \in [0, 1]$ ,

$$\varphi_{\rho \cdot (f \circ \pi)}^t(q, p) = (q, p + t \cdot df_q)$$

and hence  $\varphi_{\rho \cdot (f \circ \pi)}(X) = \Gamma_f(X)$ . This finishes the proof of Lemma 4.4.  $\square$

To prove Lemma 4.3, we use a generalized version of Contreras' argument [Co, Proposition 8.2].

*Proof of Lemma 4.3.* Let  $g$  be a Riemannian metric on  $N$ . By condition  $(\star)_\Sigma$ , for each  $q \in N$  the restricted function  $H|_{T_q^*N \cap \Sigma}$  is constant and let  $c_q$  denote that constant. Then,  $m_H = \max_{q \in N} c_q$ .

Choose  $c \in \mathbb{R}$  such that  $c < m_H$ . By Lemma 4.4, it is sufficient to prove that there exists a function  $f': N \rightarrow \mathbb{R}$  such that  $\Gamma_{f'}(\Sigma) \cap H^{-1}(c) = \emptyset$ .

Take a non-empty open subset  $U$  of  $N$  so that  $\{c_q\}_{q \in U} \subset (c, m_H]$ . Choose a smooth function  $f: N \rightarrow \mathbb{R}$  whose critical points are contained in  $U$ . Since  $N \setminus U$  is compact and  $df_q \neq 0$  for any  $q \in N \setminus U$ , the number  $R_1 = \min_{q \in N \setminus U} \|df_q\|_g$  is positive. We set  $\Sigma|_{N \setminus U} = \Sigma \cap T^*N|_{N \setminus U}$  where  $T^*N|_{N \setminus U} \subset T^*N$  is the subbundle restricted to  $N \setminus U$ . By condition  $(\star)_\Sigma$ , the sets  $\Sigma|_{N \setminus U}$  and  $H^{-1}((-\infty, c])$  are compact. Hence there exists a positive number  $R_2$  such that

$$\Sigma|_{N \setminus U} \cup H^{-1}((-\infty, c]) \subset B_{g, R_2}^*N|_{N \setminus U}.$$

We set  $R_3 = 2R_2/R_1$ . Now we claim that

$$(8) \quad \Gamma_{R_3 f}(\Sigma|_{N \setminus U}) \cap H^{-1}((-\infty, c]) = \emptyset.$$

By the choice of  $R_2$ , it is enough to show that

$$(9) \quad \Gamma_{R_3 f} \left( B_{g, R_2}^*N|_{N \setminus U} \right) \cap B_{g, R_2}^*N|_{N \setminus U} = \emptyset.$$

Arguing by contradiction, assume that there exists a point  $(q_0, p_0)$  in the left hand side of (9). Recall that

$$\Gamma_{R_3 f} \left( B_{g, R_2}^*N|_{N \setminus U} \right) = \left\{ (q, p + R_3 \cdot df_q) \in T^*N \mid (q, p) \in B_{g, R_2}^*N|_{N \setminus U} \right\}.$$

Since  $(q_0, p_0) \in \Gamma_{R_3 f} \left( B_{g, R_2}^*N|_{N \setminus U} \right)$ , we have  $\|R_3 \cdot df_{q_0} - p_0\|_g < R_2$ . Since  $(q_0, p_0) \in B_{g, R_2}^*N|_{N \setminus U}$ , we have  $\|p_0\|_g < R_2$ . Thus, by the triangle inequality,

$$\|R_3 \cdot df_{q_0}\|_g \leq \|R_3 \cdot df_{q_0} - p_0\|_g + \|p_0\|_g < R_2 + R_2 = 2R_2.$$

Therefore, by the choice of  $R_1$  and the definition of  $R_3$ , we have

$$R_1 \leq \|df_{q_0}\|_g < \frac{2R_2}{R_3} = R_1,$$

and we obtain a contradiction. Therefore, (8) holds.

Let  $q \in U$ . By condition  $(\star)_\Sigma$  and  $\{c_q\}_{q \in U} \subset (c, m_H]$ , for any  $p \in T_q^*N \cap \Sigma$  we have

$$H(q, p + R_3 \cdot df_q) \geq H(q, p) = c_q > c.$$

Namely,  $\Gamma_{R_3 f}(\Sigma|_U) \cap H^{-1}((-\infty, c]) = \emptyset$ .

Combining with (8), we conclude that  $\Gamma_{R_3 f}(\Sigma) \cap H^{-1}((-\infty, c]) = \emptyset$ . In particular,  $H^{-1}(c)$  is displaceable from  $\Gamma_{R_3 f}(\Sigma)$ . By Lemma 4.4,  $H^{-1}(c)$  is displaceable from  $\Sigma$ . This completes the proof of Lemma 4.3.  $\square$

*Remark 4.5.* When the authors first found and proved Lemma 4.3, they did not know Contreras' argument. Seongchan Kim pointed out that Contreras had already used a similar technique. They would like to thank his pointing out.

Now we are in a position to prove Theorem 4.2.

*Proof of Theorem 4.2.* Let  $y = (y_1, \dots, y_k) \in \mathbb{R}^k$ . If  $y \in \mathbb{R}^k \setminus \Phi(\Sigma)$ , then  $\Phi^{-1}(y) \cap \Sigma = \emptyset$ . In particular, the fiber  $\Phi^{-1}(y)$  is displaceable from  $\Sigma$ .

Assume that  $y \in \Phi(\Sigma)$ . Then, in particular,  $y_1 \in \Phi_1(\Sigma)$ . Since  $\Phi_1$  satisfies condition  $(\star)_\Sigma$ ,

$$(10) \quad y_1 \leq \max_{q \in N} \Phi_1|_{T_q^*N \cap \Sigma} = m_{\Phi_1}.$$

If  $y_1 \neq m_{\Phi_1}$ , then (10) and Lemma 4.3 imply that  $\Phi_1^{-1}(y_1)$  is displaceable from  $\Sigma$  and hence so is  $\Phi^{-1}(y) \subset \Phi_1^{-1}(y_1)$ .

If  $y_1 = m_{\Phi_1}$ , then  $\Phi^{-1}(y) \subset \Phi_1^{-1}(y_1) = S_{\Phi_1}$ . Since  $\Phi(S_{\Phi_1}) = \{y_0\}$ , we have  $y = y_0$ .

Therefore, the above argument implies that every fiber of  $\Phi$ , other than  $\Phi^{-1}(y_0)$ , is displaceable from  $\Sigma$ . By condition  $(\star)_\Sigma$ , the sublevel set  $\Phi_1^{-1}((-\infty, m_{\Phi_1}])$  is compact and hence so is the fiber  $\Phi^{-1}(y_0) \subset \Phi_1^{-1}((-\infty, m_{\Phi_1}])$ . Therefore,  $\Phi^{-1}(y_0)$  is a  $\Sigma$ -stem. This finishes the proof of Theorem 4.2.  $\square$

#### 4.2. Proof of Theorem 1.11.

*Proof.* Take  $y = (y_1, \dots, y_k) \in \Phi(T^*N) \subset \mathbb{R}^k$ , where  $\Phi = (H_1, \dots, H_k): T^*N \rightarrow \mathbb{R}^k$ . If  $y_i > m_{H_i}$  for some  $i \in \{1, \dots, k\}$ , then  $H_i^{-1}(y_i)$  is disjoint from the zero-section  $0_N$  and hence so is  $\Phi^{-1}(y) \subset H_i^{-1}(y_i)$ . If  $y_i < m_{H_i}$  for some  $i \in \{1, \dots, k\}$ , then applying Lemma 4.3 for  $\Sigma = 0_N$ ,  $H_i^{-1}(y_i)$  is displaceable from  $0_N$  and hence so is  $\Phi^{-1}(y) \subset H_i^{-1}(y_i)$ .

The above argument then implies that every fiber of  $\Phi$ , other than  $\Phi^{-1}(m_\Phi)$ , is displaceable from  $0_N$ , where  $m_\Phi = (m_{H_1}, \dots, m_{H_k}) \in \mathbb{R}^k$ . By Corollary 3.5,  $\Phi^{-1}(m_\Phi)$  is non-displaceable from  $0_N$ . Thus,

$$\bigcap_{i=1}^k S_{H_i} = \Phi^{-1}(m_\Phi) \cap 0_N \neq \emptyset.$$

This completes the proof of Theorem 1.11.  $\square$

**4.3. Proof of Proposition 1.13.** Proposition 1.13 immediately follows from Theorem 1.3 (iii), Proposition 1.6 and the following assertion.

**Proposition 4.6.** *Let  $X$  be a compact subset of  $T^*N$ . If  $\pi(X) \neq N$ , then  $X$  is displaceable from the zero-section  $0_N$ .*

*Proof.* By Lemma 4.4, it is enough to show that  $\Gamma_f(X)$  is displaceable from  $0_N$  for some smooth function  $f: N \rightarrow \mathbb{R}$ . Let  $f: N \rightarrow \mathbb{R}$  be a smooth function whose critical points are all contained in  $N \setminus \pi(X)$ . Then  $df_q \neq 0$  for any  $(q, p) \in X$ . Since  $X$  is compact, there exists a positive number  $R_0 > 0$  such that for any  $(q, p) \in X$ ,  $R_0 \cdot df_q \neq -p$ . It means that

$$\Gamma_{R_0 f}(X) \cap 0_N = \emptyset.$$

This completes the proof of Proposition 4.6.  $\square$

## 5. PROOF OF PROPOSITION 2.14

In this section, we prove Proposition 2.14 and provide another corollary (Corollary 5.1) of Theorem 4.2. Under the assumption of Proposition 2.14, there are many disjoint superheavy subsets in  $T^*N$ . We use these superheavy subsets to prove the existence of many non-displaceable fibers. This idea comes from [KO19b].

*Proof of Proposition 2.14.* Arguing by contradiction, assume that the moment map  $\Phi$  has finitely many non-displaceable fibers. Let  $\Phi^{-1}(y_1), \dots, \Phi^{-1}(y_\ell)$  be all the non-displaceable fibers of  $\Phi$ , where  $y_1, \dots, y_\ell \in \mathbb{R}^k$ . By the assumption on  $H$ , the fibers  $\Phi^{-1}(y_i)$ ,  $i = 1, \dots, \ell$ , are compact. Then there exists a positive number  $r$  such that

$$(11) \quad \bigcup_{i=1}^{\ell} \Phi^{-1}(y_i) \subset B_{g,r}^*N.$$

By assumption, there exist a positive number  $R$  with  $R > r$  and a partial symplectic quasi-state  $\zeta_R: C_c(T^*N) \rightarrow \mathbb{R}$  such that  $S_{g,R}^*N$  is  $\zeta_R$ -superheavy. Then, by (11),

$$(12) \quad \left( \bigcup_{i=1}^{\ell} \Phi^{-1}(y_i) \right) \cap S_{g,R}^*N = \emptyset.$$

Since  $S_{g,R}^*N$  is  $\zeta_R$ -superheavy, by Corollary 3.5, there exists  $y_0 \in \Phi(T^*N)$  such that the fiber  $\Phi^{-1}(y_0)$  is non-displaceable from itself and from  $S_{g,R}^*N$ . Therefore,  $y_0 \in \{y_1, \dots, y_\ell\}$  and  $\Phi^{-1}(y_0) \cap S_{g,R}^*N \neq \emptyset$ . It contradicts (12) and we complete the proof of Proposition 2.14.  $\square$

Moreover, we have the following corollary of Theorem 4.2.

**Corollary 5.1.** *Let  $N$  be a closed manifold,  $\Sigma$  a compact subset of  $T^*N$ , and  $H: T^*N \rightarrow \mathbb{R}$  a Hamiltonian satisfying condition  $(\star)_\Sigma$ . Assume that there exists a partial symplectic quasi-state  $\zeta: C_c(T^*N) \rightarrow \mathbb{R}$  on  $(T^*N, \omega_0)$  such that  $\Sigma$  is  $\zeta$ -superheavy. Then, the level set  $H^{-1}(m_H) \subset T^*N$  is non-displaceable from itself and from  $\Sigma$ .*

*Proof.* By Theorem 4.2, the level set  $H^{-1}(m_H)$  is a  $\Sigma$ -stem. By Corollary 3.5,  $H^{-1}(m_H)$  is non-displaceable from itself and from  $\Sigma$ .  $\square$

We provide an example of Corollary 5.1.

**Example 5.2.** Let  $(N, g)$  be a closed Riemannian manifold and  $r$  a non-negative number. Let  $H: T^*N \rightarrow \mathbb{R}$  be a Hamiltonian of the form

$$H(q, p) = \rho(\|p\|^2) + U(q),$$



where  $\rho: [0, \infty) \rightarrow \mathbb{R}$  is a smooth function which attains its minimum value at  $r^2$  and  $U: N \rightarrow \mathbb{R}$  is a smooth potential. Then  $H$  satisfies condition  $(\star)_\Sigma$  where  $\Sigma = S_{g,r}^*N$ . Assume that there exists a partial symplectic quasi-state  $\zeta: C_c(T^*N) \rightarrow \mathbb{R}$  on  $(T^*N, \omega_0)$  such that  $S_{g,r}^*N$  is  $\zeta$ -superheavy. Then, by Corollary 5.1, the level set  $H^{-1}(m_H) \subset T^*N$  is non-displaceable from itself and from  $S_{g,r}^*N$ .

## ACKNOWLEDGMENTS

The authors cordially thank Felix Schlenk for reading a preliminary version very carefully and for giving very helpful comments. They also thank Seongchan Kim for giving them the trigger to study the present topic. When they talked with him about a different mathematical topic, he explained how interesting the spherical pendulum is to them. This started the first author to consider a superheavy fiber of the spherical pendulum. He also suggested some other integrable systems (Examples 2.4, 2.5 and 2.8) and gave remarks (Remarks 2.7 and 4.5). They also sincerely thank Mitsuaki Kimura, Takahiro Matsushita, and Yuhei Suzuki for fruitful discussions and warmhearted advices.

## REFERENCES

- [AFOOO] M. Abouzaid, K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Quantum cohomology and split generation in Lagrangian Floer theory*, in preparation.
- [ABM] M. Abreu, M. Borman and D. McDuff, *Displacing Lagrangian toric fibers by extended probes*, *Algebr. Geom. Topol.* **14** (2014), no. 2, 687–752.
- [AM] M. Abreu and L. Macarini, *Remarks on Lagrangian intersections in toric manifolds*, *Trans. Amer. Math. Soc.* **365** (2013), no. 7, 3851–3875.
- [AF08] P. Albers and U. Frauenfelder, *A nondisplaceable Lagrangian torus in  $T^*S^2$* , *Comm. Pure Appl. Math.* **61** (2008), no. 8, 1046–1051.
- [AF10] P. Albers and U. Frauenfelder, *Spectral invariants in Rabinowitz-Floer homology and global Hamiltonian perturbations*, *J. Mod. Dyn.* **4** (2010), no. 2, 329–357.
- [Ar] V. I. Arnol'd, *Mathematical methods of classical mechanics*, Translated from the Russian by K. Vogtmann and A. Weinstein. Second edition. Graduate Texts in Mathematics, 60. Springer-Verlag, New York, 1989.
- [BEP] P. Biran, M. Entov and L. Polterovich, *Calabi quasimorphisms for the symplectic ball*, *Commun. Contemp. Math.* **6** (2004), no. 5, 793–802.
- [BKS] J. Brendel, S. Kim and F. Schlenk, *private communication*, 2019.
- [Ch] C.-H. Cho, *Holomorphic discs, spin structures, and Floer cohomology of the Clifford torus*, *Int. Math. Res. Not.* **2004** (2004), no. 35, 1803–1843.
- [CKO] Y. Cho, Y. Kim and Y.-G. Oh, *Lagrangian fibers of Gelfand–Cetlin systems*, Preprint (2017), [arXiv:1704.07213](https://arxiv.org/abs/1704.07213).
- [CFO] K. Cieliebak, U. Frauenfelder and A. Oancea, *Rabinowitz Floer homology and symplectic homology*, *Ann. Scient. Éc. Norm. Sup. (4)* **43** (2010), no. 6, 957–1015.
- [CFP] K. Cieliebak, U. Frauenfelder and G. Paternain, *Symplectic topology of Mañé’s critical values*, *Geom. Topol.* **14** (2010), no. 3, 1765–1870.
- [Cl] A. Clebsch, *Ueber die Bewegung eines Körpers in einer Flüssigkeit*, *Math. Ann.* **3** (1870), no. 2, 238–262.
- [Co] G. Contreras, *The Palais-Smale condition on contact type energy levels for convex Lagrangian systems*, *Calc. Var.* **27** (2006), no. 3, 321–395.
- [CB] R. Cushman and L. Bates, *Global aspects of classical integrable systems*, Second edition. Birkhäuser/Springer, Basel, 2015.
- [En] M. Entov, *Quasi-morphisms and quasi-states in symplectic topology*, The Proceedings of the International Congress of Mathematicians (Seoul, 2014).
- [EP06] M. Entov and L. Polterovich, *Quasi-states and symplectic intersections*, *Comment. Math. Helv.* **81** (2006), no. 1, 75–99.
- [EP09] M. Entov and L. Polterovich, *Rigid subsets of symplectic manifolds*, *Compos. Math.* **145** (2009), no. 3, 773–826.
- [Eu] L. Euler, *Du mouvement de rotation des corps solides autour d’un axe variable*, *Mémoires l’académie des sciences de Berlin* **14** (1758), 154–193.
- [Fl] A. Floer, *Morse theory for Lagrangian intersections*, *J. Differential Geom.* **28** (1988), no. 3, 513–547.

- [FOOO10] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian Floer theory on compact toric manifolds. I*, Duke Math. J. **151** (2010), no. 1, 23–174.
- [FOOO11] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian Floer theory on compact toric manifolds II: bulk deformations*, Selecta Math. (N.S.) **17** (2011), no. 3, 609–711.
- [FOOO12] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Toric degeneration and nondisplaceable Lagrangian tori in  $S^2 \times S^2$* , Int. Math. Res. Not. IMRN **2012** (2012), no. 13, 2942–2993.
- [FOOO19] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Spectral invariants with bulk, quasimorphisms and Lagrangian Floer theory*, Mem. Amer. Math. Soc. **260** (2019), no. 1254.
- [Gr] M. Gromov, *Pseudo holomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), no. 2, 307–347.
- [Ho] H. Hofer, *Lagrangian embeddings and critical point theory*, Ann. Inst. H. Poincaré Anal. Non Linéaire **2** (1985), no. 6, 407–462.
- [HLS] S. Hu, E. Leandro and M. Santoprete, *On the topology of the double spherical pendulum*, Regul. Chaotic Dyn. **17** (2012), no. 1, 36–53.
- [Ja] C. Jacobi, *Note von der geodätischen Linie auf einem Ellipsoid und den verschiedenen Anwendungen einer merkwürdigen analytischen Substitution*, J. Reine Angew. Math. **19** (1839), 309–313.
- [Ka17] M. Kawasaki, *Bavard’s duality theorem on conjugation-invariant norms*, Pacific J. Math. **288** (2017), no. 1, 157–170.
- [Ka18] M. Kawasaki, *Function theoretical applications of Lagrangian spectral invariants*, Preprint (2018), [arXiv:1811.00527](https://arxiv.org/abs/1811.00527).
- [KO19a] M. Kawasaki and R. Orita, *Disjoint superheavy subsets and fragmentation norms*, J. Topol. Anal., Online Ready, doi:10.1142/S179352532050017X.
- [KO19b] M. Kawasaki and R. Orita, *Existence of pseudo-heavy fibers of moment maps*, Preprint (2019), [arXiv:1901.09395](https://arxiv.org/abs/1901.09395).
- [Ki] S. Kim, *private communication*, 2019.
- [KLS] Y. Kim, J. Lee and F. Sanda, *Detecting non-displaceable toric fibers on compact toric manifolds via tropicalizations*, Internat. J. Math. **30** (2019), no. 1, 1950003.
- [Ko] S. Kowalewski, *Sur le problème de la rotation d’un corps solide autour d’un point fixe*, Acta. Math. **12** (1889), 177–232.
- [La] J. Lagrange, *Mécanique analytique*, Paris, 1788.
- [LS] F. Laudenbach and J.-C. Sikorav, *Persistence d’intersection avec la section nulle au cours d’une isotopie hamiltonienne dans un fibré cotangent*, Invent. Math. **82** (1985), no. 2, 349–357.
- [MS] J. E. Marsden and J. Scheurle, *Lagrangian reduction and the double spherical pendulum*, Z. angew. Math. Phys. **44** (1993), no. 1, 17–43.
- [Mc] D. McDuff, *Displacing Lagrangian toric fibers via probes*, Low-dimensional and symplectic topology, 131–160, Proc. Sympos. Pure Math. **82**, Amer. Math. Soc., Providence, RI, 2011.
- [MVZ] A. Monzner, N. Vichery and F. Zapolsky, *Partial quasimorphisms and quasistates on cotangent bundles, and symplectic homogenization*, J. Mod. Dyn. **6** (2012), no. 2, 205–249.
- [Neu] C. Neumann, *De problemate quodam mechanico, quod ad primam integralium ultraellipticorum classem revocatur*, J. Reine Angew. Math. **56** (1859), 46–63.
- [NNU] T. Nishinou, Y. Nohara and K. Ueda, *Toric degenerations of Gelfand-Cetlin systems and potential functions*, Adv. Math. **224** (2010), no. 2, 648–706.
- [Oh97] Y.-G. Oh, *Symplectic topology as the geometry of action functional. I. Relative Floer theory on the cotangent bundle.*, J. Differential Geom. **46** (1997), no. 3, 499–577.
- [Oh99] Y.-G. Oh, *Symplectic topology as the geometry of action functional. II. Pants product and cohomological invariants.*, Comm. Anal. Geom. **7** (1999), no. 1, 1–54.
- [Oh05] Y.-G. Oh, *Construction of spectral invariants of Hamiltonian paths on closed symplectic manifolds*, in The Breadth of Symplectic and Poisson Geometry, eds. J. E. Marsden and T. Ratiu (Birkhäuser/Springer, 2005), 525–570.
- [PPS] G. Paternain, L. Polterovich and K. Siburg, *Boundary rigidity for Lagrangian submanifolds, non-removable intersections, and Aubry–Mather theory*, Mosc. Math. J. **3** (2003), no. 2, 593–619.
- [Po98] L. Polterovich, *Hofer’s diameter and Lagrangian intersections*, Int. Math. Res. Not. **1998** (1998), no. 4, 217–223.
- [Po14] L. Polterovich, *Symplectic intersections and invariant measures*, Ann. Math. Qu. Not. **38** (2014), no. 1, 81–93.
- [PR] L. Polterovich and D. Rosen, *Function theory on symplectic manifolds*, CRM Monograph Series, Vol. 34 (American Mathematical Society, 2014).
- [Sch] M. Schwarz, *On the action spectrum for closed symplectically aspherical manifolds*, Pacific J. Math. **193** (2000), no. 2, 419–461.
- [Vi] R. Vianna, *On exotic Lagrangian tori in  $\mathbb{C}\mathbb{P}^2$* , Geom. Topol. **18** (2014), no. 4, 2419–2476.

[Wu] W. Wu, *On an exotic Lagrangian torus in  $CP^2$* , Compos. Math. **151** (2015), no. 7, 1372–1394.

(Morimichi Kawasaki) RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN

*E-mail address:* [kawasaki@kurims.kyoto-u.ac.jp](mailto:kawasaki@kurims.kyoto-u.ac.jp)

(Ryuma Orita) DEPARTMENT OF MATHEMATICAL SCIENCES, TOKYO METROPOLITAN UNIVERSITY, TOKYO 192-0397, JAPAN

*E-mail address:* [ryuma.orita@gmail.com](mailto:ryuma.orita@gmail.com)

*URL:* <https://ryuma-orita.github.io/>