

題目

Causes of liver steatosis influence the severity of ischemia reperfusion injury and survival after liver transplantation in rats

脂肪肝の成因が肝移植における虚血再灌流障害に与える影響

申請者

宮地洋介¹,

¹京都大学大学院医学研究科 肝胆膵・移植外科

*"This is the peer reviewed version of the following article:
[Causes of liver steatosis influence the severity of ischemia reperfusion injury
and survival after liver transplantation in rats],
which has been published in final form at [<https://doi.org/10.1002/lt.25814>].
This article may be used for non-commercial purposes
in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions."*

Liver Transplantation**Causes of liver steatosis influence the severity of ischemia reperfusion injury and survival after liver transplantation in rats**

Journal:	<i>Liver Transplantation</i>
Manuscript ID	LT-20-040.R1
Wiley - Manuscript type:	Original Articles
Date Submitted by the Author:	n/a
Complete List of Authors:	Miyachi, Yosuke; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Yagi, Shintaro; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Hirata, Masaaki; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Iwamura, Sena; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Yao, Siyuan; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Shirai, Hisaya; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Okumura, Shinya; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Iida, Taku; Kyoto University Graduate School of Medicine, Department of Hepatobiliary Pancreas and Transplant surgery; Ito, Takashi; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Uozumi, Ryuji; Kyoto University Graduate school of medicine54 Kawaracho, Shogoin, Sakyo-ku, Kyoto city, Kyoto prefecture, 606-8507, Department of Biomedical Statistics and Bioinformatics Kaido, Toshimi; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery Uemoto, Shinji; Kyoto Daigaku Igakubu Fuzoku Byoin, Department of Hepatobiliary, Pancreas and Transplant Surgery
Keywords:	primary non-function, hemeoxygenase-1, liver sinusoidal endothelial cell, microcirculation, methionine and choline deficient diet

SCHOLARONE™
Manuscripts

1
2
3
4
5
6 **Original article**
7

8
9 **Title: Causes of liver steatosis influence the severity of ischemia reperfusion injury and**
10
11
12 **survival after liver transplantation in rats**
13
14

15
16
17
18 **Authors**
19

20
21 Yosuke Miyachi, MD¹, Shintaro Yagi, MD, PhD¹, Masaaki Hirata, MD¹, Sena Iwamura, MD¹,
22
23
24 Siyuan Yao, MD¹, Hisaya Shirai, MD¹, Shinya Okumura, MD, PhD¹, Taku Iida, MD, PhD¹,
25
26
27 Takashi Ito, MD, PhD¹, Ryuji Uozumi, PhD², Toshimi Kaido, MD, PhD¹, Shinji Uemoto, MD,
28
29
30 PhD¹
31
32
33
34
35

36 **Affiliations**
37

- 38
39 1. Division of Hepato-Biliary-Pancreatic and Transplant Surgery, Department of Surgery,
40
41
42 Graduate School of Medicine, Kyoto University, Kyoto, Japan
43
44
45 2. Biomedical Statistics and Bioinformatics, Graduate School of Medicine, Kyoto University,
46
47
48 Kyoto, Japan
49
50
51
52
53

54 **Key words**
55
56
57
58
59
60

1
2
3
4
5
6 primary non-function; hemoxygenase-1; liver sinusoidal endothelial cell; microcirculation;
7
8

9 methionine and choline deficient diet
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Footnote page**1. Abbreviations**

ALT; alanine aminotransferase

AST; aspartate aminotransferase

ATP; adenosine triphosphate

eNOS; endothelial nitric oxide synthase

ET-1; endothelin-1

FHA; fasting and hyper alimentation

H & E; hematoxylin-eosin stain

HO-1; hemeoxygenase-1

IL; interleukin

iNOS; inducible nitric oxide synthase

IRI; ischemia reperfusion injury

LSEC; liver sinusoidal endothelial cell

MCDD; methionine and choline deficient diet

MDA; malondialdehyde

NAFLD; nonalcoholic fatty liver disease

Nqo1; NAD(P)H Quinone Dehydrogenase -1

1
2
3
4
5
6 OLT; orthotopic liver transplantation
7
8

9 ORO; oil-red-O stain
10
11

12 PNF; primary non-function
13
14

15 RP1, RP3 and RP24; 1, 3 and 24 hours after reperfusion
16
17

18 SEM; scanning electron microscopy
19
20

21 TEM; transmission electron microscopy
22
23

24 TNF; Tumor necrosis factor
25
26

27 TG; triglyceride
28
29
30
31
32

33 **2. Grants and financial support**

34 Japan Society for the Promotion of Science; grant number 17K10519
35
36
37
38
39
40

41 **3. Conflict of interest; None**

42
43
44
45
46

47 **4. Corresponding author email**

48 Shintaro Yagi, MD, PhD
49
50
51

52 **ORCID;** 0000-0001-7465-5761
53
54
55
56
57
58
59
60

1
2
3
4
5
6 Division of Hepato-Biliary-Pancreatic and Transplant Surgery, Department of Surgery,
7

8
9 Graduate School of Medicine, Kyoto University, 54 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto
10

11
12 606-8507, Japan
13

14
15 Phone number: +81-75-751-4323
16

17
18 Fax number: +81-75-751-4348, E-mail: shintaro@kuhp.kyoto-u.ac.jp
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Abstract

Liver steatosis is a leading cause of graft disposal in liver transplantation; however, the degree of steatosis has been so far the almost single factor determining whether the grafts is acceptable or not. We investigated how the cause of liver steatosis affects graft function in rat orthotopic liver transplantation (OLT). OLT was performed using two types of steatotic liver grafts; the fasting and hyper alimentation (FHA) and methionine and choline deficient diet (MCDD) models. The FHA and 4-week MCDD feeding (MCDD4wk) groups showed similar liver triglyceride levels without signs of steatohepatitis; therefore, the two groups were compared in the following experiment. With 6-hour cold storage, 7-days survival rate after OLT was far worse in the FHA than the MCDD4wk group (0% vs. 100%, $P = 0.002$). With 1-hour cold storage, the FHA group showed higher aspartate and alanine aminotransferase levels and histological injury score in zone 1 and 2 at 24 hours after reperfusion than the Normal liver and MCDD4wk groups. Intrahepatic microcirculation and tissue ATP level were significantly lower in the FHA group after reperfusion. Hepatocyte necrosis, sinusoidal endothelial cell injury and abnormal swelling of mitochondria were also found in the FHA group after reperfusion. Tissue malondialdehyde levels were higher in the MCDD4wk group before and after reperfusion; however, the grafts upregulated several antioxidant enzymes soon after reperfusion.

Conclusion: Even though the degree of steatosis was equivalent, the two liver steatosis models

1
2
3
4
5
6 possessed quite unique basal characteristics and showed completely different response against
7
8
9 ischemia reperfusion injury and survival after transplantation. Our results demonstrated that the
10
11
12 degree of fat accumulation is not a single determinant for the fate of steatotic liver graft.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1. Introduction

Because of the fear of post-transplant primary non-function (PNF), liver steatosis is a leading cause of graft disposal in liver transplantation.¹ While livers with less than 30% steatosis have no increased risk of PNF, the risk increases if graft steatosis exceeds more than 30%¹; therefore, grafts with more than 60% steatosis are deemed unacceptable in US or UK.^{2,3} On the other hand, as the prevalence of patients with overweight, obesity or diabetes has increased,^{4,5} the rate of liver steatosis in donor candidates will almost certainly rise. However, considering the severe donor shortage, using as many liver grafts as possible, even though they may have severe steatosis, is essential to help patients who suffering from end-stage liver disease.

Some authors argued that even severe steatotic liver grafts can be used safely under certain clinical conditions. Wong et al. reported that none of 19 patients who received severe (more than 60%) steatotic liver grafts experienced PNF nor early allograft dysfunction.⁶ McCormack et al. also reported that PNF did not occur in the 20 cases of liver steatosis group (median 90%, range 65-100%) and postoperative survival rate was equivalent to those in the control group.⁷

Considering these results,^{6,7} steatosis itself seems to be a superficial phenomenon and there may be another key factor that determine whether the graft is acceptable for transplantation or not.

Liver steatosis is not a homogenous disease.⁸ Various factors can cause fat accumulation and its pathophysiological process is also not uniform.⁸ Therefore, it seems natural that two different

1
2
3
4
5
6 types of steatotic liver grafts, even though they have the same degree of steatosis, show different
7
8
9 graft function and survival after transplantation. Currently, various rat liver steatosis models
10
11
12 have been established worldwide,⁹⁻¹² some of these are now available in our laboratory.
13
14
15 Therefore, we can now precisely mimic the aforementioned clinical situation. From the above,
16
17
18 the aims of this study are to i) create the same degree of liver steatosis using two different
19
20
21 methods, ii) compare the graft functions using rat orthotopic liver transplantation (OLT) and iii)
22
23
24 investigate the factors affecting function or survival of steatotic liver grafts.
25
26
27
28
29

30 **2. Material & methods**

31 **2.1 Animals**

32
33
34 A total of 114 male LEW/CrIcrlj rats weighting 250 to 350g were purchased from Charles
35
36
37 River Laboratories Japan, Inc., Yokohama, Japan. The animals were housed under specific
38
39
40 pathogen-free conditions in a temperature- and humidity-controlled environment with a 12-hour
41
42
43 light/dark cycle. Except for the following liver steatosis models, rats were fed with a standard
44
45
46 diet (F-2; Oriental Bio Service, Kyoto, Japan) and tap water ad libitum. The experimental
47
48
49 protocol was approved by the institutional ethics committee of Kyoto University (Medkyo
50
51
52 18536), which met the ethical guidelines of the Declaration of Helsinki. And all animals
53
54
55
56
57 received humane care according to criteria outlined in the “Guide for the Care and Use of
58
59
60

1
2
3
4
5
6 Laboratory Animals'' prepared by the National Academy of Sciences and published by the
7
8
9 National Institutes of Health (NIH Publication No 86-23, revised in 1985).
10
11
12
13
14

15 **2.2. Experimental protocol**

16 17 18 **2.2.1 Experiment I: Comparison of hepatic steatosis models.**

19
20
21 The following two types of liver steatosis models were compared.
22
23

24 i) Fasting and hyperalimentation (FHA) model

25
26
27 The modified protocol by Delzenne et al. was used.^{9, 13} In short, the rats (n = 5) were fasted for 2
28
29 days, followed by refeeding with a fat-free, carbohydrate-rich diet (saccharose 38%, starch
30
31 38%, casein 16%, and a mineral and vitamin mix 8%, CLEA Japan, Inc., Tokyo, Japan) for 3
32
33 days. During this period, access to tap water was not limited.
34
35
36
37
38

39 ii) Methionine and choline deficient diet (MCDD) model

40
41
42 The rats were fed a methionine and choline deficient diet (sucrose 10%, starch 39%, amino acid
43
44 mix without methionine 17%, and a mineral and vitamin mix without choline 1%, Oriental
45
46 Yeast co., Ltd, Tokyo, Japan) for 2-, 4-, 6-weeks (n = 3, 5, 3 respectively). As with the previous
47
48 model, access to tap water was not limited during the feeding period.
49
50
51
52
53
54
55
56

57 **2.2.2 Experiment II-A: Comparison of survival rates with 6-hour cold storage**

58
59
60

1
2
3
4
5
6 Whole liver grafts from the FHA and MCDD groups (n = 5, each) were preserved in University
7
8
9 of Wisconsin solution (Belzer, Astellas Pharma Inc., Tokyo, Japan) for 6-hours at 4°C. After
10
11
12 cold storage, OLT was performed using a modified method by Kamada.¹⁴ Briefly, the portal
13
14
15 vein was reconstructed using the cuff method, and anastomosis of the infra-hepatic vena cava
16
17
18 was hand-sewn using 8-0 nylon. The hepatic artery was also reconstructed using a modified
19
20
21 sleeve method.¹⁵
22
23
24
25
26

27 **2.2.3 Experiment II-B: Evaluation of liver graft with 1-hour cold storage**

28
29
30 Whole liver grafts from the Normal liver, FHA and MCDD group were preserved in University
31
32
33 of Wisconsin solution for 1 hours at 4°C. After cold storage, OLT was performed as previously
34
35
36 described. Subsequently, 3, and 24 hours after portal reperfusion, serum and liver graft tissue
37
38
39 samples were collected prior to euthanasia by exsanguination (RP-3 and RP-24, respectively, n
40
41
42 = 5, at each time point for each group). The samples of 3 groups before procurement were also
43
44
45 collected and presented as controls (n = 5, for each group).
46
47
48
49
50

51 **2.3 Assay procedure**

52 **2.3.1 Measurement of serum transaminase and albumin levels**

53
54
55
56
57
58
59
60

1
2
3
4
5
6 Serum aspartate aminotransferase (AST), alanine aminotransferase (ALT) and albumin levels
7
8
9 were measured using a standard spectrophotometric method using an automated clinical
10
11
12 analyzer (JCA- BM9030, JEOL Ltd., Tokyo, Japan).
13
14
15
16
17

18 **2.3.2 Measurement of liver tissue triglyceride, adenosine triphosphate and** 19 20 21 **malondialdehyde**

22
23
24 The left lateral segment of liver in each sample was snap frozen and stored at -80°C until
25
26 subsequent assays. Triglyceride (TG) was extracted from liver tissues according to the method
27
28 of Folch et al.¹⁶ Briefly, snap frozen liver (more than 100mg) was homogenized and extracted
29
30 using a chloroform-methanol solution. The organic phase was dried and resolubilized in 2-
31
32 propanol. TG was then determined using a commercial kit (Sekisui. Medical Co., Tokyo,
33
34 Japan). The liver tissue adenosine triphosphate (ATP) was measured using AMERIC-ATP(T)
35
36 kit (Applied Medical Enzyme Research Institute Co., Tokushima, Japan) according to the
37
38 manufacturer's directions. Liver tissue malondialdehyde (MDA) was measured using the
39
40 thiobarbituric acid assay (NWLSS malondialdehyde assay, Northwest Life Science Specialties,
41
42 LLC, WA, USA) according to the manufacturer's protocol.
43
44
45
46
47
48
49
50
51
52
53
54
55
56

57 **2.3.3 Measurement of tissue microcirculation**

58
59
60

1
2
3
4
5
6 The tissue microcirculation was evaluated at the left lateral segment and middle lobe (left and
7
8
9 right side separately) using an advanced laser flow meter (Advance Bio Research Center Co.,
10
11
12 Ltd, Nagoya, Japan). The time points were before procurement, RP-1 (before abdominal
13
14
15 closure) and RP-3. Hepatic microcirculations before procurement and at RP-1 of rats euthanized
16
17
18 at RP-24 were also recorded and included as the result.
19
20
21
22
23

24 **2.3.4 Histological analysis**

25
26
27 Formalin-fixed, paraffin-embedded sections (4 μ m thickness) were stained with hematoxylin-
28
29 eosin (H & E) and oil-red-O (ORO) stain. Two independent investigators examined all tissue
30
31
32 section in a blinded fashion. The nonalcoholic fatty liver disease (NAFLD) activity score was
33
34
35 calculated using the existing scoring system.¹⁷ The severity of ischemia reperfusion injury (IRI)
36
37
38 was quantified at RP-24 with hepatocyte swelling, vacuolization, necrosis, apoptosis, and
39
40
41 neutrophil infiltration, and degree of each factor was graded as 1 = no change or negligible
42
43
44 lesions, affecting 0%–10% of the field; 2 = mild, lesions affecting 10%–40% of the field; 3 =
45
46
47 moderate, lesions affecting 40%–70% of the field; 4 = severe, lesion affecting > 70% of the
48
49
50 field.¹⁸ The scores were evaluated in 10 random fields (magnification \times 200) per slide and
51
52
53
54 averaged for each slide.¹⁸
55
56
57
58
59
60

2.3.5 Electron microscopy

The rat livers were perfused through the portal vein with a mixture of 2% glutaraldehyde and 4% paraformaldehyde and then extracted. The livers were cut into small pieces and stored overnight at 4°C. The sections were stained with saturated uranyl acetate and lead citrate and were observed with a Hitachi H-7650 electron microscope (Hitachi, Tokyo, Japan) for transmission electron microscopy (TEM). Additionally, the samples were ion-sputter-coated and observed with a Hitachi S-4700 scanning electron microscope for scanning electron microscopy (SEM). The area of sinusoidal space was identified in 10 randomly selected TEM images ($\times 700$) in each group and calculated using Image J version 1.46r (National Institutes of Health, USA). The time points were before procurement and at RP-24 in three groups. Two independent investigators including an expert pathologist of the liver evaluated the sample findings.

2.3.6 Real-time polymerase chain reaction

For the analysis of gene expression, real-time polymerase chain reaction, using TaqMan technology, was performed. The probe and primers for interleukin-1 β (IL-1 β , assay ID Rn00580432_m1), IL-6 (assay ID Rn01410330_m1), tumor necrosis factor- α (TNF- α , assay ID Rn01525859_g1), endothelin-1 (ET-1, assay ID Rn00561129_m1), endothelial nitric oxide

1
2
3
4
5
6 synthase (eNOS, assay ID Rn02132634_s1), inducible nitric oxide synthase (iNOS, assay ID
7
8
9 Rn00561646_m1), heme oxygenase-1 (HO-1, assay ID Rn01536933_m1), NAD(P)H Quinone
10
11
12 Dehydrogenase 1 (Nqo1, assay ID Rn00566528_m1) and beta-actin (β -actin, assay ID
13
14
15 Rn00667869_m1) were obtained for TaqMan gene expression assays from Applied Biosystems,
16
17
18 Life Technologies Japan Ltd., Japan.
19

20 21 22 23 24 **2.4 Statistics**

25
26
27 Data are expressed as means \pm standard error and are compared statistically using Student's t-
28
29
30 test (Experiment I, Table 2) or Tukey-Kramer analysis. For the survival study, a log-rank test
31
32
33 was performed. $P < 0.05$ was considered as a threshold; however, according to the previous
34
35
36 recommendation of American Statistical Association,^{19,20} we avoided describing $P < 0.05$ as
37
38
39 “statistically significant”; instead, we described the measured values and P values as continuous
40
41
42 quantities in the text, tables or figures as possible. All statistical analyses were performed using
43
44
45 JMP Pro, version 14.0.0 (SAS Institute, Inc., Cary, NC, USA).
46
47
48
49
50

51 **3. Results**

52 53 54 **3.1 Experiment I: Comparison of hepatic steatosis models**

55
56
57
58
59
60

1
2
3
4
5
6 According to the definition of liver steatosis,⁸ liver tissue TG was quantified and used to
7
8
9 compare the degree of steatosis between the FHA and MCDD models (Table 1). Compared with
10
11
12 the Normal liver, both the FHA and MCDD protocols created considerable fat accumulation in
13
14
15 the liver. Among them, the FHA and 4-week feeding of MCDD (MCDD4wk) showed the
16
17
18 nearest TG accumulation ($P = 0.64$). The two groups also showed almost equivalent serum
19
20
21 AST, ALT and albumin level ($P = 0.28, 0.24$ and 0.49 respectively, Table 2).

22
23
24 H & E and ORO staining of the FHA group showed mixed (micro- and macro-vesicular)
25
26
27 steatosis exceeding 60%; on the other hand, the MCDD4wk group showed pure macro-vesicular
28
29
30 steatosis exceeding 80% (Fig. 1). The NAFLD activity scores of the two liver steatosis were
31
32
33 equivalent ($P = 0.99$) and did not reach the category of steatohepatitis (Table 2).¹⁷ Accordingly,
34
35
36 we compared the FHA and MCDD4wk groups in the following experiments.

3.2 Experiment II-A: Comparison of survival rates under 6-hour cold storage

37
38
39
40
41
42 Fig. 2 showed the survival rate of the FHA and MCDD4wk group under 6-hour cold storage.
43
44
45
46 While all five rats in the MCDD4wk group survived 7 days after transplantation, only one
47
48
49 survived 1 day and none survived more than 2 days after transplantation in the FHA group ($P =$
50
51
52
53
54
55
56
57
58
59
60
0.002). With this 6-hour cold storage model, we could not obtain the samples of 24-hour after

1
2
3
4
5
6 reperfusion in the FHA group; therefore, we conducted the experiment II-B using 1-hour cold
7
8 storage. The 24-hour survival rate of the FHA group with 1-hour cold storage improved to 80%.
9
10
11
12
13
14

15 **3.3 Experiment II-B: Evaluation of liver grafts under 1-hour cold storage**

16 **3.3.1 Results of serological study, liver tissue ATP and MDA (Table 3)**

17
18
19
20
21 The FHA group showed higher AST level at RP-3 and RP-24 than the Normal liver (both $P <$
22
23 0.001) and MCDD4wk groups ($P = 0.001$ and < 0.001 , respectively). The FHA group also
24
25 showed higher ALT levels at RP-3 and RP-24 than the Normal liver ($P = 0.001$ and < 0.001 ,
26
27 respectively) and MCDD4wk group ($P = 0.09$ and < 0.001 , respectively).
28
29
30
31
32

33 Liver tissue ATP level was lower in the MCDD4wk group before procurement (vs the Normal
34
35 liver and FHA groups, both $P = 0.02$, respectively); however, after reperfusion, while the
36
37 MCDD4wk group showed a gradual increase in the ATP level, the FHA group showed a
38
39 significant decrease in ATP (vs. the Normal liver and MCDD4wk groups, at RP-3, $P < 0.001$
40
41 and 0.17 and at RP-24, $P = 0.005$ and 0.04, respectively). MDA level was higher in the
42
43 MCDD4wk group before procurement (vs. the Normal liver and FHA groups, $P = 0.15$ and
44
45 0.03), at RP-3 (both $P < 0.001$) and RP-24 ($P = 0.11$ and 0.04, respectively).
46
47
48
49
50
51
52
53
54
55
56

57 **3.3.2 Intrahepatic microcirculation (Table 4)**

1
2
3
4
5
6
7 At the left lateral segment, the FHA group showed the lowest microcirculation before
8
9 procurement (vs. the Normal liver and MCDD4wk groups, $P < 0.001$ and 0.01), at RP-3 ($P =$
10
11 0.04 and 0.02) and RP-24 ($P < 0.001$ and 0.004 , respectively). Before procurement, the
12
13 MCDD4wk group also showed lower micro circulation than the Normal liver group ($P = 0.02$).
14
15
16
17
18 The same trend was also observed at the left and right sides of the middle lobe.
19
20
21
22
23

24 **3.3.3 Histological analysis of liver ischemia reperfusion injury**

25
26
27 Unlike the Normal liver and MCDD4wk groups, the FHA group showed a patchy necrotic area
28
29 in zones 1 and 2 (Fig. 3A). The FHA group showed a higher IRI score in zones 1 (vs. the
30
31 Normal liver and MCDD4wk groups, both $P < 0.001$, Fig. 3B) and zone 2 (both $P < 0.001$,
32
33 respectively). In zone 3, the score was also higher in the FHA group than in the Normal liver
34
35 group ($P = 0.04$). The MCDD4wk group showed a higher score than the Normal liver group in
36
37 zone 1 ($P = 0.04$).
38
39
40
41
42
43
44
45
46
47
48

49 **3.3.4 Electron microscopy**

50
51 Unlike the Normal liver and MCDD4wk group, the FHA group showed necrotic cell death at
52
53 RP-24 (Fig. 4). Hepatocyte mitochondria of the FHA group showed marked swelling and an
54
55 obscured silhouette at RP-24 (Fig. 4). Liver sinusoidal endothelial cells (LSECs) were also
56
57
58
59
60

1
2
3
4
5
6 evaluated. Before procurement (Fig. 5), while rats in the Normal liver groups showed an intact
7
8 sinusoidal structure, the FHA group showed abnormal LSEC swelling, widening of the space of
9
10 Disse and sparse hepatocyte microvilli. The MCDD4wk group also showed a subendothelial
11
12 basal lamina and collagen deposition in the space of Disse. Sinusoidal space was quantified
13
14 using the aforementioned method. The FHA and MCDD4wk group showed a decrease in
15
16 sinusoidal space compared with the Normal liver group ($P < 0.001$ and 0.008 , Supplementary
17
18 Figure 1), even before procurement. The FHA group also showed a trend toward a smaller
19
20 sinusoidal space than the MCDD4wk group ($P = 0.30$). At RP-24 (Fig. 6), LSEC injury, that is,
21
22 swelling or detachment of the LSEC, was apparent in the FHA group. SEM also showed an
23
24 irregular surface and diminished fenestrae and sieve plate structure in the FHA group. The
25
26 MCDD4wk groups also showed slightly disrupted fenestrae.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

3.3.5 Real-Time Polymerase Chain Reaction (Table 5)

43
44
45 The mRNA expression of IL-1 β and IL-6 showed a trend toward a higher level in the FHA
46
47 group at RP-24 compared with the Normal liver ($P = 0.12$ and 0.17) and MCDD4wk group ($P =$
48
49 0.16 and 0.20 , respectively). TNF- α expression at RP-24 was also higher in the FHA group than
50
51 in the Normal liver and MCDD4wk groups ($P = 0.002$ and 0.008 , respectively). Before
52
53 procurement, the MCDD4wk group showed higher ET-1 expression than the Normal liver and
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FHA groups ($P = 0.01$ and 0.003 , respectively); however, after reperfusion, the value became higher in the FHA group at RP-3 (vs. the Normal liver and MCDD4wk groups, $P = 0.01$ and 0.07) and RP-24 ($P = 0.003$ and 0.009 , respectively). The mRNA expression of eNOS was almost equivalent among the three groups at all time points. The MCDD4wk group showed a higher iNOS expression at RP-3 (vs. the Normal liver and FHA groups, $P = 0.006$ and 0.06 , respectively). In contrast, the FHA group showed a lower expression of HO-1 (vs. the Normal liver and MCDD4wk groups, $P = 0.03$ and 0.30) and Nqo1 ($P = 0.10$ and 0.04 , respectively) at RP-3. The MCDD4wk group also showed higher Nqo1 expression before procurement (vs. the Normal liver and FHA groups, both $P < 0.001$, respectively).

4 Discussion

The results of this study provide us fundamental knowledge about transplantation using steatotic liver grafts. The FHA and MCDD4wk groups had similar liver tissue TG contents; however, the two liver steatosis models possessed quite unique basal characteristics and showed completely different response against IRI and survival after transplantation.

The FHA protocol was first introduced by Delzenne et al. and increased synthesis of free fatty acid after refeeding is thought to be the main mechanism of fatty liver formation.⁹ This model mimic a liver steatosis acutely formed by short-term fasting during resuscitation and subsequent

1
2
3
4
5
6 high caloric refeeding; e.g. the situation sometimes seen in traumatic brain-dead donors.²¹ On
7
8
9 the other hand, methionine and choline are essential to create very-low-density-lipoprotein in rat
10
11
12 liver; therefore, if these elements are lacking, the rat liver cannot create the lipoprotein, which
13
14
15 leads to the accumulation of fat droplets in the hepatocytes.^{22, 23} Although some discrepancies in
16
17
18 the pathophysiological process were suggested,²⁴ this liver steatosis has long been regarded as a
19
20
21 model for NAFLD.^{25, 26} As shown in Table 1, the TG level of the MCDD4wk group was almost
22
23
24 comparable with that of the FHA group. With H & E or ORO staining, the MCDD4wk group
25
26
27 had an equal or even more prominent fat accumulation than the FHA group (Fig. 1). The type of
28
29
30 steatosis was also disadvantageous for the MCDD4wk group because in clinical situations, it is
31
32
33 a macro- (not micro-) vesicular steatosis that increases the risk of PNF.²⁷ However, it was the
34
35
36 FHA group that showed far worse graft function and survival after transplantation.

37
38
39 As described by several authors,²⁸⁻³¹ impaired microcirculation is an important factor
40
41
42 affecting IRI and worse graft outcome after liver transplantation. Our results showed that rats in
43
44
45 the FHA group had lower microcirculation than those in the Normal liver and MCDD4wk
46
47
48 groups before and after reperfusion. We must discuss the two points separately. First, the
49
50
51 microcirculation of rats in the FHA group was already compromised before procurement. In the
52
53
54 steatotic liver, fat droplets in the hepatocytes increase the cell volume, which leads to the
55
56
57 obstruction of the hepatic sinusoidal space and impaired microcirculation.²⁸ This is consistent
58
59
60

1
2
3
4
5
6 with our result that the FHA and MCDD4wk groups had a smaller sinusoidal space than the
7
8
9 Normal liver group before procurement. Although, a subendothelial basal lamina and collagen
10
11
12 deposition in space of Disse, which are the signs of microvascular change in liver steatosis,³²
13
14
15 were apparent in the MCDD4wk group, the FHA group also showed a trend toward a smaller
16
17
18 sinusoidal space and more severe LSEC swelling. Previous study demonstrated that LSEC can
19
20
21 responsively act like sphincters by swelling or contracting and narrow the sinusoidal lumen.³³
22
23
24 Therefore, the changes in LSEC seen in the FHA group might also contribute to lower
25
26
27 microcirculation than in the MCDD4wk group.
28
29

30
31 Second, the disturbance of microcirculation became more prominent in the FHA group after
32
33
34 reperfusion. This is also consistent with the results of TEM and SEM at RP-24 demonstrating
35
36
37 severe LSEC injury and capillarization in the FHA group. Additionally, tissue ATP was
38
39
40 considerably lower after reperfusion in the FHA group. The lack of ATP leads to a dysfunction
41
42
43 of the sodium/potassium ATP-dependent plasma membrane pump and results in further
44
45
46 swelling of hepatocytes or LSEC.³⁴ The expression of ET-1, which is an initiator of IRI and also
47
48
49 a strong vasoconstrictor induced by microcirculatory disturbance, was higher in the FHA group
50
51
52 after reperfusion and this also could lead to further impaired microcirculation.^{35, 36} Our results
53
54
55 not only reinforce the previous theory that impaired microcirculation and IRI aggravate each
56
57
58
59
60

1
2
3
4
5
6 other,^{28,29} but also add a new insight that this vicious cycle might be more strongly emphasized
7
8
9 in some types of liver steatosis.

10
11
12 Liver tissue MDA levels, the marker of lipid peroxidation, were higher in the MCDD4wk
13
14
15 group at all time points; in contrast, despite the same degree of fat accumulation, the FHA group
16
17
18 showed MDA levels almost comparable with the Normal liver group. This seems natural
19
20
21 because previous studies showed that the oxidative stress is strongly associated with the
22
23
24 pathophysiology of NAFLD³⁷ and the MCDD model has been regarded as a mimic of the
25
26
27 disease. Besides iNOS, both HO-1 and Nqo1 are essential antioxidant enzymes expressed
28
29
30 against various kinds of cellular stresses.^{38,39} Therefore, higher perioperative iNOS, HO-1 and
31
32
33 Nqo1 expressions in the MCDD4wk group should be regarded as a proper countermeasure
34
35
36 against higher oxidative stress. In contrast, the result of tissue MDA showed that oxidative
37
38
39 stress does not seem to weigh heavily on the FHA group; however, HO-1 and Nqo1 expression
40
41
42 were lower in the FHA group than in the Normal liver group; this might suggest the presence of
43
44
45 some cellular disturbances in the FHA group.

46
47
48 Currently, several experimental strategies have been proposed for increased utilization of
49
50
51 extended-criteria donors; e.g. statins,⁴⁰ venous systemic oxygen persufflation⁴¹ or ex situ
52
53
54 machine perfusion.⁴² However, the results of our study suggest that a uniform strategy might not
55
56
57 fully address the complex pathophysiology of steatotic liver graft. Severely impaired
58
59
60

1
2
3
4
5
6 microcirculation and acutely depleted ATP levels are thought to be the main cause of higher IRI
7
8
9 in the FHA group and oxidative stress seems to have only a low impact on this process.

10
11
12 Abnormally swollen hepatocyte mitochondria at RP-24 might be the sign of permeabilization
13
14
15 and the precursor to subsequent ATP depletion and necrotic cell death in the FHA group.⁴³
16
17

18 Therefore, protecting LSEC and hepatocyte mitochondria and maintaining microcirculation and
19
20
21 tissue ATP levels would be an effective strategy for this type of liver steatosis. On the other
22
23
24 hand, considering the relatively well-maintained postoperative microcirculation and tissue ATP,
25
26
27 higher oxidative stress might be the main cause of IRI in the MCDD4wk group; therefore,
28
29
30 ameliorating oxidative stress and underpinning antioxidant capacity might be an adequate
31
32
33 strategy for the MCDD type steatotic liver graft.
34
35

36 This study has several limitations. Among them, translatability to clinical practice is the most
37
38
39 important and difficult question to address. Despite the efforts of several researchers, there
40
41
42 remain some ambiguities over the correspondence of liver steatosis between rats and humans.⁴⁴
43
44

45 The study using discarded grafts in liver transplantation would be a bridge connecting the gap;
46
47
48 however, this type of experiment is now ethically prohibited in Japan. Therefore, our study
49
50
51 focused on demonstrating the difference between two types of liver steatosis being offered as a
52
53
54 graft for liver transplantation. In addition, although our results showed that the survival of the
55
56
57 FHA grafts dramatically worsened after 6-hour cold storage, it is still unclear whether the same
58
59
60

1
2
3
4
5
6 mechanism discussed in 1-hour cold storage model could be applied in 6-hour cold storage. The
7
8
9 FHA and MCDD groups supposedly are affected differently by cold storage. This difference
10
11
12 would be crucial for an understanding of steatotic liver graft transplantation and will be our next
13
14
15 research target.
16

17
18 In conclusion, the results of this study showed that the FHA and MCDD4wk groups, even
19
20
21 though they have almost equivalent degree of steatosis, had quite different graft function and
22
23
24 survival after transplantation. It should be remembered that the degree of fat accumulation is not
25
26
27 a single determinant for the fate of steatotic liver graft.
28
29
30
31
32

33 **Acknowledgement**

34
35
36 The authors thank Ms. Keiko Okamoto-Furuta and Mr. Haruyasu Kohda in the Division of
37
38
39 Electron Microscopy, Center for Anatomical Studies, Kyoto University, and Dr Masato
40
41
42 Nagahama in Department of Pathology, Handa City Hospital for the evaluation of the
43
44
45 specimens of electron microscopy.
46
47
48
49
50

51 **Disclosure**

52
53
54 All authors declared no conflict of interest.
55
56
57
58
59
60

References

1. Chu MJ, Dare AJ, Phillips AR, Bartlett AS. Donor Hepatic Steatosis and Outcome After Liver Transplantation: a Systematic Review. *J Gastrointest Surg* 2015;19:1713-1724.
2. Imber CJ, St Peter SD, Lopez I, Guiver L, Friend PJ. Current practice regarding the use of fatty livers: a trans-Atlantic survey. *Liver Transpl* 2002;8:545-549.
3. Doyle MB, Vachharajani N, Wellen JR, Anderson CD, Lowell JA, Shenoy S et al. Short- and long-term outcomes after steatotic liver transplantation. *Archives of surgery (Chicago, Ill : 1960)* 2010;145:653-660.
4. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A et al. Health Effects of Overweight and Obesity in 195 Countries over 25 Years. *N Engl J Med* 2017;377:13-27.
5. Mavrogiannaki AN, Migdalis IN. Nonalcoholic Fatty liver disease, diabetes mellitus and cardiovascular disease: newer data. *Int J Endocrinol* 2013;2013:450639.
6. Wong TC, Fung JY, Chok KS, Cheung TT, Chan AC, Sharr WW et al. Excellent outcomes of liver transplantation using severely steatotic grafts from brain-dead donors. *Liver Transpl* 2016;22:226-236.
7. McCormack L, Petrowsky H, Jochum W, Mullhaupt B, Weber M, Clavien PA. Use

1
2
3
4
5
6 of severely steatotic grafts in liver transplantation: a matched case-control study. *Ann*
7
8
9 *Surg* 2007;246:940-946; discussion 946-948.

10
11
12 8. Bedogni G, Nobili V, Tiribelli C. Epidemiology of fatty liver: an update. *World J*
13
14
15 *Gastroenterol* 2014;20:9050-9054.

16
17
18 9. Delzenne NM, Hernaux NA, Taper HS. A new model of acute liver steatosis
19
20
21 induced in rats by fasting followed by refeeding a high carbohydrate-fat free diet.
22
23
24 *Biochemical and morphological analysis. J Hepatol* 1997;26:880-885.

25
26
27 10. Rogers AE, Newberne PM. Animal model: fatty liver and cirrhosis in lipotrope-
28
29
30 deficient male rats. *The American journal of pathology* 1973;73:817-820.

31
32
33 11. He S, Rehman H, Wright GL, Zhong Z. Inhibition of inducible nitric oxide
34
35
36 synthase prevents mitochondrial damage and improves survival of steatotic partial liver
37
38
39 grafts. *Transplantation* 2010;89:291-298.

40
41
42 12. Anderson CD, Upadhya G, Conzen KD, Jia J, Brunt EM, Tiriveedhi V et al.
43
44
45 Endoplasmic reticulum stress is a mediator of posttransplant injury in severely steatotic
46
47
48 liver allografts. *Liver Transpl* 2011;17:189-200.

49
50
51 13. Okamura Y, Hata K, Tanaka H, Hirao H, Kubota T, Inamoto O et al. Impact of
52
53
54 Subnormothermic Machine Perfusion Preservation in Severely Steatotic Rat Livers: A
55
56
57 Detailed Assessment in an Isolated Setting. *Am J Transplant* 2017;17:1204-1215.
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
14. Kamada N, Calne RY. Orthotopic liver transplantation in the rat. Technique using cuff for portal vein anastomosis and biliary drainage. *Transplantation* 1979;28:47-50.
 15. Liu T, Freise CE, Ferrell L, Ascher NL, Roberts JP. A modified vascular "sleeve" anastomosis for rearterialization in orthotopic liver transplantation in rats. *Transplantation* 1992;54:179-180.
 16. Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipides from animal tissues. *The Journal of biological chemistry* 1957;226:497-509.
 17. Kleiner DE, Brunt EM, Van Natta M, Behling C, Contos MJ, Cummings OW et al. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology* 2005;41:1313-1321.
 18. Yagi S, Doorschodt BM, Afify M, Klinge U, Kobayashi E, Uemoto S et al. Improved preservation and microcirculation with POLYSOL after partial liver transplantation in rats. *J Surg Res* 2011;167:e375-383.
 19. Wasserstein RL, Lazar NA. The ASA Statement on p-Values: Context, Process, and Purpose. *The American Statistician* 2016;70:129-133.
 20. Wasserstein RL, Schirm AL, Lazar NA. Moving to a World Beyond "p < 0.05".

1
2
3
4
5
6 The American Statistician 2019;73:1-19.
7

8
9 21. Hata K, Tolba RH, Wei L, Doorschodt BM, Buttner R, Yamamoto Y et al. Impact
10 of polysol, a newly developed preservation solution, on cold storage of steatotic rat
11 livers. Liver Transpl 2007;13:114-121.
12
13
14
15

16
17
18 22. Yao ZM, Vance DE. The active synthesis of phosphatidylcholine is required for
19 very low density lipoprotein secretion from rat hepatocytes. The Journal of biological
20 chemistry 1988;263:2998-3004.
21
22
23
24
25

26
27 23. Yao ZM, Vance DE. Head group specificity in the requirement of
28 phosphatidylcholine biosynthesis for very low density lipoprotein secretion from cultured
29 hepatocytes. The Journal of biological chemistry 1989;264:11373-11380.
30
31
32
33
34

35
36 24. Rinella ME, Green RM. The methionine-choline deficient dietary model of
37 steatohepatitis does not exhibit insulin resistance. J Hepatol 2004;40:47-51.
38
39
40

41
42 25. Oz HS, Chen TS, Neuman M. Methionine deficiency and hepatic injury in a
43 dietary steatohepatitis model. Dig Dis Sci 2008;53:767-776.
44
45
46

47
48 26. Ma C, Kesarwala AH, Eggert T, Medina-Echeverez J, Kleiner DE, Jin P et al.
49 NAFLD causes selective CD4(+) T lymphocyte loss and promotes hepatocarcinogenesis.
50 Nature 2016;531:253-257.
51
52
53
54

55
56
57 27. Selzner N, Selzner M, Jochum W, Amann-Vesti B, Graf R, Clavien PA. Mouse
58
59
60

livers with macrosteatosis are more susceptible to normothermic ischemic injury than those with microsteatosis. *J Hepatol* 2006;44:694-701.

28. Ijaz S, Yang W, Winslet MC, Seifalian AM. Impairment of hepatic microcirculation in fatty liver. *Microcirculation* 2003;10:447-456.

29. Cheng Q, Ng KT, Xu A, Li CX, Liu XB, Guo DY et al. The roles of lipocalin-2 in small-for-size fatty liver graft injury. *Ann Surg* 2014;260:1062-1072.

30. El-Badry AM, Moritz W, Contaldo C, Tian Y, Graf R, Clavien PA. Prevention of reperfusion injury and microcirculatory failure in macrosteatotic mouse liver by omega-3 fatty acids. *Hepatology* 2007;45:855-863.

31. Kuroda S, Tashiro H, Igarashi Y, Tanimoto Y, Nambu J, Oshita A et al. Rho inhibitor prevents ischemia-reperfusion injury in rat steatotic liver. *J Hepatol* 2012;56:146-152.

32. McCuskey RS, Ito Y, Robertson GR, McCuskey MK, Perry M, Farrell GC. Hepatic microvascular dysfunction during evolution of dietary steatohepatitis in mice. *Hepatology* 2004;40:386-393.

33. Sorensen KK, Simon-Santamaria J, McCuskey RS, Smedsrod B. Liver Sinusoidal Endothelial Cells. *Compr Physiol* 2015;5:1751-1774.

34. Abu-Amara M, Yang SY, Tapuria N, Fuller B, Davidson B, Seifalian A. Liver

1
2
3
4
5
6 ischemia/reperfusion injury: processes in inflammatory networks--a review. *Liver*
7
8
9 *Transpl* 2010;16:1016-1032.

10
11
12 35. Pulitano C, Joseph D, Sandroussi C, Verran D, Ho P, Debiasio A et al.
13
14
15 Postreperfusion microcirculatory derangements after liver transplantation: Relationship
16
17
18 to hemodynamics, serum mediators, and outcome. *Liver Transpl* 2017;23:527-536.

19
20
21 36. Liu B, Qian JM. Cytoprotective role of heme oxygenase-1 in liver ischemia
22
23
24 reperfusion injury. *International journal of clinical and experimental medicine*
25
26
27 2015;8:19867-19873.

28
29
30 37. Seki S, Kitada T, Yamada T, Sakaguchi H, Nakatani K, Wakasa K. In situ
31
32
33 detection of lipid peroxidation and oxidative DNA damage in non-alcoholic fatty liver
34
35
36 diseases. *J Hepatol* 2002;37:56-62.

37
38
39 38. Amersi F, Buelow R, Kato H, Ke B, Coito AJ, Shen XD et al. Upregulation of
40
41
42 heme oxygenase-1 protects genetically fat Zucker rat livers from ischemia/reperfusion
43
44
45 injury. *J Clin Invest* 1999;104:1631-1639.

46
47
48 39. Ross D, Siegel D. Functions of NQO1 in Cellular Protection and CoQ10
49
50
51 Metabolism and its Potential Role as a Redox Sensitive Molecular Switch. *Front Physiol*
52
53
54 2017;8:595.

55
56
57 40. Gracia-Sancho J, Garcia-Caldero H, Hide D, Marrone G, Guixe-Muntet S, Peralta
58
59
60

1
2
3
4
5
6 C et al. Simvastatin maintains function and viability of steatotic rat livers procured for
7
8
9 transplantation. *J Hepatol* 2013;58:1140-1146.
10

11
12 41. Yagi S, Nagai K, Kadaba P, Afify M, Teramukai S, Uemoto S et al. A novel organ
13
14 preservation for small partial liver transplantations in rats: venous systemic oxygen
15
16 persuf lation with nitric oxide gas. *Am J Transplant* 2013;13:222-228.
17
18

19
20
21 42. Boteon YL, Boteon A, Attard J, Mergental H, Mirza DF, Bhogal RH et al. Ex situ
22
23 machine perfusion as a tool to recondition steatotic donor livers: Troublesome features
24
25 of fatty livers and the role of defatting therapies. A systematic review. *Am J Transplant*
26
27 2018;18:2384-2399.
28
29

30
31
32 43. Malhi H, Gores GJ, Lemasters JJ. Apoptosis and necrosis in the liver: a tale of
33
34 two deaths? *Hepatology* 2006;43:S31-44.
35
36

37
38
39 44. Sanches SC, Ramalho LN, Augusto MJ, da Silva DM, Ramalho FS. Nonalcoholic
40
41 Steatohepatitis: A Search for Factual Animal Models. *Biomed Res Int*
42
43 2015;2015:574832.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure legends

Fig. 1 Histology of the Normal liver, FHA and MCDD4wk group.

Above; H & E stain, below; ORO stain, $\times 400$ magnification. The FHA group showed mixed (micro- and macro-vesicular) steatosis more than 60%. The MCDD4wk group showed pure macro-vesicular steatosis more than 80%. Both type of steatotic liver grafts showed no signs of steatohepatitis. FHA, fasting and hyper alimentation; H & E, hematoxylin-eosin stain; MCDD, methionine and choline deficient diet; ORO, oil-red-O stain.

Fig. 2 The survival rate of the FHA and MCDD4wk group under 6-hour cold storage.

While all five rats of the MCDD4wk group survived 7-days after transplantation, only one survived 1-day and none survived more than 2 days after transplantation in the FHA group ($n = 5$ each, log-rank test, $P = 0.002$). FHA, fasting and hyper alimentation; MCDD, methionine and choline deficient diet.

Fig. 3 Histology of liver graft at RP-24.

A; H & E stain. Compared with the Normal liver and MCDD4wk group, the FHA group showed patchy necrotic area (dotted line) through zone 1 to zone 2. B: Result of histological score. The FHA group had significantly higher injury score at zones 1 and zone 2 than the

1
2
3
4
5
6 Normal liver and MCDD4wk group. At zone 3, in contrast, the scores were not so different
7
8
9 between the FHA and MCDD4wk group ($P = 0.36$). ** $P < 0.001$, * $P < 0.05$. FHA, fasting and
10
11 hyper alimentation; H & E, hematoxylin-eosin stain; MCDD, methionine and choline deficient
12
13 diet.
14
15
16
17
18
19

20
21 Fig. 4 Evaluation of hepatocyte injury and mitochondria using TEM at RP-24.

22
23
24 Upper row: While hepatocytes of the Normal liver and MCDD4wk groups seemed unaffected,
25
26 the FHA group showed necrotic cell death characterized by rupture of the cell membrane and
27
28 release of intracellular organelles. TEM: magnification $\times 1000$
29
30
31

32
33 Lower row: Hepatocyte mitochondria. While the Normal liver and MCDD4wk group showed
34
35 almost intact hepatocyte mitochondria, the FHA group showed marked swelling and obscured
36
37 silhouette of mitochondria.
38
39
40

41
42 FHA, fasting and hyper alimentation; MCDD, methionine and choline deficient diet. TEM,
43
44 Transmission electron microscopy.
45
46
47
48
49

50
51 Fig. 5 Evaluation of LSEC using TEM; before procurement.

52
53
54 While the Normal liver group showed an intact sinusoidal structure, the FHA group showed
55
56 marked swelling and contraction of LSEC and widening of the space of Disse. Hepatocyte
57
58
59
60

1
2
3
4
5
6 microvilli were also sparse in the FHA group. The MCDD4wk group showed a subendothelial
7
8
9 basal lamina (white arrow) and collagen deposition (black arrow) in space of Disse. TEM:
10
11
12 magnification $\times 2500$. E: endothelial cell; D: Space of Disse.

13
14
15 FHA, fasting and hyper alimentation; LSEC, liver sinusoidal endothelial cell; MCDD,
16
17
18 methionine and choline deficient diet; TEM, transmission electron microscopy.
19
20

21
22
23
24 Fig. 6 Evaluation of LSEC using TEM and SEM; at RP-24.

25
26
27 Upper row: While the Normal liver and MCDD4wk groups maintained an almost smooth LSEC
28
29
30 lining, the FHA group showed disruption and detachment of LSEC. TEM: $\times 2500$.

31
32
33 Lower row: While the Normal liver group showed an intact sinusoidal structure, the FHA
34
35
36 group showed loss of fenestrae and sieve plate structures. The MCDD4wk group maintained
37
38
39 fenestrae in sieve plate fairly well; however, these structures were disrupted compared with the
40
41
42 Normal liver group. Scales are indicated on the figure. SEM: $\times 3500$.

43
44
45 FHA, fasting and hyper alimentation; LSEC, liver sinusoidal endothelial cell; MCDD,
46
47
48 methionine and choline deficient diet; SEM, scanning electron microscopy; TEM, transmission
49
50
51 electron microscopy.

52
53
54
55
56
57 Supplementary Figure 1 Quantification of sinusoidal space.
58
59
60

1
2
3
4
5
6 The area of sinusoidal space before procurement was identified in 10 randomly selected TEM
7
8
9 images ($\times 700$) in each group and calculated using Image J version 1.46r (National Institutes of
10
11
12 Health, USA). The Normal liver group showed a larger sinusoidal space than the FHA and
13
14
15 MCDD4wk groups ($P < 0.001$ and 0.008). The FHA group also showed a trend toward a
16
17
18 smaller sinusoidal space than the MCDD4wk group ($P = 0.30$).
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1, Tissue TG contents in each liver steatosis models.

	Normal liver	FHA	MCDD2wk	MCDD4wk	MCDD6wk
	n = 3	n = 5	n = 3	n = 5	n = 3
Tissue triglyceride (mg/g tissue)	2.6 ± 0.0	116.7 ± 10.1	76.3 ± 7.1	137.2 ± 18.0*	252.8 ± 4.9* **
<i>P</i> value (vs. FHA)		reference	0.23	0.64	< 0.001

The data are shown as mean ± standard error.

* *P* < 0.05 vs. MCDD2wk. ** *P* < 0.05 vs. MCDD4wk

FHA, fasting and hyperalimantation; TG, triglyceride; MCDD methionine and choline deficient diet.

Table 2, Comparison of the FHA and MCDD4wk group.

Factors	Normal liver	FHA	MCDD4wk	<i>P</i>
	n = 3	n = 5	n = 5	(FHA vs. MCDD4wk)
Serum study				
AST (IU/L)	82 ± 3	90 ± 5	102 ± 8	0.28
ALT (IU/L)	40 ± 1	60 ± 8	89 ± 22	0.24
Albumin (g/dL)	3.7 ± 0.0	3.7 ± 0.0	3.7 ± 0.1	0.49

NAFLD activity score	0 ± 0	2.8 ± 0.2	2.8 ± 0.2	0.99
----------------------	-------	-----------	-----------	------

The data are shown as mean ± standard error. Student's t-test was used to directly compare the FHA and MCDD4wk group.

ALT, alanine aminotransferase; AST, aspartate aminotransferase; FHA, fasting and hyperalimantation; MCDD4wk, 4-week feeding of methionine and choline deficient diet. NAFLD, non-alcoholic fatty liver disease.

Table 3; Summary of serum studies, ATP and MDA analysis in Experiment IIB.

		Before procurement	RP-3	RP-24
		n = 5	n = 5	n = 5
AST	Normal liver	75 ± 4	488 ± 88*	505 ± 59*
(IU/L)	FHA	90 ± 5	2221 ± 352 ^{ab*}	2610 ± 256 ^{ab*}
	MCDD4wk	102 ± 8 ^a	686 ± 135*	601 ± 68*
ALT	Normal liver	40 ± 1	281 ± 61*	258 ± 53*
(IU/L)	FHA	60 ± 8	1753 ± 263 ^{a*}	1709 ± 140 ^{ab*}
	MCDD4wk	89 ± 22	1035 ± 272*	567 ± 58
ATP	Normal liver	1.8 ± 0.3	2.0 ± 0.2	1.9 ± 0.3
(µmol/g)	FHA	1.9 ± 0.1 ^b	0.6 ± 0.1 ^{ab*}	0.7 ± 0.2 ^{ab*}

tissue)	MCDD4wk	0.8 ± 0.2^a	1.1 ± 0.2	1.6 ± 0.2
MDA	Normal liver	2.5 ± 0.7	1.2 ± 0.2	2.6 ± 0.6
($\mu\text{mol/g}$	FHA	1.9 ± 0.3^b	1.9 ± 0.3^b	2.0 ± 0.5^b
tissue)	MCDD4wk	3.9 ± 0.3	7.5 ± 1.5^a	5.3 ± 1.2

The data are shown as mean \pm standard error.

^a $P < 0.05$, vs. the Normal liver group. ^b $P < 0.05$ vs. the MCDD4wk group. * $P < 0.05$ vs.

before procurement, ** $P < 0.05$ vs RP3. ALT, alanine aminotransferase; AST, aspartate

aminotransferase; ATP, adenosine triphosphate; FHA, fasting and hyperalimantation;

MCDD4wk, 4-week feeding of methionine and choline deficient diet. MDA, malondialdehyde;

RP-3 and RP-24, 3- and 24-hour after reperfusion.

Table 4, Summary of intrahepatic microcirculation in Experiment IIB.

		Before	RP-1	RP-3
		procurement		
		n = 10	n = 10	n = 5
Left lateral segment	Normal liver	15.6 ± 0.5	$7.1 \pm 0.9^*$	$11.6 \pm 0.3^* **$
(ml/min/100g tissue)	FHA	8.5 ± 0.6^{ab}	$4.4 \pm 0.4^{ab*}$	$4.2 \pm 0.8^{ab*}$
	MCDD4wk	12.2 ± 1.2^a	$7.3 \pm 0.8^*$	8.9 ± 1.1

Left side of middle lobe	Normal liver	16.4 ± 1.1	9.8 ± 1.0*	11.0 ± 1.4*
(ml/min/100g tissue)	FHA	12.1 ± 0.7 ^a	6.0 ± 0.5 ^{b*}	5.4 ± 0.7 ^{ab*}
	MCDD4wk	15.4 ± 1.3	11.1 ± 1.5	11.9 ± 0.6
Right side of middle lobe	Normal liver	15.9 ± 1.3	8.3 ± 1.0*	13.5 ± 1.8**
(ml/min/100g tissue)	FHA	11.3 ± 0.7	5.9 ± 0.5 ^{b*}	5.1 ± 0.7 ^{ab*}
	MCDD4wk	15.9 ± 2.1	9.6 ± 1.0*	11.6 ± 2.2

The data are shown as mean ± standard error.

^a $P < 0.05$, vs. the Normal liver group. ^b $P < 0.05$ vs. the MCDD4wk group. * $P < 0.05$ vs.

before procurement, ** $P < 0.05$ vs RP3. FHA, fasting and hyperalimentation; MCDD4wk, 4-week feeding of methionine and choline deficient diet. RP-1 and RP-3, 1- and 3-hour after reperfusion.

Table 5, Summary of mRNA expression in Experiment IIB.

		Before procurement	RP-3	RP-24
		n = 5	n = 5	n = 5
IL-1 β	Normal liver	1.5 ± 0.2	4.0 ± 0.9*	1.8 ± 0.4**
(/ β -actin)	FHA	1.5 ± 0.2	3.5 ± 0.5	7.8 ± 3.7
	MCDD4wk	1.8 ± 0.3	4.4 ± 0.4*	1.2 ± 0.1**

IL-6	Normal liver	2.2 ± 0.5	$102.5 \pm 23.1^*$	$18.0 \pm 6.7^{**}$
(/ β -actin)	FHA	15.9 ± 13.3	203.9 ± 54.5	278.8 ± 165.6
	MCDD4wk	11.0 ± 6.7	673.7 ± 323.2	29.6 ± 12.6
TNF- α	Normal liver	0.8 ± 0.1	$2.1 \pm 0.4^*$	$0.7 \pm 0.1^{**}$
(/ β -actin)	FHA	0.6 ± 0.1	$2.6 \pm 0.5^*$	$2.8 \pm 0.5^{ab*}$
	MCDD4wk	1.1 ± 0.2	$3.5 \pm 0.7^*$	$1.1 \pm 0.1^{**}$
ET-1	Normal liver	1.0 ± 0.1	$3.6 \pm 0.8^*$	1.7 ± 0.4
(/ β -actin)	FHA	0.8 ± 0.1^b	$10.4 \pm 2.0^{a*}$	$5.8 \pm 1.1^{ab*}$
	MCDD4wk	1.9 ± 0.3^a	$5.6 \pm 0.9^*$	$2.2 \pm 0.4^{**}$
eNOS	Normal liver	0.9 ± 0.0	$0.6 \pm 0.1^*$	$0.6 \pm 0.1^*$
(/ β -actin)	FHA	0.7 ± 0.1	0.7 ± 0.1	0.5 ± 0.1
	MCDD4wk	0.8 ± 0.1	0.5 ± 0.1	$0.4 \pm 0.1^*$
iNOS	Normal liver	0.7 ± 0.2	$24.4 \pm 6.0^*$	$6.3 \pm 3.1^{**}$
(/ β -actin)	FHA	0.5 ± 0.2	42.2 ± 7.8	430.0 ± 386.9
	MCDD4wk	1.4 ± 0.3	$78.7 \pm 14.2^{a*}$	$7.7 \pm 1.8^{**}$
HO-1	Normal liver	1.5 ± 0.4	$36.3 \pm 3.5^*$	$4.4 \pm 0.6^{**}$
(/ β -actin)	FHA	2.5 ± 0.3	$16.1 \pm 5.6^{a*}$	3.8 ± 0.8
	MCDD4wk	2.0 ± 0.4	$26.5 \pm 4.9^*$	$1.7 \pm 0.1^{a**}$

Nqo1	Normal liver	1.2 ± 0.2	1.4 ± 0.3	3.2 ± 0.4* **
(/β-actin)	FHA	0.8 ± 0.1 ^b	0.7 ± 0.1 ^b	0.7 ± 0.2 ^a
	MCDD4wk	2.5 ± 0.2 ^a	1.6 ± 0.2	1.1 ± 0.2 ^{a* **}

The data are shown as mean ± standard error.

^a $P < 0.05$, vs. the Normal liver group. ^b $P < 0.05$ vs. the MCDD4wk group. * $P < 0.05$ vs.

before procurement, ** $P < 0.05$ vs RP3. eNOS, endothelial nitric oxide synthase; ET-1,

endothelin-1; FHA, fasting and hyper alimentation; HO-1, heme oxygenase-1; IL interleukin;

iNOS, inducible nitric oxide synthase; MCDD4wk, 4-week feeding of methionine and choline

deficient diet; Nqo1, NAD(P)H Quinone Dehydrogenase 1; RP-3 and RP-24, 3- and 24-hour

after reperfusion; TNF, Tumor necrosis factor.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 1




Fig. 1 Histology of the Normal liver, FHA and MCDD4wk group. Above; H & E stain, below; ORO stain, ×400 magnification. The FHA group showed mixed (micro- and macro-vesicular) steatosis more than 60%. The MCDD4wk group showed pure macro-vesicular steatosis more than 80%. Both type of steatotic liver grafts showed no signs of steatohepatitis. FHA, fasting and hyper alimentation; H & E, hematoxylin-eosin stain; MCDD, methionine and choline deficient diet; ORO, oil-red-O stain.

209x297mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 2 The survival rate of the FHA and MCDD4wk group under 6-hour cold storage. While all five rats of the MCDD4wk group survived 7-days after transplantation, only one survived 1-day and none survived more than 2 days after transplantation in the FHA group (n = 5 each, log-rank test, P = 0.002). FHA, fasting and hyper alimentation; MCDD, methionine and choline deficient diet.

209x297mm (300 x 300 DPI)




Fig. 3 Histology of liver graft at RP-24.

A; H & E stain. Compared with the Normal liver and MCDD4wk group, the FHA group showed patchy necrotic area (dotted line) through zone 1 to zone 2. B: Result of histological score. The FHA group had significantly higher injury score at zones 1 and zone 2 than the Normal liver and MCDD4wk group. At zone 3, in contrast, the scores were not so different between the FHA and MCDD4wk group ($P = 0.36$). ** $P < 0.001$, * $P < 0.05$. FHA, fasting and hyper alimentation; H & E, hematoxylin-eosin stain; MCDD, methionine and choline deficient diet.

209x297mm (300 x 300 DPI)

Fig. 4




Fig. 4 Evaluation of hepatocyte injury and mitochondria using TEM at RP-24. Upper row: While hepatocytes of the Normal liver and MCDD4wk groups seemed unaffected, the FHA group showed necrotic cell death characterized by rupture of the cell membrane and release of intracellular organelles. TEM: magnification $\times 1000$. Lower row: Hepatocyte mitochondria. While the Normal liver and MCDD4wk group showed almost intact hepatocyte mitochondria, the FHA group showed marked swelling and obscured silhouette of mitochondria. FHA, fasting and hyper alimentation; MCDD, methionine and choline deficient diet. TEM, Transmission electron microscopy.

209x297mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 5




Fig. 5 Evaluation of LSEC using TEM; before procurement. While the Normal liver group showed an intact sinusoidal structure, the FHA group showed marked swelling and contraction of LSEC and widening of the space of Disse. Hepatocyte microvilli were also sparse in the FHA group. The MCDD4wk group showed a subendothelial basal lamina (white arrow) and collagen deposition (black arrow) in space of Disse. TEM: magnification $\times 2500$. E: endothelial cell; D: Space of Disse. FHA, fasting and hyperalimentation; LSEC, liver sinusoidal endothelial cell; MCDD, methionine and choline deficient diet; TEM, transmission electron microscopy.

209x297mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 6





Fig. 6 Evaluation of LSEC using TEM and SEM; at RP-24.
 Upper row: While the Normal liver and MCDD4wk groups maintained an almost smooth LSEC lining, the FHA group showed disruption and detachment of LSEC. TEM: $\times 2500$.
 Lower row: While the Normal liver group showed an intact sinusoidal structure, the FHA group showed loss of fenestrae and sieve plate structures. The MCDD4wk group maintained fenestrae in sieve plate fairly well; however, these structures were disrupted compared with the Normal liver group. Scales are indicated on the figure. SEM: $\times 3500$.
 FHA, fasting and hyper alimentation; LSEC, liver sinusoidal endothelial cell; MCDD, methionine and choline deficient diet; SEM, scanning electron microscopy; TEM, transmission electron microscopy.

209x297mm (300 x 300 DPI)

Supplementary Figure 1



Supplementary Figure 1 Quantification of sinusoidal space.

The area of sinusoidal space before procurement was identified in 10 randomly selected TEM images (×700) in each group and calculated using Image J version 1.46r (National Institutes of Health, USA). The Normal liver group showed a larger sinusoidal space than the FHA and MCDD4wk groups ($P < 0.001$ and 0.008). The FHA group also showed a trend toward a smaller sinusoidal space than the MCDD4wk group ($P = 0.30$).

209x297mm (300 x 300 DPI)