TITLE:

Protective Effects of a Hydrogen-Rich
Preservation Solution in a Canine Lung
Transplantation Model(Dissertation_全文)

AUTHOR(S):

Kayawake, Hidenao

CITATION:

Kayawake, Hidenao. Protective Effects of a Hydrogen-Rich Preservation Solution in a Canine Lung Transplantation Model. 京都大学, 2021, 博士(医学)

ISSUE DATE:

2021-03-23

URL:

<https://doi.org/10.14989/doctor.k23107>

RIGHT:

<https://doi.org/10.1016/j.athoracsur.2020.05.076>

Protective Effects of a Hydrogen-Rich Preservation Solution in a Canine Lung Transplantation Model

Hidenao Kayawake, MD, Toyofumi F. Chen-Yoshikawa, MD, PhD, Masao Saito, MD, PhD, Hiroya Yamagishi, MD, Akihiko Yoshizawa, MD, PhD, Shin-ich Hirano, DVM, PhD, Ryosuke Kurokawa, and Hiroshi Date, MD, PhD

Department of Thoracic Surgery, Kyoto University Graduate School of Medicine, Kyoto, Japan; Department of Thoracic Surgery, Nagoya University Graduate School of Medicine, Nagoya, Japan; Department of Thoracic Surgery, Shimada Municipal Hospital, Shimada, Japan; Department of Diagnostic Pathology, Kyoto University Hospital, Kyoto, Japan; and MiZ Company Ltd, Kamakura, Japan

Background. Molecular hydrogen (H_2) has protective effects against ischemia-reperfusion injury in various organs. Because they are easier to transport and safer to use than inhaled H_2 , H_2 -rich solutions are suitable for organ preservation. In this study, we examined the protective effects of an H_2 -rich solution for lung preservation in a canine left lung transplantation (LTx) model.

Methods. Ten beagles underwent orthotopic left LTx after 23 hours of cold ischemia followed by reperfusion for 4 hours. Forty-five minutes after reperfusion, the right main pulmonary artery was clamped to evaluate the function of the implanted graft. The beagles were divided into two groups: control group ($n = 5$), and H_2 group ($n = 5$). In the control group, the donor lungs were flushed and immersed during cold preservation at $4^\circ C$ using ET-Kyoto solution, and in the H_2 group, these were flushed and immersed using H_2 -rich ET-Kyoto solution. Physiologic assessments were performed during

reperfusion. After reperfusion, the wet-to-dry ratios were determined, and histology examinations were performed.

Results. Significantly higher partial pressure of arterial oxygen and significantly lower partial pressure of carbon dioxide were observed in the H_2 group than in the control group ($P = .045$ and $P < .001$, respectively). The wet-to-dry ratio was significantly lower in the H_2 group than in the control group ($P = .032$). Moreover, in histology examination, less lung injury and fewer apoptotic cells were observed in the H_2 group ($P < .001$ and $P < .001$, respectively).

Conclusions. Our results demonstrated that the H_2 -rich preservation solution attenuated ischemia-reperfusion injury in a canine left LTx model.

(Ann Thorac Surg 2021;111:246-52)

© 2021 by The Society of Thoracic Surgeons

Lung transplantation (LTx) is the final option to save the lives of patients with end-stage pulmonary diseases.^{1,2} However, severe donor shortage remains a serious problem. Another important issue is that lungs are vulnerable to ischemia, resulting in the limitation of the ischemic time. In deceased-donor LTx, the graft lungs are exposed to ischemia during transportation. A long ischemic time often leads to ischemia-reperfusion (I/R) injury after LTx.^{3,4} Therefore, developing better methods of lung preservation during the ischemic time is required to improve the treatment outcome after LTx.

Since Ohsawa and associates⁵ first reported the beneficial effects of molecular hydrogen (H_2) for brain I/R

injury in a mouse brain infarction model in 2007, H_2 has been reported to have protective effects against I/R injury in various organs.⁶⁻⁹ Although the precise mechanism remains uncertain, the antioxidative, antiinflammatory, and antiapoptosis effects of H_2 have been reported.¹⁰⁻¹² Our group has previously reported the protective effects of an H_2 -rich solution against lung I/R injury in a rat left hilar clamp model¹³ and those of an H_2 -rich lung preservation solution in a rat orthotopic left LTx model.¹⁴ To examine the diffusion of H_2 in lungs of larger animals and evaluate the protective effects of H_2 -rich preservation solution in larger animals having much more complicated immune systems than rats, we performed this preclinical study using a canine orthotopic left LTx model.

Material and Methods

Animals

Ten pairs of weight-matched Toyo beagles (Kitayama Labes, Hongo Farm, Yamaguchi, Japan) weighing 8.95 to 11.1 kg were used in this experiment. All animals

Accepted for publication May 11, 2020.

Presented at the Thirty-ninth Annual Meeting and Scientific Sessions of the International Society for Heart and Lung Transplantation, Orlando, FL, Apr 3-6, 2019.

Address correspondence to Dr Date, Department of Thoracic Surgery, Kyoto University, 54 Shogoin-Kawahara-cho, Sakyo-ku, Kyoto 606-8507, Japan; email: hdate@kuhp.kyoto-u.ac.jp.

Abbreviations and Acronyms

cDNA	= complementary DNA
H ₂	= molecular hydrogen
IL	= interleukin
I/R	= ischemia-reperfusion
LTx	= lung transplantation
mRNA	= messenger RNA
PCR	= polymerase chain reaction
ssDNA	= single-strand DNA
TNF	= tumor necrosis factor

received humane care in accordance with the *Guide for the Care and Use of Laboratory Animals* prepared by the National Institutes of Health (publication no. 86-23, revised 1996). The study protocol of this experiment was approved by the Ethics Committee of the Graduate School of Medicine at Kyoto University, Japan (MedKyo 19276).

Hydrogen-rich Preservation Solution

In this study, the ET-Kyoto solution (Otsuka Pharmaceutical, Tokushima, Japan), which is clinically used for LTx at Kyoto University Hospital, was used as the preservation solution.¹⁵ The H₂-rich preservation solution was prepared using a commercially available hydrogen-generating agent (Aquila 7.0; MiZ Company, Kamakura, Japan) consisting of metal aluminum grains and calcium hydroxide.^{14,16} The concentration of hydrogen was quantified with a titrant consisting of methylene blue and platinum.¹⁷

Experimental Design

Figure 1 shows the experimental design of this study. After the retrieval of lung grafts from a donor dog, cold preservation at 4°C for 23 hours was performed. Thereafter, the left lung graft was prepared and orthotopic left LTx for a recipient dog was performed. We set 60 minutes as the transplantation time. After the orthotopic left LTx, the graft lungs were reperfused for 4 hours. Forty-five minutes after transplantation, the right main pulmonary artery was clamped to evaluate the function of the implanted graft.

Ten donor dogs were randomly divided into two groups: H₂ group (n = 5) and control group (n = 5). In the control group, normal ET-Kyoto solution was used both for flushing and immersing the graft lungs, whereas in the H₂ group, H₂-rich ET-Kyoto solution was used for both flushing and immersing.

Donor Procurement

The operative technique has been precisely described elsewhere.¹⁸⁻²⁰ Briefly, the beagles were premedicated with an intramuscular injection of midazolam (0.5 mg/kg), xylazine (2.0 mg/kg), and atropine sulfate (0.05 mg/kg). They were intubated and mechanically ventilated with fraction of inspired oxygen of 0.5, tidal volume of 20 mL/kg, frequency of 15 breaths per minute, and positive

end-expiratory pressure of 5 cm H₂O, and maintained on 0.8% to 2.0% sevoflurane and administered an intravenous injection of rocuronium bromide (0.5 mg/kg). After median sternotomy, the main pulmonary artery was cannulated through the right ventricular outflow tract and the superior and inferior vena cavae were ligated. After the incision of the left atrial appendage, 1000 mL cold normal ET-Kyoto solution (control group) or cold H₂-rich ET-Kyoto solution (H₂ group) was perfused antegradely and retrogradely from a height of 30 cm above the heart. Ventilation was continued until the retrieval of lung grafts and recovered heart-lung block, which were semi-inflated with 50% oxygen, immersed in 2500 mL normal (control group) or H₂-rich (H₂ group) ET-Kyoto solution, and preserved at 4°C for 23 hours in an aluminum bag (Lamizip AL-34L; Seisannipponsha, Tokyo, Japan).

Recipient Preparation and Orthotopic Left Lung Transplantation

Recipient dogs were anesthetized and maintained in the same manner as the donors except for fraction of inspired oxygen of 1.0. A Swan-Ganz catheter and an arterial catheter were inserted before the operation. Before the orthotopic left LTx, the left lung graft was separated from the recovered heart-lung block and flushed again antegradely and retrogradely with 500 mL normal (control group) or H₂-rich (H₂ group) ET-Kyoto solution. After left pneumonectomy, orthotopic left LTx was performed as described elsewhere.¹⁸⁻²¹ After reperfusion, a 2.0-mm catheter was directly inserted into the left atrium to monitor the left atrial pressure. Thereafter, the right main pulmonary artery was clamped 45 minutes after reperfusion to evaluate the function of the implanted graft, and the implanted lung grafts were reperfused for 4 hours in total.

Lung Function After Transplantation

Systemic arterial pressure, pulmonary arterial pressure, and peak airway pressure were continuously monitored throughout the experiment. Arterial blood gas analysis was performed before the recipient operation and at 30, 60, 120, 180, and 240 minutes after the implantation using an automatic blood gas analyzer (ABL 80 FLEX; Radiometer, Tokyo, Japan). At the same time, cardiac output was also measured. Pulmonary dynamic compliance was defined as tidal volume divided by ([peak airway pressure] minus [positive end-expiratory pressure]).^{18,21}

Evaluation of Lung Edema

At the end of the experiment, partial resection of the left lower lobe was performed to calculate the wet-to-dry lung weight ratio for evaluating lung edema. To calculate the wet-to-dry weight ratio, wet samples were first weighed to obtain the wet lung weight; after heating the samples at 100°C for 48 hours, they were weighed again to obtain the dry lung weight.

Figure 1. Design of the experiment. After retrieval of lung grafts, cold preservation for 23 hours was performed. After orthotopic left lung transplantation (LTx), graft lungs were reperfused for 4 hours. Forty-five minutes after transplantation, the right main pulmonary artery (PA) was clamped for evaluating function of implanted graft.

Macroscopic and Histologic Findings

Macroscopic appearance of the implanted lung grafts at 4 hours after reperfusion was recorded. The left middle lobe was resected at the end of reperfusion and was fixed in 10% buffered formalin, embedded in paraffin, and stained with hematoxylin and eosin. To evaluate the degree of lung injury, the lung injury score was calculated based on the lung injury scoring system defined in the official American Thoracic Society workshop report.^{22,23} Two investigators (H.K. and H.Y.) blindly evaluated lung injury at 20 randomly chosen high-power fields per section at a magnification of $\times 400$ under the supervision of a pulmonary pathologist (A.Y.).

The evaluation of apoptotic cells was performed by immunostaining for single-strand DNA (ssDNA) using rabbit polyclonal antibodies against ssDNA (IBL, Takasaki, Japan).²⁴ The number of ssDNA-positive cells was quantified using the average number in 10 randomly chosen fields per section at a magnification of $\times 200$.

Quantitative Real-Time Polymerase Chain Reaction

The left upper lobe was partially resected at the end of reperfusion for quantitative real-time polymerase chain reaction (PCR) to evaluate the gene expression of interleukin-1 β messenger RNA (mRNA), tumor necrosis factor α (TNF- α) mRNA, and interleukin-6 (IL-6) mRNA. The lung tissue was stabilized with RNAlater (Qiagen, Hilden, Germany), and RNA was extracted using an RNeasy Plus Mini Kit (Qiagen) following the manufacturer's protocol. Thereafter, the total RNA was reversely transcribed to complementary DNA (cDNA) with the Ready-To-Go You-Prime First-Strand Beads (GE Healthcare, Pittsburgh, PA), and each cDNA sample was diluted to 10 μ g/mL. Furthermore, 2 μ L cDNA was mixed with the Thunderbird probe qPCR Mix (Toyobo, Osaka, Japan) and TaqMan Gene Expression Assay probe set (Thermo Fisher Scientific, Waltham, MA). The PCR was performed using the StepOnePlus Real-Time PCR system (Thermo Fisher Scientific). The relative quantity of each sample was calculated using the comparative $\Delta\Delta$ CT method. Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as the reference gene. Each sample was

analyzed in triplicate. Probes used for PCR assays were as follows: GAPDH (NM_001003142.2), IL-1 β mRNA (NM_001037971.1), TNF- α mRNA (NM_001003244.4), and IL-6 mRNA (NM_001003301.1).

Statistical Analysis

All statistical analyses were performed using the EZR software (Saitama Medical Center, Jichi Medical University, Saitama, Japan), a graphical user interface for R (The R Foundation for Statistical Computing, Vienna, Austria).^{25,26} All values are expressed as mean $\pm$ SD. The Mann-Whitney U test was performed for the comparison of two groups. Repeated measures analysis of variance was performed for the comparison of PaO₂ and PaCO₂, pulmonary vascular resistance, and pulmonary dynamic compliance. Statistical significance was defined as *P* less than .05.

Results

There were no significant differences between the control and H₂ groups in donor weight (control group, 9.9 ± 0.4 kg; H₂ group, 9.9 ± 0.7 kg) or recipient weight (control group, 9.6 ± 0.8 kg; H₂ group, 9.7 ± 0.5 kg). The H₂ concentration of the H₂-rich ET-Kyoto solution was greater than 1.0 ppm before being used for immersing and flushing. The H₂ concentration of the H₂-rich ET-Kyoto solution used for immersing was maintained at more than 0.6 ppm for 23 hours after retrieval.

Lung Function and Pulmonary Hemodynamics After Transplantation

All five recipients in both groups survived until 4 hours after reperfusion. Significantly higher PaO₂ was observed in the H₂ group than in the control group (*P* = .045; Figure 2A). Furthermore, PaCO₂ was significantly lower in the H₂ group than in the control group (*P* < .001; Figure 2B). There was no significant difference between the two groups in pulmonary vascular resistance (*P* = .412; Figure 2C); however, pulmonary dynamic compliance in the H₂ group was significantly better than that in the control group (*P* = .029; Figure 2D).

Wet-to-Dry Lung Weight Ratio

The wet-to-dry lung weight ratio at 240 minutes after reperfusion was significantly lower in the H₂ group than in the control group (H₂ group, 9.2 ± 1.4 ; control group, 11.4 ± 1.6 ; *P* = .032; Figure 3A).

Macroscopic and Histologic Findings at 4 Hours After Transplantation

The graft lungs of the H₂ group appeared less damaged than those of the control group (Figure 3B). With respect to histologic findings, lesser neutrophil infiltration was observed in the H₂ group than in the control group (Figure 4A). Consequently, the lung injury score in the H₂ group was significantly lower than that in the control group (H₂ group, 0.30 ± 0.13 ; control group, 0.42 ± 0.14 ; *P* < .001; Figure 4B). Furthermore, significantly fewer

Figure 2. Physiologic data of the experiment: hydrogen-rich (H_2) group (blue line [$n = 5$]); and control (CON) group (red line [$n = 5$]). (A) Significantly higher PaO_2 was observed in the H_2 group than in the control group ($P = .045$). (B) The PaCO_2 was significantly lower in the H_2 group than in the control group ($P < .001$). (C) There was no significant difference in pulmonary vascular resistance between the two groups ($P = .412$). (D) Significantly better pulmonary dynamic compliance was observed in the H_2 group than in the control group ($P = .029$). (LTx, lung transplantation; PA, pulmonary artery.)

ssDNA-positive cells were observed in the H_2 group than in the control group (H_2 group, 2.3 ± 1.6 ; control group, 6.6 ± 4.6 ; $P < .001$; Figure 5).

Expression of mRNA of Proinflammatory Cytokines

The expression of IL-1 β mRNA in the H_2 group was lower than that in the control group, although the difference did not reach statistical significance ($P = .056$; Figure 6A). The expression of TNF- α mRNA and IL-6 mRNA was comparable between the two groups ($P = .222$ and $P = .222$, respectively; Figures 6B, 6C).

Comment

In this study, an H_2 -rich solution was used for lung preservation during cold ischemia in a canine left orthotopic LTx model. By using this preservation solution, both more effective ventilation and better pulmonary compliance during 4 hours of reperfusion were achieved than with the normal preservation solution. Moreover, lung edema after LTx was attenuated with the use of the H_2 -rich preservation solution, and lung injury was also attenuated macroscopically and histologically. Furthermore, this preservation solution decreased the number of

Figure 3. Wet-to-dry lung weight ratio and macroscopic findings: hydrogen-rich (H_2) group (blue [$n = 5$]); and control (CON) group (red [$n = 5$]). (A) Wet-to-dry weight ratio was measured to evaluate degree of lung edema. Wet-to-dry weight ratio in H_2 group was significantly lower than in control group ($P = .032$). (B) Macroscopic appearance of implanted lung grafts at 4 hours after reperfusion. These two figures were representative cases. Graft lungs of the H_2 group appeared less damaged than those of the control group.

Figure 4. Histologic evaluation using the lung injury score based on the official American Thoracic Society workshop report. (A) Representative histologic findings of the two groups (original magnification $\times 400$). Less neutrophil infiltration was observed in the hydrogen-rich (H_2) group than in the control (CON) group. (B) The lung injury score in the H_2 group (blue [$n = 5$]) was significantly lower than that in the control group (red [$n = 5$]; $P < .001$).

apoptotic cells and suppressed the production of proinflammatory cytokines in lung tissue. These results indicated the protective effects of the H_2 -rich preservation solution against lung I/R injury after prolonged cold ischemia in a canine orthotopic left LTx model.

Since the report by Ohsawa and colleagues,⁵ H_2 has been used as a therapeutic antioxidant for I/R injury attenuation and organ preservation in various medical fields.^{6-11,27-32} Our group previously reported the protective effects of an H_2 -rich preservation solution during cold ischemia in a rat LTx model. As a further research for clinical application, we used larger animals in this study.

In the present study, corresponding to better physiologic function of the implanted grafts, attenuation of lung edema after LTx was observed when an H_2 -rich preservation solution was used for lung preservation during cold ischemia. In previous reports, when an H_2 -rich preservation solution was used, the regular lining of vascular endothelial cells was well preserved based on scanning electron microscope findings in a rat LTx

model¹⁴ and a rat ex vivo liver perfusion model.²⁷ Furthermore, the protective effects of an H_2 -rich medium on vascular endothelial permeability by the inhibition of signal pathways and downregulated expression of the adhesion molecules causing injury of the vascular endothelium in vitro were also observed in other studies.^{33,34} Based on these reports and our results, we speculated that the effects of an H_2 -rich preservation solution on the attenuation of lung edema in this study may be partly attributed to the reduction of vascular endothelial cell injury and vascular endothelial permeability.

This study also showed that posttransplantation lung injury and production of proinflammatory cytokines were attenuated by the use of an H_2 -rich preservation solution. Previous reports suggested that H_2 inhibits the activation of nuclear factor-kappa B and p38 mitogen-activated protein kinase pathway, thereby reducing the expression of proinflammatory cytokines and infiltration of neutrophils.^{12,28,29,35} Furthermore, this study showed the

Figure 5. Evaluation of apoptotic cells using immunostaining for single strand DNA (ssDNA). (A) Representative histologic findings of the two groups (original magnification $\times 200$). Arrows show ssDNA-positive cells. (B) Significantly fewer ssDNA-positive cells were observed in the hydrogen-rich (H_2) group (blue [$n = 5$]) than in the control (CON) group (red [$n = 5$]); $P < .001$).

Figure 6. Quantitative real-time polymerase chain reaction results of proinflammatory cytokines in grafts at end of reperfusion. (A) Gene expression of interleukin-1 β (IL-1 β) in the hydrogen-rich (H₂) group (blue [n = 5]) was lower than that in the control (CON) group (red [n = 5]), although the difference did not reach statistical significance (P = .056). (B) Gene expression of tumor necrosis factor- α (TNF- α) was comparable between the two groups (P = .222). (C) Gene expression of interleukin-6 was also comparable between the two groups (P = .222). (mRNA, messenger RNA.)

antiapoptosis effects of the H₂-rich preservation solution by revealing that fewer ssDNA-positive cells were observed in the H₂ group. Apoptosis is considered a factor for early graft failure in solid organ transplantation. It was previously reported that the antiapoptosis effects of H₂ are partly derived from the suppression of proapoptosis factors such as caspase-3 and caspase-8 and upregulation of antiapoptosis factors such as B-cell lymphoma-2.^{8,28,30,35,36} However, details of the mechanism remain unknown. As we could not reveal the details in this experiment, further studies using in vitro and LTx experiments in a mouse model are required.

Some of the methods used to deliver H₂ to organs include inhalation of H₂ and dissolution of H₂ into the preservation solution. In this study, we dissolved H₂ into the organ preservation solution because we considered that the use of an H₂-rich solution has some advantages over inhalation. First, an H₂-rich solution is easy to transport. Therefore, it can be used both for flushing during donor operation and for immersing the grafts before they are transported. Second, an H₂-rich solution is safer to use than H₂ gas, as more than 4% H₂ gas has flammable and explosive properties. In this study, the H₂-rich preservation solution obviously attenuated lung I/R injury in the LTx model after prolonged cold ischemia even with a low concentration of H₂, which indicated that the H₂-rich solution was efficient in delivering H₂ to

organ tissues. As a future perspective, we plan to apply lung preservation with an H₂-rich preservation solution to clinical LTx.

This study has several limitations. First, the number of subjects in both groups was relatively small (n = 5) because more animal sacrifices should be avoided from the viewpoint of animal welfare. Second, we did not use a sterile method for preparing the H₂-rich preservation solution. Therefore, techniques for producing a sterile and stable H₂-rich solution are required for clinical application.

In conclusion, we demonstrated that an H₂-rich preservation solution attenuated lung I/R injury after prolonged cold ischemia in a canine LTx model. The H₂-rich preservation solutions could be a useful and effective option for cold preservation and transportation of lung grafts.

The authors wish to thank the remaining members of their department for discussions about this experiment; the staff members of the Institute of Laboratory Animals, Graduate School of Medicine, Kyoto University, for their humane animal care; and the staff members of the Center for Anatomical, Pathological and Forensic Medical Researches, Graduate School of Medicine, Kyoto University, for their technical support. This work was supported by the Japan Society for the Promotion of Science KAKENHI (17H04295) to H.D. Financial support and the hydrogen generator were provided by MiZ Co, Ltd.

References

- Chambers DC, Cherikh WS, Goldfarb SB, et al. The International Thoracic Organ Transplant Registry of the International Society for Heart and Lung Transplantation: thirty-fifth adult lung and heart-lung transplant report-2018; focus theme: multiorgan transplantation. *J Heart Lung Transplant*. 2018;37:1169-1183.
- Kayawake H, Chen-Yoshikawa TF, Aoyama A, et al. Surgical management of bronchial stumps in lobar lung transplantation. *J Thorac Cardiovasc Surg*. 2018;156:451-460.
- Gelman AE, Fisher AJ, Huang HJ, et al. Report of the ISHLT working group on primary lung graft dysfunction part III: mechanisms: a 2016 consensus group statement of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. 2017;36:1114-1120.
- Chen F, Date H. Update on ischemia-reperfusion injury in lung transplantation. *Curr Opin Organ Transplant*. 2015;20:515-520.
- Ohsawa I, Ishikawa M, Takahashi K, et al. Hydrogen acts as a therapeutic antioxidant by selectively reducing cytotoxic oxygen radicals. *Nat Med*. 2007;13:688-694.
- Hayashida K, Sano M, Ohsawa I, et al. Inhalation of hydrogen gas reduces infarct size in the rat model of myocardial ischemia-reperfusion injury. *Biochem Biophys Res Commun*. 2008;373:30-35.
- Fukuda K, Asoh S, Ishikawa M, et al. Inhalation of hydrogen gas suppresses hepatic injury caused by ischemia/reperfusion through reducing oxidative stress. *Biochem Biophys Res Commun*. 2007;361:670-674.
- Kawamura T, Huang CS, Peng X, et al. The effect of donor treatment with hydrogen on lung allograft function in rats. *Surgery*. 2011;150:240-249.
- Haam S, Lee JG, Paik HC, et al. Hydrogen gas inhalation during ex vivo lung perfusion of donor lungs recovered after cardiac death. *J Heart Lung Transplant*. 2018;37:1271-1278.
- Chen M, Zhang J, Chen Y, et al. Hydrogen protects lung from hypoxia/re-oxygenation injury by reducing hydroxyl radical production and inhibiting inflammatory responses. *Sci Rep*. 2018;8:8004.
- Abe T, Li XK, Yazawa K, et al. Hydrogen-rich University of Wisconsin solution attenuates renal cold ischemia-reperfusion injury. *Transplantation*. 2012;94:14-21.
- Zhang G, Li Z, Meng C, et al. The anti-inflammatory effect of hydrogen on lung transplantation model of pulmonary microvascular endothelial cells during cold storage period. *Transplantation*. 2018;102:1253-1261.
- Takahashi M, Chen-Yoshikawa TF, Saito M, et al. Immersing lungs in hydrogen-rich saline attenuates lung ischaemia-reperfusion injury. *Eur J Cardiothorac Surg*. 2017;51:442-448.
- Saito M, Chen-Yoshikawa TF, Takahashi M, et al. Protective effects of a hydrogen-rich solution during cold ischemia in rat lung transplantation. *J Thorac Cardiovasc Surg*. 2020;159:2110-2118.
- Ikedo M, Bando T, Yamada T, et al. Clinical application of ET-Kyoto solution for lung transplantation. *Surg Today*. 2015;45:439-443.
- Ozeki N, Yamawaki-Ogata A, Narita Y, et al. Hydrogen water alleviates obliterative airway disease in mice. *Gen Thorac Cardiovasc Surg*. 2020;68:158-163.
- Seo T, Kurokawa R, Sato B. A convenient method for determining the concentration of hydrogen in water: use of methylene blue with colloidal platinum. *Med Gas Res*. 2012;2:1.
- Ohsumi A, Chen F, Sakamoto J, et al. Protective effect of pre-recovery surfactant inhalation on lungs donated after cardiac death in a canine lung transplantation model. *J Heart Lung Transplant*. 2012;31:1136-1142.
- Sakamoto J, Chen F, Nakajima D, et al. The effect of β -2 adrenoreceptor agonist inhalation on lungs donated after cardiac death in a canine lung transplantation model. *J Heart Lung Transplant*. 2012;31:773-779.
- Hijiya K, Chen-Yoshikawa TF, Kondo T, et al. Bronchodilator inhalation during ex vivo lung perfusion improves post-transplant graft function after warm ischemia. *Ann Thorac Surg*. 2017;103:447-453.
- Nakajima D, Chen F, Okita K, et al. Reconditioning lungs donated after cardiac death using short-term hypothermic machine perfusion. *Transplantation*. 2012;94:999-1004.
- Matute-Bello G, Downey G, Moore BB, et al. An official American Thoracic Society workshop report: features and measurements of experimental acute lung injury in animals. *Am J Respir Cell Mol Biol*. 2011;44:725-738.
- Cosgun T, Iskender I, Yamada Y, et al. Ex vivo administration of trimetazidine improves post-transplant lung function in pig model. *Eur J Cardiothorac Surg*. 2017;52:171-177.
- Saito M, Chen-Yoshikawa TF, Suetsugu K, et al. Pirfenidone alleviates lung ischemia-reperfusion injury in a rat model. *J Thorac Cardiovasc Surg*. 2019;158:289-296.
- Kanda Y. Investigation of the freely available easy-to-use software 'EZ' for medical statistics. *Bone Marrow Transplant*. 2013;48:452-458.
- Kayawake H, Chen-Yoshikawa TF, Hamaji M, et al. Acquired recipient pulmonary function is better than lost donor pulmonary function in living-donor lobar lung transplantation. *J Thorac Cardiovasc Surg*. 2019;158:1710-1716.
- Tamaki I, Hata K, Okamura Y, et al. Hydrogen flush after cold storage as a new end-ischemic ex vivo treatment for liver grafts against ischemia/reperfusion injury. *Liver Transpl*. 2018;24:1589-1602.
- Zhou H, Fu Z, Wei Y, et al. Hydrogen inhalation decreases lung graft injury in brain-dead donor rats. *J Heart Lung Transplant*. 2013;32:251-258.
- Noda K, Shigemura N, Tanaka Y, et al. Hydrogen preconditioning during ex vivo lung perfusion improves the quality of lung grafts in rats. *Transplantation*. 2014;98:499-506.
- Shi J, Yao F, Zhong C, et al. Hydrogen saline is protective for acute lung ischaemia/reperfusion injuries in rats. *Heart Lung Circ*. 2012;21:556-563.
- Noda K, Shigemura N, Tanaka Y, et al. A novel method of preserving cardiac grafts using a hydrogen-rich water bath. *J Heart Lung Transplant*. 2013;32:241-250.
- Buchholz BM, Masutani K, Kawamura T, et al. Hydrogen-enriched preservation protects the isogeneic intestinal graft and amends recipient gastric function during transplantation. *Transplantation*. 2011;92:985-992.
- Yu Y, Wang WN, Han HZ, et al. Protective effects of hydrogen-rich medium on lipopolysaccharide-induced monocytic adhesion and vascular endothelial permeability through regulation of vascular endothelial cadherin. *Genet Mol Res*. 2015;14:6202-6212.
- Xie K, Wang W, Chen H, et al. Hydrogen-rich medium attenuated lipopolysaccharide-induced monocyte-endothelial cell adhesion and vascular endothelial permeability via rho-associated coiled-coil protein kinase. *Shock*. 2015;44:58-64.
- Ge L, Yang M, Yang NN, et al. Molecular hydrogen: a preventive and therapeutic medical gas for various diseases. *Oncotarget*. 2017;8:102653-102673.
- Matei N, Camara R, Zhang JH, et al. Emerging mechanisms and novel applications of hydrogen gas therapy. *Med Gas Res*. 2018;8:98-102.