

## Advanced Plasma Energy Research Section

K. Nagasaki, Professor  
S. Kobayashi, Associate Professor

## 1. Introduction

The current subjects of this research section are to study the properties of high temperature plasmas in order to control and improve the plasma energy confinement from the physical viewpoint of nuclear fusion research. The experimental and theoretical investigations for the optimization of the helical-axis heliotron configuration are in progress under the collaboration with other groups of the international/national institutes and also groups of other universities under the auspices of the Collaboration Program of the Lab. Complex Energy Processes, IAE and the Collaborative Research Program of NIFS (National Institute for Fusion Science).

In this report, a remarkable result obtained in the Heliotron J experimental study in FY2020 is reported focusing on (1) O-X mode conversion analysis for the electron Bernstein wave heating and diagnostic, (2) development of interferometer system for the electron density measurement and (3) X-ray spectrum measurement in non-resonant microwave heated plasmas.

## 2. O-X mode conversion analysis using finite element method for EBW heating and diagnostics

Electron cyclotron wave (ECW) is used as one of the methods for heating and diagnostics in magnetic confinement fusion devices. The propagation modes, ordinary (O)-mode and extraordinary (X)-mode, are reflected when the plasma density reaches the cutoff density, resulting that we are not able to access the core region. To solve this problem, O-X-B (from O-mode to X-mode and then electron Bernstein wave (EBW)) mode conversion is proposed. In this study, we have analyzed the optimal conditions for O-X conversion and its conversion efficiency by the finite element method based on the wave equation using COMSOL Multiphysics and compared them with ray-tracing calculations.

We have used a two-dimensional model with a uniform magnetic field and a density gradient in the x-axis direction. we have analyzed the injection condition for maximal O-X conversion efficiency for three parameters, that is, the beam size, the focusing diameter of the injection port, and the angle of incidence, and we have obtained more than 80 % con-version efficiency for the entire injection beam. The conversion

from O-mode to X-mode can be seen near the optimum incident angle in Fig. 1(b). In comparison with ray-tracing, there is no significant difference in the optimal angle, and the trajectories of both were consistent as shown in Fig. 1 (d). This suggests that the ray-tracing calculation is useful for finding the optimal incidence angle. In the finite element analysis, the beam width is seen to expand after conversion to X-mode, and the trajectory of the finite width EC beam can be traced.

## 3. Development of 320 GHz interferometer system in Heliotron J

Measurement of the density profile with high time resolution is required to understand the particle transport in fusion plasma experiments. A new multi-channel heterodyne Michelson interferometer with 320 GHz solid-state sources has been designed and is being constructed in Heliotron J for high-density




Fig. 1. The electric field norm of O-mode injection (42 GHz). The incidence angle is (a) 25 deg, (b) 35.7 deg and (c) 45deg to the magnetic field (1.8 T), and (d) Trajectory of ECW at optimal incidence conditions. COMSOL (color), ray-tracing (black line).

plasma operation.

To reduce the influence of the plasma refraction effect and ensure that the injected beam returns back in the original direction after reflection, an array of retroreflectors was mounted on the facing wall of the vacuum chamber. To make sure the influence on the beam profile with the retroreflector, an optical test experiment has been done to compare the reflection effects between the metal mirror and the retroreflector.

Figure 2 shows the experimental setup in the optical test experiment. The beam profile was measured by moving a detector in the horizontal and vertical directions. Figure 3 shows the measured beam profiles in the metal mirror and the retroreflector. The beam profile in the retroreflector is more peaked than that in the flat metal mirror, and the signal intensity is high even at large scanning angles, indicating that the retroreflector works for beam transmission. The retroreflector was installed on the inner wall of the vacuum chamber with the angle of less than 40 deg.




Fig. 2. The schematic view of the test bench experiment for 320 GHz interferometer system




Fig. 3. (a) Beam profile in the horizontal direction and (b) dependence on the scanning angle obtained by the test experiment.

#### 4. Behavior of high-energy electrons in non-resonant microwave heated plasma in Heliotron J

High-energy electrons have been generated by non-

resonant 2.45 GHz microwaves in a magnetic field configuration of Heliotron J. To study the production mechanism, we measured the X-ray energy spectrum produced by Bremsstrahlung processes of the high energy electrons. Three sets of  $\text{LaBr}_3(\text{Ce})$  scintillator covered with shields for stray radiation and magnetic field) have been installed to obtain the X-ray spectrum emitted in three (clockwise, perpendicular and counter-clockwise) directions to the magnetic field line. Each scintillator signal is measured with a multi-channel analyzer (MCA) through a pre-amplifier and a shaping amplifier.

Even though the shielding effect by the vacuum vessel was not negligible, we confirmed the existence of MeV-class high energy electrons by the scintillator signal when the detector observes the X-ray flux in the perpendicular direction (see Fig. 4). The peak of the X-ray spectrum around 0.3 MeV may be due to the shielding effect by the Stainless-steel vacuum vessel whose thickness of 20 mm. The strong signal below than 0.1 MeV is attributed to the noise produced by the magnetic coil for the plasma confinement. We also found that the microwave power had a threshold to produce the high energy electrons and that the spectrum weakly depends on the microwave power once the high energy electrons were produced. These results indicate that (1) the observed high energy X-ray flux might be produced by the reactions when the high energy electrons hit the vacuum vessel and (2) the feature is consistent to the production mechanism that the high-energy electrons are produced by the stochastic acceleration.




Fig. 4. (a) Schematic view of X-ray measurement system using  $\text{LaBr}_3$  scintillator and (b) X-ray energy spectrum obtained in non-resonant microwave heated plasmas.

## Collaboration Works

長崎百伸, 西南物理研究所 (中華人民共和国), IPP, Greifswald (ドイツ), University of Wisconsin (アメリカ), 反射計を用いた電子密度分布・揺動解析

長崎百伸, IPP, Greifswald (ドイツ), 電子サイクロトロン電流駆動の理論解析

長崎百伸, 岡田浩之, 小林進二, 南貴司, 大島慎介, Univ. Wisconsin (アメリカ), Oak Ridge National Laboratory (アメリカ), Max Plank Institute (ドイツ), Stuttgart Univ (ドイツ), CIEMAT (スペイン), Australian National Univ., (オーストラリア), Kharkov Institute (ウクライナ), Southwest Institute of Physics (中華人民共和国), ヘリカル型装置における SOL/ダイバータプラズマに関する研究

長崎百伸, 大島慎介, 岡田浩之, 南貴司, 小林進二, Stuttgart Univ., CIEMAT, 先進閉じ込め配位

小林進二, 長崎百伸, 大島慎介, CIEMAT (スペイン), Kurchatov Institute (ロシア), ORNL (アメリカ), 低磁気シアヘリカル装置における高速イオン励起 MHD 不安定性に関する研究

小林進二, IPP, Greifswald (ドイツ), Kharkov Institute (ウクライナ), Electron acceleration by non-resonant microwave launch and its application to assist NBI plasma startup

大島慎介, 長崎百伸, 岡田浩之, 南貴司, 小林進二, Wisconsin University (アメリカ), CIEMAT (スペイン), Max-Planck Institute (ドイツ), ヘリカル磁場配位における周辺揺動研究

## Financial Support

### 1. Grant-in-Aid for Scientific Research

長崎百伸, 基盤研究(B), 高密度プラズマ輸送解析に向けた電子バーンスタイン放射計測に関する研究

小林進二, 基盤研究(B), 先進ヘリカル配位のベータ効果が対称性と熱・乱流輸送に与える影響の実験的検証

### 2. Others

長崎百伸, 自然科学研究機構, 2020 年度双方向型共同研究

長崎百伸, (株) 日立製作所, 核融合研究に対する

## 研究助成

長崎百伸, 研究拠点形成事業 (A), 磁場の多様性が拓く超高温プラズマダイナミクスと構造形成の国際研究拠点形成

## Publications

T. Minami, D. Qiu, T. Nishide, M. Miyoshi, Y. Yamana, N. Kenmochi, R. Yasuhara, C. Takahashi, S. Kobayashi, H. Okada, S. Kado, S. Ohshima, T. Mizuuchi, K. Nagasaki, Current status and plan of development of Nd:YAG laser Thomson scattering system in Heliotron J, Journal of Instrumentation, 15, 2, C02011, 2020

S. Yamamoto, K. Nagasaki, K. Nagaoka, J. Varela, 'A. Cappa, E. Ascasibar, F. Castej'on, J.M. Fontdecaba, J.M. García-Regaña, 'A. Gonz'alez-Jerez5, K. Ida, A. Ishizawa, M. Isobe, S. Kado, S. Kobayash, M. Liniers, D. L'opez-Bruna, N. Marushchenko, F. Medina, A. Melnikov, T. Minami, T. Mizuuchi, Y. Nakamura, M. Ochando, K. Ogawa, S. Ohshima, H. Okada, M. Osakabe, M. Sanders, J.L. Velasco, G.M. Weirm, M. Yoshinuma, Effect of ECH/ECCD on energetic-particle-driven MHD modes in helical plasmas, Nuclear Fusion, 60, 6, 066018, 2020

S. Kobayashi, K. Nagasaki, T. Stange, T. Mizuuchi, H. Okada, T. Minami, S. Kado, S. Yamamoto, S. Ohshima, K. Hada, G. Weir, S. Konoshima, Y. Naamura, N. Kenmochi, Y. Otani, X.X. Lu, A. Panith, K. Toki, Y. Suzuki, Study of seed plasma generation for NBI plasma start-up using non-resonant microwave launch in Heliotron J, Plasma Physics and Controlled Fusion, 62, 065009, 2020

Y. Suzuki, S. Purohit, S. Odachi, S. Yamamoto, K. Nagasaki, New tomographic reconstruction technique based on Laplacian eigenfunction, Plasma Science & Technology, 22, 10, 102002, 2020

M. Luo, K. Nagasaki, G. Weir, H. Okada, T. Minami, S. Kado, S. Kobayashi, S. Yamamoto, S. Ohshima, T. Mizuuchi, S. Konoshima, Y. Nakamura, A. Ishizawa, Measurement of Electron Temperature Profile and Fluctuation with ECE Radiometer System in Heliotron J, Plasma and Fusion Research, 15, SI, 2402038, 2020

D. Qiu, T. Minami, T. Nishide, M. Miyoshi, Y. Yamana, S. Kado, S. Ohshima, K. Nagasaki, H. Okada, S. Kobayashi, T. Mizuuchi, S. Konoshima, R. Yasuhara, Design of Compact Multi-Path Thomson Scattering Diagnostic with Signal Delay System in Heliotron J, Plasma and Fusion Research, 15, SI, 2401044, 2020

J. Varela, K. Nagasaki, K. Nagaoka, S. Yamamoto, K.Y. Watanabe, D.A. Spong, L. Garcia, A. Cappa, A. Aze-gami, Modeling of the ECCD injection effect on the Heliotron J and LHD plasma stability, Nuclear Fusion, 60, 11, 112015, 2020

N. Smith, K. Nagasaki, J.H.E. Proll, H. Okada, T. Minami, S. Kado, S. Kobayashi, S. Ohshima, S. Konoshima, T. Mizuuchi, Y. Nakamura, A. Ishizawa, T. Tomita, T. Fukuda, Y. Kondo, Measurement of Radial Correlation Lengths of Electron Density Fluctuations in Heliotron J Using O-Mode Reflectometry, Plasma and Fusion Research, 15, 2054, 2020

P. Adulsiriswad, Y. Todo, S. Yamamoto, S. Kado, S. Kobayashi, S. Ohshima, H. Okada, T. Minami, Y. Nakamura, A. Ishizawa, S. Konoshima, T. Mizuuchi, K. Nagasaki, Magnetohydrodynamic hybrid simulation of Alfvén eigenmodes in Heliotron J, a low shear helical axis stellarator/heliotron, Nuclear Fusion, 60, 9, 096005, 2020

Y. Kondo, T. Fukuda, K. Nagasaki, H. Okada, T. Minami, S. Kado, S. Kobayashi, S. Yamamoto, S. Ohshima, T. Mizuuchi, T. Tomita, Measurement of radial electric field using doppler reflectometer in high-density plasma of heliotron J, Plasma and Fusion Research, 15, 2064, 2020

M. Luo, K. Nagasaki, T. Kobayashi, H. Okada, T. Minami, S. Kado, S. Kobayashi, S. Ohshima, S. Konoshima, T. Mizuuchi, Study of electron thermal transport using ECH modulation in Heliotron J, 4th Asia-Pacific Conference on Plasma Physics, MF-P14, 2020

S.J.G. Voermans, J.H.E. Proll, C. Mora Moreno, M.J. Pueschel, G.M. Weir, A. Ishizawa, S. Kobayashi, K. Nagasaki, Linear microinstabilities in the standard and the high-bumpiness configuration of Heliotron-J, 47th EPS Conference on Plasma Physics, 16212, 2020

長崎百伸, 水内亨, 岡田浩之, 南貴司, 門信一郎, 小林進二, 大島慎介, 岸本泰明, 中村祐司, 石澤明宏, 四竈泰一, 木島滋, 的池遼太, 横山雅之, 本島巖, 小林達哉, 西野信博, 山本聡, Gavin M. Weir, Ling Zang, Heliotron J 実験, Plasma and Fusion Research, 96, 9, 475-518, 2020

## Presentations

的池遼太, 河村学思, 大島慎介, 鈴木康浩, 小林政弘, 増崎貴, 小林進二, 門信一郎, 南貴司, 岡田浩之, 水内亨, 木島滋, Y. Feng, H. Frerichs, 長崎百伸, Heliotron J における三次元磁場構造を背景とし

た周辺プラズマ輸送特性, 日本物理学会 2020 年秋季大会, オンライン開催, 2020.9.8-11

松浦寛人, ブイスアンニャットソン, 村岡英樹, 的池遼太, 大島慎介, ヘリオトロンJ複合ダイバータープローブアレイの熱電対信号解析, 日本物理学会 2020 年秋季大会, オンライン開催, 2020.9.8-11

小林進二, 永岡賢一, 長崎百伸, 徳原圭一, 大垣英明, 紀井俊輝, 全炳俊, 岡田浩之, 村田駿介, 大島慎介, 門信一郎, 南貴司, 木島滋, 水内亨, ヘリオトロン型磁場配位における非共鳴マイクロ波を利用した確率的静電加速, 日本物理学会 2020 年秋季大会, オンライン開催, 2020.9.8-11

A. Sanpei, H. Himura, Y. Ishida, T. Inoue, T. Inoue, S. Masamune, K. Nagasaki, H. Okada, S. Kado, Analysis of transition from axisymmetric torus to helical axis toroidal plasma, 第 11 回エネルギー理工学研究所国際シンポジウム, Online, 2020.9.15-16

T. Shikama, Y. Ueno, T. Chatani, S. Kado, H. Kawazome, H. Okada, L. Matsuoka, T. Minami, S. Kobayashi, S. Ohshima, A. Ishizawa, Y. Nakamura, S. Konoshima, T. Mizuuchi, K. Nagasaki, M. Hasuo, Spatially resolved measurement of atomic emission line spectra using NIR Zeeman spectroscopy, 第 11 回エネルギー理工学研究所国際シンポジウム, Online, 2020.9.15-16

H. Kawazome, S. Kado, R. Matoike, S. Ohshima, Development of HeI image reconstruction technique using neural network in Heliotron J, 第 11 回エネルギー理工学研究所国際シンポジウム, Online, 2020.9.15-16

S. Inagaki, K. Nagasaki, G.M. Weir, T. Minami, S. Kado, S. Kobayashi, S. Yamamoto, S. Ohshima, H. Okada, T. Mizuuchi, M. Sasaki, Time-series data analysis of Heliotron-J plasma by statistical modeling, 第 11 回エネルギー理工学研究所国際シンポジウム, Online, 2020.9.15-16

M. Yoshikawa, S. Kobayashi, Y. Nakashima, J. Kohagura, T. Iijima, Y. Ariizumi, Y. Takeda, H. Okada, S. Ohshima, S. Murata, K. Tokuhara, M. Luo, R. Fukushima, Transport analysis for neutral beam heated plasmas in advanced heliotron configuration, 第 11 回エネルギー理工学研究所国際シンポジウム, Online, 2020.9.15-16

S. Ohshima, H. Okada, L. Zang, S. Kobayashi, S. Yamamoto, T. Minami, S. Kado, Y. Nakamura, S. Konoshima, T. Mizuuchi, N. Kenmochi, Y. Ohtani, M. Motoshima, P. Adlsriiswad, R. Matoike, D. Qiu, M. Luo, K. Nagasaki, Isotope Effects on Nonlinear Turbulence System in a Torus Plasma, 4th Asia-Pacific Conference on Plasma Physics, Online, 2020.10.26-31

D. Qiu, T. Minami, Y. Yamanaka, M. Miyoshi, R. Shinotsuka, M. Luo, S. Kado, N. Kenmochi, S. Ohshima, S. Kobayashi, H. Okada, S. Konoshima, T. Mizuuchi, K. Nagasaki, Development of Double Pockels Cells Multi-path Thomson Scattering System in Heliotron J based on Gauss Beam Analysis, 4th Asia-Pacific Conference on Plasma Physics, Online, 2020.10.26-31

P. Adulsiriswad, Y. Todo, S. Kado, S. Yamamoto, S. Kobayashi, S. Ohshima, H. Okada, T. Minami, Y. Nakamura, A. Ishizawa, S. Konoshima, T. Mizuuchi, K. Nagasaki, Simulation Study of Energetic Particle-driven MHD Modes and Energetic, 4th Asia-Pacific Conference on Plasma Physics, Online, 2020.10.26-31

岡佑旗, 長崎百伸, 小林進二, 大島慎介, 笠原寛史, 伊神弘恵, 門信一郎, 南貴司, 中村祐司, 石澤明宏, 木島滋, 水内亨, 岡田浩之, 加藤悠, 磁場閉じ込め核融合プラズマにおけるミリ波帯域電磁波モード変換シミュレーション, COMSOL Conference 2020 Tokyo, オンライン開催, 2020.12.4-11

森敦樹, 門信一郎, 本島巖, 岩田晃拓, 岡野竜成, 馮超, 岡田浩之, 南貴司, 大島慎介, 小林進二, 中村祐司, 石澤明宏, 木島滋, 水内亨, 長崎百伸, ヘリオトロン J におけるペレット溶発雲の 2 次元計測のための変倍分光器の開発, 第 37 回プラズマ・核融合学会年会, オンライン開催, 2020.12.1-4

村田駿介, 小林進二, 門信一郎, 大島慎介, 小林達哉, 南貴司, 岡田浩之, 吉川正志, 中村祐司, 石澤明宏, 水内亨, 木島滋, 長崎百伸, ヘリオトロン J におけるビーム放射分光法を用いた密度揺動計測, 第 37 回プラズマ・核融合学会年会, オンライン開催, 2020.12.1-4

山中雄太, 南貴司, 大島慎介, Dechuan Qiu, 三好正博, 篠塚凌我, 南雲竜也, 釧持尚輝, 門信一郎, 岡田浩之, 小林進二, 木島滋, 水内亨, 長崎百伸, ヘリオトロン J における異なる磁場配位での ECH プラズマの実効電子熱拡散係数の比較II, 第 37 回プラズマ・核融合学会年会, オンライン開催, 2020.12.1-4

岡佑旗, 長崎百伸, 小林進二, 大島慎介, 笠原寛史, 岡田浩之, 門信一郎, 南貴司, 中村祐司, 石澤明宏, 木島滋, 水内亨, 加藤悠, ヘリオトロン J における電子バーンシュタイン波加熱・計測のための有限要素法を用いた O-X モード変換解析, 第 37 回プラズマ・核融合学会年会, オンライン開催, 2020.12.1-4

徳原圭一, 小林進二, 大垣英明, 紀井俊輝, 全炳俊, 永岡賢一, 岡田浩之, 南貴司, 門信一郎, 大島慎介, 水内亨, 木島滋, 長崎百伸, ヘリオトロン J における非共鳴マイクロ波加熱プラズマ中の高エネルギー X 線スペクトル計測, 第 37 回プラズマ・核融合

学会年会, オンライン開催, 2020.12.1-4

上野陽平, 四竈泰一, 茶谷智樹, 門信一郎, 川染勇人, 岡田浩之, 松岡雷士, 南貴司, 小林進二, 大島慎介, 石澤明宏, 中村祐司, 木島滋, 水内亨, 長崎百伸, 蓮尾昌裕, 核融合プラズマ中原子輝線スペクトルのゼーマン効果高感度計測に向けた近赤外分光システム開発, 第 37 回プラズマ・核融合学会年会, オンライン開催, 2020.12.1-4

X.N.S. Bui, H. Matsuura, R. Matoike, S. Ohshima, T. Mizuuchi, Heat Flux Estimation with Divertor Probes Array in Heliotron J, 第 37 回プラズマ・核融合学会年会, Online, 2020.12.1-4

P. Zhang, S. Ohshima, H. Kume, S. Kobayashi, H. Okada, T. Minami, S. Kado, S. Konoshima, T. Mizuuchi, K. Nagasaki, Development of a320 GHz interferometer system in Heliotron J, 第 37 回プラズマ・核融合学会年会, Online, 2020.12.1-4

宮下顕, 向井清史, 大島慎介, 的池遼太, 小林進二, 岡田浩之, 門信一郎, 南貴司, 水内亨, 木島滋, 長崎百伸, ヘリオトロン J におけるイメージングボロメータの開発, 第 37 回プラズマ・核融合学会年会, オンライン開催, 2020.12.1-4

岡野竜成, 門信一郎, 岩田晃拓, 森敦樹, 馮超, 岡田浩之, 南貴司, 大島慎介, 小林進二, 本島巖, 中村祐司, 石澤明宏, 木島滋, 水内亨, 長崎百伸, ヘリオトロン J の固体水素ペレット入射実験における真空紫外スペクトルの時間発展計測, 第 37 回プラズマ・核融合学会年会, オンライン開催, 2020.12.1-4

三好正博, 南貴司, 釧持尚輝, Dechuan Qiu, 山中雄太, 篠塚凌我, 南雲竜也, 高橋千尋, 岡田浩之, 門信一郎, 大島慎介, 小林進二, 木島滋, 水内亨, 長崎百伸, ヘリオトロン J における Nd:YAG レーザトムソン散乱計測のためのイベントトリガシステムの開発, 第 37 回プラズマ・核融合学会年会, オンライン開催, 2020.12.1-4

R. Matoike, G. Kawamura, S. Ohshima, Y. Suzuki, M. Kobayashi, S. Masuzaki, S. Kobayashi, S. Kado, T. Minami, H. Okada, T. Mizuuchi, S. Konoshima, Y. Feng, H. Frerichs, K. Nagasaki, Peripheral Plasma Transport Characteristics arising from 3D Magnetic Field Structure in Heliotron J, The 24th International Conference on Plasma Surface Interactions in Controlled Fusion Devices, オンライン開催, 2021.1.25-29

K. Nagasaki, Y. Sakamoto, T. Maekawa, A. Fukuyama, H. Idei, Y. Oda, ECCD Calculation for JA DEMO Reactors, Japan-Korea Workshop on Physics and Technology of Heating and Current Drive, オンライン開催,

2021.2.25-26

岡野竜成, 門信一郎, 大島慎介, 山中雄太, 鈴木琢士, 岩田晃拓, 森敦樹, 馮超, 南貴司, 小林進二, 中村祐司, 石澤明宏, 岡田浩之, 木島滋, 水内亨, 長崎百伸, ヘリオトロン J における真空紫外分光法を用いた低価数輝線の空間分布計測, 日本物理学会第 76 回年次大会, オンライン開催, 2021.3.12-15

茶谷智樹, 四竈泰一, 上野陽平, 門信一郎, 川染勇人, 南貴司, 岡田浩之, 松岡雷士, 小林進二, 大島慎介, 石澤明宏, 中村祐司, 木島滋, 水内亨, 長崎百伸, 蓮尾昌裕, ヘリオトロン J における近赤外原子輝線のゼーマン効果計測, 日本物理学会第 76 回年次大会, オンライン開催, 2021.3.12-15