

## **Static stretching time required to reduce iliacus muscle stiffness**

Shusuke Nojiri<sup>a</sup>, Masahide Yagi<sup>a</sup>, Yu Mizukami<sup>a</sup>, Noriaki Ichihashi<sup>a</sup>

*<sup>a</sup>Human Health Sciences, Graduate School of Medicine, Kyoto University*

*53 Shogoin-Kawahara-cho, Sakyo-ku, Kyoto, 606-8507, Japan*

### Corresponding Author

Shusuke Nojiri

E-mail: [nojiri.shusuke.35v@st.kyoto-u.ac.jp](mailto:nojiri.shusuke.35v@st.kyoto-u.ac.jp)

Human Health Sciences, Graduate School of Medicine, Kyoto University

53 Shogoin-Kawahara-cho, Sakyo-ku, Kyoto, 606-8507, Japan

# 1    **Static stretching time required to reduce iliacus muscle stiffness**

2

## 3    ABSTRACT

4    Static stretching (SS) is an effective intervention to reduce muscle stiffness and is also  
5    performed for the iliopsoas muscle. The iliopsoas muscle consists of the iliacus and  
6    psoas major muscles, among which the former has a greater physiological cross-  
7    sectional area and hip flexion moment arm. Static stretching time required to reduce  
8    muscle stiffness can differ among muscles, and the required time for the iliacus muscle  
9    remains unclear. The purpose of this study was to investigate the time required to reduce  
10    iliacus muscle stiffness. Twenty-six healthy men participated in this study. A 1-min hip  
11    extension SS was performed five times. Shear elastic modulus, an index of muscle  
12    stiffness, of the iliacus muscle was measured using ultrasonic shear wave elastography  
13    before SS and immediately after each SS. One-way repeated analysis of variance  
14    showed a statistical effect of time on the shear elastic modulus. A paired *t*-test with  
15    Holm adjustment revealed that the shear elastic moduli after 1-5 SS were statistically  
16    lower than that before SS. In addition, the shear elastic modulus after 5 SS was  
17    statistically lower than that after 1 SS. The results suggested that the stiffness of the  
18    iliacus muscle decreased with 1-min SS and further decreased with 5-min SS.

19 (200 words)

20

21 **KEYWORDS**

22 Iliacus muscle

23 Static stretching

24 Ultrasonic shear wave elastography

25

26 **Introduction**

27 Limited hip extension range of motion (ROM) owing to increased stiffness or shortening of the  
28 iliopsoas muscle is one of the functional impairments observed in athletes and patients (Ferber,  
29 Kendall, & McElroy, 2010; Harvey, 1998; Roach et al., 2015). Limited hip extension ROM can  
30 be a risk factor for various musculoskeletal disorders (Delp, Hess, Hungerford, & Jones, 1999;  
31 Krivickas & Feinberg, 1996). Limited hip extension ROM reduces peak hip extension angle  
32 during gait (Tsukagoshi et al., 2015), which leads to changes in gait such as shortened step length,  
33 decreased gait velocity, and increased pelvic motion (Kerrigan, Lee, Collins, Riley, & Lipsitz,  
34 2001; Miki et al., 2004; Perron, Malouin, Moffet, & McFadyen, 2000).

35         The iliopsoas muscle consists of the iliacus and psoas major muscles. The iliacus muscle  
36 has greater physiological cross-sectional area (PCSA) and hip flexion moment arm than the psoas  
37 major muscle (Blemker & Delp, 2005; Klein Horsman, Koopman, van der Helm, Prosé, & Veeger,  
38 2007). Therefore, increased stiffness or shortening of the iliacus muscle affects hip extension  
39 ROM more strongly than similar changes in the psoas major muscle.

40         Static stretching (SS) is an effective intervention to reduce muscle stiffness. Many  
41 previous studies have used ROM (Boyce & Brosky, 2008; Ryan et al., 2008), passive torque, and  
42 passive stiffness (Fowles, Sale, & MacDougall, 2000; S. Peter Magnusson, Simonsen, Aagaard,  
43 & Kjaer, 1996) as indices of SS effects. However, ROM is inadequate as an index of muscle

44 stiffness because it is influenced by not only muscle stiffness but also pain and stretch tolerance  
45 (Weppeler & Magnusson, 2010). Passive torque and passive stiffness reflect the stiffness of many  
46 tissues other than the muscle (e.g., ligaments and joint capsule).

47           Recently, shear elastic modulus, assessed using ultrasonic shear wave elastography  
48 (SWE), has been used as an index of muscle stiffness (Kusano et al., 2017; Umegaki et al., 2015;  
49 Umehara et al., 2017). SWE estimates muscle stiffness by calculating shear elastic modulus from  
50 shear wave speed (Bercoff, Tanter, & Fink, 2004). Several studies reported a high correlation  
51 between the shear elastic modulus and passive muscle force (Eby et al., 2013; Koo, Guo, Cohen,  
52 & Parker, 2013). Therefore, the stiffness of an individual muscle can be evaluated using SWE.

53           Investigating the time required to decrease muscle stiffness is important to perform  
54 effective stretching, and is useful in time-limited situations such as clinical and athletic situations.  
55 A few studies have investigated the time required to reduce muscle stiffness and reported different  
56 results (Kusano et al., 2017; Nakamura, Ikezoe, Takeno, & Ichihashi, 2013). One of the potential  
57 reasons for the different results could be the innate differences in the targeted muscles, especially  
58 muscle size. With regard to muscle size, the iliacus muscle has a much smaller volume compared  
59 to the hamstring muscles or the gastrocnemius (Klein Horsman et al., 2007). Therefore, if the time  
60 required to reduce muscle stiffness is related to muscle size, the time required to reduce the  
61 stiffness of the iliacus muscle would be shorter than that required for the hamstring muscles or

62 the gastrocnemius. In addition, it was reported that passive torque decreased gradually even after  
63 a statistically significant reduction in passive torque occurred compared with before SS  
64 (Nakamura et al., 2013). Therefore, it is also important to investigate the time course of muscle  
65 stiffness after the first statistical difference is observed to perform effective SS.

66 Thus far, no study has investigated the effect of SS on the iliacus muscle. While several  
67 studies have performed a long-term intervention by using hip extension SS (Kerrigan,  
68 Xenopoulos-Oddsson, Sullivan, Lelas, & Riley, 2003; Watt et al., 2011), its effect on muscle  
69 stiffness or the time course remains unclear.

70 The purpose of the present study was to investigate the time required for hip extension  
71 SS to reduce the stiffness of the iliacus muscle. We hypothesised that the time required to reduce  
72 muscle stiffness of the iliacus muscle would be shorter than that of the hamstring muscles or the  
73 gastrocnemius reported in previous studies.

74

## 75 **Methods**

### 76 *Participants*

77 The sample size required for multiple comparisons after a one-way repeated analysis of variance  
78 (ANOVA) (effect size = 0.58,  $\alpha$  error = 0.05, and power = 0.80) was calculated using G\* power  
79 software (Heinrich Heine University, Düsseldorf, Germany). The effect size was determined

80 based on a previous study that investigated the acute effect of SS using SWE (Kusano et al., 2017).  
81 The calculated sample size was 26. Twenty-six men (age:  $23.2 \pm 2.9$  years; height:  $170.5 \pm 5.9$   
82 cm; mass:  $63.7 \pm 6.3$  kg) were recruited for this study. None of the participants had  
83 musculoskeletal injury or neuromuscular disease in the hip or lumbar region. The exclusion  
84 criteria were (1) difficulty in taking the position at which the shear elastic modulus was measured  
85 owing to limited hip extension ROM, (2) no stretch sensation in their upper leg at maximal hip  
86 extension, and (3) pain or numbness in the right leg during SS.

87 This study was approved by the ethics committee of the Kyoto University Graduate  
88 School and the Faculty of Medicine (R0233-3). Each participant provided written informed  
89 consent for participation in the study.

90

### 91 *Experimental protocol*

92 Hip extension SS was performed for 1 min; this was repeated five times with 1-min rest  
93 intervals, corresponding to the time for measurement of shear elastic modulus. We used 1 min of  
94 SS to test the hypothesis that the time required to reduce the iliacus muscle stiffness would be  
95 shorter than 2 min. Also, we performed a total of 5 min of SS based on a previous study  
96 (Nakamura et al., 2013). The shear elastic modulus of the iliacus muscle was measured before  
97 SS (bSS) and immediately after each round of SS (SS1–SS5), corresponding to a total of six

98 measurements.

99           The participants were instructed to relax and not to activate their lower limb muscles  
100 throughout the experiment. Each participant lay supine with the hip joint positioned at the edge  
101 of the bed. The left hip was passively flexed as much as possible to tilt the pelvis backward  
102 maximally by an investigator (YM), and thereafter, the pelvis was fixed to the bed with a non-  
103 elastic belt. The right hip was held at 5° extension by another investigator (SN) and the shear  
104 elastic modulus was measured (Figure 1). All six measurements of the shear elastic modulus  
105 were performed at this position. We confirmed via a preliminary experiment that the shear  
106 elastic modulus of the iliacus muscle did not decrease by maintaining this position for 1 min. In  
107 hip extension SS, the left hip was maintained at maximal flexion by an investigator (YM), and  
108 the right hip was extended by another investigator (SN) to the maximal angle that could be  
109 achieved without the participants feeling any discomfort or pain (Figure 2). The right knee was  
110 maintained in full extension to avoid elongation of the rectus femoris. The maximal hip  
111 extension angle was measured during each round of SS and after all rounds of SS, using a 1°-  
112 scale goniometer. The hip extension angle was defined as the angle between the trunk and the  
113 femur. All measurements were obtained by the same three examiners, one of whom (MY)  
114 performed the measurement of the shear elastic modulus and the hip extension angle, and two of  
115 whom (YM and SN) fixed the limb position.



116

117 ***Measurement of shear elastic modulus***

118 Shear elastic modulus was measured to assess the muscle stiffness. Ultrasonic SWE (Aixplorer;  
119 SuperSonicImagine, Aix-en-Provence, France) with a SuperLinear SL 10-2 probe was used to  
120 measure the shear elastic modulus. The shear elastic modulus of the iliacus muscle was measured  
121 in the right limb. The measurement site was defined as a level 4 cm distal from anterior superior  
122 iliac spines, because it was reported that the iliopsoas muscle was located most superficially at  
123 this level (Jiroumaru, Kurihara, & Isaka, 2014). The iliacus muscle belly was identified at this  
124 level using a B-mode ultrasonic image. Subsequently, the measurement site was determined and  
125 marked on the skin. The probe was placed parallel to the muscle fiber on the mark, and it was  
126 confirmed that the muscle fiber was uninterrupted on the ultrasonic image. Subsequently, the  
127 shear elastic modulus was measured in ultrasonic SWE mode. The shear elastic modulus was  
128 measured twice at each time point, and the mean value was used for statistical analysis. The total  
129 time required for the two measurements in each round was < 1 min.

130 A region of interest (ROI), a square of side 1.5 cm, was set at the center of the iliacus  
131 muscle belly. A circle was drawn in full size within the ROI. The mean shear wave speed in the  
132 circle was calculated automatically (Figure 3). The shear elastic modulus (G) was calculated from  
133 the shear wave speed (V) using the following equation:

134  $G \text{ (kPa)} = \rho V^2,$

135 where  $\rho$  is the muscle mass density, which is assumed to be  $1000 \text{ kg/m}^3$  (Gennisson, Cornu,  
136 Catheline, Fink, & Portero, 2005). The calculation of shear elastic modulus values was performed  
137 by an investigator (SN), who was different from the investigator who measured the shear elastic  
138 modulus.

139 The intraclass correlation coefficient (ICC) was calculated in accordance with Shrout &  
140 Fleiss (1979) for the two measurements at bSS as an index of the reliability of shear elastic  
141 modulus values. ICC<sub>1,1</sub> was 0.85 (95% confidence interval [CI]: 0.69–0.93), and ICC<sub>1,2</sub> was  
142 0.92 (95% CI: 0.82–0.96), and therefore good reliability was observed (Portney & Watkins, 2000;  
143 Shrout & Fleiss, 1979).

144

#### 145 ***Statistical analysis***

146 Statistical analysis was performed using SPSS Statistics (version 22; IBM, Armonk, NY, USA).

147 A one-way repeated measures ANOVA was performed to assess the effect of time on the shear  
148 elastic modulus. When a statistical effect was observed, a post hoc test was performed. A paired  
149 *t*-test was performed between the shear elastic modulus at bSS and that at SS1–SS5.

150 Furthermore, the shear elastic moduli were compared between the time when the first statistical  
151 difference compared with bSS was observed and afterward, by using a paired *t*-test. The level of

152 statistical rareness was set at  $P < 0.05$ . In post hoc tests,  $P$  values were corrected with Holm  
153 adjustment in each  $t$ -test. We estimated the effect size using partial  $\eta^2$  and  $r$  for the one-way  
154 repeated measures ANOVA and post hoc test, respectively. The partial  $\eta^2$  value is considered  
155 moderate and large when it is  $\geq 0.07$  and  $\geq 0.14$ , respectively (Cohen, 1988).

156

## 157 **Results**

158 The shear elastic modulus at each time point is shown in Table 1 as a mean  $\pm$  standard deviation.

159 The maximal hip extension angle during each round of SS is shown in Table 2 as a mean  $\pm$   
160 standard deviation.

161 The one-way repeated measures ANOVA showed a statistical effect of time (effect size  
162 partial  $\eta^2 = 0.31$ ). The post hoc test revealed that the shear elastic moduli at SS1–SS5 were  
163 statistically lower than at bSS. Moreover, from a comparison of the shear elastic moduli using a  
164 paired  $t$ -test between SS1 and SS2–SS5, the shear elastic modulus at SS5 was observed to be  
165 statistically lower than at SS1.

166

## 167 **Discussion and implications**

168 In this study, we investigated the effect of hip extension SS on the stiffness of the iliacus muscle  
169 using SWE. The shear elastic moduli at measurements SS1–SS5 were statistically lower than that

170 at bSS. This result suggests that the stiffness of the iliacus muscle decreased with 1 min of SS,  
171 and is consistent with our hypothesis. Furthermore, the shear elastic modulus at SS5 was  
172 statistically lower than that at SS1. This result suggests that the stiffness of the iliacus muscle  
173 further decreased with 5 min of SS compared with 1 min of SS. To the best of our knowledge,  
174 this is the first study to demonstrate the time required for hip extension SS to reduce the stiffness  
175 of the iliacus muscle.

176           Previous studies reported that passive torque or passive stiffness decreased after 2–2.5  
177 min of SS (Nakamura et al., 2013; Nordez, Cornu, & McNair, 2006) and did not decrease after  
178 1–1.5 min of SS (S. P. Magnusson, Aagard, Simonsen, & Bojsen-Møller, 1998; McNair,  
179 Dombroski, Hewson, & Stanley, 2001). Therefore, more than 2 min of SS has been considered  
180 necessary to reduce muscle stiffness (Akagi & Takahashi, 2013; Nakamura et al., 2014, 2013).  
181 However, the shear elastic modulus of the iliacus muscle decreased after 1 min of SS in this study.  
182 The reasons for the shorter time in this study could be explained by the difference in the muscle  
183 size and the index of muscle flexibility.

184           Previous studies investigated the time to reduce muscle stiffness in hamstring muscles  
185 (S. P. Magnusson et al., 1998; Nordez et al., 2006) or the gastrocnemius (McNair et al., 2001;  
186 Nakamura et al., 2013). Kusano et al. (2017) reported that the stiffness of the infraspinatus muscle  
187 decreased after 20 s of SS. They explained that the smaller muscle size could be the reason for

188 the shorter time required. With regards to muscle size, the volume of the iliacus muscle is smaller  
189 than that of the hamstring muscles or the gastrocnemius (Klein Horsman et al., 2007). Therefore,  
190 the shorter time in this study could be explained by the smaller size of the iliacus muscle compared  
191 with that of hamstring or the gastrocnemius muscles.

192           The difference in the index of muscle stiffness could also be the reason for the shorter  
193 time required in the current study. The referred studies used passive torque or passive stiffness as  
194 an index of muscle stiffness (S. P. Magnusson et al., 1998; McNair et al., 2001; Nakamura et al.,  
195 2013; Nordez et al., 2006). While those indices reflect the stiffness of not only the muscle but also  
196 the entire joint complex, we evaluated the stiffness of the iliacus muscle solely by using SWE. By  
197 using shear elastic modulus as an index of muscle stiffness, Kusano et al. (2017) reported much  
198 shorter time than the referred studies that used passive torque and passive stiffness as an index of  
199 muscle stiffness (S. P. Magnusson et al., 1998; McNair et al., 2001; Nakamura et al., 2013; Nordez  
200 et al., 2006). In other words, it is indicated that the stiffness of muscle decreases earlier than that  
201 of the entire joint complex.

202           Furthermore, the shear elastic modulus of the iliacus muscle decreased gradually over  
203 every SS and a statistically significant difference was observed with SS5 compared with SS1.  
204 This result suggests that the stiffness of the iliacus muscle decreased further with 5 min of SS than  
205 1 min of SS. Nakamura et al. (2013) reported a gradual decrease in passive torque over every

206 minute during 5 min of SS, which was similar to the result of this study. They showed that passive  
207 torque decreased statistically after 2 min of SS compared with before SS and decreased  
208 statistically after 5 min of SS compared with 2 min of SS. The mechanism of gradual decrease of  
209 passive torque was reported to be viscoelastic stress relaxation, which is a decline in the stress or  
210 force of the tissues when held at an extended position (Taylor, Dalton, Seaber, & Garrett, 1990).  
211 It has been reported that the force declines rapidly in the first few tens of seconds and thereafter  
212 declines gradually until 5 min (McNair et al., 2001; Toft, Sinkjaer, Kålund, & Espersen, 1989). In  
213 this study, five repetitions of SS could cause viscoelastic stress relaxation as well as 5 consecutive  
214 min of SS in the previous study (Nakamura et al., 2013).

215           In this study, a gradual decrease in muscle stiffness similar to that in consecutive SS was  
216 observed in repeated SS. This result could be clinically beneficial. This is because repeating 1  
217 min of SS five times may be much easier for therapists than performing 5 consecutive min of SS.

218           There are a few limitations to this study. First, 1 min of SS might not necessarily be  
219 required to reduce the shear elastic modulus of the iliacus muscle because the effect of SS shorter  
220 than 1 min is unclear. However, we confirmed that the shear elastic modulus of the iliacus muscle  
221 hardly decreased in a preliminary experiment in which 30 s of SS was repeated. Therefore, we  
222 chose to repeat 1 min of SS. Second, we investigated only the acute effect of SS, and the duration  
223 of the effect and the long-term effect are unclear. Therefore, the acute effect of SS for shorter time

224 intervals (i.e., 30–60 s), the effect of long-term intervention, and the effect on performance will  
225 be further investigated. Third, the effects of SS on the psoas major remain unclear, although we  
226 chose the iliacus muscle rather than the psoas major, based on the greater PCSA and hip flexion  
227 moment arm.

228

## 229 **Conclusion**

230 In this study it was suggested that the stiffness of the iliacus muscle decreased with 1 min of hip  
231 extension SS and further decreased with 5 min of SS.

232

233 **Acknowledgments**

234 This work was supported by Grant-in-Aid for Scientific Research (B) 15H03043. We would like  
235 to thank Ms. Ibuki and Editage (www.editage.jp) for English language editing.

236

237 **Disclosure statement**

238 The authors declare no conflicts of interest.

239

240 **Funding**

241 This work was supported by Grant-in-Aid for Scientific Research (B) (15H03043).

242

243 **References**

244 Akagi, R., & Takahashi, H. (2013). Acute effect of static stretching on hardness of the  
245 gastrocnemius muscle. *Medicine and Science in Sports and Exercise*, 45(7), 1348–1354.

246 <https://doi.org/10.1249/MSS.0b013e3182850e17>

247 Bercoff, J., Tanter, M., & Fink, M. (2004). Supersonic shear imaging: a new technique for soft  
248 tissue elasticity mapping. *IEEE Transactions on Ultrasonics, Ferroelectrics, and*

249 *Frequency Control*, 51(4), 396–409. <https://doi.org/10.1109/TUFFC.2004.1295425>

250 Blemker, S. S., & Delp, S. L. (2005). Three-dimensional representation of complex muscle



251 architectures and geometries. *Annals of Biomedical Engineering*, 33(5), 661–673.

252 <https://doi.org/10.1007/s10439-005-1433-7>

253 Boyce, D., & Brosky, J. A. (2008). Determining the minimal number of cyclic passive stretch

254 repetitions recommended for an acute increase in an indirect measure of hamstring length.

255 *Physiotherapy Theory and Practice*, 24(2), 113–120.

256 <https://doi.org/10.1080/09593980701378298>

257 Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed). New York:

258 Lawrence Erlbaum Associates.

259 Delp, S. L., Hess, W. E., Hungerford, D. S., & Jones, L. C. (1999). Variation of rotation

260 moment arms with hip flexion. *Journal of Biomechanics*, 32(5), 493–501.

261 [https://doi.org/10.1016/S0021-9290\(99\)00032-9](https://doi.org/10.1016/S0021-9290(99)00032-9)

262 Eby, S. F., Song, P., Chen, S., Chen, Q., Greenleaf, J. F., & An, K.-N. (2013). Validation of

263 shear wave elastography in skeletal muscle. *Journal of Biomechanics*, 46(14), 2381–2387.

264 <https://doi.org/10.1016/j.jbiomech.2013.07.033>

265 Ferber, R., Kendall, K. D., & McElroy, L. (2010). Normative and critical criteria for iliotibial

266 band and iliopsoas muscle flexibility. *Journal of Athletic Training*, 45(4), 344–348.

267 <https://doi.org/10.4085/1062-6050-45.4.344>

268 Fowles, J. R., Sale, D. G., & MacDougall, J. D. (2000). Reduced strength after passive stretch

269 of the human plantarflexors. *Journal of Applied Physiology (Bethesda, Md. : 1985)*, 89(3),  
270 1179–1188. <https://doi.org/10.1152/jappl.2000.89.3.1179>

271 Gennisson, J. L., Cornu, C., Catheline, S., Fink, M., & Portero, P. (2005). Human muscle  
272 hardness assessment during incremental isometric contraction using transient elastography.  
273 *Journal of Biomechanics*, 38(7), 1543–1550.  
274 <https://doi.org/10.1016/j.jbiomech.2004.07.013>

275 Harvey, D. (1998). Assessment of the flexibility of elite athletes using the modified Thomas  
276 test. *British Journal of Sports Medicine*, 32(1), 68–70.  
277 <https://doi.org/10.1136/bjism.32.1.68>

278 Jiroumaru, T., Kurihara, T., & Isaka, T. (2014). Establishment of a recording method for surface  
279 electromyography in the iliopsoas muscle. *Journal of Electromyography and Kinesiology :  
280 Official Journal of the International Society of Electrophysiological Kinesiology*, 24(4),  
281 445–451. <https://doi.org/10.1016/j.jelekin.2014.02.007>

282 Kerrigan, D. C., Lee, L. W., Collins, J. J., Riley, P. O., & Lipsitz, L. A. (2001). Reduced hip  
283 extension during walking: healthy elderly and fallers versus young adults. *Archives of  
284 Physical Medicine and Rehabilitation*, 82(1), 26–30.  
285 <https://doi.org/10.1053/apmr.2001.18584>

286 Kerrigan, D. C., Xenopoulos-Oddsson, A., Sullivan, M. J., Lelas, J. J., & Riley, P. O. (2003).

287 Effect of a hip flexor-stretching program on gait in the elderly. *Archives of Physical*  
288 *Medicine and Rehabilitation*, 84(1), 1–6. <https://doi.org/10.1053/apmr.2003.50056>

289 Klein Horsman, M. D., Koopman, H. F. J. M., van der Helm, F. C. T., Prosé, L. P., & Veeger,  
290 H. E. J. (2007). Morphological muscle and joint parameters for musculoskeletal modelling  
291 of the lower extremity. *Clinical Biomechanics (Bristol, Avon)*, 22(2), 239–247.  
292 <https://doi.org/10.1016/j.clinbiomech.2006.10.003>

293 Koo, T. K., Guo, J.-Y., Cohen, J. H., & Parker, K. J. (2013). Relationship between shear elastic  
294 modulus and passive muscle force: an ex-vivo study. *Journal of Biomechanics*, 46(12),  
295 2053–2059. <https://doi.org/10.1016/j.jbiomech.2013.05.016>

296 Krivickas, L. S., & Feinberg, J. H. (1996). Lower extremity injuries in college athletes: relation  
297 between ligamentous laxity and lower extremity muscle tightness. *Archives of Physical*  
298 *Medicine and Rehabilitation*, 77(11), 1139–1143. [https://doi.org/10.1016/S0003-](https://doi.org/10.1016/S0003-9993(96)90137-9)  
299 [9993\(96\)90137-9](https://doi.org/10.1016/S0003-9993(96)90137-9)

300 Kusano, K., Nishishita, S., Nakamura, M., Tanaka, H., Umehara, J., & Ichihashi, N. (2017).  
301 Acute effect and time course of extension and internal rotation stretching of the shoulder  
302 on infraspinatus muscle hardness. *Journal of Shoulder and Elbow Surgery*, 26(10), 1782–  
303 1788. <https://doi.org/10.1016/j.jse.2017.04.018>

304 Magnusson, S. P., Aagard, P., Simonsen, E., & Bojsen-Møller, F. (1998). A biomechanical

305 evaluation of cyclic and static stretch in human skeletal muscle. *International Journal of*  
306 *Sports Medicine*, 19(5), 310–316. <https://doi.org/10.1055/s-2007-971923>

307 Magnusson, S. P., Simonsen, E. B., Aagaard, P., & Kjaer, M. (1996). Biomechanical responses  
308 to repeated stretches in human hamstring muscle in vivo. *The American Journal of Sports*  
309 *Medicine*, 24(5), 622–628. <https://doi.org/10.1177/036354659602400510>

310 McNair, P. J., Dombroski, E. W., Hewson, D. J., & Stanley, S. N. (2001). Stretching at the  
311 ankle joint: viscoelastic responses to holds and continuous passive motion. *Medicine and*  
312 *Science in Sports and Exercise*, 33(3), 354–358. [https://doi.org/10.1097/00005768-](https://doi.org/10.1097/00005768-200103000-00003)  
313 [200103000-00003](https://doi.org/10.1097/00005768-200103000-00003)

314 Miki, H., Sugano, N., Hagio, K., Nishii, T., Kawakami, H., Kakimoto, A., ... Yoshikawa, H.  
315 (2004). Recovery of walking speed and symmetrical movement of the pelvis and lower  
316 extremity joints after unilateral THA. *Journal of Biomechanics*, 37(4), 443–455.  
317 <https://doi.org/10.1016/j.jbiomech.2003.09.009>

318 Nakamura, M., Ikezoe, T., Kobayashi, T., Umegaki, H., Takeno, Y., Nishishita, S., & Ichihashi,  
319 N. (2014). Acute effects of static stretching on muscle hardness of the medial  
320 gastrocnemius muscle belly in humans: an ultrasonic shear-wave elastography study.  
321 *Ultrasound in Medicine & Biology*, 40(9), 1991–1997.  
322 <https://doi.org/10.1016/j.ultrasmedbio.2014.03.024>

323 Nakamura, M., Ikezoe, T., Takeno, Y., & Ichihashi, N. (2013). Time course of changes in  
324 passive properties of the gastrocnemius muscle-tendon unit during 5 min of static  
325 stretching. *Manual Therapy, 18*(3), 211–215. <https://doi.org/10.1016/j.math.2012.09.010>

326 Nordez, A., Cornu, C., & McNair, P. (2006). Acute effects of static stretching on passive  
327 stiffness of the hamstring muscles calculated using different mathematical models.  
328 *Clinical Biomechanics (Bristol, Avon), 21*(7), 755–760.  
329 <https://doi.org/10.1016/j.clinbiomech.2006.03.005>

330 Perron, M., Malouin, F., Moffet, H., & McFadyen, B. J. (2000). Three-dimensional gait analysis  
331 in women with a total hip arthroplasty. *Clinical Biomechanics (Bristol, Avon), 15*(7), 504–  
332 515. [https://doi.org/10.1016/S0268-0033\(00\)00002-4](https://doi.org/10.1016/S0268-0033(00)00002-4)

333 Portney, L., & Watkins, M. (2000). *Foundations of clinical research: application to practice*  
334 (2nd ed). New Jersey: Prentice Hall Health.

335 Roach, S. M., San Juan, J. G., Suprak, D. N., Lyda, M., Bies, A. J., & Boydston, C. R. (2015).  
336 Passive hip range of motion is reduced in active subjects with chronic low back pain  
337 compared to controls. *International Journal of Sports Physical Therapy, 10*(1), 13–20.  
338 Retrieved from  
339 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4325283&tool=pmcentrez&re](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4325283&tool=pmcentrez&rendertype=abstract)  
340 [ndertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4325283&tool=pmcentrez&rendertype=abstract)

341 Ryan, E. D., Beck, T. W., Herda, T. J., Hull, H. R., Hartman, M. J., Stout, J. R., & Cramer, J. T.  
342 (2008). Do practical durations of stretching alter muscle strength? A dose-response study.  
343 *Medicine and Science in Sports and Exercise*, 40(8), 1529–1537.  
344 <https://doi.org/10.1249/MSS.0b013e31817242eb>

345 Shrout, P. E., & Fleiss, J. L. (1979). Intraclass correlations: uses in assessing rater reliability.  
346 *Psychological Bulletin*, 86(2), 420–428. Retrieved from [http://rokwa.x-y.net/Shrout-Fleiss-](http://rokwa.x-y.net/Shrout-Fleiss-ICC.pdf)  
347 [ICC.pdf](http://rokwa.x-y.net/Shrout-Fleiss-ICC.pdf)

348 Taylor, D. C., Dalton, J. D., Seaber, A. V., & Garrett, W. E. (1990). Viscoelastic properties of  
349 muscle-tendon units. The biomechanical effects of stretching. *The American Journal of*  
350 *Sports Medicine*, 18(3), 300–309. <https://doi.org/10.1177/036354659001800314>

351 Toft, E., Sinkjaer, T., Kålund, S., & Espersen, G. T. (1989). Biomechanical properties of the  
352 human ankle in relation to passive stretch. *Journal of Biomechanics*, 22(11–12), 1129–  
353 1132. [https://doi.org/10.1016/0021-9290\(89\)90214-5](https://doi.org/10.1016/0021-9290(89)90214-5)

354 Tsukagoshi, R., Tateuchi, H., Fukumoto, Y., Akiyama, H., So, K., Kuroda, Y., ... Ichihashi, N.  
355 (2015). Factors associated with restricted hip extension during gait in women after total  
356 hip arthroplasty. *Hip International : The Journal of Clinical and Experimental Research*  
357 *on Hip Pathology and Therapy*, 25(6), 543–548. <https://doi.org/10.5301/hipint.5000286>

358 Umegaki, H., Ikezoe, T., Nakamura, M., Nishishita, S., Kobayashi, T., Fujita, K., ... Ichihashi,

359 N. (2015). Acute effects of static stretching on the hamstrings using shear elastic modulus  
360 determined by ultrasound shear wave elastography: Differences in flexibility between  
361 hamstring muscle components. *Manual Therapy*, 20(4), 610–613.

362 <https://doi.org/10.1016/j.math.2015.02.006>

363 Umehara, J., Hasegawa, S., Nakamura, M., Nishishita, S., Umegaki, H., Tanaka, H., ...

364 Ichihashi, N. (2017). Effect of scapular stabilization during cross-body stretch on the  
365 hardness of infraspinatus, teres minor, and deltoid muscles: An ultrasonic shear wave  
366 elastography study. *Musculoskeletal Science & Practice*, 27, 91–96.

367 <https://doi.org/10.1016/j.math.2016.10.004>

368 Watt, J. R., Jackson, K., Franz, J. R., Dicharry, J., Evans, J., & Kerrigan, D. C. (2011). Effect of  
369 a supervised hip flexor stretching program on gait in elderly individuals. *PM & R : The  
370 Journal of Injury, Function, and Rehabilitation*, 3(4), 324–329.

371 <https://doi.org/10.1016/j.pmrj.2010.11.012>

372 Wepler, C. H., & Magnusson, S. P. (2010). Increasing muscle extensibility: a matter of  
373 increasing length or modifying sensation? *Physical Therapy*, 90(3), 438–449.

374 <https://doi.org/10.2522/ptj.20090012>

375

376

**Table 1 Shear elastic modulus of the iliacus muscle at each time point**

	Shear elastic modulus (kPa)	Vs. bSS		Vs. SS1	
		<i>P</i> value	effect size (r)	<i>P</i> value	effect size (r)
bSS	22.1 ± 3.5	-	-	-	-
SS1	20.5 ± 4.2	0.008	0.50	-	-
SS2	20.1 ± 4.4	0.008	0.54	0.49	0.14
SS3	19.8 ± 3.7	<0.001	0.71	0.28	0.29
SS4	19.4 ± 3.5	<0.001	0.69	0.19	0.36
SS5	18.2 ± 2.4	<0.001	0.85	0.006	0.58

The shear elastic modulus is expressed as a mean ± standard deviation.

SS: static stretching

377

378

379

**Table 2 Maximal hip extension angle during each round of SS and after SS**

Maximal hip extension angle (°)	
1st SS	19 ± 4
2nd SS	21 ± 5
3rd SS	23 ± 5
4th SS	25 ± 5
5th SS	26 ± 5
After SS	26 ± 6

Results are expressed as a mean ± standard deviation.

The angle was measured during each round of SS and after all rounds of SS. The angle during 2nd SS was indicated as the maximal angle, which was a result of 1st SS, for example.

380



381 **Figure captions**

382



383

384

385 **Figure 1 Position at which the shear elastic modulus was measured**

386 The left hip was maintained at maximal flexion and the right hip was maintained at 5° extension.

387

388

389



390


391

392 **Figure 2 Position of static stretching**

393 The left hip was maintained at maximal flexion and the right hip was extended to the maximal

394 angle at which there was no pain or discomfort.

395



396

397

398 **Figure 3 Typical example of measuring the shear wave speed**

399 An ROI, a square of side 1.5 cm, was set at the center of the iliacus muscle belly. A circle was  
400 drawn in full size within the ROI. The mean shear wave speed in the circle was calculated  
401 automatically.

402