

SOLUTIONS FOR A FRACTIONAL-ORDER DIFFERENTIAL EQUATION WITH BOUNDARY CONDITIONS

MEIJI UNIVERSITY
TOSHIKAZU WATANABE

1. INTRODUCTION

In [5], the authors considered the fractional-order boundary value problem, we consider existence and uniqueness of solutions of the fractional-order boundary value problem

$$(1.1) \quad \begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t)), & 0 < t < 1, \\ u(0) = u'(0) = u''(1) = u'''(1) = 0. \end{cases}$$

Inspiring the result of Ma and Silva [4], we consider the problem (1.2). The problem (1.1) is the case that $g \equiv 0$ of the problem (1.2). Due to the restriction of g , results in [5] cannot deal with the problem (1.3). where $3 < \alpha \leq 4$, f is a continuous function of $[0, 1] \times \mathbb{R}$ into \mathbb{R} , g is a function of \mathbb{R} into itself and D_{0+}^{α} denotes the Riemann-Liouville fractional derivative of order α which is defined in Section 2. A function $u \in C[0, 1]$, where $C[0, 1]$ is the set of all continuous functions of $[0, 1]$ into \mathbb{R} , is called a solution of the problem (1.1) if $D_{0+}^{\alpha} u \in C[0, 1]$ and u satisfies (1.1).

$$(1.2) \quad \begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t)), & 0 < t < 1, \\ u(0) = u'(0) = u''(1) = 0, \quad u'''(1) = g(u(1)), \end{cases}$$

A function $u \in C[0, 1]$, where $C[0, 1]$ is the set of all continuous functions of $[0, 1]$ into \mathbb{R} , is called a solution of the problem (1.2) if $D_{0+}^{\alpha} u \in C[0, 1]$ and u satisfies (1.2).

When $\alpha = 4$, the problem (1.1) is the boundary value problems for cantilever beam equations and the problem (1.2) is the following fourth-order boundary value problem

$$(1.3) \quad \begin{cases} u''''(t) = f(t, u(t)), & 0 < t < 1, \\ u(0) = u'(0) = u''(1) = 0, \quad u'''(1) = g(u(1)). \end{cases}$$

In the problem (1.3), since u'' represents the shear force at $t = 1$, the conditions $u''(1) = 0, u'''(1) = g(u(1))$ means that the vertical force is equal to $g(u(1))$, which denotes a relation, possibly nonlinear, between the vertical force and the displacement u . Furthermore, since $u''(1) = 0$ indicates that there is no bending moment at $t = 1$, the beam is resting on the bearing g . Existence and iterative schemes to solve the problem (1.3) were studied by Ma and da Silva [4]. The purpose of this article is to follow the results of the fractional-order problems (1.1) and (1.2).

2010 *Mathematics Subject Classification.* Primary 46A40, 47B50, 47H10.

Key words and phrases. Fixed point theorem, fractional boundary value problem.

2. PRELIMINARIES

In this section, we introduce preliminary facts. Especially, we construct the Green function $G(s, t)$ for the boundary value problems (1.1) and (1.2), and we discuss some properties of the function.

We start with the definition of the Riemann-Liouville fractional integral and fractional derivative. Let $\alpha > 0$ and u be a continuous function of $[0, 1]$ into \mathbb{R} . The Riemann-Liouville fractional integral of order α of u , denoted $I_{0+}^\alpha u$, is defined by

$$I_{0+}^\alpha u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} u(s) ds$$

for $0 \leq t \leq 1$. The Riemann-Liouville fractional derivative of order α of u , denoted $D_{0+}^\alpha u$, is defined by

$$D_{0+}^\alpha u(t) = \frac{1}{\Gamma(n-\alpha)} \frac{d^n}{dt^n} \int_0^t (t-s)^{n-\alpha-1} u(s) ds$$

for $0 \leq t \leq 1$, where n denotes a positive integer such that $n-1 < \alpha \leq n$. For $\alpha \geq 0$ and $\beta > -1$, we have

$$D_{0+}^\alpha t^\beta = \frac{\Gamma(\beta+1)}{\Gamma(\beta-\alpha+1)} t^{\beta-\alpha};$$

see [3]. For the case of (1.2), let h be a continuous mapping of $[0, 1]$ into \mathbb{R} . Let $3 < \alpha \leq 4$. Then the unique solution of the boundary value problem is

$$u(t) = \int_0^1 G(t, s)h(s)ds,$$

where

(2.1)

$$G(t, s) = \begin{cases} \frac{1}{\Gamma(\alpha)} ((t-s)^{\alpha-1} + t^{\alpha-1}(1-s)^{\alpha-4}(4s-\alpha s-1) + (\alpha-1)t^{\alpha-2}(1-s)^{\alpha-4}s) & (0 \leq s \leq t < 1), \\ \frac{1}{\Gamma(\alpha)} (t^{\alpha-1}(1-s)^{\alpha-4}(4s-\alpha s-1) + (\alpha-1)t^{\alpha-2}(1-s)^{\alpha-4}s) & (0 \leq t \leq s < 1); \end{cases}$$

see [5]. Also for the case of (1.2)

$$\begin{cases} D_{0+}^\alpha u(t) = h(t), & 0 < t < 1, \\ u(0) = u'(0) = u''(1) = 0, & u'''(1) = \gamma \end{cases}$$

if and only if u is a solution of the integral equation

$$(2.2) \quad u(t) = \int_0^1 G(t, s)h(s)ds + \frac{\gamma t^{\alpha-1}}{(\alpha-1)(\alpha-2)} - \frac{\gamma t^{\alpha-2}}{(\alpha-2)(\alpha-3)}$$

for $0 \leq t \leq 1$, where $G(t, s)$ is defined by (2.1); see [6].

3. MAIN RESULTS

In this section, we consider the boundary value problems (1.1) and (1.2). By the Banach fixed point theorem, we obtain a sufficient condition for uniqueness and existence of solutions of the problems.

Theorem 1. *Let $3 < \alpha \leq 4$. Let f be a continuous function of $[0, 1] \times \mathbb{R}$ into \mathbb{R} . Let g be a Lipschitz continuous function of \mathbb{R} into itself with a nonnegative constant L . Assume that there exists a nonnegative constant λ with*

$$\lambda\Lambda + \frac{2L}{(\alpha - 1)(\alpha - 2)(\alpha - 3)} < 1$$

such that for any $0 \leq t \leq 1$ and $u_1, u_2 \in \mathbb{R}$,

$$|f(t, u_1) - f(t, u_2)| \leq \lambda|u_1 - u_2|,$$

where Λ is the constant

$$\Lambda = \sup_{0 \leq t \leq 1} \int_0^1 |G(t, s)| ds.$$

Then the boundary value problem (1.2) has a unique solution.

Proof. See [6, Theorem 1]. □

For the case that $g = 0$ in Theorem 1, we have the following; see [5, Theorem 3.1].

Corollary 2. *Let $3 < \alpha \leq 4$. Let f be a continuous mapping of $[0, 1] \times \mathbb{R}$ into \mathbb{R} . Assume that there exists $\lambda \in [0, \frac{1}{\Lambda})$ such that for any $u, v \in [0, \infty)$ and $t \in [0, 1]$,*

$$|f(t, u) - f(t, v)| \leq \lambda|u - v|,$$

where

$$\Lambda = \sup_{0 \leq t \leq 1} \int_0^1 |G(t, s)| ds$$

and G is the function given by (2.1). Then the boundary value problem represented by (1.1) has a unique solution.

For the case that $\alpha = 4$ in Theorem 1, we have the following; see Theorem 1 in [4].

Corollary 3. *Let f be a continuous function of $[0, 1] \times \mathbb{R}$ into \mathbb{R} with bounded partial derivative with respect to the second variable. Let g be a Lipschitz continuous function of \mathbb{R} into itself with a nonnegative constant L . Let*

$$\lambda = \max_{(t, u) \in [0, 1] \times \mathbb{R}} \left| \frac{\partial f}{\partial u}(t, u) \right|.$$

If

$$\frac{\lambda}{8} + \frac{L}{3} < 1,$$

then the boundary value problem (1.3) has a unique solution.

Proof. By the mean value theorem, we have for any $0 \leq t \leq 1$ and $u_1, u_2 \in \mathbb{R}$,

$$|f(t, u_1) - f(t, u_2)| \leq \lambda|u_1 - u_2|.$$

In the case that $\alpha = 4$, the function $G(t, s)$ reduces to

$$G(t, s) = \begin{cases} \frac{1}{6}s^2(3t - s) & (s < t), \\ \frac{1}{6}t^2(3s - t) & (t \leq s). \end{cases}$$

Since

$$\Lambda = \sup_{0 \leq t \leq 1} \int_0^1 |G(t, s)| ds = \frac{1}{8},$$

we obtain the conclusion by Theorem 1. □

Remark 4. Let $3 < \alpha \leq 4$. Since the function $G(t, s)$ satisfies

$$(3.1) \quad \int_0^1 |G(t, s)| ds \leq \frac{1}{\Gamma(\alpha)} \left(\frac{1}{\alpha} + \frac{\alpha}{\alpha - 3} \right)$$

for all $0 \leq t \leq 1$, $\sup_{0 \leq t \leq 1} \int_0^1 |G(t, s)| ds$ is finite. The function $G(t, s)$ satisfies

$$l(t, s) \leq G(t, s) \leq m(t, s)$$

for $0 \leq t \leq 1$ and $0 \leq s < 1$. where

$$l(t, s) = \begin{cases} \frac{1}{\Gamma(\alpha)} t^{\alpha-2} (1-s)^{\alpha-4} (2s + st - t) & (s < t), \\ \frac{\alpha-2}{\Gamma(\alpha)} t^{\alpha-2} (1-s)^{\alpha-4} s & (t \leq s) \end{cases}$$

and

$$m(t, s) = \begin{cases} \frac{\alpha-1}{\Gamma(\alpha)} t^{\alpha-2} (1-s)^{\alpha-4} s & (s < t), \\ \frac{3}{\Gamma(\alpha)} t^{\alpha-2} (1-s)^{\alpha-4} s & (t \leq s); \end{cases}$$

see [6].

To conclude the paper, we present an example demonstrating an application of Theorem 1.

Example 5. Let us consider the boundary value problem

$$(3.2) \quad \begin{cases} D_{0+}^{3.1} u(t) = \frac{3}{(54e^t + 1)(1 + |u(t)|)}, & 0 < t < 1, \\ u(0) = u'(0) = u''(1) = 0, & u'''(1) = g(u(1)), \end{cases}$$

where

$$g(t) = \frac{1}{100} \sin t.$$

By (3.1), the constant Λ in Theorem 1 satisfies

$$\Lambda \leq \frac{1}{\Gamma(\alpha)} \left(\frac{1}{\alpha} + \frac{\alpha}{\alpha - 3} \right) = 14.2530 \dots < 15.$$

Moreover we have, for any $0 \leq t \leq 1$ and $u_1, u_2 \in \mathbb{R}$,

$$|f(t, u_1) - f(t, u_2)| \leq \frac{3}{55} |u_1 - u_2|,$$

5 where

$$f(t, u) = \frac{3}{(54e^t + 1)(1 + |u|)}$$

for $0 \leq t \leq 1$ and $u \in \mathbb{R}$; see Section 4 in [2]. Since the constants $\lambda = \frac{3}{55}$ and $L = \frac{1}{100}$ in Theorem 1, we have

$$\lambda\Lambda + \frac{2L}{(\alpha-1)(\alpha-2)(\alpha-3)} \leq \frac{3}{55} \times 15 + \frac{2 \times \frac{1}{100}}{2.1 \times 1.1 \times 0.1} = 0.90476\bar{1} < 1.$$

It follows from Theorem 1 that the problem (3.2) has a unique solution.

Example 6. We also consider the following.

$$(3.3) \quad \begin{cases} D_{0+}^{3.1}u(t) = \frac{1}{50e^t(1+u^2)}, & 0 < t < 1, \\ u(0) = u'(0) = u''(1) = 0, \quad u'''(1) = g(u(1)), \end{cases}$$

where

$$g(t) = \frac{1}{100} \sin t.$$

For any $0 \leq t \leq 1$ and $u_1, u_2 \in \mathbb{R}$,

$$|f(t, u_1) - f(t, u_2)| \leq \frac{3}{50} |u_1 - u_2|,$$

where

$$f(t, u) = \frac{3}{50e^t(1+u^2)}$$

for $0 \leq t \leq 1$ and $u \in \mathbb{R}$. In this case we also have

$$\lambda\Lambda + \frac{2L}{(\alpha-1)(\alpha-2)(\alpha-3)} \leq \frac{3}{50} \times 15 + \frac{2 \times \frac{1}{100}}{2.1 \times 1.1 \times 0.1} \approx 0.9866 < 1.$$

REFERENCES

- [1] Z. Bai and H. Lü, Positive solutions for boundary value problem of nonlinear fractional differential equation, *J. Math. Anal. Appl.* **311**, No.2 (2005), 495–505.
- [2] M. Benchohra, N. Hamidi and J. Henderson, Fractional differential equations with anti-periodic boundary condition, *Numer. Funct. Anal. Optim.* **34**, No.4 (2013), 404?–414.
- [3] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, In North-Holland Mathematics Studies, 204, Elsevier Science B.V., Amsterdam, 2006.
- [4] T. F. Ma and J. da Silva, Iterative solutions for a beam equation with nonlinear boundary conditions of third order, *Appl. Math. Comput.* **159**, No.1 (2004), 11?–18.
- [5] M. Toyoda and T. Watanabe, Note on solutions of boundary value problems involving a fractional differential equation, *Linear Nonlinear Anal.* **3**, No.3 (2017), 449?–455.
- [6] M. Toyoda and T. Watanabe, Solutions for a fractional-order differential equation with boundary conditions of third order, *International Journal of Applied Mathematics* Volume **33** No. 3 (2020), 403–412.

(Toshikazu Watanabe) MEIJI UNIVERSITY

Email address: wa-toshi@mti.biglobe.ne.jp