

A framework for national scenarios with varying emission reductions

Shinichiro Fujimori^{1,2,3#}, Volker Krey³, Detlef van Vuuren^{4,5}, Ken Oshiro¹, Masahiro Sugiyama⁶, Puttipong Chunark⁷, Bundit Limmeechokchai⁷, Shivika Mittal^{8,9}, Osamu Nishiura¹, Chan Park¹⁰, Salony Rajbhandari⁷, Diego Silva Herran^{2,11}, Tran Thanh Tu¹², Shiya Zhao¹, Yuki Ochi¹³, Priyadarshi R. Shukla⁹, Toshihiko Masui², Phuong V.H. Nguyen¹⁴, Anique-Marie Cabardos³, Keywan Riahi^{3,15}

Author Affiliations

1. Department of Environmental Engineering, Kyoto University, C1-3 361, Kyotodaigaku Katsura, Nishikyoku, Kyoto city, Japan
2. Center for Social and Environmental Systems Research, National Institute for Environmental Studies (NIES), 16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan
3. International Institute for Applied System Analysis (IIASA), Schlossplatz 1, A-2361 Laxenburg, Austria
4. PBL Netherlands Environmental Assessment Agency, The Hague, Netherlands
5. Copernicus Institute for Sustainable Development, Utrecht University, Utrecht, the Netherlands
6. University of Tokyo, Tokyo, Japan
7. Sustainable Energy and Low Carbon Research Unit (SELC), Sirindhorn International Institute of Technology (SIIT), Thammasat University (TU), Thailand
8. Imperial College London, Grantham Institute—Climate Change and the Environment, London, UK
9. Global Center for Environment and Energy, Ahmedabad University, India
10. Department of Landscape Architecture, College of Urban Science, University of Seoul, University of Seoul, Korea
11. Institute for Global Environmental Strategies (IGES), Japan
12. Department of Environmental Engineering, International University - Vietnam National University Ho Chi Minh City, Vietnam
13. E-Konzal, Osaka, Japan
14. Faculty of Environment-Resources and Climate Change, Ho Chi Minh City University of Food Industry, Vietnam
15. Graz University of Technology, Graz, Austria

#: Corresponding author; Shinichiro Fujimori (fujimori.shinichiro.8a@kyoto-u.ac.jp)

Editor's summary

Currently there is no common structure to how national emissions scenarios are created, hindering efforts for comparison and analysis at the larger scale. This Perspective presents a framework to guide individual national scenario creation in a standardized way.

National-level climate actions will be vital for achieving global temperature goals in the coming decades. Near-term (2025-2030) plans are laid out in Nationally Determined Contributions (NDCs); the next step is submission of long-term strategies (LTS) for 2050. Currently, national scenarios underpinning LTSs are poorly coordinated and incompatible across countries, preventing assessment of individual nations' climate policy. Here we present a systematic and standardised, yet flexible, scenario framework varying 2050 emissions to build long-term national energy and climate mitigation scenarios. Applying the framework to six major Asian countries reveals individual challenges in energy system transformation and investment needs in comparable scenarios. This framework could be a starting point for comprehensive assessments as input to the global stocktake over the coming years.

53 **Main text**

54 **Introduction**

55 The Paris Agreement¹ defines a long-term temperature goal for international climate policy:
56 “holding the increase in the global average temperature to well below 2°C above pre-
57 industrial levels and pursuing efforts to limit the temperature increase to 1.5°C above pre-
58 industrial levels”. While this global goal defines the fundamental direction of international
59 climate policy, its achievement critically depends on national actions and policy making at
60 the national level. As part of the agreement, countries are required to submit Nationally
61 Determined Contributions (NDCs) outlining their greenhouse gas (GHG) emissions reduction
62 efforts at the national scale. Within the Paris Agreement, there are several mechanisms to
63 ensure that national actions align with the global goals; first, parties are required to regularly
64 report on their progress towards implementing their NDCs. These reports form part of the so-
65 called global stocktake, i.e., an assessment of the sum of the national contributions. Second,
66 countries are also required to submit long-term strategies (LTS) to the UNFCCC (also called
67 mid-century strategies). Some countries have already submitted them, and others are
68 preparing to do so (at September 2020). Third, the Paris Agreement contains provisions to
69 increase efforts over time, through what has been dubbed the “ratchet mechanism”. In
70 summary, one objective of the agreement is to have national actions aligned with long-term
71 goals, with routine checks and revisions of short- to medium-term national goals and policies.

72 Model-based climate and emissions scenarios are pivotal instruments for determining
73 whether proposed actions are in line with the long-term goals^{2,3}. In the fifth assessment
74 report of the IPCC (AR5), over a thousand scenarios were summarized in the database,
75 assessed and classified by both assumption and mitigation levels⁴. The special report on
76 1.5°C was also accompanied by a large set of global scenarios that depict emissions pathways
77 through the 21st century⁵. A large number of global model scenarios have been developed
78 under specific model inter-comparison projects (MIPs) by sharing scenario implementation
79 protocols that prescribe the characteristics of the scenarios (e.g. carbon budgets, technological
80 availability). This allows for a systematic assessment of a set of research questions and to
81 identify robust insights on climate change mitigation^{6,7,8,9}.

82 Similarly to the role of global emissions scenarios in international negotiations, national
83 scenarios have widely contributed to national policy making². In several countries and
84 regions, this is done by national modeling teams, but results are largely disseminated in
85 governmental reports or as internal information, and only occasionally shared in academic
86 papers^{10,11,12}. Taking Japan as an example, a task force was established to determine the
87 2020 emission target in 2008, and its recommendations were published in a book¹⁰ (only
88 available in Japanese) while for the NDCs submission, there was no official scenario
89 assessment. MIPs exist not only at the national level (for the US, China, Brazil and Japan^{13,14,}
90 ^{15,16}), but also for specific regions such as the EU¹⁷, Asia¹⁸, Latin America¹⁹. There have been
91 a few attempts to collect national scenarios such as in the CD-LINKS (Linking Climate and
92 Development Policies – Leveraging International Networks and Knowledge Sharing)^{20,21}
93 which also includes individual national scenarios^{22,23,24,25,26,27,28}, COMMIT (Climate Policy
94 assessment and Mitigation Modeling to Integrate national and global Transition pathways):
95 <https://themasites.pbl.nl/commit/>²⁹ and DDPP (The Deep Decarbonization Pathways Project)
96 ^{30,31} projects. Moreover, there have been various studies assessing national NDC implications
97 from sectoral perspectives^{32,33,34,35} to the broader SDGs context^{36,37,38,39}. It should also be
98 noted that many countries do not have publicly available national energy or emissions
99 scenarios.

100 Some major emitting countries rely on the scientific basis of existing national scenarios
101 for national climate policymaking^{36,40}, while many others do not. Furthermore, the emissions
102 reduction targets of national scenarios are either determined by their own countries’

103 interpretation of global goals (e.g. such as taking 2 °C consistent pathways and judging these
104 by themselves) or are derived from global scenarios such as those based on either cost-
105 optimal scenarios or effort sharing schemes⁴¹ (see left in Table 1). While recent efforts made
106 by national MIPs (e.g. CD-LINKS) have shared a scenario protocol across countries based
107 on global IAM results, these allow only an assessment of quite specific conditions (.e.g cost-
108 optimal and global uniform carbon price). Moreover, the modeling capability and the main
109 strategies of GHG emissions reduction can be diverse across nations. Consequently, the level
110 of emissions mitigation in national scenarios varies, which implies challenges for comparing
111 mitigation costs, and the degree of energy system changes across countries and scenarios.
112 Apart from scenarios, real national emissions targets for both the near-term and long-term
113 have often changed and will continue to do so in the future as well under various political and
114 social circumstances. If only scenarios under specific but limited emissions reduction targets
115 are available, the national scenarios are quickly outdated and become irrelevant (complexity
116 of national scenarios are discussed more in the section “**Complexity in the assessment of
117 national scenarios**”).

118 Given this current situation, what if there were a standardized scenario framework which
119 covers a wide range of emissions targets under the same reduction targets that are shared and
120 implemented by many countries? For example, suppose that there were publicly available
121 scenarios to reduce national emissions by 80% or 100% (not cumulative emissions) in 2050
122 for dozens of countries. What would be the benefits for national policy making of such a
123 scientific basis standardized scenario framework? There are at least four key benefits. One, it
124 would reveal the dynamics of each nation’s energy, land-use, and agricultural systems as well
125 as economic implications if the selected countries were to reduce emissions by similar levels.
126 For example, a Japanese energy model comparison study was conducted, which found that
127 even under the 80% reduction target in 2050, Japan would still have relatively high industrial
128 sector energy demands because of its large dependency on heavy industry and limited
129 renewable energy sources due to the small area of the country in comparison with the EU and
130 the United State¹⁴. This kind of assessment would become available on a broader scale. Two,
131 the transparent publication in scientific literature of the scenarios, the simulation models, and
132 how the scenarios were generated would contribute to ensuring that the scientific basis and
133 quality of models and scenarios are maintained, to some extent, although it happens more
134 frequently than before²¹. This is critical for evidence-informed policymaking. Three, this
135 might allow a direct comparison of the challenges that countries are facing in achieving
136 emissions reduction targets, which would be valuable when assessing the forthcoming
137 national long-term climate targets from the perspective of social transition. Finally, four, this
138 would allow climate policymakers to compare each country’s emissions targets and assess
139 whether their own national targets are compatible with other countries’ multiple reduction
140 targets possibilities or sufficient to reach the global long-term goals. Ultimately,
141 policymakers may want to update the national targets; these are already supposed to undergo
142 routine reviews as part of the global stocktake under the Paris Agreement. Although
143 individual nations would have their own interests and priorities and the standardized simple
144 scenario may not be completely sufficient to assess the national climate policies, such
145 scenarios could at least be an entry point to communicate with policymakers in many
146 countries. From there, each country could build their own specialized and customized
147 scenarios. We summarized the global and national modeling and scenarios circumstances in
148 Table 1.

149 Here, we present the issues with current national scenarios, propose a systematic and
150 standardized scenario framework, and demonstrate the implementation of such a framework
151 for a few selected countries. Our proposal can ultimately contribute to the establishment of a
152 central national scenario datahub for further national scenario assessments, similar to what

153 has already been done for global scenarios (see Table 1). Next, we discuss the complexity
 154 and expected criteria of national scenarios.

155
 156

Table 1 Summary of the characteristics of global and national scenarios

	Global scenarios	National scenarios
Producers	Integrated Assessment Models	National energy/Integrated Assessment Models
Main users of the research outcomes	IPCC, UNEP, UNFCCC, international and national policymakers	National policymakers, private companies, stakeholders and IPCC
Main study target	Global climate goals and associated implications for climate, energy, economy and land-use etc.	Individual national climate goals/targets and their implications for energy, economy, land-use, etc.
Scenario implementation	Individual studies or standardized modeling protocols implemented by multiple models	Some standardization in projects, but mostly specific and varied
Community organization	Well established as Integrated Assessment Modeling Consortium (IAMC)	Partially organized in different communities, often as part of a modeling framework (e.g., The Energy Technology Systems Analysis Program (ETSAP)), but also to an extent in IAMC

157
 158

Complexity in the assessment of national scenarios

160 For short- to medium-term perspectives, focusing on the next ten years, national policies
 161 and policy options as well as stakeholder interests are the primary concerns. In contrast, from
 162 a long-term perspective, a simple, comparable and systematic approach has clear benefits,
 163 facilitating a reassessment of the option space. It should be recognized that there are many
 164 determinants that are relevant for the specification of national emissions pathways, such as i)
 165 global climate targets in the context of international commitments; ii) how to select global
 166 pathways in line with global long-term goals (e.g. multi-IAMs uncertainty and physical
 167 climate science uncertainty); iii) selection of effort sharing schemes; iv) economic
 168 development stages in individual countries; v) other societal and development priorities that
 169 may be critical factors to determining the challenges of emissions reductions. The emissions
 170 reduction levels and challenges to achieving them naturally vary across countries and
 171 scenarios, and there is no need to have identical reduction levels across countries.
 172 Additionally, the current NDCs, which are based upon each nation's voluntary actions, are in
 173 many cases ambiguous, leading to significant uncertainty regarding the actual level of
 174 emission reduction targets⁴². This may be or may not be because nations would prefer to keep
 175 some flexibility in the interpretation of their target statements, resulting in a remarkable
 176 degree of flexibility significant enough to change long-term global implications⁴². Either way,
 177 this would imply that, in principle, it is inevitable to have some degree of uncertainty in the
 178 actual national targets, and we should eventually develop strategies to cope with such
 179 uncertainties. (See more explanation for each uncertainty in Supplementary Note.)

180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229

Expected criteria for upcoming national scenarios

Given the above-mentioned uncertainties, here we discuss the expected characteristics of the national scenarios, as listed below:

- Cross-national comparability
- Compatibility and cohesion with global climate goals
- Policy relevance
- Ability to address critical national target uncertainties
- Simple implementation without ambiguities in the interpretation of the modeling protocol

The comparability, which enables exploration of the comparative stringency of national targets, is particularly beneficial for assessing national scenarios. One possible way to achieve this is fixed reduction rates across countries (e.g., 80% reduction compared to a base year). The implications for the energy, land-use, and economic transitions can reveal the associated challenges. Regarding the cohesion with global emissions pathways, global emissions scenarios with the climate goal specifications (e.g., 2 and 1.5 °C) in conjunction with effort sharing assumptions⁴¹ can bridge national and global scenarios. There can be a large variation in national emissions pathways derived from the combination of effort sharing and global pathways, but we will show how our proposed framework in this paper can be easily mapped with global scenarios. For scenarios to be policy relevant, the emission reduction levels should not be far from the targets laid out in forthcoming national LTSs. Exploring multiple mitigation levels has the advantage of identifying potential ambiguities in forthcoming LTSs, as well as enabling sensitivity analyses around the eventual LTSs. For example, supposing that the LTS for a country does not specify the GHG coverage but declares a 50% reduction target in 2050, multiple scenarios would be mapped with full Kyoto gases cases and only CO₂ cases. A similar approach can be applied to other ambiguities. Finally, the simplicity of a modeling protocol that avoids ambiguities in its interpretation and the ease of implementing scenarios are of key importance to (i) allow such exercises to be performed in a decentralized manner, and (ii) keep the barrier for joining such an effort as low as possible. The simplicity would facilitate the updating of these scenario exercises on a regular basis, which will be discussed in the next section in more detail.

Proposal for a systematic national scenario framework

We, thus, propose a systematic and standardized approach for national scenarios that appropriately cover plausible future ranges of mitigation pathways and enables comparison across countries. Here we refer to this framework as “National Long-term Pathways” (NLPs), which comprises a set of national scenarios explained below. The role of NLPs could resemble Representative Concentration Pathways (RCPs)⁴³ in formulating the ambition and range of climate targets. As defined by the earlier study, we use the term scenario to describe a plausible, comprehensive, integrated and consistent description of how the future might unfold whereas the term pathway is used for a set of scenarios⁴⁴. This NLPs approach permits hedging against future national target uncertainties by not specifying a single emission reduction target, instead, exploring multiple systematic scenarios associated with percentages of emissions reductions in 2050, the commonly considered target year for LTS, as a default set.

We classify two kinds of scenarios. One is the so-called baseline, which excludes climate change mitigation policy but can include currently implemented and planned policies as implemented in earlier literature². Other socioeconomic assumptions are up to the individual modeler’s choice but are encouraged to be without unusual specific assumptions such as without CCS⁴⁵ and low energy demand⁴⁶. Although this modeler’s choice might

230 sometimes make the assessment and interpretation of the results difficult because
231 socioeconomic backgrounds can differ among countries, there is an advantage in being able
232 to skip a process to discuss what socioeconomic assumptions should be used and reach an
233 agreement. More importantly, globally standardized socioeconomic scenarios such as the
234 Shared Socioeconomic Pathways (SSPs) ⁴⁷ would not be best for individual countries and
235 thus, the selection and assumption of socioeconomic conditions would depend on each
236 country. If the national modellers cannot access national socioeconomic perspectives, the use
237 of globally standardized SSPs would be recommended. The second kind are climate scenarios
238 which target 10 to 100% of emissions reduction in 2050 compared to base year emissions,
239 with 10%-point increments covering the space between them. This can also be mapped with
240 intensity targets, such as carbon intensity with GDP assumptions. For 2030, NDC targets can
241 be adopted but these may have variations associated with conditional/unconditional targets.
242 Considering the current political situation, in which many countries are announcing carbon
243 neutrality targets for different years, which are not always 2050, our proposed emissions
244 pathways can be easily extrapolated linearly beyond 2050 and can be assessed from the
245 timing of zero emissions and the required transition towards that goal. If a model is unable to
246 get feasible solutions for specific scenarios because emissions reductions are too strict, this
247 information would also be reported. Energy-related CO₂ emissions are the default emissions
248 coverage. As we will discuss later, although there can be multiple options in the coverage of
249 species and sectors (e.g. full GHG, including land sector), we chose specific emissions as
250 defined above for two main reasons. First, energy-related CO₂ emissions are currently the
251 major source of emissions in most countries. Second, national modeling concerned with
252 climate change mitigation policy is, in many cases, initiated from energy modeling, and
253 considering developing countries whose modelling capability is relatively low, limiting the
254 scope of coverage would be effective for enhancing participation. Incorporation of other CO₂
255 and non-CO₂ emissions is not limited because they are critical elements that determine the
256 total GHG emissions. Additionally, it is important to design a holistic human system from the
257 energy, land-use, and economic perspectives. The reduction percentages are relative to the
258 specific base year (e.g., 2010) for which the national emissions inventory is available for
259 most countries and can thus exclude unnecessary uncertainties in the current NDCs. In this
260 way, the NLPs proposal meets the criteria stated above, with comparability across countries,
261 compatibility and cohesion with global climate goals, policy relevance and a relatively simple
262 implementation protocol, and a strategy to address uncertainties.

263 There should be flexibility in this proposal regarding at least the following two points.
264 First, there are several options for emissions gas coverage. Full Kyoto GHG would yield the
265 best coverage, but sectoral and gas coverages can vary. For the gas coverage, this could
266 include only CO₂ or three major GHGs (CO₂, N₂O and CH₄). The sectoral coverage would be
267 either full-sector or energy related emissions only. This coverage should be considered
268 depending on the availability of the information, composition of gases (e.g. Brazil could have
269 a large portion of emissions from land-use sector) and model capability for each country. For
270 non-CO₂ emissions, the Global Warming Potential (GWP) should be standardized and a
271 GWP100 metric should be used, as applied by UNFCCC and IPCC as the default choice in
272 their reporting. Second, the reduction levels can be changed depending on country. For
273 example, baseline emissions would not be increased for developed countries, while most
274 developing countries can have much higher emissions in the future than now and starting the
275 reduction percentage from 0% could still be deemed ambitious. For developed countries,
276 more granularity might be needed for the range of deep reductions, and thus 5%-point
277 increments between 70-100% could be also attractive. The base year can also be flexible if
278 needed (see further flexibility options in Supplementary Table 1).

279 We also propose to routinely and periodically run this systematic scenario framework. In
280 the global IAM community, there are series of almost routine-basis MIPs (e.g. Energy
281 Modeling Forum (EMF)), which now have a large influence on global climate policies. In
282 contrast, national scenarios are not yet so well-established and can derive much more benefit
283 from a scenario generation routine. There are multiple options for the routine intervals, such
284 as every five years, every IPCC assessment cycle, or international political milestones (e.g.,
285 every global stocktake). The pros and cons of these choices can be considered later, but here
286 we emphasize the advantages of having a regular scenario exercise under a similar protocol.
287 First, the research community would be able to routinely provide policy-relevant information,
288 tracing the model development history and tracking how the scenarios have changed over the
289 period. Second, these regular exercises would allow individual countries' researchers to
290 anticipate the forthcoming exercises and prepare a plan for model development as well as
291 take advantage of funding opportunities. In particular, this would be useful for developing
292 countries where the energy models/IAMs are not yet fully developed. Note that it might be
293 challenging to have completely harmonized protocols over time as political circumstances
294 change (e.g., NDC and its updates). The need for the routine exercise can also be extended to
295 the global integrated assessment modeling community; the climate modeling community has
296 such an experimental design, namely the Diagnostic, Evaluation and Characterization of
297 Klima (DECK), under the umbrella of so-called Coupled Model Intercomparison Project
298 (CMIP)⁴⁸.

299 Keeping the scenario protocol simple is important, which would enable modelers to
300 implement the scenarios in regular intervals. In the meantime, in theory, tens of scenario
301 variations depending on socioeconomic, technological availability/cost, and policy
302 assumptions could be developed. For example, SSPs in the global modeling community
303 allows us to explore the variation of future socioeconomic assumptions⁴⁷. Concerning
304 variations in technological availability and cost, there are well-known examples in the global
305 study carried out under EMF27 and AMPERE (Assessment of Climate Change Mitigation
306 Pathways and Evaluation of the Robustness of Mitigation Cost Estimates) projects^{45, 49} (e.g.
307 none carbon capture and storage (CCS) scenario). Furthermore we can see similar national or
308 regional implementations^{50, 51, 52}. These scenario variations can be added to the standard set as
309 supplementary (extended) scenarios which are similar as proposed in the SSPs⁵³.

310 Regarding the relationships with policymakers, there are at least three main roles. First,
311 for those countries that have not yet developed national scenarios, obviously, NLPs can
312 provide opportunities to generate national scenarios, which would create dialogue between
313 modellers and policymakers. Second, regardless of the existence of the national scenarios,
314 comparable multi-national scenarios can provide meaningful insights for each national
315 policymaker because national climate policy cannot be independent of the international
316 context. These two benefits are valid for both short-term and long-term. Third, while it would
317 be valuable to continue routine-based standardized scenario making, more customization of
318 the scenarios for each country might be needed in terms of socioeconomic assumptions and
319 some specific national interests in the long-term (e.g. no more nuclear power in Japan). The
320 role of NLPs would then become an entry point for shifting from the standardized and
321 systematic approach to creating such individual and unique national scenarios. Eventually,
322 NLPs would be a platform to maintain the national scenario modeling community which can
323 enhance a dialogue among modelers and policymakers similar to CMIP as mentioned earlier.
324

325 **Demonstration of the proposal scenario design**

326 To explore how this newly proposed scenario set can be used, we have implemented the
327 proposed framework in selected Asian countries which have a large diversity in economic
328 development stages, economy size and energy consumption patterns: China, India, Japan,


329 Korea, Thailand, and Vietnam. Each country individually runs national models, which means
330 that countries do not change international market conditions. For scenario quantifications, we
331 used AIM (Asia-Pacific Integrated Model) which has been extensively applied in global and
332 national climate change mitigation studies (see **Methods**).

333 We first focus on an assessment of a single country, in this case, Japan (Fig. 1). The
334 emissions in the baseline scenario (BaU) are almost unchanged throughout the period,
335 whereas climate mitigation scenarios, named CM30, CM40 to CM100, meet the NDC
336 emission reduction target of 26% in 2030 and hit incremental 10% reductions levels as
337 prescribed in the protocol in 2050 (Fig. 1a). Then, we compare projected emissions in 2050
338 with the global emissions pathways in conjunction with effort sharing schemes (Fig. 1bcd)
339 (see **Methods**). Since we consider the multi-scenario uncertainties of global IAMs emissions
340 pathways for 1.5°C and 2°C climate stabilization, at the national level there is a large range
341 of emissions levels associated with various effort-sharing schemes (Fig. 1d). Here we also
342 illustrate emissions target space with the long-term national goal for 2050, which in case of
343 Japan is an 80% reduction, but there is a range because the reference year and the GHG
344 coverage is unspecified.

345 Then, the energy system and economic implications for each emission reduction level
346 are presented which depend on the emissions reduction target levels (Fig. 1efg). For example,
347 the total energy supply is almost constant under a 30-60% reduction, while the scenario with
348 a 100% reduction in emissions implies a drop in supply by around half of the baseline. In
349 other words, beyond 60-70% of emission reductions a significant contribution of demand-
350 side measures, including both energy efficiency improvements and behavioral change, are
351 needed. Regarding the composition of energy sources, the contribution from low carbon
352 energy technologies sources such as CCS and renewable energy sources gradually increase as
353 reduction levels rise. Macro-economic costs of mitigation increase remarkably with more
354 ambitious targets (Fig. 1f) and could rise to 3%, 4%, and 4.5% of GDP losses with emission
355 reduction targets of 80, 90, and 100% in 2050, respectively. Carbon prices are much more
356 sensitive to reduction levels, increasing sharply to over 5,000\$/tCO₂ in a 100% reduction
357 scenario, and to around 2,000 and 1,000\$/tCO₂ in 90 and 80% reduction scenarios,
358 respectively. The carbon price would become extremely high under stringent reduction
359 targets, but this is due to the availability of negative emissions in Japan where only a small
360 area is left for energy crops and BECCS. Below target reductions of 60%, prices are lower
361 than 200\$/tCO₂ over the period. More indicators are presented in Supplementary Figure 1 for
362 Japan and Supplementary Figures 2 to 6 for other countries, and several basic trends in many
363 variables can be observed. There are gradual changes, with carbon price reductions in most
364 cases, but it should be noted that there are some variables and countries where convergences
365 are apparent. For example, carbon prices and GDP losses in India and Vietnam display a
366 trend that is due to the availability of CCS, including BECCS. Once CCS becomes widely
367 available, the carbon price is reduced. Final energy consumption in China, India and Vietnam
368 therefore decreases along with the increasing rates of carbon price reduction in the 2020s and
369 2030s, but then converges in the 2040s. These results are due to the enhancement of
370 electrification under mitigation⁵⁴, which offsets the energy efficiency improvements.

371 Applying the framework to a country that submitted a LTS, the scenario outcomes
372 could provide policymakers and analysts with an independent sensitivity around the LTS
373 which allows judgement on whether the targets are plausible or feasible from the energy and
374 economic perspectives. In addition, putting the LTS into the context of different equity
375 principles sheds some light on the fairness of the target. However, policymakers need to
376 interpret the results of model estimates carefully because they include uncertainties. The
377 socioeconomic conditions were prescribed as SSP2 in this case, but the implications would
378 change substantially if other conditions were assumed. Population and GDP are such key

379 socioeconomic drivers, but technological availability and national energy policies are also
 380 sources of uncertainty. For example, unavailability of CCS pushes the policy cost much
 381 higher than usual⁴⁵, whereas low energy demand substantially mitigates the cost⁴⁶. Finally,
 382 periodic reviews and assessments of the LTS, and the forthcoming 2035 or 2040 emissions
 383 reduction targets will provide opportunities to revise and update the goals.
 384
 385



386
 387 *Fig. 1 Illustrative example of the interpretation of the national scale long-term scenarios*
 388 *(NLPs) using Japan as a case study. Panel a shows national emissions pathways, b shows*
 389 *emissions global pathways, and c is a list of effort sharing schemes. d shows emissions in*
 390 *2050 considering global pathways (all available scenarios are considered), effort sharing*

391 *schemes and the long-term national goal of an 80% reduction, with an uncertainty range (red*
392 *shaded area) associated with an unspecified reference year and gas coverages. The dashed*
393 *red lines are emissions in 2010, and black dashed lines correspond to incremental 10%*
394 *reduction levels from base year emissions. e shows the energy system implications*
395 *represented by primary energy supply in 2050, while f,g indicate policy cost implications*
396 *represented by GDP loss rates and carbon prices.*


397

398 Regarding comparative assessments of multiple countries, in Fig. 2 we show selected
399 indicators, namely mean annual rate of energy intensity change (Fig. 2a) and carbon intensity
400 change (Fig. 2b), share of low carbon energy sources (Fig. 2c), electrification rates which is
401 electricity final energy consumption divided by total final energy consumption (Fig. 2d),
402 carbon price (Fig. 2e) and GDP loss rates (Fig. 2f). These indicators were chosen because
403 they are fundamentally critical variables for assessing climate change mitigation, and since
404 the scale of economy, energy consumption, and emissions of the countries assessed in this
405 demonstration vary substantially, indicators that take percentages or relative rather than
406 absolute values are more suitable for this analysis. We also carried out a regression analysis
407 to clarify the common characteristics and the extent to which the reduction target rates in
408 2050 would change each indicator with country dummy parameters, as shown in the
409 **Methods** section, and results are summarized in Supplementary Table 2 and Fig. 2.

410 We see a strong correlation with emissions reduction rates in most indicators except for
411 the mean annual rate of energy intensity change. The mean annual rate of carbon intensity
412 change indicates 0.025% improvements per incremental 1% of emissions reduction. In
413 contrast, the response of mean annual rate of energy intensity change to reduction levels
414 varies across countries, and the regressed slope is statistically insignificant. Japan's behavior,
415 in which energy intensity rises when increasing mitigation ambitions, is normal whereas
416 some other countries like India, China, and Vietnam appear to respond inversely. This is due
417 to the requirement for negative emissions associated with bioenergy combined with CCS
418 (BECCS). This result would imply that improvements in carbon intensity are a common and
419 effective strategy to reduce CO₂ emissions, while energy efficiency improvements do not
420 always yield the expected reduction in emissions. The share of low carbon energy sources
421 also shows a clear correlation with emissions reduction levels and a 0.56% increase is
422 expected per 1% of incremental emissions reduction. Electrification is a well-known and
423 critical strategy for decarbonizing the energy system, and 0.36% is the regressed slope for
424 change in electrification rates. Note that in Korea and Vietnam (see Supplementary Figure 4
425 and 6), the time series of electrification crosses over in 2030s. In the near-term, with modest
426 emissions constraints, the electricity generation cost increases, which lowers electricity
427 consumption while gas consumption increases. In the long term, under tighter emissions
428 constraints, electrification needs to be enhanced. Carbon prices vary substantially by country,
429 while the slope of regression is statistically significant at 12.50\$/tCO₂. Finally, the GDP loss
430 rates would increase by 0.055% per 1% of additional emissions reductions. GDP loss rates
431 also show variations across countries; Vietnam shows relatively high GDP loss rates, over
432 10%, while Japan presents small values, less than 5% even in a 100% emissions reduction
433 scenario. This variation comes from socioeconomic conditions such as the share of energy
434 and food expenditures which is largely influenced by abatement of non-CO₂ emissions from
435 agricultural sector and carbon tax imposition on them (if these are large, the relative influence
436 on industrial structures and household consumption patterns would be large) and GDP per
437 capita (if low, the carbon price intervention effects would be large) as well as assumptions on
438 the availability of technology. It could be argued that this regression analysis would be
439 affected by extreme country data. To test this, we conducted a sensitivity analysis to

440 determine the robustness by withdrawing one country from the regression and then iterating
 441 the results for all countries. The results indicated that the carbon price and some other
 442 indicators were affected by the Japanese data (see Supplementary figure 7).

443 Note that this study uses single model results. The use of multiple models, including
 444 multiple types of models (e.g. top-down and bottom-up, or CGE and energy system models)
 445 could lead to different results⁵⁵, which would enrich the implications of the study by
 446 introducing diversity in future prospects, and in particular, might not indicate the clear
 447 relationships shown here.
 448
 449



450
 451 *Fig. 2 Cross-national comparison of national long-term scenarios (NLPs). Six scenario*
 452 *indicators for 2050 are plotted against reduction targets. Panels a, b, c, d, e, and f represent*
 453 *mean annual rate of energy intensity change (%), mean annual rate of carbon intensity*
 454 *change (%), share of low carbon energy sources in primary energy supply (%),*
 455 *electrification rates in final energy consumption (%), carbon prices (\$/tCO₂) and GDP loss*
 456 *rates (%). The solid lines indicate regression results using the derived slope and intercept +*
 457 *mean of dummy country results shown in Supplementary Table 2.*

458
 459 **Caveats to the proposal and discussion**

460 We recognize that there are potential limitations to our proposal. First, policy relevance
 461 is the primary concern for this approach. This scenario set with its incremental 10% reduction
 462 levels might not exactly match the forthcoming LTS. As discussed, even if one of the
 463 scenario values of reduction rates hits a target, there will still be uncertainty in the inventory
 464 of the base year and coverage of GHGs. Second, there need to be several model runs (around
 465 10 or more). However, in contrast to existing large scale global models, national models tend
 466 to have relatively small computational loads, which could allow them to run relatively a large
 467 number of scenarios. In this sense, it is crucial to keep the simplicity of the scenario
 468 requirements as the simple scenario protocol allows researchers to systematically deal with
 469 scenarios running in the programming codes. To manage these issues, we view this proposal as
 470 a default core standard set, to which supplementary scenarios can be added, such as using

471 varying technological availability taking into account individual countries' circumstances⁴⁵.
472 Moreover, NDCs can be updated and ambitious LTSs may motivate countries to achieve
473 more reductions in the near-term, which would pose the question of whether more variations
474 should be added in near-term reduction targets. Although such scenarios are excluded from
475 this study, the updated NDC scenario could also be another set of supplementary scenarios. It
476 would be worth noting that such additional scenarios would have different roles from the
477 above-proposed scenario set, and requires additional work to check and maintain the quality
478 of results. Third, the protocol ignores possible interactions with the rest of the world.
479 Increasing ambitions in one country might go in hand with actions in other countries. This
480 could lead to impacts across countries. For example, fossil fuel prices could be low if many
481 major countries decarbonized their economies. International price scenarios derived from
482 global IAMs could be used as boundary conditions for national models, and in such a case,
483 global models should also provide multi-level mitigation scenarios which could be prescribed
484 by carbon budgets⁵. Still, the most direct impacts of more ambitious targets are nearly always
485 felt simply within each country – and thus should serve as a caveat in the light of proposed
486 simplification. Future study will be needed to investigate cross-border impacts. Fourth, the
487 proposed scenarios always come with the risk of being outdated at some point, which can be
488 critical in some cases. For example, long-term strategies were supposed to be submitted by
489 the year 2020, and our proposal may not be able to keep up with them. Another possibility is
490 that some extreme economic, social and political events may completely change the relevant
491 energy-economic system. The the disaster at the Fukushima nuclear power plant was one
492 such turning point, and the COVID-19 pandemic has the potential to be another one. A
493 financial crisis, in general, could also result in structural change, which may imply that
494 additional scenarios may be needed to take these extreme (or simply outlier) events into
495 account. However, this depends on individual events and national circumstances. It may not
496 be able to generalize and will probably need to generate specific scenarios to address such
497 events.

498 Finally, the current proposal can be a first step to have systematic national scenarios,
499 much as global scenarios are currently stored and utilized effectively. Meanwhile, even if the
500 scenarios are developed by many countries, building up a valuable database, there would be
501 still the need for better communication with policymakers. This is obvious from global IAM
502 exercises. Even though there have been efforts to create transparent models⁵⁶ and
503 socioeconomic assumptions behind scenarios⁵⁷, as well as making code open-source^{58, 59}
504 consistent with the recent demand for transparency, there is still an increasing demand to
505 explain scenarios to decision makers. Furthermore, the misinterpretation of current scenarios
506 is an ongoing problem, for example, in the lack of climate change impacts⁶⁰. Therefore, just
507 developing national scenarios is not sufficient, and better translation and communication of
508 the scenarios to the policymakers is still needed.

509

510 **Community and capacity development**

511 The development of national scenarios fundamentally needs the involvement of
512 researchers from each country. Many countries, including developing countries, have national
513 models, but there are also many countries still missing national energy or integrated
514 assessment models. Even if national models exist, a certain portion of models need to
515 improve their systematic model output reporting, model validation (including diagnostics and
516 documentation), and will require significant work to reach state-of-the-art modeling
517 representation. In many cases, global integrated assessment modeling activities and
518 experiences accumulated in the Integrated Assessment Modeling Consortium (IAMC)⁶¹
519 community should greatly help national modeling capacity development^{56, 59, 62, 63, 64, 65}.

520 Note that global models are themselves not always the best; some national models have much

521 more granularity in the representation of geographical and temporal resolutions, taking
522 advantage of relatively smaller model coverage^{55, 66}. IAMC members have been actively
523 involved in capacity development (e.g., for Asian⁶⁷ and Latin American¹⁵ capacity building
524 activities, National Institute for Environmental Studies Japan (NIES) and Pacific Northwest
525 National Laboratory USA (PNNL) have taken part in some exercises) and the IAMC itself
526 sometimes coordinating them so far. However, this proposed standardized scenario exercise
527 can be a more meaningful and practical catalyst for enhancing capacity building activities
528 within the climate mitigation modeling community.

529
530

531 **Conclusions**

532 In this Perspective, we propose a new systematic and standardized scenario framework for
533 long-term national scenarios and discuss its rationale, the advantages, and possible
534 disadvantages. We believe that this proposal is valid and useful for policymaking and
535 building a research community. National climate change mitigation modelling and scenario
536 implementations might inherently have had relatively little motivation for building up a
537 research community and conducting cross-national comparisons in the past. However, the
538 political and societal conditions have changed over the last decade, and we believe that
539 national countermeasures are now a necessity for combatting climate change. The climate
540 policy circumstances and the need for national modeling and scenarios are expected to
541 continue for at least the next couple of decades until emissions drop to sufficiently low levels.
542 This research community should, therefore, devote much more attention and resources to
543 national scenarios that guide or enhance the actual societal transformative movement. We
544 envisage that the proposed framework could be a great milestone for national climate policy
545 research and many countries and models would engage with it. Thus, we call for community-
546 level activities that will let a wide range of researchers involved in national climate policy
547 assessment consider dedicating efforts to these important new activities.

548

549 **References**

550

- 551 1. United Nations Framework Convention on Climate Change, (UNFCCC). Adoption of the Paris
552 Agreement. Proposal by the President (1/CP.21). 2015 [cited 2016 02, Feb.] Available from:
553 <http://unfccc.int/resource/docs/2015/cop21/eng/10a01.pdf>
- 554 2. Roelfsema M, van Soest HL, Harmsen M, van Vuuren DP, Bertram C, den Elzen M, *et al.* Taking
555 stock of national climate policies to evaluate implementation of the Paris Agreement. *Nature*
556 *Communications* 2020, **11**(1): 2096.
- 557 3. United Nations Environment Programme, (UNEP). *Emissions Gap Report 2018*. UNEP, 2019.
- 558 4. Clarke L, Jiang K, Akimoto K, Babiker M, Blanford G, Fisher-Vanden K, *et al.* Assessing
559 Transformation Pathways. In: Edenhofer O, Pichs-Madruga R, Sokona Y, Farahani E, Kadner S,
560 Seyboth K, *et al.* (eds). *Climate Change 2014: Mitigation of Climate Change. Contribution of Working*
561 *Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*.
562 Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2014, pp 413-
563 510.
- 564 5. Fujimori S, Rogelj J, Krey V, Riahi K. A new generation of emissions scenarios should cover blind
565 spots in the carbon budget space. *Nature Climate Change* 2019, **9**(11): 798-800.
- 566 6. Bauer N, Rose SK, Fujimori S, van Vuuren DP, Weyant J, Wise M, *et al.* Global energy sector
567 emission reductions and bioenergy use: overview of the bioenergy demand phase of the EMF-33 model
568 comparison. *Climatic Change* 2018.
- 569 7. Luderer G, Vrontisi Z, Bertram C, Edelenbosch OY, Pietzcker RC, Rogelj J, *et al.* Residual fossil CO2
570 emissions in 1.5–2 °C pathways. *Nature Climate Change* 2018, **8**(7): 626-633.
- 571 8. Fujimori S, Hasegawa T, Krey V, Riahi K, Bertram C, Bodirsky BL, *et al.* A multi-model assessment
572 of food security implications of climate change mitigation. *Nature Sustainability* 2019, **2**(5): 386-396.

- 573 9. McCollum DL, Zhou W, Bertram C, de Boer H-S, Bosetti V, Busch S, *et al.* Energy investment needs
574 for fulfilling the Paris Agreement and achieving the Sustainable Development Goals. *Nature Energy*
575 2018, **3**(7): 589-599.
- 576 10. Fukui T. *Explanation of the mid-term target for global warming countermeasures (Chikyū Ondanka*
577 *Taisaku Chūki Mokuhyō no Kaisetsu)*. Gyosei, 2009.
- 578 11. Commission E-E. A Clean Planet for all—A European long-term strategic vision for a prosperous,
579 modern, competitive and climate neutral economy. *depth analysis in support of the commission;*
580 *Communication COM (2018) 2018*, **773**: 2018.
- 581 12. The White House. United States mid-century strategy for deep decarbonization. *United Nations*
582 *Framework Convention on Climate Change, Washington, DC*; 2016.
- 583 13. Fawcett AA, Clarke LE, Weyant JP. Introduction to EMF 24. *The Energy Journal* 2014, **35**: 1-8.
- 584 14. Sugiyama M, Fujimori S, Wada K, Endo S, Fujii Y, Komiyama R, *et al.* Japan's long-term climate
585 mitigation policy: Multi-model assessment and sectoral challenges. *Energy* 2019, **167**: 1120-1131.
- 586 15. Lugovoy O, Feng X-Z, Gao J, Li J-F, Liu Q, Teng F, *et al.* Multi-model comparison of CO2 emissions
587 peaking in China: Lessons from CEMF01 study. *Advances in Climate Change Research* 2018, **9**(1): 1-
588 15.
- 589 16. Lucena AFP, Clarke L, Schaeffer R, Szklo A, Rochedo PRR, Nogueira LPP, *et al.* Climate policy
590 scenarios in Brazil: A multi-model comparison for energy. *Energy Economics* 2016, **56**: 564-574.
- 591 17. Weyant J, Knopf B, De Cian E, Keppo I, van Vuuren DP. Introduction to the Emf28 Study on
592 Scenarios for Transforming the European Energy System. *Climate Change Economics* 2013,
593 **04**(supp01): 1302001.
- 594 18. Calvin K, Clarke L, Krey V, Blanford G, Jiang K, Kainuma M, *et al.* The role of Asia in mitigating
595 climate change: Results from the Asia modeling exercise. *Energy Economics* 2012, **34**: S251-S260.
- 596 19. van der Zwaan BCC, Calvin KV, Clarke LE. Climate Mitigation in Latin America: Implications for
597 Energy and Land Use: Preface to the Special Section on the findings of the CLIMACAP-LAMP
598 project. *Energy Economics* 2016, **56**: 495-498.
- 599 20. Schaeffer R, Köberle A, van Soest HL, Bertram C, Luderer G, Riahi K, *et al.* Comparing
600 transformation pathways across major economies. *Climatic Change* 2020, **162**(4): 1787-1803.
- 601 21. Schaeffer R, Bosetti V, Kriegler E, Riahi K, van Vuuren D. Climatic change: CD-Links special issue
602 on national low-carbon development pathways. *Climatic Change* 2020, **162**(4): 1779-1785.
- 603 22. Köberle AC, Rochedo PRR, Lucena AFP, Szklo A, Schaeffer R. Brazil's emission trajectories in a
604 well-below 2 °C world: the role of disruptive technologies versus land-based mitigation in an already
605 low-emission energy system. *Climatic Change* 2020, **162**(4): 1823-1842.
- 606 23. Wang H, Chen W, Zhang H, Li N. Modeling of power sector decarbonization in China: comparisons of
607 early and delayed mitigation towards 2-degree target. *Climatic Change* 2020, **162**(4): 1843-1856.
- 608 24. Vishwanathan SS, Garg A. Energy system transformation to meet NDC, 2 °C, and well below 2 °C
609 targets for India. *Climatic Change* 2020, **162**(4): 1877-1891.
- 610 25. Mathur R, Shekhar S. India's energy sector choices—options and implications of ambitious mitigation
611 efforts. *Climatic Change* 2020, **162**(4): 1893-1911.
- 612 26. Feijoo F, Iyer G, Binsted M, Edmonds J. US energy system transitions under cumulative emissions
613 budgets. *Climatic Change* 2020, **162**(4): 1947-1963.
- 614 27. Safonov G, Potashnikov V, Lugovoy O, Safonov M, Dorina A, Bolotov A. The low carbon
615 development options for Russia. *Climatic Change* 2020, **162**(4): 1929-1945.
- 616 28. Oshiro K, Gi K, Fujimori S, van Soest HL, Bertram C, Després J, *et al.* Mid-century emission
617 pathways in Japan associated with the global 2 °C goal: national and global models' assessments based
618 on carbon budgets. *Climatic Change* 2019, **162**(4): 1913-1927.
- 619 29. Fragkos P, Laura van Soest H, Schaeffer R, Reedman L, Köberle AC, Macaluso N, *et al.* Energy
620 system transitions and low-carbon pathways in Australia, Brazil, Canada, China, EU-28, India,
621 Indonesia, Japan, Republic of Korea, Russia and the United States. *Energy* 2021, **216**: 119385.
- 622 30. Waisman H, Bataille C, Winkler H, Jotzo F, Shukla P, Colombier M, *et al.* A pathway design
623 framework for national low greenhouse gas emission development strategies. *Nature Climate Change*
624 2019, **9**(4): 261-268.
- 625 31. Bataille C, Waisman H, Colombier M, Segafredo L, Williams J. The Deep Decarbonization Pathways
626 Project (DDPP): insights and emerging issues. *Climate Policy* 2016, **16**(sup1): S1-S6.
- 627 32. Islas-Samperio JM, Manzini F, Grande-Acosta GK. Toward a Low-Carbon Transport Sector in
628 Mexico. *Energies* 2020, **13**(1): 84.
- 629 33. Fyson CL, Jeffery ML. Ambiguity in the Land Use Component of Mitigation Contributions Toward the
630 Paris Agreement Goals. *Earth's Future* 2019, **7**(8): 873-891.

- 631 34. Maraseni TN, Poudyal BH, Rana E, Chandra Khanal S, Ghimire PL, Subedi BP. Mapping national
632 REDD+ initiatives in the Asia-Pacific region. *Journal of environmental management* 2020, **269**:
633 110763.
- 634 35. Goes GV, Schmitz Gonçalves DN, de Almeida D'Agosto M, de Mello Bandeira RA, Grottera C.
635 Transport-energy-environment modeling and investment requirements from Brazilian commitments.
636 *Renewable Energy* 2020, **157**: 303-311.
- 637 36. Iyer G, Ledna C, Clarke L, Edmonds J, McJeon H, Kyle P, *et al.* Measuring progress from nationally
638 determined contributions to mid-century strategies. *Nature Climate Change* 2017, **7**(12): 871-874.
- 639 37. Liu J-Y, Fujimori S, Takahashi K, Hasegawa T, Wu W, Takakura Jy, *et al.* Identifying trade-offs and
640 co-benefits of climate policies in China to align policies with SDGs and achieve the 2 °C goal.
641 *Environmental Research Letters* 2019, **14**(12): 124070.
- 642 38. Dal Maso M, Olsen KH, Dong Y, Pedersen MB, Hauschild MZ. Sustainable development impacts of
643 nationally determined contributions: assessing the case of mini-grids in Kenya. *Climate Policy* 2020,
644 **20**(7): 815-831.
- 645 39. Nogueira LP, Longa FD, van der Zwaan B. A cross-sectoral integrated assessment of alternatives for
646 climate mitigation in Madagascar. *Climate Policy* 2020, **20**(10): 1257-1273.
- 647 40. Weitzel M, Vandyck T, Keramidas K, Amann M, Capros P, den Elzen M, *et al.* Model-based
648 assessments for long-term climate strategies. *Nature Climate Change* 2019, **9**(5): 345-347.
- 649 41. van den Berg NJ, van Soest HL, Hof AF, den Elzen MGJ, van Vuuren DP, Chen W, *et al.* Implications
650 of various effort-sharing approaches for national carbon budgets and emission pathways. *Climatic*
651 *Change* 2019, **162**(4): 1805-1822.
- 652 42. Rogelj J, Fricko O, Meinshausen M, Krey V, Zilliacus JJJ, Riahi K. Understanding the origin of Paris
653 Agreement emission uncertainties. *Nature Communications* 2017, **8**: 15748.
- 654 43. van Vuuren DP, Edmonds JA, Kainuma M, Riahi K, Weyant J. A special issue on the RCPs. *Climatic*
655 *Change* 2011, **109**(1-2): 1-4.
- 656 44. van Vuuren DP, Kriegler E, O'Neill BC, Ebi KL, Riahi K, Carter TR, *et al.* A new scenario framework
657 for Climate Change Research: scenario matrix architecture. *Climatic Change* 2014, **122**(3): 373-386.
- 658 45. Kriegler E, Weyant JP, Blanford GJ, Krey V, Clarke L, Edmonds J, *et al.* The role of technology for
659 achieving climate policy objectives: overview of the EMF 27 study on global technology and climate
660 policy strategies. *Climatic Change* 2014, **123**(3-4): 353-367.
- 661 46. Grubler A, Wilson C, Bento N, Boza-Kiss B, Krey V, McCollum DL, *et al.* A low energy demand
662 scenario for meeting the 1.5 °C target and sustainable development goals without negative emission
663 technologies. *Nature Energy* 2018, **3**(6): 515-527.
- 664 47. Riahi K, Kriegler E, Johnson N, Bertram C, den Elzen M, Eom J, *et al.* Locked into Copenhagen
665 pledges — Implications of short-term emission targets for the cost and feasibility of long-term climate
666 goals. *Technological Forecasting and Social Change* 2015, **90**: 8-23.
- 667 48. Eyring V, Righi M, Lauer A, Evaldsson M, Wenzel S, Jones C, *et al.* ESMValTool (v1.0) – a
668 community diagnostic and performance metrics tool for routine evaluation of Earth system models in
669 CMIP. *Geosci Model Dev* 2016, **9**(5): 1747-1802.
- 670 49. Riahi K, Kriegler E, Johnson N, Bertram C, den Elzen M, Eom J, *et al.* Locked into Copenhagen
671 pledges — Implications of short-term emission targets for the cost and feasibility of long-term climate
672 goals. *Technological Forecasting and Social Change* 2013.
- 673 50. Silva Herran D, Fujimori S, Kainuma M. Implications of Japan's long term climate mitigation target
674 and the relevance of uncertain nuclear policy. *Climate Policy* 2019, **19**(9): 1117-1131.
- 675 51. Clarke LE, Fawcett AA, Weyant JP, McFarland J, Chaturvedi V, Zhou Y. Technology and US
676 emissions reductions goals: Results of the EMF 24 modeling exercise. *The Energy Journal* 2014,
677 **35**(Special Issue).
- 678 52. Knopf B, Chen Y-HH, De Cian E, FÖRster H, Kanudia A, Karkatsouli I, *et al.* Beyond 2020 —
679 Strategies and Costs for Transforming the European Energy System. *Climate Change Economics* 2013,
680 **04**(supp01): 1340001.
- 681 53. O'Neill B, Kriegler E, Riahi K, Ebi K, Hallegatte S, Carter T, *et al.* A new scenario framework for
682 climate change research: the concept of shared socioeconomic pathways. *Climatic Change* 2014,
683 **122**(3): 387-400.
- 684 54. Sugiyama M. Climate change mitigation and electrification. *Energy Policy* 2012, **44**: 464-468.
- 685 55. Fujimori S, Oshiro K, Shiraki H, Hasegawa T. Energy transformation cost for the Japanese mid-
686 century strategy. *Nature Communications* 2019, **10**(1): 4737.
- 687 56. IAMC. The common Integrated Assessment Model (IAM) documentation. 2018 [cited]Available
688 from: https://www.iamcdocumentation.eu/index.php/IAMC_wiki

- 689 57. Riahi K, van Vuuren DP, Kriegler E, Edmonds J, O'Neill BC, Fujimori S, *et al.* The Shared
690 Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An
691 overview. *Global Environmental Change-Human and Policy Dimensions* 2017, **42**: 153-168.
- 692 58. Huppmann D, Gidden M, Fricko O, Kolp P, Orthofer C, Pimmer M, *et al.* The MESSAGEix Integrated
693 Assessment Model and the ix modeling platform (ixmp): An open framework for integrated and cross-
694 cutting analysis of energy, climate, the environment, and sustainable development. *Environmental*
695 *Modelling & Software* 2019, **112**: 143-156.
- 696 59. Calvin K, Patel P, Clarke L, Asrar G, Bond-Lamberty B, Cui RY, *et al.* GCAM v5.1: representing the
697 linkages between energy, water, land, climate, and economic systems. *Geosci Model Dev* 2019, **12**(2):
698 677-698.
- 699 60. Grant N, Hawkes A, Napp T, Gambhir A. The appropriate use of reference scenarios in mitigation
700 analysis. *Nature Climate Change* 2020, **10**(7): 605-610.
- 701 61. IAMC. IAMC. webpage [cited 2015 1. April]Available from:
702 <http://www.globalchange.umd.edu/iamc/scientific-working-groups/evaluation-and-diagnostics/>
- 703 62. Kriegler E, Petermann N, Krey V, Schwanitz VJ, Luderer G, Ashina S, *et al.* Diagnostic indicators for
704 integrated assessment models of climate policy. *Technological Forecasting and Social Change* 2014.
- 705 63. Emmerling J, Drouet L, Aleluia Reis L, Bevione M, Berger L, Bosetti V, *et al.* The WITCH 2016
706 Model - Documentation and Implementation of the Shared Socioeconomic Pathways. FEEM Nota di
707 Lavoro 42.2016. Milano, Italy: FEEM; 2016.
- 708 64. Fujimori S, Dai H, Masui T, Matsuoka Y. Global energy model hindcasting. *Energy* 2016, **114**: 293-
709 301.
- 710 65. Calvin K, Wise M, Kyle P, Clarke L, Edmonds J. A hindcast experiment using the gcam 3.0 agriculture
711 and land-use module. *Climate Change Economics* 2017, **08**(01): 1750005.
- 712 66. Feijoo F, Iyer GC, Avraam C, Siddiqui SA, Clarke LE, Sankaranarayanan S, *et al.* The future of natural
713 gas infrastructure development in the United states. *Applied Energy* 2018, **228**: 149-166.
- 714 67. Fujimori S, Kainuma M, Masui T. *Post-2020 Climate Action: Global and Asian Perspective*. Springer,
715 2017.
- 716
717
718

719 **Methods**

720 **Overview**

721 We carried out scenario analysis for selected Asian countries, namely China, India,
722 Indonesia, Japan, Korea, Thailand, and Vietnam. As stated in the main text, we implemented
723 8 scenarios for each country. They all have different reduction rates relative to the base year
724 of 2010, and, furthermore, Vietnam and Thailand indicated conditional and unconditional
725 target statements in their NDCs. We thus additionally simulated variations for these
726 conditional statements. In scenario implementations, we considered currently planned
727 national policies as much as possible. We used AIM/Hub (formerly AIM/CGE) for the
728 proposed scenario design implementation and as the core tool of this study. It is a computable
729 general equilibrium model and has been intensively applied to assessments of Asian national
730 climate policies in past years^{68, 69, 70, 71, 72}. In the scenario implementations, we used three
731 major GHG gases (CO₂, CH₄ and N₂O) for emission coverage, considering that the countries
732 have relatively large emissions of non-CO₂ gases. The reductions start from 30% in all
733 countries because we took into account that Japan's baseline emissions have been quite stable
734 over time and thus it may not be meaningful to see the lower reduction levels such as 10%.
735 Finally, a regression analysis has been conducted on the scenario results. Note that the
736 scenarios in this study excluded climate change impacts, because global emissions scenarios
737 are needed for each national emission scenario to determine such impacts, which is an
738 important factor for national policymakers to consider^{60, 73, 74, 75}.

739

740 **Simulation model and data**

741 AIM/Hub is a one-year-step recursive-type dynamic general equilibrium model covering all
742 regions of the world. The AIM/Hub model includes 42 industrial classifications. For
743 assessing bioenergy and land use competition, agricultural sectors are disaggregated⁷⁶. The
744 details of the model structure and mathematical formulae have been described previously^{77, 78}.
745 Version 2.2 of the AIM/Hub model was used, and the main revisions from the previous
746 version are described below.

747 Production sectors are assumed to maximise profits using multi-nested constant
748 elasticity substitution (CES) functions and input prices. For energy transformation sectors, to
749 handle energy conversion efficiency appropriately in these sectors, input energy and value
750 added are fixed coefficients of the output. Power generation values from several energy
751 sources are combined with a logit function⁷⁹. This functional form was used to ensure energy
752 balance, as it was not guaranteed by the CES function. Electricity and bioenergy are produced
753 by multiple sectors (e.g. coal-fired, nuclear and solar, agricultural residue, energy crops and
754 sugarcane), which are aggregated by the logit function so that energy production by
755 individual sectors is balanced to match total generation. Household expenditures on each
756 commodity are described with a linear expenditure system (LES) function. The parameters
757 adopted in the LES function are recursively updated in accordance with income elasticity
758 assumptions. The savings ratio is endogenously determined to balance savings and
759 investment, and capital formation for each good is assigned a fixed coefficient as an
760 exogenous assumption. The Armington assumption is used for trade (using CES and the
761 constant elasticity of transformation function), and the current account is assumed to be
762 balanced.

763 In addition to energy-related CO₂, CO₂ from other sources, CH₄, N₂O, and fluorinated
764 gases (F-gases) are treated as GHGs in the model. Energy-related emissions are associated
765 with fossil fuel combustion. Non-energy-related CO₂ emissions consist of changes in land-
766 use and industrial processes. Emissions from changes in land-use are derived from the change
767 in forest area relative to the previous year multiplied by the carbon stock density, which is
768 differentiated into AEZs (Agro-Ecological Zones). Non-energy-related emissions other than

769 those associated with changes in land-use are assumed to be proportional to the level of each
770 activity (e.g. based on output). CH₄ emissions arise from a range of sources, mainly rice
771 production, livestock, fossil fuel mining, and waste management. N₂O is emitted as a result of
772 fertiliser application and livestock manure management, as well as by the chemical industry.
773 F-gases are emitted mainly from refrigerants used in air conditioners and industrial cooling
774 devices. Air pollutant gases (black carbon, CO, NH₃, non-methane volatile organic
775 compounds, NO_x, organic carbon, and SO₂) are also associated with fuel combustion and
776 activity levels. Emission factors change over time with the implementation of air pollutant
777 removal technologies and other regulations⁸⁰.

778 The implementation of mitigation actions in the model is represented by constraints on
779 CO₂ emissions. The carbon price is imposed on CO₂ as well as other GHG types, such as CH₄
780 and N₂O, arising from every sector. The carbon price increases the price of fossil fuel-based
781 goods when emissions are constrained and promotes energy savings and substitution away
782 from fossil fuels to sources and transport methods with lower GHG emissions. The carbon
783 tax also functions as an incentive to reduce non-energy-related emissions. Gases other than
784 CO₂ are weighted based on their global warming potential and summed as total GHG
785 emissions. Further parameter settings and changes under the future scenarios are documented
786 in Fujimori et al. (2017)⁸¹.

787 The main revisions from version 2.0, which was used in SSP quantification⁷², to version
788 2.2 are described in Fujimori et al. (2020)⁸² and the most relevant one for this study is the
789 reflection of historical energy data (2005 to 2015). This methodology is the same as model
790 integration with an energy system model where we exogenously provide the final energy,
791 transport energy share and power energy technological share, while the corresponding
792 parameters in the production function and household consumption are endogenized.
793 Consequently, the autonomous energy efficiency in energy consumption and logit share
794 parameters used to determine the share of power generation by different technologies were
795 calibrated during that period and then used for the future scenarios (for more methodological
796 details, see Fujimori et al.⁵⁵). We used the IEA Energy Balances as the historical energy
797 information⁸³.

798

799 **National policies**

800 We adopted current national policies that can be considered relevant for the scenarios as
801 much as possible. The NDCs are all taken into account as emissions constraints for the year
802 2030. For all countries, population and GDP projections are based on the national perspective
803 until either 2030 or 2035. Rates of SSP2 annual change are extrapolated afterward. There are
804 some vital energy and climate mitigation-related policies at national levels which are
805 reflected as either model constraints or as reference information to serve as a check that the
806 scenarios are not far from the corresponding national perspectives. For example, in China, the
807 next five-year plan, to be implemented in 2021, is scheduled to be published in late 2020 to
808 early 2021 and thus we decided not to use the latest available five-year plan but have
809 incorporated the best available current energy information. Another example is Thailand,
810 where the power development plan has been established by the Ministry of Environment^{84, 85,}
811 ⁸⁶ as is being used for model constraints. The full list of national policy information
812 considered in this study is shown in Supplementary Table 3.

813

814 **Effort sharing**

815 To map the national scenarios with global goals, we used multiple effort sharing schemes
816 shown by van den Berg *et al.*⁴¹. For the global scenarios, we adopted the latest global
817 scenarios from the IPCC Special Report on 1.5 °C database⁸⁷ by taking minimum, median,
818 and maximum ranges of IAMs pathways categorized as 1.5°C or 1.5°C-consistent and 2°C or

819 2°C-consistent for 1.5 and 2 °C goals, respectively, regardless of the scale of global mean
820 temperature overshoot.

821

822 **Regression analysis of the scenario indicators**

823 A regression analysis was carried out for the cross-country comparative assessment. The aim
824 of this regression is to derive the general relationships which can be observed in multiple
825 countries between each indicator and reduction levels. The equation applied is shown below.

$$Y_{r,s} = aX_{r,s} + b_r + c + \epsilon$$

826 Where:

827 $Y_{r,s}$ is an individual six indicators (annual mean rate of energy intensity change, annual rate of
828 mean carbon intensity change, share of low carbon energy sources, electrification rate, carbon
829 price and GDP loss rates) in country r and scenario s , $X_{r,s}$ is the emissions reduction
830 percentage relative to those of 2010. a , b_r and c represent estimated parameters and they are
831 the slope of the reduction levels, dummy countries, and intercept respectively. ϵ is an error
832 term.

833

834 All figures in this paper is generated by the code at a Github repository⁸⁸.

835

836 **Methods references**

837

- 838 41. van den Berg NJ, van Soest HL, Hof AF, den Elzen MGJ, van Vuuren DP, Chen W, *et al.* Implications
839 of various effort-sharing approaches for national carbon budgets and emission pathways. *Climatic*
840 *Change* 2019, **162**(4): 1805-1822.
- 841 55. Fujimori S, Oshiro K, Shiraki H, Hasegawa T. Energy transformation cost for the Japanese mid-
842 century strategy. *Nature Communications* 2019, **10**(1): 4737.
- 843 60. Grant N, Hawkes A, Napp T, Gambhir A. The appropriate use of reference scenarios in mitigation
844 analysis. *Nature Climate Change* 2020, **10**(7): 605-610.
- 845 68. Fujimori S, Kainuma M, Masui T, Hasegawa T, Dai H. The effectiveness of energy service demand
846 reduction: A scenario analysis of global climate change mitigation. *Energy Policy* 2014,
847 **75**(Supplement C): 379-391.
- 848 69. Fujimori S, Masui T, Matsuoka Y. Gains from emission trading under multiple stabilization targets and
849 technological constraints. *Energy Economics* 2015, **48**: 306-315.
- 850 70. Thepkhun P, Limmeechokchai B, Fujimori S, Masui T, Shrestha RM. Thailand's Low-Carbon Scenario
851 2050: The AIM/CGE analyses of CO2 mitigation measures. *Energy Policy* 2013, **62**(0): 561-572.
- 852 71. Hasegawa T, Fujimori S, Shin Y, Tanaka A, Takahashi K, Masui T. Consequence of Climate
853 Mitigation on the Risk of Hunger. *Environmental science & technology* 2015, **49**(12): 7245-7253.
- 854 72. Fujimori S, Hasegawa T, Masui T, Takahashi K, Herran DS, Dai H, *et al.* SSP3: AIM implementation
855 of Shared Socioeconomic Pathways. *Global Environmental Change-Human and Policy Dimensions*
856 2017, **42**: 268-283.
- 857 73. Burke M, Hsiang SM, Miguel E. Global non-linear effect of temperature on economic production.
858 *Nature* 2015, **527**: 235.
- 859 74. Hsiang S, Kopp R, Jina A, Rising J, Delgado M, Mohan S, *et al.* Estimating economic damage from
860 climate change in the United States. *Science* 2017, **356**(6345): 1362-1369.
- 861 75. Glanemann N, Willner SN, Levermann A. Paris Climate Agreement passes the cost-benefit test. *Nature*
862 *Communications* 2020, **11**(1): 110.
- 863 76. Fujimori S, Hasegawa T, Masui T, Takahashi K. Land use representation in a global CGE model for
864 long-term simulation: CET vs. logit functions. *Food Security* 2014, **6**(5): 685-699.
- 865 77. Fujimori S, Masui T, Matsuoka Y. AIM/CGE [basic] manual: Center for Social and Environmental
866 Systems Research, National Institute Environmental Studies; 2012. Report No.: 2012-01.
- 867 78. Fujimori S, Masui T, Matsuoka Y. AIM/CGE V2.0 model formula. In: Fujimori S, Kainuma M, Masui
868 T (eds). *Post-2020 Climate Action: Global and Asian Perspective*. Springer, 2017, pp 201-303.
- 869 79. Sands RD. Dynamics of carbon abatement in the Second Generation Model. *Energy Economics* 2004,
870 **26**(4): 721-738.
- 871 80. Rao S, Klimont Z, Smith SJ, Van Dingenen R, Dentener F, Bouwman L, *et al.* Future air pollution in
872 the Shared Socio-economic Pathways. *Global Environmental Change-Human and Policy Dimensions*
873 2017, **42**: 346-358.

- 874 81. Fujimori S, Hasegawa T, Masui T. AIM/CGE V2.0: Basic feature of the model. In: Fujimori S,
875 Kainuma M, Masui T (eds). *Post-2020 Climate Action: Global and Asian Perspective*. Springer, 2017,
876 pp 305-328.
- 877 82. Shinichiro F, Tomoko H, Kiyoshi T, Hancheng D, Jing-Yu L, Haruka O, *et al*. Measuring the
878 sustainable development implications of climate change mitigation. *Environmental Research Letters*
879 2020.
- 880 83. International Energy Agency, (IEA). World Energy balances. In: OECD/IEA, editor. Paris, France;
881 2018.
- 882 84. MOE. Thailand Power Development Plan 2015-2036 (PDP2015). In: Energy Policy and Planning
883 Office, (EPPO), editor. Bangkok: Ministry of Energy; 2015.
- 884 85. MOE. Guidelines for upgrading Thai electricity generation plans. In: Energy Mo, editor. Bangkok:
885 Ministry of Energy,, (MoE); 2018.
- 886 86. MOE. Power Development Plan 2561 - 2580 (PDP2018). Bangkok: Ministry of Energy; 2019.
- 887 87. Huppmann D, Kriegler E, Krey V, Riahi K, Rogelj J, Rose SK, *et al*. IAMC 1.5°C Scenario Explorer
888 and Data hosted by IIASA. Integrated Assessment Modeling Consortium & International Institute for
889 Applied Systems Analysis; 2018.
- 890 88. Fujimori S. Code for A framework for national scenarios with varying emission reductions. 2021
891 [cited]Available from: <https://github.com/shinichirofujimoriKU/AsianMCSAnalysis>

892

893 **Data Availability**

894 Scenario data is accessible online via the ENGAGE Scenario Database at
895 <https://data.ene.iiasa.ac.at/engage/>. Data derived from the original scenario database, which is
896 shown as figures but is not in the above database, is available upon reasonable request from
897 the corresponding author. The scenario name mapping table between this paper and the
898 database are shown in Supplementary Table 3.
899 URL: <https://data.ene.iiasa.ac.at/engage/>
900 DOI: 10.5281/zenodo.4653341

901

902 **Code availability**

903 All code used for data analysis and creating the figures is available at
904 <https://github.com/shinichirofujimoriKU/AsianMCSAnalysis>
905 DOI: 10.5281/zenodo.4677638

906

907 Correspondence and requests for materials should be addressed to Shinichiro Fujimori
908 (fujimori.shinichiro.8a@kyoto-u.ac.jp)

909

910 **Acknowledgements**

911 S.F, K.O and T.M are supported by the Environment Research and Technology Development
912 Fund (JPMEERF20192008 and JPMEERF20211001) of the Environmental Restoration and
913 Conservation Agency of Japan. S.F and K.O are supported by Sumitomo Foundation. S.F.,
914 K.O., V.K. D.vV. K.R, B.L., and T.T.T received funding from the European Union’s Horizon
915 2020 research and innovation programme under grant agreement no. 821471 (ENGAGE).
916 C.P. was supported by the International Research & Development Program of the National
917 Research Foundation of Korea (NRF) funded by the Ministry of Science and ICT
918 (NRF2019K1A3A1A7811297311).

919

920 **Conflict of interest**

921 The authors declare no competing interests.

922

923 **Author contributions**

924 S.F, V.K., DvV, M.S and K.O. designed the research; S.F and Y.O carried out analysis of the
925 modelling results; S.F and Y.O created figures; S.F wrote the draft of the paper; S.F, K.O,
926 O.N. and D.H.S. set up the model; P.C., S.M., C.P., D.H.S., T.T.T., and S.Z. simulated the

927 model and P.C., S.M., C.P., D.H.S., T.T.T., P.N. and S.Z. provided national policy
928 information; and all authors contributed to writing the entire manuscript.
929

