

# International Research Center for Elements Science – Organometallic Chemistry –

<http://om.kuicr.kyoto-u.ac.jp/en/index.html>



Prof  
OHKI, Yasuhiro  
(D Eng)



Assist Prof  
WAKIOKA, Masayuki  
(D Eng)



Assist Prof  
TANIFUJI, Kazuki  
(D Sc)

## Assist Techn Staff

FUJISAKI, Yoshie

## Students

SAKAI, Yuta (M2)\*  
KATO, Itsuki (UG)  
MATSUOKA, Yuto (UG)


\*Special Research Students

## Scope of Research

Developing efficient energy storage systems and innovative material production processes is a significant challenge for chemists to contribute to a sustainable society. We plan to approach these problems by using *transition metal clusters* of which multiple metal atoms work together as catalysts and functional materials. Our laboratory focuses explicitly on creating a new method to synthesize the clusters with atomic precision and applying the obtained clusters to difficult reactions such as the reduction of CO<sub>2</sub> and N<sub>2</sub>.

### KEYWORDS

Transition Metal Clusters  
Homogeneous Catalysis  
Nitrogen Fixation  
Bioinorganic Chemistry



## Recent Selected Publications

Wakioka, M.; Yamashita, N.; Mori, H.; Murdey, R.; Shimoaka, T.; Shioya, N.; Wakamiya, A.; Nishihara, Y.; Hasegawa, T.; Ozawa, F., Formation of *trans*-Poly(thienylenevinylene) Thin Films by Solid-State Thermal Isomerization, *Chem. Mater.*, **33**, 5631-5638 (2021).

Wakioka, M.; Torii, N.; Saito, M.; Osaka, I.; Ozawa, F., Donor–Acceptor Polymers Containing 4,8-Dithienylbenzo[1,2-*b*:4,5-*b'*]dithiophene via Highly Selective Direct Arylation Polymerization, *ACS Appl. Polym. Mater.*, **3**, 830-836 (2021).


Tanifuji, K.; Jasnowski, A. J.; Villarreal, D.; Stiebritz, M. T.; Lee, C. C.; Wilcoxon, J.; Ohki, Y.; Chatterjee, R.; Bogacz, I.; Yano, J.; Kern, J.; Hedman, B.; Hodgson, K. O.; Britt, R. D.; Hu, Y.; Ribbe, M. W., Tracing the Incorporation of the “Ninth Sulfur” into the Nitrogenase Cofactor Precursor with Selenite and Tellurite, *Nat. Chem.*, **13**, 1228-1234 (2021).

Ohki, T.; Ishihara, K.; Yaoi, M.; Tada, M.; Sameera, W. M. C.; Cramer, R. E., A Dinuclear Mo<sub>2</sub>H<sub>8</sub> Complex Supported by Bulky C<sub>5</sub>H<sub>2</sub>Bu<sub>3</sub> Ligands, *Chem. Commun.*, **56**, 8035-8038 (2020).

Ohki, Y.; Uchida, K.; Tada, M.; Cramer, R. E.; Ogura, T.; Ohta, T., N<sub>2</sub> Activation on a Molybdenum–Titanium–Sulfur Cluster, *Nat. Commun.*, **9**, 3200 (2018).

## Catalytic N<sub>2</sub> Silylation by the Fe Sites of Cuboidal [Mo<sub>3</sub>S<sub>4</sub>Fe] Clusters


Biological N<sub>2</sub> fixation is conducted by nitrogenase that employs a unique Fe/Mo-S-C cluster as its catalytic site (FeMoco, [(*R*-homocitrate)MoFe<sub>7</sub>S<sub>9</sub>C]). Synthetic counterparts of the FeMoco, metal-sulfur clusters, demonstrated capturing N<sub>2</sub> on rare occasions; nevertheless, the catalytic conversion of this stable molecule has not been achieved despite its relevance to the biological N<sub>2</sub> fixation. This study focuses on capture, activation, and catalytic conversion of N<sub>2</sub> by an Fe atom incorporated into our [Mo<sub>3</sub>S<sub>4</sub>] incomplete-cubane platform bearing bulky Cp ligands. Treatment of these clusters with excess Na and ClSiMe<sub>3</sub> under a N<sub>2</sub> atmosphere gave N(SiMe<sub>3</sub>)<sub>3</sub> with up to 248 eq. per cluster. This work exemplifies the N<sub>2</sub>-reducing capability of Fe atoms in a S-rich environment, which biological systems have selected to achieve a similar purpose.



**Figure 1.** Proposed N<sub>2</sub> binding state of FeMoco and a N<sub>2</sub>-bound Mo-Fe-S cluster.

## Tracing the S Incorporation into the Nitrogenase Cofactor Precursor

FeMoco is arguably one of the most complex metallocofactors in Nature. Its biosynthetic pathway is correspondingly complicated and remains unclear, which hampers applications of this enzyme toward artificial N<sub>2</sub> fixation. In this study, we investigated an enzymatic process by which FeMoco precursor (L-cluster, [Fe<sub>8</sub>S<sub>9</sub>C]) is generated from two [Fe<sub>4</sub>S<sub>4</sub>] clusters on the protein by using a *semi-synthetic* approach. The study revealed that this process includes a S atom uptake from SO<sub>3</sub><sup>2-</sup> and that the S atom is replaceable with homologous elements (Se, Te). Moreover, we succeeded in selective observation of the incorporated elements and theoretical simulations supporting the reactions' feasibility. These results show that nitrogenase requires an S source as an external substrate for its function.



**Figure 2.** Schematic description of a sulfur uptake from sulfite (SO<sub>3</sub><sup>2-</sup>) in the biosynthetic pathway of FeMoco.