

Triplet of Fibonacci Duals

— Identical Duality —

Seiichi Iwamoto
Professor emeritus, Kyushu University

Yutaka Kimura
Department of Management Science and Engineering
Faculty of Systems Science and Technology
Akita Prefectural University

Abstract

We consider three pairs of quadratic optimization problems from a view point of identical duality. An identity

$$(CI) \quad \sum_{k=1}^{n-1} [(x_{k-1} - x_k)\mu_k + x_k(\mu_k - \mu_{k+1})] + (x_{n-1} - x_n)\mu_n + x_n\mu_n = x_0\mu_1$$

is called complementary. We show that a complementary identity via *conditional complementarity* produces a pair of conditional minimization (primal) problem and conditional maximization (dual) problem, together with an equality condition. It is shown that both the problems have an identical optimal solution (point and value). Moreover, we show that a primal and its dual satisfy *Fibonacci Identical Duality*.

1 Introduction

Bellman and others [1–12, 26] have analyzed a wide class of quadratic optimization problems. Dynamic programming has solved its partial class [2, 17, 18, 29]. Further a dual approach has been discussed through convex-concavity [14, 16, 28].

Recently some dual approaches — (1) extended Lagrangean method, (2) plus-minus method, (3) inequality method, (4) complementary method and others — have been proposed in [18, 20–25].

In this paper, we show an *identical duality* for three pairs of minimization (primal) problems and maximization (dual) problems — (P_1^*) vs (D_1^*) , (P_2^*) vs (D_2^*) and (P_3^*) vs (D_3^*) —. These three are the respective identical versions of (P_1) vs (D_1) , (P_2) vs (D_2) and (P_3) vs (D_3) , which have analyzed the *complementary duality* [23].

It is shown that each pair is dual to each other. It turns out that the duality is based upon the complementary identity and an elementary inequality with equality

$$2xy \leq x^2 + y^2 \quad \text{on } R^2; \quad x = y. \quad (1)$$

2 Identical Duality

Let $x = \{x_k\}_0^n$, $\mu = \{\mu_k\}_1^n$ be any two sequences of real number with $x_0 = c$.

2.1 (P₁^{*}) vs (D₁^{*})

Then a complementary identity

$$(C_1) \quad c\mu_1 = \sum_{k=1}^{n-1} [(x_{k-1} - x_k)\mu_k + x_k(\mu_k - \mu_{k+1})] + (x_{n-1} - x_n)\mu_n + x_n\mu_n$$

holds true.

Let us define two sequences $y = \{y_k\}_1^{2n}$, $\nu = \{\nu_k\}_1^{2n}$ from $x = \{x_k\}_0^n$, $\mu = \{\mu_k\}_1^n$ through

$$\begin{aligned} y_1 &= c - x_1, \quad y_2 = x_1, \quad y_3 = x_1 - x_2, \quad y_4 = x_2, \quad y_5 = x_2 - x_3 \\ &\dots, \quad y_{2n-2} = x_{n-1}, \quad y_{2n-1} = x_{n-1} - x_n, \quad y_{2n} = x_n \\ \nu_1 &= \mu_1, \quad \nu_2 = \mu_1 - \mu_2, \quad \nu_3 = \mu_2, \quad \nu_4 = \mu_2 - \mu_3, \quad \nu_5 = \mu_3 \\ &\dots, \quad \nu_{2n-2} = \mu_{n-1} - \mu_n, \quad \nu_{2n-1} = \mu_n, \quad \nu_{2n} = \mu_n \end{aligned} \quad (2)$$

, respectively. Then an identity

$$(C_1^*) \quad c\nu_1 = \sum_{k=1}^{2n} y_k \nu_k \quad (3)$$

holds *under a constraint* – a linear system of $4n$ -variable (y, ν) on $2n$ -equation – :

$$\begin{aligned} (C^{y\nu}) \quad & \begin{array}{ll} c = y_1 + y_2 & \nu_1 = \nu_2 + \nu_3 \\ y_2 = y_3 + y_4 & \nu_3 = \nu_4 + \nu_5 \\ \vdots & \vdots \\ y_{2n-4} = y_{2n-3} + y_{2n-2} & \nu_{2n-3} = \nu_{2n-2} + \nu_{2n-1} \\ y_{2n-2} = y_{2n-1} + y_{2n} & \nu_{2n-1} = \nu_{2n}. \end{array} \end{aligned} \quad (4)$$

An equality (3) with constraint (4) is called a $2n$ -variable *conditional complementarity*. This is simply written as (C_1^*) under $(C^{y\nu})$.

Now let $y = \{y_k\}_1^{2n}$, $\nu = \{\nu_k\}_1^{2n}$ satisfy $(C_1^{y\nu})$. Then an elementary inequality (1) yields

$$2c\nu_1 \leq \sum_{k=1}^{2n} (y_k^2 + \nu_k^2).$$

Thus we have an inequality

$$2c\nu_1 - \sum_{k=1}^{2n} \nu_k^2 \leq \sum_{k=1}^{2n} y_k^2.$$

The sign of equality holds iff

$$(EC_1^*) \quad y_k = \nu_k \quad 1 \leq k \leq 2n. \quad (5)$$

Hence we have a pair of conditional minimization problem:

$$\begin{aligned}
 & \text{minimize } y_1^2 + y_2^2 + \cdots + y_{2n-1}^2 + y_{2n}^2 \\
 & \text{subject to } (1) \ y_1 + y_2 = c \\
 & \qquad \qquad \qquad (2) \ y_3 + y_4 = y_2 \\
 (P_1^*) \quad & \qquad \qquad \qquad \vdots \\
 & (n-1) \ y_{2n-3} + y_{2n-2} = y_{2n-4} \\
 & \qquad \qquad \qquad (n) \ y_{2n-1} + y_{2n} = y_{2n-2} \\
 & (n+1) \ y \in R^{2n}
 \end{aligned}$$

and conditional maximization problem:

$$\begin{aligned}
 & \text{Maximize } 2c\nu_1 - (\nu_1^2 + \nu_2^2 + \cdots + \nu_{2n-1}^2 + \nu_{2n}^2) \\
 & \text{subject to } [1] \ \nu_2 + \nu_3 = \nu_1 \\
 & \qquad \qquad \qquad [2] \ \nu_4 + \nu_5 = \nu_3 \\
 (D_1^*) \quad & \qquad \qquad \qquad \vdots \\
 & [n-1] \ \nu_{2n-2} + \nu_{2n-1} = \nu_{2n-3} \\
 & \qquad \qquad \qquad [n] \ \nu_{2n} = \nu_{2n-1} \\
 & [n+1] \ \nu \in R^{2n}.
 \end{aligned}$$

Let (AC_1) be an *augmentation* of the system $(C_1^{y\nu})$ with the additional equality condition (EC_1^*) :

$$\begin{aligned}
 (AC_1) \quad & \begin{array}{ll} c = y_1 + y_2 & \nu_1 = \nu_2 + \nu_3 \\ y_2 = y_3 + y_4 & \nu_3 = \nu_4 + \nu_5 \\ \vdots & \vdots \\ y_{2n-4} = y_{2n-3} + y_{2n-2} & \nu_{2n-3} = \nu_{2n-2} + \nu_{2n-1} \\ y_{2n-2} = y_{2n-1} + y_{2n} & \nu_{2n-1} = \nu_{2n} \end{array} \\
 & y_k = \nu_k \quad 1 \leq k \leq 2n.
 \end{aligned}$$

The linear system (AC_1) is of $4n$ -variable on $4n$ -equation. Let (y, ν) satisfy (AC_1) . Then both sides become a common value with five expressions.

$$\begin{aligned}
 (5V_1) \quad & y_1^2 + y_2^2 + \cdots + y_{2n}^2 \\
 & = cy_1 \\
 & = 2c\nu_1 - (\nu_1^2 + \nu_2^2 + \cdots + \nu_{2n}^2) \\
 & = \nu_1^2 + \nu_2^2 + \cdots + \nu_{2n}^2 \\
 & = c\nu_1.
 \end{aligned}$$

The system (AC₁) has indeed a unique common solution:

$$\begin{aligned} y &= (y_1, y_2, \dots, y_k, \dots, y_{2n-1}, y_{2n}) \\ &= \frac{c}{F_{2n+1}}(F_{2n}, F_{2n-1}, \dots, F_{2n-k+1}, \dots, F_2, F_1), \\ \nu &= (\nu_1, \nu_2, \dots, \nu_k, \dots, \nu_{2n-1}, \nu_{2n}) \\ &= \frac{c}{F_{2n+1}}(F_{2n}, F_{2n-1}, \dots, F_{2n-k+1}, \dots, F_2, F_1) \end{aligned}$$

where $\{F_n\}$ is the *Fibonacci sequence* [13,15,27,30]. This is defined as the solution to the second-order linear difference equation

$$x_{n+2} - x_{n+1} - x_n = 0, \quad x_1 = 1, \quad x_0 = 0. \quad (6)$$

n	\dots	-2	-1	0	1	2	3	4	5	6	7	8	9	10	11
F_n	\dots	-1	1	0	1	1	2	3	5	8	13	21	34	55	89

Table 1 Fibonacci sequence $\{F_n\}$

The primal (P₁^{*}) has a minimum value $m_1 = \frac{F_{2n}}{F_{2n+1}}c^2$ at a path

$$\begin{aligned} \hat{y} &= (\hat{y}_1, \hat{y}_2, \dots, \hat{y}_k, \dots, \hat{y}_{2n-1}, \hat{y}_{2n}) \\ &= \frac{c}{F_{2n+1}}(F_{2n}, F_{2n-1}, \dots, F_{2n-k+1}, \dots, F_2, F_1). \end{aligned}$$

The dual (D₁^{*}) has a maximum value $M_1 = \frac{F_{2n}}{F_{2n+1}}c^2$ at a path

$$\begin{aligned} \nu^* &= (\nu_1^*, \nu_2^*, \dots, \nu_k^*, \dots, \nu_{2n-1}^*, \nu_{2n}^*) \\ &= \frac{c}{F_{2n+1}}(F_{2n}, F_{2n-1}, \dots, F_{2n-k+1}, \dots, F_2, F_1). \end{aligned}$$

Both optimal solutions (point and value) are identical:

$$\hat{x} = \mu^*, \quad m_1 = M_1.$$

Further both are Fibonacci:

$$\begin{aligned} \hat{x} = \mu^* &= \frac{c}{F_{2n+1}}(F_{2n}, F_{2n-1}, \dots, F_{2n-k+1}, \dots, F_2, F_1), \\ m_1 = M_1 &= \frac{F_{2n}}{F_{2n+1}}c^2. \end{aligned}$$

Thus *Fibonacci Identical Duality* (FID) [18–20,22,24,25] holds between (P₁^{*}) and (D₁^{*}).

We remark that the $2n$ -variable pair is a transliteration from n -variable one

$$(P_1) \quad \begin{aligned} & \text{minimize} \quad \sum_{k=1}^{n-1} [(x_{k-1} - x_k)^2 + x_k^2] + (x_{n-1} - x_n)^2 + x_n^2 \\ & \text{subject to} \quad (i) \ x \in R^n, \quad (ii) \ x_0 = c \end{aligned}$$

$$(D_1) \quad \begin{aligned} & \text{Maximize} \quad 2c\mu_1 - \sum_{k=1}^{n-1} [\mu_k^2 + (\mu_k - \mu_{k+1})^2] - \mu_n^2 - \mu_n^2 \\ & \text{subject to} \quad (i) \ \mu \in R^n. \end{aligned}$$

2.2 (P₂^{*}) vs (D₂^{*})

On the other hand, we assume that $\underline{\mu_n = 0}$. Then an identity

$$(C_2) \quad c\mu_1 = \sum_{k=1}^{n-1} [(x_{k-1} - x_k)\mu_k + x_k(\mu_k - \mu_{k+1})] + (x_{n-1} - x_n)\mu_n$$

holds true.

Let us define two sequences $y = \{y_k\}_1^{2n-1}$, $\nu = \{\nu_k\}_1^{2n-1}$ from $x = \{x_k\}_0^n$, $\mu = \{\mu_k\}_1^n$ through

$$\begin{aligned} & y_1 = c - x_1, \ y_2 = x_1, \ y_3 = x_1 - x_2, \ y_4 = x_2, \ y_5 = x_2 - x_3, \ \dots \\ & , \ y_{2n-4} = x_{n-2}, \ y_{2n-3} = x_{n-2} - x_{n-1}, \ y_{2n-2} = x_{n-1}, \ y_{2n-1} = x_{n-1} - x_n \\ & \nu_1 = \mu_1, \ \nu_2 = \mu_1 - \mu_2, \ \nu_3 = \mu_2, \ \nu_4 = \mu_2 - \mu_3, \ \nu_5 = \mu_3, \ \dots \\ & , \ \nu_{2n-4} = \mu_{n-2} - \mu_{n-1}, \ \nu_{2n-3} = \mu_{n-1}, \ \nu_{2n-2} = \mu_{n-1} - \mu_n, \ \nu_{2n-1} = \mu_n \end{aligned} \tag{7}$$

, respectively. Then an identity

$$(C_2^*) \quad c\nu_1 = \sum_{k=1}^{2n-1} y_k\nu_k$$

holds under a constraint – a linear system of $(4n - 2)$ -variable on $(2n - 1)$ -equation – :

$$(C_2^{y\nu}) \quad \begin{array}{ll} c = y_1 + y_2 & \nu_1 = \nu_2 + \nu_3 \\ y_2 = y_3 + y_4 & \nu_3 = \nu_4 + \nu_5 \\ \vdots & \vdots \\ y_{2n-6} = y_{2n-5} + y_{2n-4} & \nu_{2n-5} = \nu_{2n-4} + \nu_{2n-3} \\ y_{2n-4} = y_{2n-3} + y_{2n-2} & \nu_{2n-3} = \nu_{2n-2} + \nu_{2n-1} \\ & \underline{\nu_{2n-1} \equiv 0}. \end{array}$$

Thus we have a $(2n - 1)$ -variable *conditional complementarity* (C_2^*) under $(C_2^{y\nu})$.

Let $y = \{y_k\}_1^{2n-1}$, $\nu = \{\nu_k\}_1^{2n-1}$ satisfy $(C_2^{y\nu})$. Then the elementary inequality (1) yields

$$2c\nu_1 \leq \sum_{k=1}^{2n-1} (y_k^2 + \nu_k^2).$$

Thus we have an inequality

$$2c\nu_1 - \sum_{k=1}^{2n-1} \nu_k^2 \leq \sum_{k=1}^{2n-1} y_k^2.$$

The sign of equality holds iff

$$(EC_2^*) \quad y_k = \nu_k \quad 1 \leq k \leq 2n-1. \quad (8)$$

Hence we have a pair of conditional minimization problem:

$$(P_2^*) \quad \begin{array}{ll} \text{minimize} & y_1^2 + y_2^2 + \cdots + y_{2n-2}^2 + y_{2n-1}^2 \\ \text{subject to} & (1) \quad y_1 + y_2 = c \\ & (2) \quad y_3 + y_4 = y_2 \\ & \vdots \\ & (n-2) \quad y_{2n-5} + y_{2n-4} = y_{2n-6} \\ & (n-1) \quad y_{2n-3} + y_{2n-2} = y_{2n-4} \\ & (n) \quad y \in R^{2n-1} \end{array}$$

and conditional maximization problem:

$$(D_2^*) \quad \begin{array}{ll} \text{Maximize} & 2c\nu_1 - (\nu_1^2 + \nu_2^2 + \cdots + \nu_{2n-2}^2 + \nu_{2n-1}^2) \\ \text{subject to} & [1] \quad \nu_2 + \nu_3 = \nu_1 \\ & [2] \quad \nu_4 + \nu_5 = \nu_3 \\ & \vdots \\ & [n-1] \quad \nu_{2n-2} + \nu_{2n-1} = \nu_{2n-3} \\ & [n] \quad \underline{\nu_{2n-1} = 0} \\ & [n+1] \quad \nu \in R^{2n-1}. \end{array}$$

Let (AC_2) be an *augmentation* of the system $(C_2^{y\nu})$ with the additional equality condi-

tion (EC₂^{*}):

$$\begin{array}{rcl}
 c = y_1 + y_2 & & \nu_1 = \nu_2 + \nu_3 \\
 y_2 = y_3 + y_4 & & \nu_3 = \nu_4 + \nu_5 \\
 \vdots & & \vdots \\
 (AC_2) \quad y_{2n-6} = y_{2n-5} + y_{2n-4} & & \nu_{2n-5} = \nu_{2n-4} + \nu_{2n-3} \\
 y_{2n-4} = y_{2n-3} + y_{2n-2} & & \nu_{2n-3} = \nu_{2n-2} + \nu_{2n-1} \\
 & & \underline{\nu_{2n-1} = 0} \\
 y_k = \nu_k \quad 1 \leq k \leq 2n-1.
 \end{array}$$

The linear system (AC₂) is of $(4n-2)$ -variable on $(4n-2)$ -equation. Let (y, ν) satisfy (AC₂). Then both sides become a common value with five expressions.

$$\begin{aligned}
 & y_1^2 + y_2^2 + \cdots + y_{2n-1}^2 \\
 & = cy_1 \\
 (5V_2) \quad & = 2c\nu_1 - (\nu_1^2 + \nu_2^2 + \cdots + \nu_{2n-1}^2) \\
 & = \nu_1^2 + \nu_2^2 + \cdots + \nu_{2n-1}^2 \\
 & = c\nu_1.
 \end{aligned}$$

The system (AC₂) has indeed a unique common solution:

$$\begin{aligned}
 & y = (y_1, y_2, \dots, y_k, \dots, y_{2n-2}, y_{2n-1}) \\
 & = \frac{c}{F_{2n-1}}(F_{2n-2}, F_{2n-3}, \dots, F_{2n-k-1}, \dots, F_1, F_0), \\
 & \nu = (\nu_1, \nu_2, \dots, \nu_k, \dots, \nu_{2n-2}, \nu_{2n-1}) \\
 & = \frac{c}{F_{2n-1}}(F_{2n-2}, F_{2n-3}, \dots, F_{2n-k-1}, \dots, F_1, F_0).
 \end{aligned}$$

The primal (P₂^{*}) has a minimum value $m_2 = \frac{F_{2n-2}}{F_{2n-1}}c^2$ at a path

$$\begin{aligned}
 & \hat{y} = (\hat{y}_1, \hat{y}_2, \dots, \hat{y}_k, \dots, \hat{y}_{2n-2}, \hat{y}_{2n-1}) \\
 & = \frac{c}{F_{2n-1}}(F_{2n-2}, F_{2n-3}, \dots, F_{2n-k-1}, \dots, F_1, F_0).
 \end{aligned}$$

The dual (D₂^{*}) has a maximum value $M_2 = \frac{F_{2n-2}}{F_{2n-1}}c^2$ at a path

$$\begin{aligned}
 & \nu^* = (\nu_1^*, \nu_2^*, \dots, \nu_k^*, \dots, \nu_{2n-2}^*, \nu_{2n-1}^*) \\
 & = \frac{c}{F_{2n-1}}(F_{2n-2}, F_{2n-3}, \dots, F_{2n-k-1}, \dots, F_1, F_0).
 \end{aligned}$$

Both optimal solutions (point and value) are identical:

$$\hat{x} = \mu^*, \quad m_2 = M_2.$$

Further both are Fibonacci:

$$\hat{x} = \mu^* = \frac{c}{F_{2n-1}}(F_{2n-2}, F_{2n-3}, \dots, F_{2n-k-1}, \dots, F_1, F_0),$$

$$m_2 = M_2 = \frac{F_{2n-2}}{F_{2n-1}}c^2.$$

Thus FID holds between (P_2^*) and (D_2^*) .

Note that the $(2n-1)$ -variable pair is a transliteration from n -variable one

$$(P_2) \quad \begin{aligned} & \text{minimize} \quad \sum_{k=1}^{n-1} [(x_{k-1} - x_k)^2 + x_k^2] + (x_{n-1} - x_n)^2 \\ & \text{subject to} \quad (i) \ x \in R^n, \quad (ii) \ x_0 = c \end{aligned}$$

$$(D_2) \quad \begin{aligned} & \text{Maximize} \quad 2c\mu_1 - \sum_{k=1}^{n-1} [\mu_k^2 + (\mu_k - \mu_{k+1})^2] - \mu_n^2 \\ & \text{subject to} \quad (i) \ \mu \in R^n, \quad (ii) \ \underline{\mu_n = 0}. \end{aligned}$$

2.3 (P_3^*) vs (D_3^*)

On the other hand, we assume that $x_n = 0$. Then an identity

$$(C_3) \quad c\mu_1 = \sum_{k=1}^{n-1} [(x_{k-1} - x_k)\mu_k + x_k(\mu_k - \mu_{k+1})] + (x_{n-1} - x_n)\mu_n$$

holds true.

Let us define two sequences $y = \{y_k\}_1^{2n-1}$, $\nu = \{\nu_k\}_1^{2n-1}$ from $x = \{x_k\}_0^n$, $\mu = \{\mu_k\}_1^n$ through

$$\begin{aligned} & y_1 = c - x_1, \ y_2 = x_1, \ y_3 = x_1 - x_2, \ y_4 = x_2, \ y_5 = x_2 - x_3, \ \dots \\ & , \ y_{2n-4} = x_{n-2}, \ y_{2n-3} = x_{n-2} - x_{n-1}, \ y_{2n-2} = x_{n-1}, \ y_{2n-1} = x_{n-1} - x_n \\ & \nu_1 = \mu_1, \ \nu_2 = \mu_1 - \mu_2, \ \nu_3 = \mu_2, \ \nu_4 = \mu_2 - \mu_3, \ \nu_5 = \mu_3, \ \dots \\ & , \ \nu_{2n-4} = \mu_{n-2} - \mu_{n-1}, \ \nu_{2n-3} = \mu_{n-1}, \ \nu_{2n-2} = \mu_{n-1} - \mu_n, \ \nu_{2n-1} = \mu_n \end{aligned} \tag{9}$$

, respectively. Then an identity

$$(C_3^*) \quad c\nu_1 = \sum_{k=1}^{2n-1} y_k \nu_k \tag{10}$$

Let (AC_3) be an *augmentation* of the system $(C_3^{y\nu})$ with the additional equality condition (EC_3^*) :

$$\begin{aligned}
 & c = y_1 + y_2 & \nu_1 &= \nu_2 + \nu_3 \\
 & y_2 = y_3 + y_4 & \nu_3 &= \nu_4 + \nu_5 \\
 & \vdots & \vdots & \\
 (AC_3) \quad & y_{2n-4} = y_{2n-3} + y_{2n-2} & \nu_{2n-3} &= \nu_{2n-2} + \nu_{2n-1} \\
 & \underline{y_{2n-2}} = \underline{y_{2n-1}} & & \\
 & y_k = \nu_k \quad 1 \leq k \leq 2n-1.
 \end{aligned}$$

The linear system (AC_3) is of $(4n-2)$ -variable on $(4n-2)$ -equation. Let (y, ν) satisfy (AC_3) . Then both sides become a common value with five expressions.

$$\begin{aligned}
 & y_1^2 + y_2^2 + \cdots + y_{2n-1}^2 \\
 &= cy_1 \\
 (5V_3) \quad &= 2c\nu_1 - (\nu_1^2 + \nu_2^2 + \cdots + \nu_{2n-1}^2) \\
 &= \nu_1^2 + \nu_2^2 + \cdots + \nu_{2n-1}^2 \\
 &= c\nu_1.
 \end{aligned}$$

The system (AC_3) has indeed a unique common solution:

$$\begin{aligned}
 & y = (y_1, y_2, \dots, y_k, \dots, y_{2n-2}, y_{2n-1}) \\
 &= \frac{c}{F_{2n}}(F_{2n-1}, F_{2n-2}, \dots, F_{2n-k}, \dots, F_2, F_1), \\
 & \nu = (\nu_1, \nu_2, \dots, \nu_k, \dots, \nu_{2n-2}, \nu_{2n-1}) \\
 &= \frac{c}{F_{2n}}(F_{2n-1}, F_{2n-2}, \dots, F_{2n-k}, \dots, F_2, F_1).
 \end{aligned}$$

The primal (P_3^*) has a minimum value $m_3 = \frac{F_{2n-1}}{F_{2n}}c^2$ at a path

$$\begin{aligned}
 & \hat{y} = (\hat{y}_1, \hat{y}_2, \dots, \hat{y}_k, \dots, \hat{y}_{2n-2}, \hat{y}_{2n-1}) \\
 &= \frac{c}{F_{2n}}(F_{2n-1}, F_{2n-2}, \dots, F_{2n-k}, \dots, F_2, F_1).
 \end{aligned}$$

The dual (D_3^*) has a maximum value $M_3 = \frac{F_{2n-1}}{F_{2n}}c^2$ at a path

$$\begin{aligned}
 & \nu^* = (\nu_1^*, \nu_2^*, \dots, \nu_k^*, \dots, \nu_{2n-2}^*, \nu_{2n-1}^*) \\
 &= \frac{c}{F_{2n}}(F_{2n-1}, F_{2n-2}, \dots, F_{2n-k}, \dots, F_2, F_1).
 \end{aligned}$$

Both optimal solutions (point and value) are identical:

$$\hat{x} = \mu^*, \quad m_3 = M_3.$$

Further both are Fibonacci:

$$\hat{x} = \mu^* = \frac{c}{F_{2n}}(F_{2n-1}, F_{2n-2}, \dots, F_{2n-k}, \dots, F_2, F_1),$$

$$m_3 = M_3 = \frac{F_{2n-1}}{F_{2n}}c^2.$$

Thus FID holds between (P_3^*) and (D_3^*) .

Note that the $(2n - 1)$ -variable pair is a transliteration from n -variable one

$$(P_3) \quad \begin{array}{l} \text{minimize} \quad \sum_{k=1}^{n-1} [(x_{k-1} - x_k)^2 + x_k^2] + (x_{n-1} - x_n)^2 \\ \text{subject to} \quad \text{(i) } x \in R^n, \quad \text{(ii) } x_0 = c, \underline{x_n = 0} \end{array}$$

$$(D_3) \quad \begin{array}{l} \text{Maximize} \quad 2c\mu_1 - \sum_{k=1}^{n-1} [\mu_k^2 + (\mu_k - \mu_{k+1})^2] - \mu_n^2 \\ \text{subject to} \quad \text{(i) } \mu \in R^n. \end{array}$$

References

- [1] E.F. Beckenbach and R.E. Bellman, Inequalities, Springer-Verlag, Ergebnisse 30, 1961.
- [2] R.E. Bellman, Dynamic Programming, Princeton Univ. Press, NJ, 1957.
- [3] R.E. Bellman, Introduction to the Mathematical Theory of Control Processes, Vol.I, Linear Equations and Quadratic Criteria, Academic Press, NY, 1967.
- [4] R.E. Bellman, Introduction to the Mathematical Theory of Control Processes, Vol.II, Nonlinear Processes, Academic Press, NY, 1971.
- [5] R.E. Bellman, Methods of Nonlinear Analysis, Vol.I, Nonlinear Processes, Academic Press, NY, 1972.
- [6] R.E. Bellman, Methods of Nonlinear Analysis, Vol.II, Nonlinear Processes, Academic Press, NY, 1972.
- [7] R.E. Bellman, Introduction to Matrix Analysis, McGraw-Hill, New York, NY, 1970 (Second Edition is a SIAM edition 1997).
- [8] R.E. Bellman and Wm. Karush, On a new functional transform in analysis : the maximum transform, Bull. Amer. Math. Soc., **67**(1961), 501-503.

- [9] R.E. Bellman and Wm. Karush, Mathematical programming and the maximum transform, *J. SIAM Appl. Math.*, **10**(1962), 550-567.
- [10] R.E. Bellman and Wm. Karush, On the maximum transform and semigroups of transformations, *Bull. Amer. Math. Soc.*, **68**(1962), 516-518.
- [11] R.E. Bellman and Wm. Karush, Functional equations in the theory of dynamic programming-XII : an application of the maximum transform, *J. Math. Anal. Appl.*, **6**(1963), 155-157.
- [12] R.E. Bellman and Wm. Karush, On the maximum transform, *J. Math. Anal. Appl.*, **6**(1963), 57-74.
- [13] A. Beutelspacher and B. Petri, *Der Goldene Schnitt 2, überarbeitete und erweiterte Auflage*, Elsevier GmbH, Spectrum Akademischer Verlag, Heidelberg, 1996.
- [14] J.M. Borwein and A.S. Lewis, *Convex Analysis and Nonlinear Optimization Theory and Examples*, Springer-Verlag, New York, 2000.
- [15] R.A. Dunlap, *The Golden Ratio and Fibonacci Numbers*, World Scientific Publishing Co.Pte.Ltd., 1977.
- [16] W. Fenchel, *Convex Cones, Sets and Functions*, Princeton Univ. Dept. of Math, NJ, 1953.
- [17] S. Iwamoto, *Theory of Dynamic Program*, Kyushu Univ. Press, Fukuoka, 1987 (*in Japanese*).
- [18] S. Iwamoto, *Mathematics for Optimization II Bellman Equation*, Chisen Shokan, Tokyo, 2013 (*in Japanese*).
- [19] S. Iwamoto and Y. Kimura, Nonhomogeneous Semi-Fibonacci Programming – Identical Duality – , *RIMS Kokyuroku*, Vol.2078, pp.114–120, 2018.
- [20] S. Iwamoto and Y. Kimura, Semi-Fibonacci Programming – odd-variable – , *RIMS Kokyuroku*, Vol.2158, pp.30–37, 2020.
- [21] S. Iwamoto and Y. Kimura, Semi-tridiagonal Programming – Complementary Approach – , *RIMS Kokyuroku*, Vol.2190, 2021, to appear.
- [22] S. Iwamoto and Y. Kimura, Identical Duals – Gap Function – , *RIMS Kokyuroku*, Vol.2194, pp.56–67, 2021.
- [23] S. Iwamoto and Y. Kimura, Triplet of Fibonacci Duals – with or without constraint – , *RIMS Kokyuroku*, to appear.
- [24] S. Iwamoto, Y. Kimura and T. Fujita, Complementary versus shift dualities, *J. Non-linear Convex Anal.*, **17**(2016), 1547–1555.

- [25] S. Iwamoto, Y. Kimura and T. Fujita, On complementary duals – both fixed points –, Bull. Kyushu Inst. Tech. Pure Appl. Math., **67**(2020), 1–28.
- [26] E.S. Lee, Quasilinearization and Invariant Imbedding, Academic Press, New York, 1968.
- [27] S. Nakamura, A Microcosmos of Fibonacci Numbers — Fibonacci Numbers, Lucas Numbers, and Golden Section — (Revised), Nippon Hyoronsha, 2008 (*in Japanese*).
- [28] R.T. Rockafeller, Conjugate Duality and Optimization, SIAM, Philadelphia, 1974.
- [29] M. Sniedovich, Dynamic Programming: foundations and principles, 2nd ed., CRC Press 2010.
- [30] H. Walsert, DER GOLDENE SCHNITT, B.G. Teubner, Leipzig, 1996.