

1  
2  
3  
4  
5  
6  
7 **Effect of electrical activity of the diaphragm waveform patterns on SpO<sub>2</sub> for**  
8  
9  
10 **extremely preterm infants ventilated with neurally adjusted ventilatory assist**  
11  
12  
13  
14  
15

16 Ryosuke Araki MD; Seiichi Tomotaki MD; Mitsuyo Akita MD; Kouji Motokura MD;  
17  
18

19 Yutaro Tomobe MD; Takeru Yamauchi MD; Shintaro Hanaoka MD; Hiroko Tomotaki  
20  
21

22 MD; Kogoro Iwanaga MD; Fusako Niwa MD, PhD; Junko Takita MD, PhD; Masahiko  
23  
24

25  
26 Kawai MD, PhD  
27  
28

29  
30 Department of Pediatrics, Graduate School of Medicine, Kyoto University, 54  
31  
32

33 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan.  
34  
35  
36  
37  
38  
39

40 **Corresponding Author:** Ryosuke Araki, Department of Pediatrics, Graduate School of  
41  
42

43 Medicine, Kyoto University, 54 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507,  
44  
45

46 Japan.  
47  
48

49  
50 E-mail: r\_araki@kuhp.kyoto-u.ac.jp; Tel: +81-75-751-3291; Fax: +81-75-751-2361  
51  
52  
53  
54  
55

56 **Conflicts of Interest and Source of Funding:** None declared.  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Keywords:** bronchopulmonary dysplasia; neonatal intensive care units; diaphragm;  
lung injury.

For Peer Review

## Abstract

### Objective

This study aimed to evaluate the association between electrical activity of the diaphragm (Edi) waveform patterns and peripheral oxygen saturation (SpO<sub>2</sub>) in extremely preterm infants who are ventilated with neurally adjusted ventilatory assist (NAVA).

### Study Design

We conducted a retrospective cohort study at a level III neonatal intensive care unit. Extremely preterm infants born at our hospital between November 2019 and November 2020 and ventilated with NAVA were included. We collected Edi waveform data and classified them into four Edi waveform patterns, including the phasic pattern, central apnea pattern, irregular low-voltage pattern, and tonic burst pattern. We analyzed the Edi waveform pattern for the first 15 hours of collectable data in each patient. To investigate the association between Edi waveform patterns and SpO<sub>2</sub>, we analyzed the dataset every 5 minutes as one data unit. We compared the proportion of each waveform pattern between the desaturation (Desat (+)) and non-desaturation (Desat (-)) groups.

## Results

We analyzed collected data for 105 hours (1260 data units). The proportion of the phasic pattern in the Desat (+) group was significantly lower than that in the Desat (-) group ( $p < 0.001$ ). However, the proportions of the central apnea, irregular low-voltage, and tonic burst patterns in the Desat (+) group were significantly higher than those in the Desat (-) group (all  $p < 0.05$ ).

## Conclusion

Our results indicate that proportions of Edi waveform patterns have an effect on desaturation of SpO<sub>2</sub> in extremely preterm infants who are ventilated with NAVA.

## INTRODUCTION

Recent advancement in perinatal management has improved the survival rate of extremely preterm infants. Consequently, bronchopulmonary dysplasia (BPD) has become one of the most common and severe sequelae for extremely preterm infants<sup>1,2</sup> and management for preventing BPD has been the focus of attention. Extremely preterm infants require mechanical ventilation for a relatively long period because of prematurity of their lungs. In such a long mechanical ventilation period, high ventilatory pressure and patient–ventilator asynchrony are important factors for BPD<sup>3,4</sup>. Therefore, the strategy of ventilation for extremely preterm infants has focused on minimizing inspiratory pressure and more synchronized ventilation.

Neurally adjusted ventilatory assist (NAVA) is a relatively novel mode of mechanical ventilation and provides respiratory support in response to the patient’s electrical activity of the diaphragm (Edi). Compared with other patient-triggered ventilator modes, NAVA improves patient–ventilatory synchrony and reduces peak inspiratory pressure and the work of breathing without an elevated fraction of inspiratory oxygen ( $\text{FiO}_2$ )<sup>5–9</sup>. There has been a recent increase in reports on ventilation with NAVA for

1  
2  
3  
4  
5  
6 preterm and low birth weight infants<sup>10-13</sup>. NAVA is expected to improve the prognosis  
7  
8  
9  
10 of extremely preterm infants, especially in proceeding to BPD<sup>14,15</sup>. However, despite  
11  
12  
13 appropriate ventilatory settings, some preterm infants cannot be appropriately supported  
14  
15  
16 by NAVA. These infants develop an unstable respiratory condition or need to have their  
17  
18  
19 ventilatory mode changed from NAVA<sup>16-18</sup>. However, there are limited clinical  
20  
21  
22  
23 indicators of whether infants can be appropriately ventilated by NAVA.  
24  
25

26  
27 During ventilation with NAVA, respiratory support is supplied from a ventilator on the  
28  
29  
30 basis of the Edi waveform. Preterm infants are known to represent variable Edi  
31  
32  
33 waveform patterns<sup>19,20</sup>, but there are limited data on the association between Edi  
34  
35  
36 waveform patterns and respiratory conditions. On the basis of our clinical experience,  
37  
38  
39 we consider that not only frequent central apnea, but also low inspiratory movements  
40  
41  
42  
43 prevent appropriate ventilatory support by NAVA.  
44  
45

46  
47 We hypothesize that Edi waveform patterns reflect the respiratory pattern, which  
48  
49  
50 affects appropriate ventilatory support by NAVA and the frequency of desaturation of  
51  
52  
53 peripheral oxygen saturation (SpO<sub>2</sub>).  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 In this study, we evaluated the association between Edi waveform patterns and the  
8  
9  
10 frequency of desaturation in extremely preterm infants who are ventilated with NAVA.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## MATERIALS AND METHODS

### Patients

We conducted a retrospective cohort study at a single level III neonatal intensive care unit (Kyoto University Hospital, Kyoto, Japan). Extremely preterm infants of < 28 weeks gestational age who were born at our hospital between November 2019 and November 2020 and required mechanical ventilation with NAVA delivered by Servo-n<sup>®</sup> (Maquet, Critical Care AB, Solna, Sweden) were eligible for our study. Infants who needed to have their ventilation mode changed from NAVA to another mode (including those who were extubated) within 24 consecutive hours or those who did not have more than 15 hours of data collected were excluded.

### Study procedure

According to our usual practice, the NAVA level was set and adjusted to maintain Edi peak < 15  $\mu\text{V}$  and positive end-expiratory pressure (PEEP) was set and adjusted to maintain Edi minimum at < 5  $\mu\text{V}$ . The backup support setting and apnea time were adjusted to supply proper backup ventilation to avoid desaturation. Because we

1  
2  
3  
4  
5  
6 routinely adjust the NAVA ventilator settings with this procedure, the initial settings  
7  
8  
9  
10 were usually the same as follows: NAVA level = 1.5 to 2 cmH<sub>2</sub>O/ $\mu$ V, PEEP = 7  
11  
12  
13 cmH<sub>2</sub>O, apnea time = 2 seconds, pressure level above PEEP in backup = 10 cmH<sub>2</sub>O,  
14  
15  
16 and back up respiratory rate = 40. By this procedure of adjustment, subsequent  
17  
18  
19 ventilator setting after the initial setting also did not vary in each patient. In accordance  
20  
21  
22  
23 with the manufacturer's recommendations, proper positioning of the Edi catheter was  
24  
25  
26 frequently verified by attending physicians.  
27

28  
29  
30 The waveforms of ventilator pressure, flow, and Edi were acquired from the ventilator  
31  
32  
33 and recorded continuously by using the software SERVO CONNECT (Fukuda Denshi  
34  
35  
36 Co. Ltd., Tokyo, Japan). We routinely use this software and collect data when possible.  
37  
38  
39  
40  
41  
42

### 43 **Data collection**

44  
45  
46 Demographic and clinical data of included infants were extracted from medical  
47  
48  
49 records. Valid data for analysis of Edi waveforms and SpO<sub>2</sub> were defined as when the  
50  
51  
52 infants had not been manipulated and the ventilator setting had not been changed at  
53  
54  
55  
56 least within 3 hours, except for FiO<sub>2</sub>. With SERVO CONNECT, Edi waveforms were  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 shown in one window every 10 seconds. Edi waveforms were classified into four  
7  
8  
9 patterns including those previously reported and a novel pattern<sup>19,20</sup> (Figure 1). These  
10  
11  
12 patterns were as follows: 1) a phasic pattern was defined as a regular increasing and  
13  
14 decreasing pattern, 2) a central apnea pattern was defined as continuous low Edi, which  
15  
16 induced backup mandatory ventilation, 3) a tonic burst pattern was defined as elevated  
17  
18 tonic activity without clear phasic activity or with irregular activity, and 4) an irregular  
19  
20 low-voltage pattern was defined as low Edi activity (Edi [peak-minimum] < 1.5  $\mu$ V).  
21  
22  
23  
24  
25  
26  
27  
28  
29 Because irregular low-voltage patterns did not induce backup ventilation and generated  
30  
31 low peak inspiratory pressure, we distinguished irregular low-voltage patterns from  
32  
33  
34  
35  
36  
37 central apnea patterns.

38  
39  
40 Ventilatory settings, including the NAVA level, PEEP, backup setting, and FiO<sub>2</sub>, were  
41  
42  
43 collected from the medical records. Ventilatory parameters, including Edi peak, Edi  
44  
45  
46 minimum, peak airway pressure, mean airway pressure, and time in backup ventilation  
47  
48  
49 were collected from stored data in the ventilators.  
50  
51

52  
53 SpO<sub>2</sub> data were recorded by a central monitoring system (Phillips Information  
54  
55  
56 Management System<sup>®</sup>; Phillips Japan, Japan) every minute. We extracted these data  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 recorded at the same timing of ventilator waveform analysis. We defined desaturation as  
8  
9  
10 a decrease in  $SpO_2 < 85\%$  because our setting of the lower limit of  $SpO_2$  was routinely  
11  
12  
13 85% as part of our respiratory strategy.  
14  
15  
16  
17  
18  
19

### 20 **Data analysis**

21  
22  
23 Edi waveforms every 10 seconds were assessed and classified into four patterns by  
24  
25  
26 only one investigator (blinded clinical data). When only one waveform pattern (e.g.,  
27  
28  
29 phasic pattern) was shown in a window, this window was classified in this waveform  
30  
31  
32 pattern (e.g., phasic pattern) and counted with one point. When there were multiple  
33  
34  
35 patterns in one window (e.g., central apnea pattern and irregular low-voltage pattern;  
36  
37  
38 Figure 2 online), one point was distributed to each pattern equally (e.g., 0.5 points each  
39  
40  
41 in the central apnea pattern and irregular low-voltage pattern). We summed the points of  
42  
43  
44 each waveform pattern every 5 minutes (total of 30 points) and calculated the  
45  
46  
47 proportion (%) of the points (by dividing by 30). In each patient, data for the first 15  
48  
49  
50  
51  
52  
53 hours of data collection were analyzed.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 To investigate the association between Edi waveform patterns and SpO<sub>2</sub>, we analyzed  
7  
8  
9  
10 the dataset every 5 minutes as one data unit. Some Edi waveform patterns might not  
11  
12  
13 affect SpO<sub>2</sub> immediately, but have an effect later. Therefore, the association between  
14  
15  
16 Edi waveform patterns and SpO<sub>2</sub> in 1 or 2 minutes cannot reflect the truth. One data unit  
17  
18  
19 of the dataset every 5 minutes included the proportion of each waveform pattern and  
20  
21  
22 SpO<sub>2</sub>. There were 180 data units in each extremely preterm infant because we assessed  
23  
24  
25  
26  
27 15 hours of data.

28  
29  
30 We divided all data units into the following two groups. The Desat (+) group was  
31  
32  
33 defined as when desaturation was observed in even only once. The Desat (-) group was  
34  
35  
36 defined as when no desaturation was observed in any time. We compared the proportion  
37  
38  
39 of each waveform pattern between the two groups. SpO<sub>2</sub> was also compared between  
40  
41  
42  
43 the two groups.

44  
45  
46 In addition, to evaluate the association between the Edi waveform pattern and  
47  
48  
49 ventilatory parameters, we analyzed the data using another method. We calculated the  
50  
51  
52 median proportion of all data units in each waveform pattern. In each waveform pattern,  
53  
54  
55  
56 all data units were divided into two groups as the high frequency (H-) group and the low  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 frequency (L-) group (e.g., H-phasic group or L-phasic group) by each median  
7  
8  
9  
10 proportion. The H- group and L- group of each waveform pattern were defined as data  
11  
12  
13 units that were higher and lower, respectively, than the median proportion of all data  
14  
15  
16 units. If one data unit had a higher proportion than the median proportion in the phasic  
17  
18  
19 pattern, this data unit was classified as the H-phasic group. In the same data unit, if it  
20  
21  
22 had a lower proportion than the median proportion in the central apnea pattern, this data  
23  
24  
25 unit was also classified as the L-apnea group. We compared SpO<sub>2</sub>, frequency of  
26  
27  
28 desaturation, and ventilatory parameters between the H- and L- groups in each  
29  
30  
31  
32  
33 waveform pattern.

34  
35  
36 The study protocol was approved by the local medical ethics committee (reference  
37  
38  
39 number R2154).

### 46 **Statistical analysis**

47  
48  
49 Statistical analysis was performed with R statistical software version 4.0.3 (R  
50  
51  
52 Foundation for Statistical Computing, Vienna, Austria) using the EZR application.

53  
54  
55  
56 Actual ventilator setting data are expressed as the mean  $\pm$  standard deviation (SD). The  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 other data are expressed as the median (interquartile range [IQR]) and n (%).  
7  
8

9  
10 Continuous variables were analyzed using the Mann-Whitney test. We defined  
11

12  
13 significant differences as  $p < 0.05$ .  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## RESULTS

### Patients' characteristics

Ten extremely preterm infants were included in our study. Of these, one preterm infant was excluded because she was extubated before being ventilated with NAVA for 24 consecutive hours. Another two preterm infants were excluded because more than 15 hours of data could not be collected when they changed their ventilatory support from NAVA. Therefore, we analyzed the remaining seven preterm infants. The patients' clinical data are shown in Table 1. The preterm infants were born at 24 or 26 gestational weeks and weighed between 503 g and 1128 g. They were ventilated with NAVA at approximately 2 or 3 weeks after birth. In total, we analyzed collected data for 105 hours (1260 data units). The actual ventilatory settings for all data units are also shown in Table 1.

### Edi waveform patterns and desaturation

Comparison of the proportion of each waveform pattern between the Desat (+) and (–) groups is shown in Figure 3. The Desat (+) group included 261 data units and the Desat

1  
2  
3  
4  
5  
6  
7 (-) group included 999 data units. The proportion of the phasic pattern in the Desat (+)  
8  
9  
10 group was significantly lower than that in the Desat (-) group ( $p < 0.001$ ). However, the  
11  
12 proportions of the central apnea, irregular low-voltage, and tonic burst patterns in the  
13  
14 Desat (+) group were significantly higher than those in the Desat (-) group ( $p < 0.001$ ,  $p$   
15  
16  $< 0.001$ , and  $p = 0.039$ , respectively). SpO<sub>2</sub> in the Desat (+) group were significantly  
17  
18 lower than that in the Desat (-) group (Desat (+) group: 88% (86%, 89%), Desat (-)  
19  
20 group: 95% (92%, 97%),  $p < 0.001$ ).  
21  
22  
23  
24  
25  
26  
27  
28  
29

### 30 **Edi waveform patterns and ventilatory parameters**

31  
32  
33 The median proportions in all data units were 53% (42%, 71%) for the phasic pattern,  
34  
35  
36 22% (12%, 37%) for the central apnea pattern, 8.9% (5%, 15%) for the irregular low-  
37  
38  
39 voltage pattern, and 9.4% (4.7%, 16%) for the tonic burst pattern.  
40  
41  
42

43 In every waveform pattern, these median proportions were used as thresholds for  
44  
45  
46 dividing data units into the H- and L- groups. (e.g., in one unit, if the phasic pattern was  
47  
48  
49 62%, the central apnea pattern was 13%, the irregular low-voltage pattern was 12%, and  
50  
51  
52 the tonic burst pattern was 13%, they were classified into the H-phasic group, the L-  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 apnea group, the H-irregular low-voltage group, and the H-tonic burst group,  
8  
9  
10 respectively).

11  
12  
13 SpO<sub>2</sub> in the H- and L- groups in each waveform pattern are shown in Table 2. SpO<sub>2</sub> in  
14  
15  
16 the H-phasic group was significantly higher than that in the L-phasic group ( $p < 0.001$ ).

17  
18  
19 However, SpO<sub>2</sub> was significantly lower in the H-apnea group than in the L-apnea group

20  
21  
22 ( $p < 0.001$ ) and also lower in the H-irregular low-voltage group than in the L-irregular

23  
24  
25 low-voltage group ( $p < 0.001$ ). The presence of desaturation between the H- and L-

26  
27  
28 groups in each waveform pattern is also shown in Table 2. The frequency of

29  
30  
31 desaturation was significantly lower in the H-phasic group than in the L-phasic group ( $p$

32  
33  
34  $< 0.001$ ). However, the frequency of desaturation was significantly higher in the H-

35  
36  
37 apnea group than in the L-apnea group ( $p < 0.001$ ), and higher in the H-irregular low-

38  
39  
40 voltage group than in the L-irregular low-voltage group ( $p < 0.001$ ).

41  
42  
43 Comparison of ventilatory parameters and proportions of each waveform pattern

44  
45  
46 between the H- and L- groups of each waveform pattern are also shown in Table 2. Edi

47  
48  
49 peak in the H-phasic group was significantly higher than that in the L-phasic group ( $p <$

50  
51  
52  $0.001$ ). Time in backup ventilation in the H-phasic group were significantly lower than

1  
2  
3  
4  
5  
6 those in the L-phasic group ( $p < 0.001$ ). On the other hand, Edi peak in the H-irregular  
7  
8  
9 low-voltage group was significantly lower than that in the L-irregular low-voltage group  
10  
11  
12 ( $p < 0.001$ ). Peak airway pressure in the H-irregular low-voltage group was significantly  
13  
14  
15 lower than that in the L-irregular low-voltage group ( $p < 0.001$ ). Time in backup  
16  
17  
18 ventilation was not different between the H- and L-irregular low-voltage groups.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## DISCUSSION

Several studies have reported characterization of breathing patterns as shown by an Edi catheter in preterm infants<sup>19,20</sup>. However, the association between breathing patterns and respiratory conditions was not shown in these reports. This is the first report to evaluate the effect of Edi waveform patterns on SpO<sub>2</sub> and the frequency of desaturation. Our results indicated that Edi waveform patterns were associated with the frequency of desaturation. Therefore, analysis of Edi waveform patterns could be an important indicator for appropriately using NAVA in extremely preterm infants.

We found that patients had a stable SpO<sub>2</sub> when the phasic pattern was frequently observed. However, when an irregular low-voltage, central apnea, or tonic burst pattern was frequently observed, patients showed unstable SpO<sub>2</sub>. These results suggest that a high frequency of an irregular low-voltage, central apnea, or tonic burst pattern is a cause of failure to ventilate with NAVA.

In this study, we defined the central apnea pattern as continuous flat Edi, which induces backup mandatory ventilation, to appropriately assess the effect of central apnea. In contrast, we defined a low Edi peak waveform as an irregular low-voltage

1  
2  
3  
4  
5  
6 pattern. This waveform pattern has not been defined in previous reports. A low Edi peak  
7  
8  
9 waveform (Edi [peak-min] < 1.5  $\mu$ V) prevents backup ventilation, but it supplies low  
10  
11  
12 peak inspiratory pressure <sup>21</sup>. Peak airway pressure supplied by NAVA is calculated  
13  
14 according to the following formula: NAVA level (cmH<sub>2</sub>O/ $\mu$ V)×(Edi peak [ $\mu$ V]-Edi  
15  
16 minimum [ $\mu$ V])+PEEP (cmH<sub>2</sub>O). During this study, an efficient observed peak  
17  
18  
19 inspiratory pressure was approximately 15 cmH<sub>2</sub>O (IQR: 14–16 cmH<sub>2</sub>O). Because we  
20  
21  
22 set the NAVA level at approximately  $1.6 \pm 0.4$  and PEEP at 7 cmH<sub>2</sub>O, peak airway  
23  
24  
25 pressure supplied by an irregular low-voltage pattern was calculated as < 10 cmH<sub>2</sub>O  
26  
27  
28 (1.6×1.5+PEEP). In reality, peak airway pressure with a low Edi peak waveform  
29  
30  
31 observed in this study was < 10 cmH<sub>2</sub>O. This low peak inspiratory pressure appeared to  
32  
33  
34 provide insufficient support. Our study indicated that a high frequency of the irregular  
35  
36  
37 low-voltage pattern lead to low peak airway pressure generated with low Edi peak. Our  
38  
39  
40 study also showed that time in backup ventilation was not different between H- and L-  
41  
42  
43 irregular low-voltage groups. These results indicate that the irregular low-voltage  
44  
45  
46 pattern may prevent backup ventilation, but supply insufficient peak airway pressure.  
47  
48  
49  
50 Therefore, irregular low-voltage pattern could lead to desaturation.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 In addition, a high frequency of the tonic burst pattern was associated with slightly  
8  
9  
10 higher peak airway pressure generated with slightly higher Edi peak. In contrast, the  
11  
12  
13 tonic burst pattern was more frequently observed in the Desat (+) group than in the  
14  
15  
16 Desat (-) group. This contradiction arose because the tonic burst pattern is an irregular  
17  
18  
19 waveform and is considered to be an asynchronized pattern. Therefore, the tonic burst  
20  
21  
22 pattern was also considered to result in desaturation. On the other hand, as tonic burst  
23  
24  
25 pattern was known to be an essential respiratory pattern for premature infants to recruit  
26  
27  
28 their lung<sup>22,23</sup>, desaturation may lead infants to breath with tonic burst pattern. Further  
29  
30  
31 research about the effect of tonic burst pattern for respiratory condition would be  
32  
33  
34  
35  
36  
37 needed.

38  
39  
40 Ventilatory parameters from the ventilator did not include information of the  
41  
42  
43 waveform pattern. Therefore, the ventilatory parameters especially of Edi peak and peak  
44  
45  
46 airway pressure could include all waveform patterns. Even if these parameters were  
47  
48  
49 indicated as the same value, the meaning of the value was different among the different  
50  
51  
52 Edi waveform patterns. When the Edi peak was 5  $\mu$ V in the phasic pattern, the  
53  
54  
55 ventilator may provide stable support, but when that in the irregular low-voltage pattern  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 or tonic burst pattern the ventilator may supply ineffective or asynchronized support and  
7  
8  
9  
10 cause desaturation. Therefore, desaturation could not be predicted by only using these  
11  
12  
13 ventilatory parameters. Consequently, the Edi waveform pattern is considered to be  
14  
15  
16 important for assessing desaturation.  
17  
18

19  
20 To exclude the effect of changing the ventilator setting for the results, our data were  
21  
22  
23 collected when the ventilator setting had not been changed within at least 3 hours to  
24  
25  
26 confirm the stability of the respiratory condition with the ventilator setting at that time.  
27  
28  
29 In addition, because the ventilator setting did not vary much, we consider that our  
30  
31  
32 results did not depend on the ventilator setting, but on the variability of Edi waveform  
33  
34  
35 patterns instead.  
36  
37  
38

39  
40 There are some limitations of our study. The main limitation was the limited number  
41  
42  
43 of cases. Additionally, because the gestational age and duration of postconceptional age  
44  
45  
46 with ventilation by NAVA of the included infants had a limited range, our results might  
47  
48  
49 not be able to be applied for more mature infants. However, because we were able to  
50  
51  
52 evaluate many data units, the similarity of the patients' characteristics was a strength of  
53  
54  
55 this study. Further studies are required to validate our results.  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 In future studies, we need to identify the appropriate proportion of Edi waveform  
8  
9  
10 patterns that result in stable SpO<sub>2</sub> of infants. The criterion that indicates when the  
11  
12  
13 attending physician should change the ventilator mode from NAVA to another mode  
14  
15  
16 also needs to be investigated.  
17  
18  
19  
20  
21  
22

## 23 **CONCLUSION**

24  
25  
26 We assessed the association between Edi waveform patterns and SpO<sub>2</sub>. Proportions of  
27  
28  
29 Edi waveform patterns affect SpO<sub>2</sub> in extremely preterm infants. Our results indicate  
30  
31  
32 that analysis of Edi waveform patterns is an important indicator for assessing whether  
33  
34  
35 preterm infants are appropriately ventilated with NAVA.  
36  
37  
38  
39  
40  
41  
42

## 43 **Acknowledgment**

44  
45  
46 We thank Ellen Knapp, PhD, from Edanz Group ([https://en-author-](https://en-author-services.edanzgroup.com/ac)  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
services.edanzgroup.com/ac) for editing a draft of this manuscript.

**REFERENCES**

1. Stoll BJ, Hansen NI, Bell EF, Walsh MC, Carlo WA, Shankaran S, Laptook AR, Sánchez PJ, Van Meurs KP, Wyckoff M, et al. Trends in care practices, morbidity, and mortality of extremely preterm neonates, 1993-2012. *JAMA* 2015;314:1039–1051.
2. Jensen EA, Dysart K, Gantz MG, McDonald S, Bamat NA, Keszler M, Kirpalani H, Laughon MM, Poindexter BB, Duncan AF, et al. The diagnosis of bronchopulmonary dysplasia in very preterm infants. An evidence-based approach. *Am J Respir Crit Care Med* 2019;200:751–759.
3. Walsh MC, Morris BH, Wrage LA, Vohr BR, Poole WK, Tyson JE, Wright LL, Ehrenkranz RA, Stoll BJ, Fanaroff AA. Extremely low birthweight neonates with protracted ventilation: mortality and 18-month neurodevelopmental outcomes. *J Pediatr* 2005;146:798–804.
4. Mally PV, Beck J, Sinderby C, Caprio M, Bailey SM. Neural breathing pattern and patient-ventilator interaction during neurally adjusted ventilatory assist and conventional ventilation in newborns. *Pediatr Crit Care Med* 2018;19:48–55.

- 1  
2  
3  
4  
5  
6  
7 5. Karikari S, Rausa J, Flores S, Loomba RS. Neurally adjusted ventilatory assist versus  
8  
9  
10 conventional ventilation in the pediatric population: Are there benefits? *Pediatr*  
11  
12  
13 *Pulmonol* 2019;54:1374–1481.  
14
- 15  
16 6. Clement KC, Thurman TL, Holt SJ, Heulitt MJ. Neurally triggered breaths reduce  
17  
18  
19 trigger delay and improve ventilator response times in ventilated infants with  
20  
21  
22 bronchiolitis. *Intensive Care Med* 2011;37:1826–1832.  
23  
24
- 25  
26 7. Alander M, Peltoniemi O, Pokka T, Kontiokari T. Comparison of pressure-, flow-,  
27  
28  
29 and NAVA-triggering in pediatric and neonatal ventilatory care. *Pediatr Pulmonol*  
30  
31  
32 2012;47:76–83.  
33  
34
- 35  
36 8. Bengtsson JA, Edberg KE. Neurally adjusted ventilatory assist in children: an  
37  
38  
39 observational study. *Pediatr Crit Care Med* 2010;11:253–257.  
40  
41  
42
- 43  
44 9. Breatnach C, Conlon NP, Stack M, Healy M, O’Hare BP. A prospective crossover  
45  
46  
47 comparison of neurally adjusted ventilatory assist and pressure-support ventilation in  
48  
49  
50 a pediatric and neonatal intensive care unit population. *Pediatr Crit Care Med*  
51  
52  
53 2010;11:7–11.  
54  
55
- 56  
57 10. Stein H, Alosch H, Ethington P, White DB. Prospective crossover comparison between  
58  
59  
60

- 1  
2  
3  
4  
5  
6 NAVA and pressure control ventilation in premature neonates less than 1500 grams.  
7  
8  
9  
10 J Perinatol 2013; 33:452–456.  
11  
12
- 13 11. Rosterman JL, Pallotto EK, Truog WE, Escobar H, Meinert KA, Holmes A, Dai H,  
14  
15  
16 Manimtim WM. The impact of neurally adjusted ventilatory assist mode on  
17  
18  
19 respiratory severity score and energy expenditure in infants: a randomized crossover  
20  
21  
22  
23 trial. J Perinatol 2018; 38:59–63.  
24  
25
- 26 12. Kallio M, Koskela U, Peltoniemi O, Kontiokari T, Pokka T, Suo-Palosaari, et al.  
27  
28  
29 Neurally adjusted ventilatory assist (NAVA) in preterm newborn infants with  
30  
31  
32  
33 respiratory distress syndrome-a randomized controlled trial. Eur J Pediatr 2016;  
34  
35  
36  
37 175:1175–1183.  
38  
39
- 40 13. Oda A, Kamei Y, Hiroma T, Nakamura T. Neurally adjusted ventilatory assist in  
41  
42  
43  
44 extremely low-birthweight infants. Pediatr Int 2018;60:844–848.  
45
- 46 14. Rong X, Liang F, Li YJ, Liang H, Zhao XP, Zou HM, Lu WN, Shi H, Zhang JH,  
47  
48  
49  
50 Guan RL, et al. Application of neurally adjusted ventilatory assist in  
51  
52  
53  
54 premature neonates less than 1,500 grams with established or evolving  
55  
56  
57  
58 bronchopulmonary dysplasia. Front Pediatr 2020;8:110. doi:  
59  
60



- 1  
2  
3  
4  
5  
6 infants. *Pediatr Res* 2011;70:607–613.  
7  
8  
9
- 10 20. García-Muñoz Rodrigo F, Urquía Martí L, Galán Henríquez G, Rivero Rodríguez,  
11  
12  
13 Hernandez Gomez A. Neural breathing patterns in preterm newborns supported with  
14  
15  
16 non-invasive neurally adjusted ventilatory assist. *J Perinatol* 2018; 38:1235–41.  
17  
18  
19
- 20 21. Narchi H, Chedid F. Neurally adjusted ventilator assist in very low birth weight  
21  
22  
23 infants: Current status. *World J Methodol* 2015;5:62–67.  
24  
25  
26
- 27 22. Emeriaud G, Beck J, Tucci M, Lacroix J, Sinderby C.  
28  
29  
30 Diaphragm electrical activity during expiration in mechanically ventilated infants.  
31  
32  
33 *Pediatr Res* 2006;59:705-710.  
34  
35  
36
- 37 23. Larouche A, Massicotte E, Constantin G, Ducharme-Crevier L, Essouri S, Sinderby  
38  
39  
40 C, Beck J, et al. Tonic diaphragmatic activity in critically ill children with and without  
41  
42  
43 ventilatory support. *Pediatr Pulmonol* 2015;50:1304-1312.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10 **Image legends**  
11  
12  
13  
14  
15

16 Figure 1. Representative waveforms for four Edi waveform patterns.

17  
18  
19  
20 Pressure curves are shown in each upper row and Edi waveforms are shown in each  
21  
22  
23 lower row. A: Phasic pattern, B: central apnea pattern, C: tonic bursts pattern, D:  
24  
25  
26 irregular low-voltage pattern.  
27  
28  
29  
30  
31  
32

33 Figure 2. Example of one window of servo connect including multiple waveform  
34  
35  
36 patterns.  
37  
38  
39

40 A pressure curve is in the upper row and an Edi waveform is in the lower row.

41  
42  
43 Waveforms on the left side were classified as central apnea and waveforms on the right  
44  
45  
46 side were classified as an irregular low-voltage pattern. Each pattern was calculated as  
47  
48  
49 0.5 points.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 Figure 3. Comparison of the proportions each waveform pattern between the

8  
9  
10 Desaturation (+) and (-) groups.

11  
12  
13 Data are expressed as median and interquartile range. The Desat (+) group included 261

14  
15  
16 data units and the Desat (-) group included 999 data units. \*,  $p < 0.05$ .

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

Table 1. Patients' clinical data and ventilatory settings

Clinical data	
Gestational Age (wks)	25.3 (24.5, 25.5)
Birth Weight (g)	626 (549, 790)
Apgar score 1 minute	4 (3, 5)
Apgar score 5 minute	7(6, 7)
DOL at NAVA started	19 (18, 20)
PCA at NAVA started (wks)	28.2 (27.4, 28.4)
Ventilator settings	
NAVA level (cmH <sub>2</sub> O/ $\mu$ V)	1.6 $\pm$ 0.4
PEEP (cmH <sub>2</sub> O)	7.2 $\pm$ 0.4
Apnea time (s)	2.0 $\pm$ 0.6
Pressure level above PEEP in backup (cmH <sub>2</sub> O)	10 $\pm$ 1.5
Backup RR	40 $\pm$ 4.3
FiO <sub>2</sub> (%)	22 $\pm$ 1.4


Patients' clinical data are expressed as the median (interquartile range). Ventilator settings are presented as mean  $\pm$  standard deviation. DOL, day of life; NAVA, neurally adjusted ventilatory assist; PCA, postconceptional age; PEEP, positive end expiratory pressure; RR, respiratory rate.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

Table 2. Comparison between Edi waveform patterns and stability of SpO<sub>2</sub> and ventilatory parameters


	Phasic pattern			Central apnea pattern			Irregular low-voltage pattern			Tonic burst pattern		
	H-	L-	P value	H-	L-	P value	H-	L-	P value	H-	L-	P value
SpO <sub>2</sub> (%)	94 (91, 97)	93 (90, 95)	<0.001	93 (90, 96)	94 (91, 96)	<0.001	93 (90, 96)	94 (91, 96)	<0.001	94 (91, 96)	93 (91, 96)	0.51
Presence of desaturation (%)	16	26	<0.001	25	17	<0.001	26	17	<0.001	22	20	0.531
Edi peak (μV)	5.2 (4.2, 6.3)	4.8 (3.8, 5.9)	<0.001	4.9 (3.9, 5.8)	5.2 (4.2, 6.4)	<0.001	4.7 (3.5, 5.7)	5.3 (4.5, 6.4)	<0.001	5.2 (4.2, 6.3)	4.9 (3.9, 5.8)	<0.001
Edi min (μV)	1.3 (1.0, 1.7)	1.4 (1.0, 1.9)	0.026	1.3 (1.0, 1.8)	1.3 (1.0, 1.8)	0.964	1.3 (0.9, 1.8)	1.4 (1.0, 1.8)	0.11	1.4 (1.1, 1.9)	1.2 (0.9, 1.7)	<0.001
Peak airway Pressure (cmH <sub>2</sub> O)	15 (15, 16)	15 (14, 16)	0.001	15 (14, 16)	15 (14, 16)	0.47	15 (14, 16)	16 (15, 17)	<0.001	15 (14, 17)	15 (14, 16)	<0.001
Mean airway Pressure (cmH <sub>2</sub> O)	10 (9.6, 10)	10 (9.6, 10)	0.078	10 (9.6, 10)	10 (9.6, 10)	0.34	10 (9.6, 10)	10 (9.7, 10)	<0.001	10 (9.7, 10)	10 (9.6, 10)	0.16
Time in backup (%)	14 (6.0, 28)	19 (9.0, 37)	<0.001	19 (8.7, 37)	15 (6.3, 28)	<0.001	16 (7.0, 35)	15 (6.7, 31)	0.055	15 (7.3, 30)	18 (7.7, 35)	0.028
FiO <sub>2</sub> (%)	21 (21, 22)	21 (21, 23)	0.014	21 (21, 22)	21 (21, 22)	0.364	21 (21, 23)	21 (21, 22)	0.019	21 (21, 23)	21 (21, 22)	<0.001
Median of Phasic pattern (%)	67 (60, 76)	35 (23, 45)	<0.001	37 (23, 48)	66 (57, 76)	<0.001	43 (29, 58)	62 (47, 74)	<0.001	52 (36, 64)	54 (34, 72)	0.003
Median of Central apnea pattern (%)	13 (7.5, 19)	37 (26, 51)	<0.001	37 (28, 52)	12 (7.2, 17)	<0.001	26 (15, 41)	19 (10, 34)	<0.001	19 (11, 30)	26 (13, 45)	<0.001
Median of Irregular low-voltage pattern (%)	6.1 (3.3, 11)	13 (7.8, 21)	<0.001	11 (6.1, 18)	7.8 (3.9, 14)	<0.001	17 (13, 22)	2.0 (2.8, 7.2)	<0.001	8.9 (4.4, 15)	10 (5.0, 17)	0.026
Median of Tonic burst pattern (%)	8.9 (5.0, 13)	9.4 (4.4, 16)	0.11	7.2 (4.2, 13)	10 (5.6, 17)	<0.001	8.3 (4.4, 14)	9.4 (5.0, 16)	0.002	15 (12, 20)	5.0 (2.8, 6.7)	<0.001

Data are expressed as the median (interquartile range). Edi, electrical activity of the diaphragm; H-, high frequency; L-, low frequency.



Caption: Representative waveforms for four Edi waveform patterns. Pressure curves are shown in each upper row and Edi waveforms are shown in each lower row. A: Phasic pattern, B: central apnea pattern, C: tonic bursts pattern, D: irregular low-voltage pattern.


254x190mm (72 x 72 DPI)



31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Caption : Example of one window of servo connect including multiple waveform patterns. A pressure curve is in the upper row and an Edi waveform is in the lower row. Waveforms on the left side were classified as central apnea and waveforms on the right side were classified as an irregular low-voltage pattern. Each pattern was calculated as 0.5 points.

254x190mm (72 x 72 DPI)



Data are expressed as median and interquartile range. The Desat (+) group included 261 data units and the Desat (-) group included 999 data units. \*, p<0.05.

338x190mm (54 x 54 DPI)

1  
2  
3  
4  
5  
6  
7 **Effect of electrical activity of the diaphragm waveform patterns on SpO<sub>2</sub> for**

8  
9  
10 **extremely preterm infants ventilated with neurally adjusted ventilatory assist**

11  
12  
13  
14  
15  
16 Ryosuke Araki MD; Seiichi Tomotaki MD; Mitsuyo Akita MD; Kouji Motokura MD;

17  
18  
19 Yutaro Tomobe MD; Takeru Yamauchi MD; Shintaro Hanaoka MD; Hiroko Tomotaki

20  
21  
22 MD; Kogoro Iwanaga MD; Fusako Niwa MD, PhD; Junko Takita MD, PhD; Masahiko

23  
24  
25  
26 Kawai MD, PhD

27  
28  
29 Department of Pediatrics, Graduate School of Medicine, Kyoto University, 54

30  
31  
32 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan.

33  
34  
35  
36  
37  
38  
39 **Corresponding Author:** Ryosuke Araki, Department of Pediatrics, Graduate School of

40  
41  
42 Medicine, Kyoto University, 54 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507,

43  
44  
45  
46 Japan.

47  
48  
49 E-mail: r\_araki@kuhp.kyoto-u.ac.jp; Tel: +81-75-751-3291; Fax: +81-75-751-2361

50  
51  
52  
53  
54  
55  
56 **Conflicts of Interest and Source of Funding:** None declared.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10 **Keywords:** bronchopulmonary dysplasia; neonatal intensive care units; diaphragm;  
11  
12  
13 lung injury.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## Abstract

### Objective

This study aimed to evaluate the association between electrical activity of the diaphragm (Edi) waveform patterns and peripheral oxygen saturation (SpO<sub>2</sub>) in extremely preterm infants who are ventilated with neurally adjusted ventilatory assist (NAVA).

### Study Design

We conducted a retrospective cohort study at a level III neonatal intensive care unit. Extremely preterm infants born at our hospital between November 2019 and November 2020 and ventilated with NAVA were included. We collected Edi waveform data and classified them into four Edi waveform patterns, including the phasic pattern, central apnea pattern, irregular low-voltage pattern, and tonic burst pattern. We analyzed the Edi waveform pattern for the first 15 hours of collectable data in each patient. To investigate the association between Edi waveform patterns and SpO<sub>2</sub>, we analyzed the dataset every 5 minutes as one data unit. We compared the proportion of each waveform pattern between the desaturation (Desat (+)) and non-desaturation (Desat (-)) groups.

## Results

We analyzed collected data for 105 hours (1260 data units). The proportion of the phasic pattern in the Desat (+) group was significantly lower than that in the Desat (-) group ( $p < 0.001$ ). However, the proportions of the central apnea, irregular low-voltage, and tonic burst patterns in the Desat (+) group were significantly higher than those in the Desat (-) group (all  $p < 0.05$ ).

## Conclusion

Our results indicate that proportions of Edi waveform patterns have an effect on desaturation of SpO<sub>2</sub> in extremely preterm infants who are ventilated with NAVA.

## INTRODUCTION

Recent advancement in perinatal management has improved the survival rate of extremely preterm infants. Consequently, bronchopulmonary dysplasia (BPD) has become one of the most common and severe sequelae for extremely preterm infants<sup>1,2</sup> and management for preventing BPD has been the focus of attention. Extremely preterm infants require mechanical ventilation for a relatively long period because of prematurity of their lungs. In such a long mechanical ventilation period, high ventilatory pressure and patient–ventilator asynchrony are important factors for BPD<sup>3,4</sup>. Therefore, the strategy of ventilation for extremely preterm infants has focused on minimizing inspiratory pressure and more synchronized ventilation.

Neurally adjusted ventilatory assist (NAVA) is a relatively novel mode of mechanical ventilation and provides respiratory support in response to the patient’s electrical activity of the diaphragm (Edi). Compared with other patient-triggered ventilator modes, NAVA improves patient–ventilatory synchrony and reduces peak inspiratory pressure and the work of breathing without an elevated fraction of inspiratory oxygen ( $\text{FiO}_2$ )<sup>5–9</sup>. There has been a recent increase in reports on ventilation with NAVA for

1  
2  
3  
4  
5  
6 preterm and low birth weight infants<sup>10-13</sup>. NAVA is expected to improve the prognosis  
7  
8  
9  
10 of extremely preterm infants, especially in proceeding to BPD<sup>14,15</sup>. However, despite  
11  
12  
13 appropriate ventilatory settings, some preterm infants cannot be appropriately supported  
14  
15  
16 by NAVA. These infants develop an unstable respiratory condition or need to have their  
17  
18  
19 ventilatory mode changed from NAVA<sup>16-18</sup>. However, there are limited clinical  
20  
21  
22  
23 indicators of whether infants can be appropriately ventilated by NAVA.  
24  
25

26  
27 During ventilation with NAVA, respiratory support is supplied from a ventilator on the  
28  
29  
30 basis of the Edi waveform. Preterm infants are known to represent variable Edi  
31  
32  
33 waveform patterns<sup>19,20</sup>, but there are limited data on the association between Edi  
34  
35  
36 waveform patterns and respiratory conditions. On the basis of our clinical experience,  
37  
38  
39 we consider that not only frequent central apnea, but also low inspiratory movements  
40  
41  
42 prevent appropriate ventilatory support by NAVA.  
43  
44  
45

46 We hypothesize that Edi waveform patterns reflect the respiratory pattern, which  
47  
48  
49 affects appropriate ventilatory support by NAVA and the frequency of desaturation of  
50  
51  
52 peripheral oxygen saturation (SpO<sub>2</sub>).  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 In this study, we evaluated the association between Edi waveform patterns and the  
8  
9  
10 frequency of desaturation in extremely preterm infants who are ventilated with NAVA.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## MATERIALS AND METHODS

### Patients

We conducted a retrospective cohort study at a single level III neonatal intensive care unit (Kyoto University Hospital, Kyoto, Japan). Extremely preterm infants of < 28 weeks gestational age who were born at our hospital between November 2019 and November 2020 and required mechanical ventilation with NAVA delivered by Servo-n<sup>®</sup> (Maquet, Critical Care AB, Solna, Sweden) were eligible for our study. Infants who needed to have their ventilation mode changed from NAVA to another mode (including those who were extubated) within 24 consecutive hours or those who did not have more than 15 hours of data collected were excluded.

### Study procedure

~~To properly use NAVA~~ According to our usual practice, the NAVA level was set and adjusted to maintain Edi peak < 15  $\mu$ V and positive end-expiratory pressure (PEEP) was set and adjusted to maintain Edi minimum at < 5  $\mu$ V. The backup support setting and apnea time were adjusted to supply proper backup ventilation to avoid desaturation.

1  
2  
3  
4  
5  
6  
7 Because we routinely adjust the NAVA ventilator settings with this procedure, the  
8  
9  
10 initial settings were usually the same as follows: NAVA level = 1.5 to 2 cmH<sub>2</sub>O/ $\mu$ V,  
11  
12  
13 PEEP = 7 cmH<sub>2</sub>O, apnea time = 2 seconds, pressure level above PEEP in backup = 10  
14  
15  
16 cmH<sub>2</sub>O, and back up respiratory rate = 40. ~~Subsequent setting with adjustment by this~~  
17  
18  
19 ~~procedure did not vary in each patient.~~ By this procedure of adjustment, subsequent  
20  
21  
22 ~~ventilator setting after the initial setting also did not vary in each patient.~~ In accordance  
23  
24  
25  
26 with the manufacturer's recommendations, proper positioning of the Edi catheter was  
27  
28  
29 frequently verified by attending physicians.  
30  
31

32  
33 The waveforms of ventilator pressure, flow, and Edi were acquired from the ventilator  
34  
35  
36 and recorded continuously by using the software SERVVO CONNECT (Fukuda Denshi  
37  
38  
39 Co. Ltd., Tokyo, Japan). We routinely use this software and collect data when possible.  
40  
41  
42  
43  
44  
45

#### 46 **Data collection**

47  
48  
49 Demographic and clinical data of included infants were extracted from medical  
50  
51  
52 records. Valid data for analysis of Edi waveforms and SpO<sub>2</sub> were defined as when the  
53  
54  
55 infants had not been manipulated and the ventilator setting had not been changed at  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 least within 3 hours, except for  $\text{FiO}_2$ . With SERVO CONNECT, Edi waveforms were  
7  
8  
9  
10 shown in one window every 10 seconds. Edi waveforms were classified into four  
11  
12  
13 patterns including those previously reported and a novel pattern<sup>19,20</sup> (Figure 1). These  
14  
15  
16 patterns were as follows: 1) a phasic pattern was defined as a regular increasing and  
17  
18  
19 decreasing pattern, 2) a central apnea pattern was defined as continuous low Edi, which  
20  
21  
22 induced backup mandatory ventilation, 3) a tonic burst pattern was defined as elevated  
23  
24  
25 tonic activity without clear phasic activity or with irregular activity, and 4) an irregular  
26  
27  
28 low-voltage pattern was defined as low Edi activity ( $\text{Edi} [\text{peak-minimum}] < 1.5 \mu\text{V}$ ).  
29  
30  
31  
32  
33 Because irregular low-voltage patterns did not induce backup ventilation and generated  
34  
35  
36 low peak inspiratory pressure, we distinguished irregular low-voltage patterns from  
37  
38  
39 central apnea patterns.  
40  
41  
42

43 Ventilatory settings, including the NAVA level, PEEP, backup setting, and  $\text{FiO}_2$ , were  
44  
45  
46 collected from the medical records. Ventilatory parameters, including Edi peak, Edi  
47  
48  
49 minimum, peak airway pressure, mean airway pressure, and time in backup ventilation  
50  
51  
52  
53 were collected from stored data in the ventilators.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 SpO<sub>2</sub> data were recorded by a central monitoring system (Phillips Information  
8  
9 Management System<sup>®</sup>; Phillips Japan, Japan) every minute. We extracted these data  
10  
11 recorded at the same timing of ventilator waveform analysis. We defined desaturation as  
12  
13 a decrease in SpO<sub>2</sub> < 85% because our setting of the lower limit of SpO<sub>2</sub> was routinely  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

85% as part of our respiratory strategy.

### Data analysis

Edi waveforms every 10 seconds were assessed and classified into four patterns by only one investigator (blinded clinical data). When only one waveform pattern (e.g., phasic pattern) was shown in a window, this window was classified in this waveform pattern (e.g., phasic pattern) and counted with one point. When there were multiple patterns in one window (e.g., central apnea pattern and irregular low-voltage pattern; Figure 2 online), one point was distributed to each pattern equally (e.g., 0.5 points each in the central apnea pattern and irregular low-voltage pattern). We summed the points of each waveform pattern every 5 minutes (total of 30 points) and calculated the

1  
2  
3  
4  
5  
6 proportion (%) of the points (by dividing by 30). In each patient, data for the first 15  
7  
8  
9 hours of data collection were analyzed.  
10  
11  
12

13 To investigate the association between Edi waveform patterns and SpO<sub>2</sub>, we analyzed  
14  
15  
16 the dataset every 5 minutes as one data unit. Some Edi waveform patterns might not  
17  
18  
19 affect SpO<sub>2</sub> immediately, but have an effect later. Therefore, the association between  
20  
21  
22 Edi waveform patterns and SpO<sub>2</sub> in 1 or 2 minutes cannot reflect the truth. One data unit  
23  
24  
25 of the dataset every 5 minutes included the proportion of each waveform pattern and  
26  
27  
28 SpO<sub>2</sub>. There were 180 data units in each extremely preterm infant because we assessed  
29  
30  
31  
32  
33 15 hours of data.  
34  
35  
36

37 We divided all data units into the following two groups. The Desat (+) group was  
38  
39  
40 defined as when desaturation was observed in even only once. The Desat (–) group was  
41  
42  
43 defined as when no desaturation was observed in any time. We compared the proportion  
44  
45  
46 of each waveform pattern between the two groups. SpO<sub>2</sub> was also compared between  
47  
48  
49 the two groups.  
50  
51  
52

53 In addition, to evaluate the association between the Edi waveform pattern and  
54  
55  
56 ventilatory parameters, we analyzed the data using another method. We calculated the  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 median proportion of all data units in each waveform pattern. In each waveform pattern,  
8  
9  
10 all data units were divided into two groups as the high frequency (H-) group and the low  
11  
12  
13 frequency (L-) group (e.g., H-phasic group or L-phasic group) by each median  
14  
15  
16 proportion. The H- group and L- group of each waveform pattern were defined as data  
17  
18  
19 units that were higher and lower, respectively, than the median proportion of all data  
20  
21  
22 units. If one data unit had a higher proportion than the median proportion in the phasic  
23  
24  
25 pattern, this data unit was classified as the H-phasic group. In the same data unit, if it  
26  
27  
28 had a lower proportion than the median proportion in the central apnea pattern, this data  
29  
30  
31 unit was also classified as the L-apnea group. We compared SpO<sub>2</sub>, frequency of  
32  
33  
34 desaturation, and ventilatory parameters between the H- and L- groups in each  
35  
36  
37 waveform pattern.  
38  
39  
40  
41  
42

43 The study protocol was approved by the local medical ethics committee (reference  
44  
45  
46 number R2154).  
47  
48  
49  
50  
51  
52

### 53 **Statistical analysis**

54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 Statistical analysis was performed with R statistical software version 4.0.3 (R  
8  
9  
10 Foundation for Statistical Computing, Vienna, Austria) using the EZR application.  
11  
12  
13 Actual ventilator setting data are expressed as the mean  $\pm$  standard deviation (SD). The  
14  
15  
16 other data are expressed as the median (interquartile range [IQR]) and n (%).  
17  
18  
19  
20 Continuous variables were analyzed using the Mann-Whitney test. We defined  
21  
22  
23 significant differences as  $p < 0.05$ .  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## RESULTS

### Patients' characteristics

Ten extremely preterm infants were included in our study. Of these, one preterm infant was excluded because she was extubated before being ventilated with NAVA for 24 consecutive hours. Another two preterm infants were excluded because more than 15 hours of data could not be collected when they changed their ventilatory support from NAVA. Therefore, we analyzed the remaining seven preterm infants. The patients' clinical data are shown in Table 1. The preterm infants were born at 24 or 26 gestational weeks and weighed between 503 g and 1128 g. They were ventilated with NAVA at approximately 2 or 3 weeks after birth. In total, we analyzed collected data for 105 hours (1260 data units). The actual ventilatory settings for all data units are also shown in Table 1.

### **Edi waveform patterns and desaturation**

Comparison of the proportion of each waveform pattern between the Desat (+) and (-) groups is shown in Figure 3. The Desat (+) group included 261 data units and the Desat (-) group included 999 data units. The proportion of the phasic pattern in the Desat (+) group was significantly lower than that in the Desat (-) group ( $p < 0.001$ ). However, the proportions of the central apnea, irregular low-voltage, and tonic burst patterns in the Desat (+) group were significantly higher than those in the Desat (-) group ( $p < 0.001$ ,  $p < 0.001$ , and  $p = 0.039$ , respectively). SpO<sub>2</sub> in the Desat (+) group were significantly lower than that in the Desat (-) group (Desat (+) group: 88% (86%, 89%), Desat (-) group: 95% (92%, 97%),  $p < 0.001$ ).

### **Edi waveform patterns and ventilatory parameters**

The median proportions in all data units were 53% (42%, 71%) for the phasic pattern, 22% (12%, 37%) for the central apnea pattern, 8.9% (5%, 15%) for the irregular low-voltage pattern, and 9.4% (4.7%, 16%) for the tonic burst pattern.

In every waveform pattern, these median proportions were used as thresholds for dividing data units into the H- and L- groups. (e.g., in one unit, if the phasic pattern was

1  
2  
3  
4  
5  
6 62%, the central apnea pattern was 13%, the irregular low-voltage pattern was 12%, and  
7  
8  
9  
10 the tonic burst pattern was 13%, they were classified into the H-phasic group, the L-  
11  
12  
13 apnea group, the H-irregular low-voltage group, and the H-tonic burst group,  
14  
15  
16 respectively).

17  
18  
19  
20 SpO<sub>2</sub> in the H- and L- groups in each waveform pattern are shown in Table 2. SpO<sub>2</sub> in  
21  
22  
23 the H-phasic group was significantly higher than that in the L-phasic group ( $p < 0.001$ ).  
24  
25  
26 However, SpO<sub>2</sub> was significantly lower in the H-apnea group than in the L-apnea group  
27  
28  
29 ( $p < 0.001$ ) and also lower in the H-irregular low-voltage group than in the L-irregular  
30  
31  
32 low-voltage group ( $p < 0.001$ ). The presence of desaturation between the H- and L-  
33  
34  
35 groups in each waveform pattern is also shown in Table 2. The frequency of  
36  
37  
38 desaturation was significantly lower in the H-phasic group than in the L-phasic group ( $p$   
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
60  
 $< 0.001$ ). However, the frequency of desaturation was significantly higher in the H-  
apnea group than in the L-apnea group ( $p < 0.001$ ), and higher in the H-irregular low-  
voltage group than in the L-irregular low-voltage group ( $p < 0.001$ ).

61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158

1  
2  
3  
4  
5  
6  
7 peak in the H-phasic group was significantly higher than that in the L-phasic group ( $p <$   
8  
9  
10 0.001). Time in backup ventilation in the H-phasic group were significantly lower than  
11  
12 those in the L-phasic group ( $p < 0.001$ ). On the other hand, Edi peak in the H-irregular  
13  
14 low-voltage group was significantly lower than that in the L-irregular low-voltage group  
15  
16 ( $p < 0.001$ ). Peak airway pressure in the H-irregular low-voltage group was significantly  
17  
18 lower than that in the L-irregular low-voltage group ( $p < 0.001$ ). Time in backup  
19  
20 ventilation was not different between the H- and L-irregular low-voltage groups.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## DISCUSSION

Several studies have reported characterization of breathing patterns as shown by an Edi catheter in preterm infants<sup>19,20</sup>. However, the association between breathing patterns and respiratory conditions was not shown in these reports. This is the first report to evaluate the effect of Edi waveform patterns on SpO<sub>2</sub> and the frequency of desaturation. Our results indicated that Edi waveform patterns were associated with the frequency of desaturation. Therefore, analysis of Edi waveform patterns could be an important indicator for appropriately using NAVA in extremely preterm infants.

We found that patients had a stable SpO<sub>2</sub> when the phasic pattern was frequently observed. However, when an irregular low-voltage, central apnea, or tonic burst pattern was frequently observed, patients showed unstable SpO<sub>2</sub>. These results suggest that a high frequency of an irregular low-voltage, central apnea, or tonic burst pattern is a cause of failure to ventilate with NAVA.

In this study, we defined the central apnea pattern as continuous flat Edi, which induces backup mandatory ventilation, to appropriately assess the effect of central apnea. In contrast, we defined a low Edi peak waveform as an irregular low-voltage

1  
2  
3  
4  
5  
6 pattern. This waveform pattern has not been defined in previous reports. A low Edi peak  
7  
8  
9  
10 waveform (Edi [peak-min] < 1.5  $\mu$ V) prevents backup ventilation, but it supplies low  
11  
12  
13 peak inspiratory pressure <sup>21</sup>. Peak airway pressure supplied by NAVA is calculated  
14  
15  
16 according to the following formula: NAVA level (cmH<sub>2</sub>O/ $\mu$ V)×(Edi peak [ $\mu$ V]-Edi  
17  
18  
19 minimum [ $\mu$ V])+PEEP (cmH<sub>2</sub>O). During this study, an efficient observed peak  
20  
21  
22 inspiratory pressure was approximately 15 cmH<sub>2</sub>O (IQR: 14–16 cmH<sub>2</sub>O). Because we  
23  
24  
25 set the NAVA level at approximately  $1.6 \pm 0.4$  and PEEP at 7 cmH<sub>2</sub>O, peak airway  
26  
27  
28 pressure supplied by an irregular low-voltage pattern was calculated as < 10 cmH<sub>2</sub>O  
29  
30  
31 (1.6×1.5+PEEP). In reality, peak airway pressure with a low Edi peak waveform  
32  
33  
34 observed in this study was < 10 cmH<sub>2</sub>O. This low peak inspiratory pressure appeared to  
35  
36  
37 provide insufficient support. Our study indicated that a high frequency of the irregular  
38  
39  
40 low-voltage pattern lead to low peak airway pressure generated with low Edi peak. Our  
41  
42  
43 study also showed that time in backup ventilation was not different between H- and L-  
44  
45  
46 irregular low-voltage groups. These results indicate that the irregular low-voltage  
47  
48  
49 pattern may prevent backup ventilation, but supply insufficient peak airway pressure.  
50  
51  
52  
53  
54  
55  
56 Therefore, irregular low-voltage pattern could lead to desaturation.  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 In addition, a high frequency of the tonic burst pattern was associated with slightly  
8  
9  
10 higher peak airway pressure generated with slightly higher Edi peak. In contrast, the  
11  
12  
13 tonic burst pattern was more frequently observed in the Desat (+) group than in the  
14  
15  
16 Desat (-) group. This contradiction arose because the tonic burst pattern is an irregular  
17  
18  
19 waveform and is considered to be an asynchronized pattern. Therefore, the tonic burst  
20  
21  
22 pattern was also considered to result in desaturation. On the other hand, as tonic burst  
23  
24  
25 pattern was known to be an essential respiratory pattern for premature infants to recruit  
26  
27  
28 their lung<sup>22,23</sup>, desaturation may lead infants to breath with tonic burst pattern. Further  
29  
30  
31 research about the effect of tonic burst pattern for respiratory condition would be  
32  
33  
34  
35  
36  
37 needed.

38  
39  
40 Ventilatory parameters from the ventilator did not include information of the  
41  
42  
43 waveform pattern. Therefore, the ventilatory parameters especially of Edi peak and peak  
44  
45  
46 airway pressure could include all waveform patterns. Even if these parameters were  
47  
48  
49 indicated as the same value, the meaning of the value was different among the different  
50  
51  
52 Edi waveform patterns. When the Edi peak was 5  $\mu$ V in the phasic pattern, the  
53  
54  
55 ventilator may provide stable support, but when that in the irregular low-voltage pattern  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 or tonic burst pattern the ventilator may supply ineffective or asynchronized support and  
7  
8  
9  
10 cause desaturation. Therefore, desaturation could not be predicted by only using these  
11  
12  
13 ventilatory parameters. Consequently, the Edi waveform pattern is considered to be  
14  
15  
16 important for assessing desaturation.  
17  
18

19  
20 To exclude the effect of changing the ventilator setting for the results, our data were  
21  
22  
23 collected when the ventilator setting had not been changed within at least 3 hours to  
24  
25  
26 confirm the stability of the respiratory condition with the ventilator setting at that time.  
27  
28

29  
30 In addition, because the ventilator setting did not vary much, we consider that our  
31  
32  
33 results did not depend on the ventilator setting, but on the variability of Edi waveform  
34  
35  
36 patterns instead.  
37  
38

39  
40 There are some limitations of our study. The main limitation was the limited number  
41  
42  
43 of cases. Additionally, because the gestational age and duration of postconceptional age  
44  
45  
46 with ventilation by NAVA of the included infants had a limited range, our results might  
47  
48  
49 not be able to be applied for more mature infants. However, because we were able to  
50  
51  
52 evaluate many data units, the similarity of the patients' characteristics was a strength of  
53  
54  
55 this study. Further studies are required to validate our results.  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 In future studies, we need to identify the appropriate proportion of Edi waveform  
8  
9  
10 patterns that result in stable SpO<sub>2</sub> of infants. The criterion that indicates when the  
11  
12  
13 attending physician should change the ventilator mode from NAVA to another mode  
14  
15  
16 also needs to be investigated.  
17  
18  
19  
20  
21  
22

## 23 **CONCLUSION**

24  
25  
26 We assessed the association between Edi waveform patterns and SpO<sub>2</sub>. Proportions of  
27  
28  
29 Edi waveform patterns affect SpO<sub>2</sub> in extremely preterm infants. Our results indicate  
30  
31  
32 that analysis of Edi waveform patterns is an important indicator for assessing whether  
33  
34  
35 preterm infants are appropriately ventilated with NAVA.  
36  
37  
38  
39  
40  
41  
42

## 43 **Acknowledgment**

44  
45  
46 We thank Ellen Knapp, PhD, from Edanz Group ([https://en-author-](https://en-author-services.edanzgroup.com/ac)  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
services.edanzgroup.com/ac) for editing a draft of this manuscript.

**REFERENCES**

1. Stoll BJ, Hansen NI, Bell EF, Walsh MC, Carlo WA, Shankaran S, Laptook AR, Sánchez PJ, Van Meurs KP, Wyckoff M, et al. Trends in care practices, morbidity, and mortality of extremely preterm neonates, 1993-2012. *JAMA* 2015;314:1039–1051.
2. Jensen EA, Dysart K, Gantz MG, McDonald S, Bamat NA, Keszler M, Kirpalani H, Laughon MM, Poindexter BB, Duncan AF, et al. The diagnosis of bronchopulmonary dysplasia in very preterm infants. An evidence-based approach. *Am J Respir Crit Care Med* 2019;200:751–759.
3. Walsh MC, Morris BH, Wrage LA, Vohr BR, Poole WK, Tyson JE, Wright LL, Ehrenkranz RA, Stoll BJ, Fanaroff AA. Extremely low birthweight neonates with protracted ventilation: mortality and 18-month neurodevelopmental outcomes. *J Pediatr* 2005;146:798–804.
4. Mally PV, Beck J, Sinderby C, Caprio M, Bailey SM. Neural breathing pattern and patient-ventilator interaction during neurally adjusted ventilatory assist and conventional ventilation in newborns. *Pediatr Crit Care Med* 2018;19:48–55.

- 1  
2  
3  
4  
5  
6  
7 5. Karikari S, Rausa J, Flores S, Loomba RS. Neurally adjusted ventilatory assist versus  
8  
9  
10 conventional ventilation in the pediatric population: Are there benefits? *Pediatr*  
11  
12  
13 *Pulmonol* 2019;54:1374–1481.  
14
- 15  
16 6. Clement KC, Thurman TL, Holt SJ, Heulitt MJ. Neurally triggered breaths reduce  
17  
18  
19 trigger delay and improve ventilator response times in ventilated infants with  
20  
21  
22 bronchiolitis. *Intensive Care Med* 2011;37:1826–1832.  
23  
24
- 25  
26 7. Alander M, Peltoniemi O, Pokka T, Kontiokari T. Comparison of pressure-, flow-,  
27  
28  
29 and NAVA-triggering in pediatric and neonatal ventilatory care. *Pediatr Pulmonol*  
30  
31  
32 2012;47:76–83.  
33  
34
- 35  
36 8. Bengtsson JA, Edberg KE. Neurally adjusted ventilatory assist in children: an  
37  
38  
39 observational study. *Pediatr Crit Care Med* 2010;11:253–257.  
40  
41  
42
- 43  
44 9. Breatnach C, Conlon NP, Stack M, Healy M, O'Hare BP. A prospective crossover  
45  
46  
47 comparison of neurally adjusted ventilatory assist and pressure-support ventilation in  
48  
49  
50 a pediatric and neonatal intensive care unit population. *Pediatr Crit Care Med*  
51  
52  
53 2010;11:7–11.  
54  
55
- 56  
57 10. Stein H, Alosch H, Ethington P, White DB. Prospective crossover comparison between  
58  
59  
60

1  
2  
3  
4  
5  
6  
7 NAVA and pressure control ventilation in premature neonates less than 1500 grams.

8  
9  
10 J Perinatol 2013; 33:452–456.

11  
12  
13 11. Rosterman JL, Pallotto EK, Truog WE, Escobar H, Meinert KA, Holmes A, Dai H,

14  
15  
16 Manimtim WM. The impact of neurally adjusted ventilatory assist mode on  
17  
18 respiratory severity score and energy expenditure in infants: a randomized crossover  
19  
20 trial. J Perinatol 2018; 38:59–63.  
21  
22  
23

24  
25  
26 12. Kallio M, Koskela U, Peltoniemi O, Kontiokari T, Pokka T, Suo-Palosaari, et al.

27  
28  
29 Neurally adjusted ventilatory assist (NAVA) in preterm newborn infants with  
30  
31 respiratory distress syndrome-a randomized controlled trial. Eur J Pediatr 2016;  
32  
33 175:1175–1183.  
34  
35  
36

37  
38  
39 13. Oda A, Kamei Y, Hiroma T, Nakamura T. Neurally adjusted ventilatory assist in

40  
41  
42 extremely low-birthweight infants. Pediatr Int 2018;60:844–848.  
43  
44  
45

46  
47 14. Rong X, Liang F, Li YJ, Liang H, Zhao XP, Zou HM, Lu WN, Shi H, Zhang JH,

48  
49 Guan RL, et al. Application of neurally adjusted ventilatory assist in  
50  
51 premature neonates less than 1,500 grams with established or evolving  
52  
53 bronchopulmonary dysplasia. Front Pediatr 2020;8:110. doi:  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7 10.3389/fped.2020.00110.  
8  
9
- 10 15. Jung YH, Kim HS, Lee J, Shin SH, Kim EK, Choi JH. Neurally adjusted ventilatory  
11  
12 assist in preterm infants with established or evolving bronchopulmonary dysplasia on  
13  
14 high-intensity mechanical ventilatory support: A single-center experience. *Pediatr*  
15  
16 *Crit Care Med* 2016;17:1142–6.  
17  
18  
19
- 20 16. McKinney RL, Keszler M, Truog WE, Norberg M, Sindelar R, Wallström L,  
21  
22 Schulman B, Gien J, Abman SH. Multicenter experience with neurally adjusted  
23  
24 ventilatory assist in infants with severe bronchopulmonary dysplasia. *Am J Perinatol*  
25  
26  
27  
28  
29  
30  
31  
32  
33 24 March 2020. doi: 10.1055/s-0040-1708559.  
34  
35
- 36 17. Stein H, Howard D. Neurally adjusted ventilatory assist in neonates weighing < 1500  
37  
38 grams: a retrospective analysis. *J Pediatr* 2012;160:786–789.  
39  
40  
41  
42
- 43 18. Lee J, Kim HS, Sohn JA, Lee JA, Choi CW, Kim EK, Kim BI, Choi JH. Randomized  
44  
45 crossover study of neurally adjusted ventilatory assist in preterm infants. *J Pediatr*  
46  
47  
48  
49  
50 2012;161:808–813.  
51  
52
- 53 19. Beck J, Reilly M, Grasselli G, Qui H, Slutsky AS, Dunn MS, Sinderby CA.  
54  
55  
56  
57  
58  
59  
60
- Characterization of neural breathing pattern in spontaneously breathing preterm

1  
2  
3  
4  
5  
6 infants. *Pediatr Res* 2011;70:607–613.  
7  
8

9  
10 20. García-Muñoz Rodrigo F, Urquía Martí L, Galán Henríquez G, Rivero Rodríguez,  
11  
12  
13 Hernandez Gomez A. Neural breathing patterns in preterm newborns supported with  
14  
15  
16 non-invasive neurally adjusted ventilatory assist. *J Perinatol* 2018; 38:1235–41.  
17  
18

19  
20 21. Narchi H, Chedid F. Neurally adjusted ventilator assist in very low birth weight  
21  
22  
23 infants: Current status. *World J Methodol* 2015;5:62–67.  
24  
25

26 22. Emeriaud G, Beck J, Tucci M, Lacroix J, Sinderby C.  
27  
28  
29 Diaphragm electrical activity during expiration in mechanically ventilated infants.  
30  
31  
32  
33 *Pediatr Res* 2006;59:705-710.  
34  
35

36 23. Larouche A, Massicotte E, Constantin G, Ducharme-Crevier L, Essouri S, Sinderby  
37  
38  
39 C, Beck J, et al. Tonic diaphragmatic activity in critically ill children with and without  
40  
41  
42  
43 ventilatory support. *Pediatr Pulmonol* 2015;50:1304-1312.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Image legends

Figure 1. Representative waveforms for four Edi waveform patterns.

Pressure curves are shown in each upper row and Edi waveforms are shown in each lower row. A: Phasic pattern, B: central apnea pattern, C: tonic bursts pattern, D: irregular low-voltage pattern.

Figure 2. Example of one window of servo connect including multiple waveform patterns.

A pressure curve is in the upper row and an Edi waveform is in the lower row.

Waveforms on the left side were classified as central apnea and waveforms on the right side were classified as an irregular low-voltage pattern. Each pattern was calculated as 0.5 points.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 3. Comparison of the proportions each waveform pattern between the

Desaturation (+) and (-) groups.

Data are expressed as median and interquartile range. The Desat (+) group included 261

data units and the Desat (-) group included 999 data units. \*,  $p < 0.05$ .

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## RESPONSES TO THE REVIEWERS' COMMENTS

We thank the reviewers for reviewing our manuscript. We have carefully considered the comments made by the reviewers and have revised our manuscript accordingly. We believe that our revised manuscript has been improved because of the reviewer's comments. We hope that our revisions and responses to the reviewers' comments are satisfactory.

Reviewer: 2

### Comments to the Author

The manuscript has been improved and I only have minor suggestions:

- Abstract line 30 : we conducted a retrospective cohort study at a level 3...

Response: We thank the reviewer for the comment. We revised the manuscript.

- Page 8, study procedure, line 46, please replace "To properly use NAVA", by something like " According to our usual practice,"...

Response: We thank the reviewer for the comment. We revised the manuscript as suggested.

-page 9, line 17 : I do not understand the sentence " Subsequent setting with adjustment by this procedure did not vary in each patient"... please rephrase.

Response: We rephrased the sentence, as suggested. We thank the reviewer for the suggestion.

- Figure 3: please indicate in the Figure the interquartile ranges (if the bars represent the median values) on the bars... (or the SD if the bars represent the mean values). Please also provide the number of periods in each desat group in the legend. Also, indicate in the legend which measures are illustrated (median or mean).

Response: We thank the reviewer for the comment. We revised the figure 3 and the legend as suggested.