

A criterion of compact sets in L^p -spaces and its application

Katsuhisa Koshino

Faculty of Engineering,
Kanagawa University

1 Introduction

In functional analysis, it is important to recognize compact subsets of function spaces, and criteria like the Ascoli-Arzelà theorem play central roles. In this article, we shall characterize subsets of an L^p -space on a metric measure space to be compact, and as its application, we will investigate the topological structure of the subspace consisting of Lipschitz functions with a bounded support. This is a résumé of the paper [10]. Assume that $X = (X, d, \mu)$ is a Borel-regular Borel metric measure space such that for every $x \in X$ and every $r > 0$, $0 < \mu(B(x, r)) < \infty$, where d is a metric, μ is a measure and $B(x, r)$ is the closed ball centered at x with radius r . Recall that a measure space is *Borel* if its Borel sets are measurable, and that a Borel measure space is *Borel-regular* if any subset is contained in some Borel set with the same measure. For $1 \leq p < \infty$, let $L^p(X) = (L^p(X), \|\cdot\|_p)$ be the L^p -space on X . It is known that $L^p(X)$ is a Banach space. Giving a criterion of compactness in L^p -spaces traces its history back to the Kolmogorov-Riesz theorem [8, 12], which is an L^p -version of the Ascoli-Arzelà one.

Theorem 1.1 (Kolmogorov-Riesz). *A bounded set $F \subset L^p(\mathbb{R}^n)$, where \mathbb{R}^n is the n -dimensional Euclidean space with the Euclidean metric and the Lebesgue measure, is relatively compact if and only if the following are satisfied.*

- (1) *For every $\epsilon > 0$, there is $\delta > 0$ such that $\|\tau_a f - f\|_p < \epsilon$ for any $f \in F$ and any $a \in \mathbb{R}^n$ with $|a| < \delta$.¹*
- (2) *For each $\epsilon > 0$, there exists $r > 0$ such that $\|f \chi_{\mathbb{R}^n \setminus B(\mathbf{0}, r)}\|_p < \epsilon$ for all $f \in F$.²*

Some generalizations in this direction are seen in [6]. Fixing $f \in L^p(X)$ and $r > 0$, define the *average function* $A_r f : X \rightarrow \mathbb{R}$ of f by

$$A_r f(x) = \frac{1}{\mu(B(x, r))} \int_{B(x, r)} f \chi_{B(x, r)}(y) d\mu(y).$$

¹The function $\tau_a f$ is defined by $\tau_a f(x) = f(x - a)$.

²For a subset $E \subset X$, we denote the characteristic function of E by χ_E .

Recently, the Kolmogorov-Riesz type criteria of compactness in L^p -spaces and more generally, in Banach function spaces, have been obtained by use of average functions, refer to [4, 5]. It is said that X is *doubling* if the following condition holds.

- There is $\gamma \geq 1$ such that $\mu(B(x, 2r)) \leq \gamma\mu(B(x, r))$ for any $x \in X$ and any $r > 0$.³

In the paper [10], the following characterization is established.

Theorem 1.2. *Let X be doubling and let F be bounded in $L^p(X)$. Suppose that for each $x \in X$ and each $r > 0$,*

$$\mu(B(x, r) \triangle B(y, r)) \rightarrow 0$$

as $y \rightarrow x$.⁴ Then F is relatively compact if and only if the following conditions are satisfied.

- (1) *For any $\epsilon > 0$, there exists $\delta > 0$ such that $\|A_r f - f\|_p < \epsilon$ for each $f \in F$ and each $r \in (0, \delta)$.*
- (2) *For every $\epsilon > 0$, there is a bounded subset $E \subset X$ such that $\|f\chi_{X \setminus E}\|_p < \epsilon$ for all $f \in F$.*

In the study of topologies on function spaces, typical convex subspaces of Hilbert space ℓ_2 and the Hilbert cube \mathbf{Q} have been detected among them. Due to the efforts by R.D. Anderson [1] and M.I. Kadec [7], we have the following:

Theorem 1.3. *If X is infinite and separable, then $L^p(X)$ is homeomorphic to ℓ_2 .*

It is known that the subspace ℓ_2^f spanned by the canonical orthonormal basis of ℓ_2 is recognized among several function spaces as a factor. For instance, the author [9] investigated the topological type of the subspace $UC(X) \subset L^p(X)$ consisting of uniformly continuous functions.⁵

Theorem 1.4 (K). *Suppose that X is a separable and locally compact metric measure space. If the subset $\{x \in X \mid \mu(\{x\}) \neq 0\}$ is not dense in X , then $UC(X)$ is homeomorphic to the countable product of ℓ_2^f .*

Let $LIP_b(X) \subset L^p(X)$ be the subspace consisting of Lipschitz functions with a bounded support. As an application of Theorem 1.2, the following corollary is obtained in [10].

Corollary 1.5. *Let X be non-degenerate, separable and doubling. If for every $x \in X$, the function*

$$(0, \infty) \ni r \mapsto \mu(B(x, r)) \in (0, \infty)$$

is continuous, then $(L^p(X), LIP_b(X))$ is homeomorphic to $(\ell_2 \times \mathbf{Q}, \ell_2^f \times \mathbf{Q})$.⁶

³The number γ is called the *doubling constant*.

⁴Given subsets $A, B \subset X$, put $A \triangle B = (A \setminus B) \cup (B \setminus A)$.

⁵R. Cauty [2] proved it in the case where $X = [0, 1]$ with the usual metric and the Lebesgue measure.

⁶For spaces $Y_1 \supset Y_2$ and $Z_1 \supset Z_2$, the pair (Y_1, Y_2) is homeomorphic to (Z_1, Z_2) provided that there exists a homeomorphism $f : Y_1 \rightarrow Z_1$ such that $f(Y_2) = Z_2$.

2 A characterization of compactness in $L^p(X)$

Combining Hölder's inequality with the Fubini-Tonelli theorem, we can prove the following lemma, which means that A_r is bounded as an operator on $L^p(X)$.

Lemma 2.1. *Suppose that X is a doubling metric measure space with the doubling constant γ and that $f \in L^p(X)$. Then $\|A_r f\|_p \leq \gamma^{1/p} \|f\|_p$ for any $r > 0$.*

We will show the “only if” part of Theorem 1.2.

Sketch of proof. Since F is relatively compact, and hence it is totally bounded, it can be approximated by finitely many bounded functions $f_i \in L^p(X)$, $1 \leq i \leq n$, with a bounded support E_i . According to Lemma 2.1, we can see that for each $f \in F$,

$$\begin{aligned} \|A_r f - f\|_p &\leq \|A_r f - A_r f_i\|_p + \|A_r f_i - f_i\|_p + \|f_i - f\|_p \\ &\leq \|A_r f_i - f_i\|_p + (\gamma^{1/p} + 1) \|f_i - f\|_p, \end{aligned}$$

where γ is the doubling constant of X . By virtue of the Lebesgue differentiation theorem and the dominated convergence theorem, $\|A_r f_i - f_i\|_p \rightarrow 0$ as $r \rightarrow 0$, and hence the condition (1) holds. On the other hand, letting $E = \bigcup_{x \in \bigcup_{i=1}^n E_i} B(x, 1)$, observe that

$$\|f \chi_{X \setminus E}\|_p \leq \|f \chi_{X \setminus E} - f_i \chi_{X \setminus E}\|_p + \|f_i \chi_{X \setminus E}\|_p \leq \|f - f_i\|_p.$$

Therefore the condition (2) follows immediately. \square

From now on, consider the following conditions between metrics and measures on X :

- (\star) For each $x \in X$, $(0, \infty) \ni r \mapsto \mu(B(x, r)) \in (0, \infty)$ is continuous;
- (\ast) For any $x \in X$ and any $r \in (0, \infty)$, $\mu(B(x, r) \Delta B(y, r)) \rightarrow 0$ as $y \rightarrow x$;
- (\dagger) For every $r \in (0, \infty)$, $X \ni x \mapsto \mu(B(x, r)) \in (0, \infty)$ is continuous.

As is easily observed, the implications (\star) \Rightarrow (\ast) \Rightarrow (\dagger) hold. Measures of closed balls with the same radius, whose centers are belonging to a bounded subset of a doubling metric measure space are lower bounded.

Lemma 2.2. *Let X be doubling. For every bounded subset $E \subset X$ and for every positive number $r > 0$, $\inf\{\mu(B(x, r)) \mid x \in E\} > 0$.*

It is a key idea in proving the “if” part of Theorem 1.2 to approximate a function by a simple function, while it is significant to approximate a function by a “line graph” in the proof of the Ascoli-Arzelà one. The following proposition implies that the operator A_r on $L^p(X)$ is compact when X is bounded and doubling, and $p > 1$.

Proposition 2.3. *Let X be a doubling metric measure space satisfying the property (\ast) and let $F \subset L^p(X)$ be a bounded subset. Suppose that $E \subset X$ is a bounded set and that $r > 0$. Then $\{(A_r f) \chi_E \mid f \in F\}$ is relatively compact in $L^p(X)$ when $p > 1$. Additionally, if for each $\epsilon > 0$, there exists $\delta > 0$ such that $\|A_r f - f\|_1 < \epsilon$ for every $f \in F$ and every $r \in (0, \delta)$, then the above holds even if $p = 1$.*

Sketch of proof. By Hölder's inequality, we get that for any $x, y \in X$,

$$|A_r f(x) - A_r f(y)| \leq \frac{1}{\mu(B(x, r))} \|f\|_p (\mu(B(x, r) \triangle B(y, r)))^{1/q} \\ + \left| \frac{1}{\mu(B(x, r))} - \frac{1}{\mu(B(y, r))} \right| \|f\|_p (\mu(B(y, r)))^{1/q}.$$

In the case where $p = 1$, if $y \in B(x, 1)$ and $s > 0$, then according to Lemma 2.2,

$$\|f\|_1 \leq \|A_s f - f\|_1 + \frac{1}{\inf\{\mu(B(z, s)) \mid z \in B(x, r+1)\}} \|f\|_1 \mu(B(x, r) \triangle B(y, r)).$$

By the assumption, $\mu(B(x, r) \triangle B(y, r)) \rightarrow 0$ as $y \rightarrow x$, and hence $\mu(B(y, r)) \rightarrow \mu(B(x, r))$. Thus the following equicontinuity of average functions on F is valid.

- For every $x \in X$ and every $\epsilon > 0$, there is $\delta > 0$ such that if $d(x, y) < \delta$, then $|A_r f(x) - A_r f(y)| < \epsilon$ for all $f \in F$.

The boundedness of average functions on F follows from the one of F .

- For each $x \in X$, $\{A_r f(x) \mid f \in F\}$ is bounded in \mathbb{R} .

Combining the above equicontinuity and the boundedness with the Vitali covering theorem (cf. [14, Theorem 6.20]), we can show that $\{A_r f \mid f \in F\}$ is approximated by a finitely many collection of simple functions, which implies that $\{A_r f \mid f \in F\}$ is totally bounded. \square

We shall prove the “if” part of Theorem 1.2.

Sketch of proof. For a bounded subset $E \subset X$ and $r > 0$,

$$\|f - (A_r f)_{\chi_E}\|_p \leq \|f_{\chi_E} - (A_r f)_{\chi_E}\|_p + \|f_{\chi_{X \setminus E}}\|_p \leq \|f - A_r f\|_p + \|f_{\chi_{X \setminus E}}\|_p.$$

It follows from the conditions (1) and (2) that F can be approximated by the subset $\{(A_r f)_{\chi_E} \mid f \in F\}$, that is totally bounded in $L^p(X)$ due to Proposition 2.3. Hence F is relatively compact. \square

3 The topological structure of $LIP_b(X)$

To prove Corollary 1.5, we shall use the following criterion, which was proven by D. Curtis, T. Dobrowolski and J. Mogilski [3].

Theorem 3.1 (Curtis-Dobrowolski-Mogilski). *Let C be a σ -compact convex set in a completely metrizable linear space, whose closure $\text{cl} C$ is an AR and not locally compact. Then $(\text{cl} C, C)$ is homeomorphic to $(\ell_2 \times \mathbf{Q}, \ell_2^f \times \mathbf{Q})$ if C contains an infinite-dimensional locally compact convex set.*

The density of $\text{LIP}_b(X) \subset \text{L}^p(X)$ follows from the doubling property and the condition (\star) of X .

Proposition 3.2. *If X is a doubling metric measure space satisfying the property (\star) , then $\text{LIP}_b(X)$ is dense in $\text{L}^p(X)$.*

Sketch of proof. To approximate a function of $\text{L}^p(X)$, by virtue of the Vitali covering theorem, we can choose a lipschitz function $g : X \rightarrow \mathbb{R}$ defined as follows:

$$g(x) = \begin{cases} a_i & \text{if } x \in B(x_i, \delta_i), i = 1, \dots, n, \\ \frac{-a_i(d(x, x_i) - \delta_i)}{r_i - \delta_i} + a_i & \text{if } x \in B(x_i, r_i) \setminus B(x_i, \delta_i), i = 1, \dots, n, \\ 0 & \text{if otherwise,} \end{cases}$$

where for each $i = 1, \dots, n$, $x_i \in X$, $a_i \neq 0$, $0 < \delta_i < r_i$, and $\{B(x_i, r_i) \mid i = 1, \dots, n\}$ is pairwise disjoint. \square

Fix any point $x_0 \in X$. Given a positive integer n , put

$$L(n) = \{f \in \text{LIP}(X) \mid \|f\|_p \leq n, \text{lip } f \leq n \text{ and } \text{supp } f \subset B(x_0, n)\}.$$
⁷

Note that $\text{LIP}_b(X) = \bigcup_{n \in \mathbb{N}} L(n)$. Applying Theorem 1.2, we can prove the following:

Proposition 3.3. *Suppose that X is a doubling metric measure space which satisfies the property (\star) . Then each $L(n)$ is compact.*

Sketch of proof. By the definition of $L(n)$, it is bounded and satisfies the condition (2) of Theorem 1.2. Observe that $L(n)$ is closed. Taking any $f \in L(n)$ and any $r \in (0, 1)$, since f is n -lipschitz, we have that for each $x \in X$,

$$\begin{aligned} |A_r f(x) - f(x)| &\leq \frac{1}{\mu(B(x, r))} \int_{B(x, r)} |f(y) - f(x)| d\mu(y) \\ &\leq \frac{1}{\mu(B(x, r))} \int_{B(x, r)} nr d(x, y) d\mu(y) \leq nr. \end{aligned}$$

Then $\|A_r f - f\|_p \leq nr(\mu(B(x_0, n+1)))^{1/p}$, which implies that the condition (1) of Theorem 1.2 holds. \square

It will be shown that each $L(n)$ is homeomorphic to \mathbf{Q} by using the following characterization [13].

Theorem 3.4 (Toruńczyk). *A compact AR Y is homeomorphic to \mathbf{Q} if and only if the following is satisfied.*

- (\sharp) *Let n be any positive integer and $f : [0, 1]^n \times \{0, 1\} \rightarrow Y$ be any continuous function. For every $\epsilon > 0$, there is a continuous function $g : [0, 1]^n \times \{0, 1\} \rightarrow Y$ such that g is ϵ -close to f and $g([0, 1]^n \times \{0\}) \cap g([0, 1]^n \times \{1\}) = \emptyset$.*⁸

⁷The symbols $\text{lip } f$ and $\text{supp } f$ stand for the lipschitz constant and the support of f , respectively.

⁸This property is called the *disjoint cells property*.

Recall that for a metric space $Y = (Y, d_Y)$, for functions $f : Z \rightarrow Y$ and $g : Z \rightarrow Y$, and for a positive number $\epsilon > 0$, f is ϵ -close to g if $d_Y(f(z), g(z)) \leq \epsilon$ for all $z \in Z$.

Lemma 3.5. *Let X be a doubling metric measure space such that $\mu(\{x_0\}) = 0$, and let n be a positive integer. For any $\epsilon > 0$, there exist maps $\Phi : L(n) \rightarrow \{f \in L(n) \mid \text{lip } f < n\}$ and $\Psi : L(n) \rightarrow \{f \in L(n) \mid \text{lip } f = n\}$ which are ϵ -close to the identity map on $L(n)$.*

Sketch of proof. Let $\Phi(f) = (1 - \epsilon/n)f$ for each $f \in L(n)$. Due to the map Φ , we may assume that there exists $m < n$ such that $\|f\|_p \leq m$ and $\text{lip } f \leq m$ for every $f \in L(n)$. Taking $0 < r_2 < r_1$ appropriately, for each $f \in L(n)$, we can define a function $\psi(f) : X \setminus (B(x_0, r_1) \setminus B(x_0, r_2)) \rightarrow \mathbb{R}$ by

$$\psi(f)(x) = \begin{cases} f(x_0) + n(r_2 - d(x, x_0)) & \text{if } x \in B(x_0, r_2), \\ f(x) & \text{if otherwise.} \end{cases}$$

Applying McShane's lipschitz extension [11], let

$$\Psi(f)(x) = \begin{cases} \psi(f)(x) & \text{if } x \in X \setminus (B(x_0, r_1) \setminus B(x_0, r_2)), \\ \inf_{a \in X \setminus (B(x_0, r_1) \setminus B(x_0, r_2))} (\psi(f)(a) + nd(a, x)) & \text{if otherwise,} \end{cases}$$

which is the desired. \square

It is a direct consequence of the above lemma that every $L(n)$ satisfies the property (\sharp) of Theorem 3.4.

Proposition 3.6. *Let X be a doubling metric measure space with $\mu(\{x_0\}) = 0$. If the condition $(*)$ holds, then $L(n)$ is homeomorphic to \mathbf{Q} for any positive integer n .*

Now we shall show Corollary 1.5.

Sketch of proof. By Propositions 3.2 and 3.6, we can prove that $\text{LIP}_b(X) \subset L^p(X)$ is a dense σ -compact subset containing topological copies of \mathbf{Q} . According to Theorem 3.1, the pair $(L^p(X), \text{LIP}_b(X))$ is homeomorphic to $(\ell_2 \times \mathbf{Q}, \ell_2^f \times \mathbf{Q})$. \square

References

- [1] R.D. Anderson, *Hilbert space is homeomorphic to the countable infinite product of lines*, Bull. Amer. Math. Soc. 72 (1966), 515–519.
- [2] R. Cauty, *Les fonctions continues et les fonctions intégrables au sens de Riemann comme sous-espaces de \mathcal{L}_1* , Fund. Math. 139 (1991), 23–36.
- [3] D.W. Curtis, T. Dobrowolski and J. Mogilski, *Some applications of the topological characterizations of the sigma-compact spaces ℓ_2^f and Σ* , Trans. Amer. Math. Soc. 284 (1984), 837–846.

- [4] P. Górká and A. Macios, *The Riesz-Kolmogorov theorem on metric spaces*, (English summary) Miskolc Math. Notes 15 (2014), 459–465.
- [5] P. Górká and H. Rafeiro, *From Arzelà-Ascoli to Riesz-Kolmogorov*, (English summary) Nonlinear Anal. 144 (2016), 23–31.
- [6] H. Hanche-Olsen and H. Holden, *The Kolmogorov-Riesz compactness theorem*, (English summary) Expo. Math. 28 (2010), 385–394.
- [7] M.I. Kadec, *A proof the topological equivalence of all separable infinite-dimensional Banach spaces* (Russian), Funkcional Anal. i Priložen, 1 (1967), 61–70.
- [8] A.N. Kolmogorov, *Über Kompaktheit der Funktionenmengen bei der Konvergenz im Mittel*, Nachr. Ges. Wiss. Göttingen 9 (1931), 60–63.
- [9] K. Koshino, *The space consisting of uniformly continuous functions on a metric measure space with the L^p norm*, Topology Appl. 282 (2020), 107303.
- [10] K. Koshino, *Characterizing compact sets in L^p -spaces and its application*, arXiv: 2205.06864 [math.GN].
- [11] E.J. McShane, *Extension of range of functions*, Bull. Amer. Math. Soc. 40 (1934), 837–842.
- [12] M. Riesz, *Sur les ensembles compacts de fonctions sommables*, Acta Szeged Sect. Math. 6 (1933), 136–142.
- [13] H. Toruńczyk, *On CE -images of the Hilbert cube and characterization of \mathbf{Q} -manifolds*, Fund. Math. 106 (1980), 31–40.
- [14] J. Yeh, *Metric in measure spaces*, World Scientific Publishing Co. Pte. Ltd., Singapore, 2020.

Faculty of Engineering
Kanagawa University
Yokohama, 221-8686, Japan
E-mail: ft160229no@kanagawa-u.ac.jp