

**WELL-POSEDNESS OF STOCHASTIC NONLINEAR HEAT AND WAVE  
EQUATIONS DRIVEN BY SUBORDINATE CYLINDRICAL BROWNIAN  
NOISES ON THE TWO-DIMENSIONAL TORUS**

HIROTATSU NAGOJI

ABSTRACT. In [12], the author studied stochastic nonlinear heat equations and stochastic nonlinear wave equations driven by subordinate cylindrical Brownian noises on the two-dimensional torus by way of renormalization. More precisely, the author studied the local well-posedness of the renormalized version of the equations. In this note, we make some remarks on the global well-posedness of the stochastic nonlinear wave equation with a cubic nonlinearity after we briefly revisit the results in [12].

1. INTRODUCTION

In [12], the author studied the local well-posedness of stochastic nonlinear heat equations and stochastic nonlinear wave equations driven by subordinate cylindrical Brownian noises on the two-dimensional torus. In this note, we briefly discuss the global well-posedness of the stochastic nonlinear wave equation with a cubic nonlinearity, which we omitted in [12].

First, we recall the setting and the main result of [12]. In that study, the author studied the following Cauchy problems of the stochastic nonlinear heat equation and stochastic nonlinear wave equation on the two-dimensional torus  $\mathbb{T}^2 = (\mathbb{R}/2\pi\mathbb{Z})^2$ :

$$(1.1) \quad \begin{cases} \partial_t u - \Delta u = \lambda u^k + \partial_t W_L \\ u(0) = u_0 \end{cases}$$

$$(1.2) \quad \begin{cases} \partial_t^2 u - \Delta u = \lambda u^k + \partial_t W_L \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

where  $k \geq 2$  is an integer and  $\lambda \in \mathbb{R}$ . The stochastic process  $W_L$  is called subordinate cylindrical Brownian motion and defined as follows: Let  $W$  be a cylindrical Brownian motion on  $L^2(\mathbb{T}^2)$  defined on the probability space  $(\Omega_1, \mathbb{P}_1)$  written by

$$(1.3) \quad W(t) = \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2} \beta^l(t) e_l$$

where  $e_l(x) = e^{\sqrt{-1}l \cdot x}$  is the Fourier basis and  $(\beta^l)_{l \in \mathbb{Z}^2}$  is an independent sequence of  $\mathbb{C}$ -valued Brownian motions conditioned with  $\bar{\beta}^l = \beta^{-l}$  and  $\text{var}(\beta^l(t)) = t$ . We also consider the  $\mathbb{R}_+$ -valued

---

2020 *Mathematics Subject Classification.* 60H15, 35K15, 35L71.

*Key words and phrases.* Subordinate cylindrical Brownian motion, stochastic nonlinear heat equation, stochastic nonlinear wave equation, renormalization.

*Statements and Declarations:* This work was supported by JST SPRING, Grant Number JPMJSP2110. The author has no conflicts of interest directly relevant to the content of this article.

stochastic process  $L$  with  $L(0) = 0$ , non-decreasing and càdlàg sample paths defined on another space  $(\Omega_2, \mathbb{P}_2)$ . Then, we define the process  $W_L$  defined on  $(\Omega, \mathbb{P}) := (\Omega_1 \otimes \Omega_2, \mathbb{P}_1 \otimes \mathbb{P}_2)$  by

$$(1.4) \quad W_L(t) = \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2} \beta^l(L(t)) e_l.$$

It is well-known that when  $L$  is a Lévy process with non-decreasing paths,  $W_L$  is also a (distribution-valued) Lévy process. See [2, Section 1.3.2] for the example of such subordinators  $L$ .

When the noise is given by space time white noise i.e. when  $L(t) = t$  in our setting, the equations (1.1) and (1.2) are related to the problem of stochastic quantization and are studied by a lot of researchers, see [5, 8, 9, 4] and references therein. In these equations, some ill-defined products of distributions appear in the equation because of the singularity of the noise, so we need to introduce renormalization in order to give meaning to the products. This kind of equations are called singular SPDEs and have attracted great attentions especially after the invention of the theory of regularity structure [10] and the theory of paracontrolled calculus [7].

Similarly to these previous works, it turns out that we need to introduce renormalization to give meaning to the nonlinearity. More precisely, we have to consider the following modified approximation of the equations:

$$(1.5) \quad \mathcal{L}u_N = \lambda H_k(u_N; C_N) + P_N \partial_t W_L$$

where  $\mathcal{L} = \partial_t - \Delta$  or  $\partial_t^2 - \Delta$ ,  $P_N$  is the approximation operator defined by

$$(1.6) \quad P_N f = \sum_{l \in \mathbb{Z}^2, |l| \leq N} \hat{f}(l) e_l \quad \text{for } f \in \mathcal{D}'(\mathbb{T}^2),$$

$H_k$  is the  $k$ -th Hermite polynomial defined via

$$(1.7) \quad e^{tx - \frac{1}{2}\sigma^2 t^2} = \sum_{k=0}^{\infty} \frac{t^k}{k!} H_k(x; \sigma^2)$$

and the renormalization constant  $C_N$  is defined by

$$(1.8) \quad C_N(t) := \begin{cases} \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2, |l| \leq N} \int_0^t e^{2(s-t)|l|^2} dL(s) & \text{when } \mathcal{L} = \partial_t - \Delta \\ \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2, |l| \leq N} \int_0^t \frac{\sin^2((t-s)|l|)}{|l|^2} dL(s) & \text{when } \mathcal{L} = \partial_t^2 - \Delta. \end{cases}$$

Note that  $C_N$  does not depend on the space variable  $x \in \mathbb{T}^2$  but depends on  $\omega_2 \in \Omega_2$ , so  $C_N$  is an  $L$ -measurable stochastic process. Then, we decompose the solution  $u_N$  as  $u_N = v_N + \Phi_N$  where  $\Phi_N$  solves the linear equation

$$(1.9) \quad \mathcal{L}\Phi_N = P_N \partial_t W_L$$

with initial condition 0, see Section 2 for more precise definition. This decomposition is called Da Prato-Debussche trick in the context of singular SPDEs. Then,  $v_N$  solves

$$(1.10) \quad \begin{aligned} \mathcal{L}v_N &= \lambda H_k(v_N + \Phi_N; C_N) \\ &= \lambda \sum_{l=0}^k \binom{k}{l} v_N^{k-l} H_l(\Phi_N; C_N) \end{aligned}$$

where we use the algebraic property of the Hermite polynomials

$$(1.11) \quad H_k(a+b; c) = \sum_{l=0}^k \binom{k}{l} a^{k-l} H_l(b; c).$$

In the argument of renormalization of singular SPDEs, we usually heavily use the Gaussianity of the noises, so similar kind of argument cannot be made in the case of non-Gaussian noises. In our setting (1.4), although  $W_L$  itself is not a Gaussian process in general, it turns out that we can deploy renormalization argument by exploiting the Gaussianity of  $W$ : We can show the convergence of the random coefficients  $\Phi_N^{\diamond l} := H_l(\Phi_N; C_N)$  of the equation (1.10), which are called the Wick powers, see Proposition 2.2. And consequently, we can show the convergence of the solution  $v_N$  (and  $u_N$ ), see Theorems 1.1 and 1.2 below.

As the driving noises of SPDEs, Gaussian noises such as space-time white noise are considered to be the most important examples, but on the other hand, SPDEs driven by Lévy noise (or jump noise) are also studied by a lot of researchers especially after publication of the monograph [14], see [1, 3, 11, 16, 18] and references therein. We also note that our setting of the noise (1.4) includes some interesting type of the noise such as stable noises [6, 15, 17] and compound Poisson noises. Now, we recall the local well-posedness result of the equations (1.1) and (1.2) in [12].

**Theorem 1.1.** (cf. [12, Theorem 1.1]) *Let  $k = 2$  and let  $v_N$  be the mild solution of*

$$(1.12) \quad \begin{cases} \partial_t v_N - \Delta v_N = \lambda \sum_{l=0}^k \binom{k}{l} v_N^{k-l} \Phi_N^{\diamond l} \\ v(0) = v_0. \end{cases}$$

*Then for any given initial condition  $v_0 \in B_{\infty, \infty}^{2/\gamma - \delta}(\mathbb{T}^2)$ , there exists  $T(\omega) > 0$  such that  $v_N$  converges to some  $v$   $\mathbb{P}$ -almost-surely in*

$$L^\gamma([0, T]; B_{\infty, \infty}^{2/\gamma - \delta}(\mathbb{T}^2)) \cap C([0, T]; B_{\infty, \infty}^{-\delta}(\mathbb{T}^2))$$

*for any  $\frac{2}{1-\epsilon} < \gamma < \frac{2}{\epsilon}$ ,  $0 < \delta < \frac{2}{\gamma} - \epsilon$ . Moreover the limit process  $v$  solves the equation*

$$(1.13) \quad \begin{cases} \partial_t v - \Delta v = \lambda \sum_{l=0}^k \binom{k}{l} v^{k-l} \Phi^{\diamond l} \\ v(0) = v_0 \end{cases}$$

*on the interval  $[0, T(\omega)]$  where  $\Phi^{\diamond k}$  are the Wick powers of  $\Phi$  defined in Proposition 2.2.*

**Theorem 1.2.** (cf. [12, Theorem 1.3]) *Let  $k \geq 2$  and let  $v_N$  be the mild solution of*

$$(1.14) \quad \begin{cases} \partial_t^2 v_N - \Delta v_N = \lambda \sum_{l=0}^k \binom{k}{l} v_N^{k-l} \Psi_N^{\diamond l} \\ (v(0), \partial_t v(0)) = (v_0, v_1). \end{cases}$$

*Then, for any given initial condition  $(v_0, v_1) \in H^{1-\epsilon}(\mathbb{T}^2) \times H^{-\epsilon}(\mathbb{T}^2)$  with sufficiently small  $\epsilon > 0$ , there exists  $T(\omega) > 0$  such that  $v_N$  converges to some  $v$   $\mathbb{P}$ -almost-surely in  $C([0, T]; H^{1-\epsilon}(\mathbb{T}^2)) \cap C^1([0, T]; H^{-\epsilon}(\mathbb{T}^2))$ . Moreover the limit process  $v$  solves the equation*

$$(1.15) \quad \begin{cases} \partial_t^2 v - \Delta v = \lambda \sum_{l=0}^k \binom{k}{l} v^{k-l} \Psi^{\diamond l} \\ (v(0), \partial_t v(0)) = (v_0, v_1) \end{cases}$$

*on the interval  $[0, T(\omega)]$ , where  $\Psi^{\diamond k}$  are the Wick powers of  $\Psi$  defined in Proposition 2.2.*

**Remark 1.3.** *When  $L(t) = t$ , it is known that  $u_N$  converges for any positive integer  $k$  both heat and wave cases, see [5, 8]. On the other hand, we have not been able to obtain the convergence result for  $k \geq 3$  in the case of the heat equation for general  $L$ . This difference between the heat case and the wave case comes from the difference in the time integrability of the Wick powers, see*

*Proposition 2.2.* This result (and the argument in the proof of the result) suggests that the heat case behaves worse than the wave case when the singular noise is of jump-type, differently from the case of space-time white noise, see [13].

As we pointed out in [12], when  $k = 3$ ,  $\lambda < 0$  and  $\mathcal{L} = \partial_t^2 - \Delta$ , it is straightforward to extend the locally-in-time convergence of  $u_N$  to the globally-in-time convergence by applying the argument with I-method in [9]. In Section 3, we will explain how to adjust their argument to our setting and we will prove the global well-posedness of the stochastic nonlinear wave equation with a cubic nonlinearity.

## 2. LINEAR EQUATIONS

In this section, we give the definition of the mild solution of the stochastic linear heat equation

$$(2.1) \quad \begin{cases} \partial_t \Phi - \Delta \Phi = \partial_t W_L \\ \Phi(0) = 0, \end{cases}$$

and the stochastic linear wave equation

$$(2.2) \quad \begin{cases} \partial_t^2 \Psi - \Delta \Psi = \partial_t W_L \\ (\Psi(0), \partial_t \Psi(0)) = (0, 0). \end{cases}$$

From the Duhamel principle, it is natural to define the solutions of these equations as the following stochastic convolutions:

$$(2.3) \quad \Phi(t) = \int_0^t e^{(t-s)\Delta} dW_L(s) = \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2} \int_0^t e^{(s-t)|l|^2} d\beta_L^l(s) e_l$$

and

$$(2.4) \quad \Psi(t) = \int_0^t \frac{\sin((t-s)|\nabla|)}{|\nabla|} dW_L(s) = \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2} \int_0^t \frac{\sin((t-s)|l|)}{|l|} d\beta_L^l(s) e_l$$

where  $\beta_L^l(t) = \beta^l(L(t))$ . Recall that  $W_L$  is defined by (1.4). The integrals  $\int_0^t e^{(s-t)|l|^2} d\beta_L^l(s)$  and  $\int_0^t \frac{\sin((t-s)|l|)}{|l|} d\beta_L^l(s)$  can be understood as the Young integrals thanks to the following lemma.

**Lemma 2.1.** (cf. [12, Lemma 3.1]) *Let  $B = B(t)$  be a one-dimensional Brownian motion. Then, the sample paths of  $B_L(t) = B(L(t))$  have the finite  $p$ -variation on any interval  $[0, T]$  for any  $p > 2$  almost surely.*

*Proof.* Let  $P[0, T] = \{D = \{0 = t_0 < t_1 < \dots < t_n = T\}\}$  be the set of all partitions of  $[0, T]$ . Because the sample paths of  $B$  is  $\frac{1}{p}$ -Hölder continuous almost surely for any  $p > 2$ , there holds

$$\sup_{D \in P[0, T]} \sum_{i=1}^n |B_L(t_i) - B_L(t_{i-1})|^p \lesssim \sup_{D \in P[0, T]} \sum_{i=1}^n |L(t_i) - L(t_{i-1})|^{p \times \frac{1}{p}} = L(T) < \infty.$$

□

From this lemma and the theory of the Young integral, we can see that the finite sums

$$(2.5) \quad \Phi_N(t) := P_N \Phi(t) = \frac{1}{2\pi} \sum_{|l| \leq N} \int_0^t e^{(s-t)|l|^2} d\beta_L^l(s) e_l$$

and

$$(2.6) \quad \Psi_N(t) := P_N \Psi(t) = \frac{1}{2\pi} \sum_{|l| \leq N} \int_0^t \frac{\sin((t-s)\langle l \rangle)}{\langle l \rangle} d\beta_L^l(s) e_l$$

are well-defined as stochastic processes. Then, from Proposition 2.2 below, we see that  $\Phi_N$  and  $\Psi_N$  converge as  $N \rightarrow \infty$ .

Now, we recall the convergence result of the Wick powers of  $\Phi_N$  and  $\Psi_N$  defined by

$$(2.7) \quad \Phi_N^{\diamond k} := H_k(\Phi_N; C_N^H)$$

and

$$(2.8) \quad \Psi_N^{\diamond k} := H_k(\Psi_N; C_N^W).$$

which appear in the equation (1.10) where

$$(2.9) \quad C_N^H := \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2, |l| \leq N} \int_0^t e^{2(s-t)|l|^2} dL(s)$$

and

$$(2.10) \quad C_N^W := \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2, |l| \leq N} \int_0^t \frac{\sin^2((t-s)|l|)}{|l|^2} dL(s).$$

Here we state the convergence result in [12].

**Proposition 2.2.** (cf. [12, Propositions 4.3 and 4.8]) *Let  $k \geq 1$  be any positive integer and let  $T > 0$ .*

- (i)  $\Phi_N^{\diamond k}$  converges to some  $\Phi^{\diamond k}$   $\mathbb{P}$ -almost-surely in  $L^p([0, T]; W^{-\alpha, \infty}(\mathbb{T}^2))$  for any  $\alpha > 0$  and  $0 < p < \frac{2}{k-1}$  with  $\frac{2}{p} + \alpha > k$  as  $N \rightarrow \infty$ .
- (ii)  $\Psi_N^{\diamond k}$  converges to some  $\Psi^{\diamond k}$   $\mathbb{P}$ -almost-surely in  $C([0, T]; W^{-\delta, \infty}(\mathbb{T}^2))$  for any  $\delta > 0$  as  $N \rightarrow \infty$ .

**Remark 2.3.** *When  $L(t) = t$ , it is well-known that both  $\Phi_N^{\diamond k}$  and  $\Psi_N^{\diamond k}$  converge in the space  $C([0, T]; W^{-\delta, \infty}(\mathbb{T}^2))$  for any  $\delta > 0$  as  $N \rightarrow \infty$ , see [5, 8] for example.*

**Remark 2.4.** *Because the noise is jump-type, we cannot expect the time-continuity for the heat case  $\Phi$ . But by working on  $L^p$ -space with respect to time variable, we can construct the Wick powers of  $\Phi$ . Note that when  $0 < p < 1$ ,  $L^p$ -space is not a norm space but a complete metric space. On the other hand, we can show the time-continuity for the wave case  $\Psi$  even when subordinator  $L$  has jumps.*

### 3. GLOBAL WELL-POSEDNESS OF THE STOCHASTIC NONLINEAR WAVE EQUATION WITH A CUBIC NONLINEARITY

In this section, we prove the global well-posedness of the stochastic nonlinear wave equation with a cubic nonlinearity by applying the globalization argument with I-method in [9].

First, we briefly recall their argument. They considered the case when  $k = 3$ ,  $\lambda < 0$ ,  $\mathcal{L} = \partial_t^2 - \Delta$  and  $L(t) = t$  in our setting. In view of Theorem 1.2, it suffices to consider the following equation:

$$(3.1) \quad \begin{cases} \partial_t^2 v - \Delta v = \lambda \sum_{l=0}^3 \binom{3}{l} v^{3-l} \Psi^{\diamond l} \\ (v(0), \partial_t v(0)) = (v_0, v_1) \end{cases}$$

where  $\lambda < 0$  is a negative constant. Let  $L(t) = t$  for a while. In order to prove the existence of the global solution of this kind of wave equation, one (or only one perhaps) known method is to utilize the formal conservation of the energy

$$(3.2) \quad E(v)(t) = \frac{1}{2} \int_{\mathbb{T}^2} v(t)^2 dx + \frac{1}{2} \int_{\mathbb{T}^2} (\partial_t v(t))^2 dx + \frac{1}{2} \int_{\mathbb{T}^2} |\nabla v(t)|^2 dx - \frac{\lambda}{4} \int_{\mathbb{T}^2} v(t)^4 dx.$$

However, in this setting, the solution  $(v, \partial_t v)$  of (3.1) lies in the space  $H^s(\mathbb{T}^2) \times H^{s-1}(\mathbb{T}^2)$  with  $s < 1$  due to the low regularity of the Wick powers  $\Psi^{\diamond l}$ , so the value of  $E(v)$  is expected to be infinite. To avoid this problem, they consider the following mollified energy

$$(3.3) \quad E^I(v) := E(Iv) = \frac{1}{2} \int_{\mathbb{T}^2} (Iv)^2 dx + \frac{1}{2} \int_{\mathbb{T}^2} (\partial_t Iv)^2 dx + \frac{1}{2} \int_{\mathbb{T}^2} |\nabla Iv|^2 dx - \frac{\lambda}{4} \int_{\mathbb{T}^2} (Iv)^4 dx$$

where  $I = I_N$  is called  $I$ -operator and defined as follows. For fixed  $0 < s < 1$  and given  $N \geq 1$ , we define the operator  $I = I_N$  on  $\mathcal{D}'(\mathbb{T}^2)$  by

$$(3.4) \quad I_N f = \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2} m_N(l) \hat{f}(l) e_l$$

where  $\hat{f}$  is the Fourier transformation of  $f$  and  $m_N \in C^\infty(\mathbb{R}_+; [0, 1])$  is a rotationally invariant function such that

$$(3.5) \quad m_N(\xi) = \begin{cases} 1 & \text{when } |\xi| \leq N \\ \left(\frac{N}{|\xi|}\right)^{1-s} & \text{when } |\xi| \geq 2N \end{cases}$$

and  $m_N(\xi)$  is non-decreasing with respect to  $|\xi|$ . Then, they successfully controlled the growth of mollified energy  $E^I(v)$  by the Gronwall-type argument and proved the global well-posedness of the equation (3.1). In that process, we need a more careful observation of the behavior of  $\Psi$  than we need in the proof of the local well-posedness. More precisely, the fact that  $\Psi^{\diamond k} \in C([0, T]; W^{-\delta, \infty}(\mathbb{T}^2))$  for any  $\delta > 0$ , which we saw in Proposition 2.2, is not enough to prove the global well-posedness and we need to use the fact that the following quantities are finite for  $\Psi$ : For given  $\Xi = (\Xi_1, \Xi_2, \Xi_3)$ , we define  $V = V(\Xi)$  and  $R = R(\Xi)$  by

$$(3.6) \quad e^{V^{\frac{1}{3}}} := \sum_{j=0}^{\infty} e^{-\theta j} e^{V_j(\Xi)^{\frac{1}{3}}}$$

and

$$(3.7) \quad R := \sum_{N=1}^{\infty} \sum_{j=1}^{\infty} e^{-\theta j \log N} \int_0^j \int_{\mathbb{T}^2} e^{|I_N \Xi_1(t, x)|} dx dt$$

where  $V_j(\Xi) := \max_{1 \leq k \leq 3} \|\Xi_k\|_{L^\infty([j, j+1]; W^{-\delta, \infty})}$  and  $\delta, \theta > 0$ . Note that the definition of  $V$  above is slightly different from the one in [9] but this does not cause any problem in the argument below. Then, it turns out that  $V(\Psi, \Psi^{\diamond 2}, \Psi^{\diamond 3})$  and  $R(\Psi, \Psi^{\diamond 2}, \Psi^{\diamond 3})$  are finite almost surely. In [9], they essentially proved the following theorem which is purely deterministic result and by applying this to the equation (3.1) pathwisely, they obtained the desired result.

**Theorem 3.1.** (cf. [9]) *Let  $s > \frac{4}{5}$ . Assume that  $\Xi = (\Xi_1, \Xi_2, \Xi_3)$  satisfies  $V(\Xi) < \infty$  and  $R(\Xi) < \infty$  for some  $\theta > 0$  and sufficiently small  $\delta > 0$ . Then, the equation*

$$(3.8) \quad \begin{cases} \partial_t^2 v - \Delta v = \lambda v^3 + \lambda \sum_{l=1}^3 \binom{3}{l} v^{3-l} \Xi_l \\ (v(0), \partial_t v(0)) = (v_0, v_1) \end{cases}$$

has a unique mild solution in the space

$$C([0, T]; H^s(\mathbb{T}^2)) \cap C^1([0, T]; H^{s-1}(\mathbb{T}^2))$$

for any given  $(v_0, v_1) \in H^s(\mathbb{T}^2) \times H^{s-1}(\mathbb{T}^2)$  and  $T > 0$ .

Now, we go back to the original setting i.e. let  $L$  be any càdlàg and non-decreasing stochastic process with  $L(0) = 0$ . Then, unfortunately,  $(\Psi, \Psi^{\diamond 2}, \Psi^{\diamond 3})$  does not satisfy the assumption of Theorem 3.1. However, it turns out that we can easily overcome this problem: Fix  $T > 0$ . We define  $\Psi_T^{\diamond k}(t) = \Psi^{\diamond k}(t \wedge T)$ . Then, we can prove that  $(\Psi_T, \Psi_T^{\diamond 2}, \Psi_T^{\diamond 3})$  satisfies the assumption of Theorem 3.1:

**Lemma 3.2.** *For any  $T > 0$ ,  $\delta > 0$  and  $\omega_2 \in \Omega_2$ , there exists some  $\theta = \theta(\omega_2, T) > 0$  such that the values of  $V(\Psi_T, \Psi_T^{\diamond 2}, \Psi_T^{\diamond 3})$  and  $R(\Psi_T, \Psi_T^{\diamond 2}, \Psi_T^{\diamond 3})$  are finite  $\mathbb{P}_1$ -almost surely.*

Before the proof we recall the following lemma.

**Lemma 3.3.** (cf. [12, Lemma 4.1]) *Let  $f$  and  $g$  be smooth deterministic functions on  $[0, T]$ . Then, for any  $l, l' \in \mathbb{Z}^2$ ,*

$$(3.9) \quad \mathbb{E}^{\mathbb{P}_1} \left[ \int_0^T f(t) d\beta_L^l(t) \int_0^T g(t) d\beta_L^{l'}(t) \right] = \delta_{l, -l'} \int_0^T f(t)g(t) dL(t).$$

*Proof of Lemma 3.2.* Because

$$V_j(\Psi_T, \Psi_T^{\diamond 2}, \Psi_T^{\diamond 3}) = \max_{1 \leq k \leq 3} \|\Psi_T^{\diamond k}\|_{L^\infty([j, j+1]; W^{-\delta, \infty})} \leq \max_{1 \leq k \leq 3} \|\Psi_T^{\diamond k}\|_{L^\infty([0, T]; W^{-\delta, \infty})} < \infty$$

uniformly in  $j \in \mathbb{N}$  from Proposition 2.2, we can see that

$$e^{V^{\frac{1}{3}}} = \sum_{j=0}^{\infty} e^{-\theta j} e^{V_j(\Psi_T, \Psi_T^{\diamond 2}, \Psi_T^{\diamond 3})^{\frac{1}{3}}} < \infty$$

for any  $\theta > 0$ . Next, we check that  $R(\Psi_T)$  is finite. We write  $\Psi_T = P_N \Psi_T + P_N^\perp \Psi_T$ . From Lemma 3.3,

$$\begin{aligned} \mathbb{E}^{\mathbb{P}_1} \left[ |I_N P_N \Psi_T(x, t)|^2 \right] &= \mathbb{E}^{\mathbb{P}_1} \left[ |P_N \Psi_T(x, t)|^2 \right] \\ &= \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2, |l| \leq N} \int_0^{t \wedge T} \frac{\sin^2((t-s)|l|)}{|l|^2} dL(s) \\ &\lesssim L(T) \log N \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}^{\mathbb{P}_1} \left[ |I_N P_N^\perp \Psi_T(x, t)|^2 \right] &= \frac{1}{2\pi} \sum_{l \in \mathbb{Z}^2, |l| > N} m_N(l)^2 \int_0^{t \wedge T} \frac{\sin^2((t-s)|l|)}{|l|^2} dL(s) \\ &\lesssim \sum_{l \in \mathbb{Z}^2, |l| > N} \frac{N^{2-2s}}{|l|^{2-2s}} \times \frac{L(T)}{|l|^2} \lesssim L(T). \end{aligned}$$

Therefore, for any fixed  $\omega_2 \in \Omega_2$ ,  $I_N \Psi_T(x, t)$  is a centered Gaussian random variable on  $(\Omega_1, \mathbb{P}_1)$  with

$$\mathbb{E}^{\mathbb{P}_1} \left[ |I_N \Psi_T(x, t)|^2 \right] \leq C_0 L(T) \log N$$

where  $C_0 > 0$  is a positive constant which is independent of  $t \in \mathbb{R}_+$  and  $x \in \mathbb{T}^2$ . Thus, we can calculate as

$$\begin{aligned} \mathbb{E}^{\mathbb{P}_1} [R(\Psi_T)] &= \mathbb{E}^{\mathbb{P}_1} \left[ \sum_{N=1}^{\infty} \sum_{j=1}^{\infty} e^{-\theta j \log N} \int_0^j \int_{\mathbb{T}^2} e^{|I_N \Psi_T(t,x)|} dx dt \right] \\ &\leq \sum_{N=1}^{\infty} \sum_{j=1}^{\infty} e^{-\theta j \log N} \int_0^j \int_{\mathbb{T}^2} e^{C_0 L(T) \log N} dx dt \\ &\lesssim \sum_{N=1}^{\infty} \sum_{j=1}^{\infty} e^{-\theta j \log N} e^{C_0 L(T) \log N} \times j \\ &= \sum_{N=1}^{\infty} \sum_{j=1}^{\infty} N^{-\theta j} N^{C_0 L(T)} \times j < \infty \end{aligned}$$

if  $\theta$  is sufficiently larger than  $L(T)$ .  $\square$

From Theorem 3.1 and Lemma 3.2, we can see that the global well-posedness of the following equation holds.

$$(3.10) \quad \begin{cases} \partial_t^2 v - \Delta v = \lambda v^3 + \lambda \sum_{l=1}^3 \binom{3}{l} v^{3-l} \Psi_T^{\diamond l} \\ (v(0), \partial_t v(0)) = (v_0, v_1). \end{cases}$$

Here, the solutions of (3.1) and (3.10) coincide on the interval  $[0, T]$  if the same initial value  $(v_0, v_1)$  is given. Therefore, we obtain the global well-posedness of the original equation (3.1).

**Acknowledgement.** The author would like to thank Professor Seiichiro Kusuoka and Professor Yoshio Tsutsumi for helpful comments. This work was supported by JST SPRING, Grant Number JPMJSP2110.

#### REFERENCES

- [1] S. Albeverio, V. Mandrekar and B. Rüdiger. Existence of mild solutions for stochastic differential equations and semilinear equations with non-Gaussian Lévy noise. *Stochastic Process. Appl.* 119 (2009), no. 3, 835-863.
- [2] D. Applebaum. *Lévy processes and stochastic calculus*. Cambridge Studies in Advanced Mathematics, 93. Cambridge University Press, Cambridge, 2004.
- [3] Z. Brzezniak and J. Zabczyk. Regularity of Ornstein-Uhlenbeck processes driven by a Lévy white noise. *Potential Anal.*, 32 (2010), 153-188.
- [4] R. Catellier and K. Chouk. Paracontrolled distributions and the 3-dimensional stochastic quantization equation. *Ann. Probab.* 46 (2018), no. 5, 2621-2679.
- [5] G. Da Prato and A. Debussche. Strong solutions to the stochastic quantization equations. *The Ann. Probab.* (2003), Vol. 31, No. 4, 1900-1916.
- [6] Z. Dong, L. Xu and X. Zhang. Exponential ergodicity of stochastic Burgers equations driven by  $\alpha$ -stable processes. *J. Stat. Phys.* 154 (2014), no. 4, 929-949.
- [7] M. Gubinelli, P. Imkeller and N. Perkowski. Paracontrolled distributions and singular PDEs. *Forum Math. Pi* 3 (2015), e6, 75 pp.
- [8] M. Gubinelli, H. Koch and T. Oh. Renormalization of the two-dimensional stochastic nonlinear wave equations. *Trans. Amer. Math. Soc.* (2018), Vol. 370, No. 10, 7335-7359.
- [9] M. Gubinelli, H. Koch, T. Oh and L. Tolomeo. Global Dynamics for the Two-dimensional Stochastic Nonlinear Wave Equations. *International Mathematics Research Notices*, rnab084, <https://doi.org/10.1093/imrn/rnab084>.
- [10] M. Hairer. A theory of regularity structures. *Invent. Math.* 198 (2014), no. 2, 269-504.
- [11] T. Meyer-Brandis and F. Proske. Explicit representation of strong solutions of SDEs driven by infinite dimensional Lévy processes. *J. Theor. Prob.*, 23:301-314, 2010.



- [12] H. Nagoji. Renormalization of stochastic nonlinear heat and wave equations driven by subordinate cylindrical Brownian noises. arXiv:2201.08010 [math.PR]
- [13] T. Oh and M. Okamoto. Comparing the stochastic nonlinear wave and heat equations: a case study. *Electron. J. Probab.* 26 (2021), Paper No. 9, 44 pp.
- [14] S. Peszat and J. Zabczyk. Stochastic partial differential equations with Lévy noise. An evolution equation approach. *Encyclopedia of Mathematics and its Applications*, 113. Cambridge University Press, Cambridge, 2007. xii+419 pp.
- [15] E. Priola and J. Zabczyk. Structural properties of semilinear SPDEs driven by cylindrical stable processes. *Probab. Theory Related Fields* 149 (2011), no. 1-2, 97-137.
- [16] B. Rüdiger and G. Ziglio. Itô formula for stochastic integrals w.r.t compensated Poisson random measures on separable Banach spaces. *Stochastics*, 78:377-410, 2006.
- [17] J. Wang and Y. Rao. A note on stability of SPDEs driven by  $\alpha$ -stable noises. *Adv. Difference Equ.* 2014, 2014:98, 11 pp.
- [18] J. H. Zhu and Z. Brzezniak. Nonlinear stochastic partial differential equations of hyperbolic type driven by Lévy-type noises. *Discrete Contin. Dyn. Syst. Ser. B*, 21 (2016), 3269-3299.

GRADUATE SCHOOL OF SCIENCE, KYOTO UNIVERSITY, KITASHIRAKAWA-OIWAKECHO, SAKYO-KU, KYOTO 606-8502, JAPAN

*Email address:* nagoji.hirotsu.63x@st.kyoto-u.ac.jp