

Definable proper quotients

Masato Fujita Japan Coast Guard Academy and
Tomohiro Kawakami Department of Mathematics, Wakayama University

Abstract

Consider a definably complete locally o-minimal expansion $\mathcal{F} = (F, +, \cdot, <, 0, 1, \dots)$ of an ordered field. We prove the existence of definable quotients of definable sets by definable equivalence relations when certain conditions are satisfied. These conditions are satisfied when X is a locally closed definable subset of F^n and there is a definable proper action of a definable group G on X . We give its application.

1 Introduction

We study definable proper quotients of definable sets by definable equivalence relations for a definably complete locally o-minimal expansion of an ordered field in this paper [3].

An expansion $\mathcal{M} = (M, <, \dots)$ of dense linear order $<$ without endpoints is *locally o-minimal* if for any point $x \in M$ and for any definable subset A of M , there exists an open interval $I \ni x$ such that $A \cap I$ is a finite union of open intervals and points [6]. We say that \mathcal{M} is *definably complete* if any nonempty subset A of M has $\sup A, \inf A \in M \cup \{-\infty, \infty\}$ [4].

This paper considers a definable proper quotients and its application.

2020 *Mathematics Subject Classification.* 03C64.

Key Words and Phrases. Locally o-minimal, definably complete, definable proper quotients.

2 History of definable quotients

Definition 2.1. A semialgebraic equivalence relation E on a semialgebraic set X is a semialgebraic subset of $X \times X$ such that the binary relation \sim_E defined by $x \sim_E y \Leftrightarrow (x, y) \in E$ is an equivalence relation defined on X .

We say that E is *proper* over X if the projection $p : E \rightarrow X$ onto the first factor is proper. That is the inverse image of every bounded closed subset C in the whole space F^n with $C \subset X$ is bounded and closed in F^{2n} .

In 1987, Brumfiel [1] considers semialgebraic proper quotients. Let X be a semialgebraic set

Theorem 2.2 (1.4 [1]). *Suppose that $E \subset X \times X$ is a closed semialgebraic equivalence relation such that the projection $p : E \rightarrow X$ onto the first factor is proper. Then there exists a topological map $f : X \rightarrow Y$ with $E = E_f$. Moreover any such f is proper, and conversely, any proper $f : X \rightarrow Y$ induces an equivalence relation $E_f \subset X \times X$ proper over X .*

Consider an o-minimal expansion $\mathcal{F} = (F, +, \cdot, <, \dots)$ of a real closed field F .

Theorem 2.3 (van den Dries [2]). *Let E be an equivalence relation on a definable set X which is definably proper over X . Then X/E exists as a definable proper quotient of X .*

Scheiderer [5] proves the following theorem.

Theorem 2.4 ([5]). *Let M be a locally complete semialgebraic space and $E \subset M \times M$ a closed semialgebraic equivalence relation. Then the following are equivalent.*

- (1) *The geometric quotient M/E exists.*
- (2) *There exists a semialgebraic subspace $K \subset M$ such that $p_1|_{p_2^{-1}(K)} : p_2^{-1}(K) \rightarrow X$ is proper and surjective.*

3 Our results

Let $\mathcal{F} = (F, +, \cdot, <, \dots)$ be a definably complete locally o-minimal expansion of an ordered field F . Suppose that $X \subset F^m, Y \subset F^n$ are definable sets

and $f : X \rightarrow Y$ is a definable continuous map.

Definition 3.1. *The map f is definably proper if for any closed bounded definable subset K in F^m with $K \subset Y$, the inverse image $f^{-1}(K)$ is closed and bounded in F^m . It is called definably indentifying if it is surjective and for any definable subset Y , K is closed in Y whenever $f^{-1}(K)$ is closed in X .*

Let $\mathcal{F} = (F, +, \cdot, <, \dots)$ be a definably complete locally o-minimal expansion of an ordered field F , X a definable set and E a definable equivalence relation. A *definably quotient* of X by E is a definable identifying map $f : X \rightarrow Y$ such that $f(x) = f(x')$ if and only if $(x, x') \in E$. A *definably proper quotient* of X by E is a definable quotient which is definably proper. The target space Y is denoted by X/E .

Let $\mathcal{F} = (F, +, \cdot, <, \dots)$ be a definably complete locally o-minimal expansion of an ordered field F , X a definable set and E a definable equivalence relation. Suppose that $p_i : E \rightarrow X$ is the restriction of the projection $X \times X \rightarrow X$ onto the i -th factor X for $i = 1, 2$. The equivalence relation E is *definably proper over X* if p_1 is a definably proper map.

Let $G \subset F^m$ be a definable set. We say that G is a *definable group* if it is a group and the group operations $G \times G \rightarrow G$ and $G \rightarrow G$ are definable continuous maps. A *definable G set* is a pair (X, ϕ) consisting of a definable set X and a group action $\phi : G \times X \rightarrow X$ is a definable continuous map. We simply write X instead of (X, ϕ) . A G invariant definable subset A of a definable G set is called *definable G subset*. The G action on X is *definably proper* if the map $G \times X \rightarrow X \times X, (g, x) \mapsto (gx, x)$ is a definable proper map.

Theorem 3.2 ([3]). *Consider a definably complete locally o-minimal expansion $\mathcal{F} = (F, +, \cdot, <, \dots)$ of an ordered field F . Suppose that X is a nonempty locally closed definable set and E is a definable equivalence relation on X which is proper over X . Then there exists a definable proper quotient $\pi : X \rightarrow X/E$.*

Theorem 3.3 ([3]). *Consider a definably complete locally o-minimal expansion $\mathcal{F} = (F, +, \cdot, <, \dots)$ of an ordered field F . Suppose that G is a definable group and X is a definable G set which is locally closed. Assume that the action on X is definable proper. Then there exists a definable quotient $X \rightarrow X/G$.*

Theorem 3.4 ([3]). *Consider a definably complete locally o -minimal expansion $\mathcal{F} = (F, +, \cdot, <, \dots)$ of an ordered field F . Suppose that G is a definable group, X is a definable G set which is locally closed and A is a closed definable G subset of X . Assume that the action on X is definable proper. Then there exists a G invariant definable continuous function $f : X \rightarrow F$ with $A = f^{-1}(0)$.*

References

- [1] G. Brunfiel, *Quotient spaces for semialgebraic equivalence relations*, Math. Z. **195** (1987), 69-78.
- [2] L. van den Dries, *Tame topology and o -minimal structures*, Lecture notes series **248**, London Math. Soc. Cambridge Univ. Press (1998).
- [3] M. Fujita and T. Kawakami, *Definable quotients in locally o -minimal structures*, arXiv:2212.06401.
- [4] C. Miller, *Expansions of dense linear orders with the intermediate value property*, J. Symbolic Logic **66** (2001), 1783-1790.
- [5] C. Scheiderer, *Quotients of semi-algebraic spaces*, Math. Z., **201** (1989), 249-271.
- [6] C. Toffalori and K. Vozoris, *Notes on local o -minimality*. MLQ Math. Log. Q. **55** (2009), 617-632.