



Research



Metabarcoding analysis provides insight into the link between prey and plant intake in a large alpine cat carnivore, the snow leopard

Cite this article: Yoshimura H, Hayakawa T, Kikuchi DM, Zhumabai Uulu K, Qi H, Sugimoto T, Sharma K, Kinoshita K. 2024 Metabarcoding analysis provides insight into the link between prey and plant intake in a large alpine cat carnivore, the snow leopard. *R. Soc. Open Sci.* **11**: 240132.

<https://doi.org/10.1098/rsos.240132>

Received: 23 January 2024

Accepted: 27 March 2024

Subject Category:

Ecology, conservation, and global change biology

Subject Areas:

molecular biology, behaviour, ecology

Keywords:

diet analysis, DNA barcoding, carnivore, felids

Author for correspondence:

Hiroto Yoshimura

e-mail: hrt.yoshimura@gmail.com

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.7162451>.

Hiroto Yoshimura¹, Takashi Hayakawa^{3,4}, Dale M. Kikuchi⁵, Kubanychbek Zhumabai Uulu⁶, Huiyuan Qi¹, Taro Sugimoto⁷, Koustubh Sharma^{6,8} and Kodzue Kinoshita²

¹Wildlife Research Center, and ²Graduate School of Asian and African Area Studies, Kyoto University, Kyoto, Japan

³Faculty of Environmental Earth Science, Hokkaido University, Sapporo, Hokkaido, Japan

⁴Japan Monkey Center, Inuyama, Aichi, Japan

⁵Department of Bioresource Development, Tokyo University of Agriculture, Kanagawa, Japan

⁶Snow Leopard Foundation in Kyrgyzstan, Bishkek, Kyrgyzstan

⁷Institute of Natural and Environmental Sciences, University of Hyogo, Tamba, Hyogo, Japan

⁸Snow Leopard Trust, Seattle, WA, USA

HY, 0000-0003-1626-0790; TH, 0000-0002-1037-0350; KZU, 0000-0002-6184-4018; HQ, 0000-0003-1337-2455; TS, 0000-0002-3217-051X; KS, 0000-0001-7301-441X; KK, 0000-0002-5804-5485

Species of the family Felidae are thought to be obligate carnivores. However, detection of plants in their faeces raises questions about the role of plants in their diet. This is particularly true for the snow leopard (*Panthera uncia*). Our study aimed to comprehensively identify the prey and plants consumed by snow leopards. We applied DNA metabarcoding methods on 90 faecal samples of snow leopards collected in Kyrgyzstan, employing one vertebrate and four plant markers. We found that argali (*Ovis ammon*) was detected only from male snow leopards. *Myricaria* sp. was the most consumed among 77 plant operational taxonomic units found in snow leopard samples. It frequently appeared in samples lacking any prey animal DNA, indicating that snow leopards might have consumed this plant especially when their digestive tracts were empty. We also observed differences in the patterns of plant consumption between male and female snow leopards. Our comprehensive overview

of prey and plants detected in the faeces of snow leopards and other sympatric mammals will help in formulating hypotheses and guiding future research to understand the adaptive significance of plant-eating behaviour in felids. This knowledge supports the enhancement of their captive environments and the conservation planning of their natural habitats.

1. Introduction

Animals interact with plants in multi-faceted ways, using them as sources of food, shelter, tools and medicine. When studying diets, researchers have predominantly focussed on plant species as primary food sources for herbivores and omnivores. Plants are not easy to digest, especially due to their cellulose-rich cell walls [1,2]. The inherent structural distinctions between plant and animal cells, compounded by potential toxicities in some plant species, necessitate specialized digestive and detoxification mechanisms for those consuming a predominantly plant-based diet [3–5].

Felids are primarily recognized as obligate carnivores. They possess specific morphological, physiological and behavioural adaptations that enable them to efficiently consume other animals. These adaptations, including specialized dentition for slicing and a shorter digestive tract, reflect their carnivorous diet [6–8]. The shorter digestive tract is an adaptation to their reduced need for fermentation [9], as animal tissue is easier to digest than plant matter [10]. Furthermore, felid taste receptors have evolved in response to their dietary needs, showing decreased sensitivity to fruit sugars but increased sensitivity to amino acids and lower tolerance to bitter compounds [11–16]. Although their morphological and physiological states are highly tuned to carnivorous diet, carnivores have been observed consuming plants, both in the wild and in captivity [17–19]. Plant occurrence in faeces was reported in 24 extant felid species [18], and camera traps captured three wild felid species eating grass in Costa Rica [17].

The reasons and implications behind interactions between felids and plants remain ambiguous. Plant-eating behaviour might be overlooked due to the presumption that plants play a minimal role in felid nutrition or might be ingested inadvertently (e.g. [20,21]). Considering captive felids thrive without consuming plant materials in zoos, little is known about their necessity for survival. Some posit plants act as supplementary food or moisture sources, as indicated by the presence of fruit seeds in felid faecal samples [22,23]. Others suggest that plants serve medicinal purposes, aiding in parasite excretion [24,25] or digestion [26]. A commonly held notion also suggests plants assist in the evacuation of hair and undigested material [27–29].

Understanding which plants wild felids use is pivotal for experimental design and hypothesis formulation to explore adaptive significance of the behaviour. It is important to use similar plants to the ones consumed in the wild when conducting experiments related to plant-eating behaviour. Given that plant-eating behaviour is both common and natural among wild felids [18], information about the plants they consume can enhance their captive environments (e.g. introduce plant species consumed by wild individuals into enclosures) and aid in the conservation planning of their natural habitats. Additionally, it enriches our understanding of the ecological interactions between felids and plants.

The snow leopard (*Panthera uncia*) is native to the high mountains of central Asia. Plant material has been reported in the faeces of 24 out of 41 extant felid species; notably, snow leopard faeces frequently contained plant materials, despite their alpine habitat where vegetation is typically sparse [18]. Previous studies have made cursory mentions of grasses and bushes; in particular, *Myricaria* spp. [30], and 45% of their faeces contained the shrub *Myricaria* spp. [31] in prey animal surveys but have not investigated the plant species further. Therefore, it is unclear if *Myricaria* spp. is more frequently consumed than other plants and if this is a phenomenon specific to snow leopards compared to other animals. We believe that investigating the plant repertoire consumed by wild snow leopards in alpine environments will deepen our understanding of the plant-eating behaviour, including which plants they consume despite limited plant resources.

The molecular approach using the next-generation sequencing is widely used in diet analysis for many animals [32]. DNA metabarcoding employs specially designed universal primer pairs to amplify standardized regions of DNA. These regions are then sequenced and matched against reference databases for taxonomic identification [33]. When combined with next-generation sequencing, this technology enables the concurrent taxonomic analysis of thousands of samples efficiently and economically [34]. DNA metabarcoding is considered suitable for identifying the potentially diverse

dietary plants in carnivores. However, application of metabarcoding method for plant identification in felids is quite limited. A study of leopard cat (*Prionailurus bengalensis*) in China is the only case at the moment [25], and there are few studies that use this method for large cat species.

The primary objective of this study was to comprehensively identify plant species in the faeces of wild snow leopards. This would enable us to infer the traits of plants they frequently consume and the function of plant-eating. Based on Illumina sequencing data, we first revealed the frequently consumed prey and plant taxa in snow leopard faeces. Additionally, we identified the dietary plant species consumed by other mammals inhabiting the same alpine ecosystem. These included ibex (*Capra sibirica*) and argali (*Ovis ammon*) that constitute the primary prey for snow leopard, wolf (*Canis lupus*) that is another apex predator species in its habitat and red fox (*Vulpes vulpes*) that functions as a mid-level predator and omnivore. By contrasting the dietary composition of the other mammal species, we can understand the characteristics of plant eating of the snow leopard. It is equally critical to consider intraspecies variations, such as differences between sexes. A study of Puma showed that sex affects the species and size of prey [35]. Additionally, the difference in reproductive roles between sexes influences their behaviour and energy requirements [36], potentially impacting plant-eating behaviour. Consequently, we tried to find out whether the dietary composition differs between sexes in snow leopards.

2. Material and methods

2.1. Ethical note

This research adhered to the legal requirements of the governments of Kyrgyzstan and Japan. All sampling procedures were non-invasive, granted by the State Agency on Environment Protection and Forestry (now Ministry of Natural Resources, Ecology and Technical Supervision) of the government of Kyrgyzstan, and carried out according to the guidelines for animal studies in the wild and ethics in animal research issued by the Wildlife Research Center of Kyoto University.

2.2. Study area

The Sarychat-Ertash Reserve (42°02'N 78°25'E) spans 1341 km² in the Central Tien-Shan Mountain range's Uch-Kol River basin. It is characterized by altitudes of 2000–5500 m and experiences a cold continental climate with mean monthly temperatures in June and January of +4.2 and –21.5°C, respectively, and annual precipitation of 295 mm. The Reserve's vegetation consists of arid grasslands, wet meadows and tundra cushion plants [37]. Snow leopard, wolf and red fox are the most common carnivores; brown bear (*Ursus arctos*), lynx (*Lynx lynx*), Palla's cat (*Otocolobus manul*) and stone marten (*Martes foina*) are the other carnivores found there. In addition to ibex and argali, potential snow leopard and wolf prey species include marmot (*Marmota baibacina*), hare (*Lepus tolai*), pika (*Ochotona roylei*) and birds such as snowcock (*Tetraogallus himalayensis*) and chukar partridge (*Alectoris chukar*). There are reports of four species of mustelids and four vole species in the area [37]. Historically affected by human activities such as livestock grazing and illegal hunting, only a small part of the Reserve's buffer zone is used for seasonal livestock grazing.

2.3. Sample collection

The faecal samples were collected in November 2017, March and September 2018, May 2019, October 2022 and May 2023. High water levels in the summer and thick snow cover in the winter prevented field-work in these seasons during the year. Faecal samples were collected opportunistically. We collected faecal droppings of ungulates in addition to those of carnivores in the autumn of 2022. Typically, whole faeces were collected into plastic bags with silica gel after photographing it in its natural setting. Geographical coordinates, altitude and sampling time were recorded. Since refrigerating facilities were not available at the study site, faecal samples were stored in a dark place at ambient temperature until they were brought to Bishkek city. Surface of each faeces was swabbed by sterile cotton swab and preserved in sterilized 2 ml plastic tubes with 1 ml lysis buffer (0.5% sodium dodecyl sulfate, 100 mM ethylenediaminetetraacetic acid (pH 8.0), 100 mM Tris-HCl (pH 8.0) and 10 mM NaCl [38]), mixed by tapping the tube, and kept in dark boxes at ambient temperature for later processing.

Table 1. List of primers used in the study.

usage	name	primer sequence (5'–3')	reference
species identification	16SrRNA_L2513_felid	GCCTGTTTACCAAAAACATCAC	this study
	16SrRNA_H2714_felid	CTCCATAGGGTCTTCTCGTCTT	
sex identification	ZFX-PF	TACCGAGCGATATAGCTCCAG	Sugimoto <i>et al.</i> [40]
	ZFX-PR	GTGTTCTACGTTAAGCTATTG	
diet analysis	DBY7-PF	CTCATGAAGCCCTATTTTGGTTG	
	DBY7-PR	ACGGCGTCCGTATCTCCA	
	12SV5F	TAGAACAGGCTCCTCTAG	Riaz <i>et al.</i> [41]
	12SV5R	TTAGATACCCCACTATGC	
	UniplantF	TGTGAATTGCARRATYCMG	Moorhouse Gann <i>et al.</i> [42]
	UniplantR	CCCGHYTGAYYTGRGGTCDC	
	rbcl-F	CTTACCAGYCTTGATCGTTACAAAGG	Erickson <i>et al.</i> [43]
	rbcl-R	GTAATAATCAAGTCCACCRCG	Kress and Erickson. [44]
	trnL-g	GGGCAATCCTGAGCAA	Taberlet <i>et al.</i> [45]
	trnL-h	CCATTGAGTCTCTGCACCTATC	
	ITS1-F	GATATCCGTTGCCGAGAGTC	Baamrane <i>et al.</i> [46]
	ITS1Poa-R	CCGAAGGGCTCAAGGAACAC	

To reduce the environmental contamination, we cut the faeces with sterile tweezers and transferred the inner parts into 2 ml sterilized plastic tubes with 1 ml RNA*later* solution (Thermo Fisher Scientific, Waltham, MA). The contents in the tube were mixed by tapping the tube, and they were kept in a dark box at ambient temperature for later processing. Samples in the lysis buffer, specifically surface swabs, were used for species and sex identification due to the expected higher concentration of host DNA. The samples taken from the inner region of faeces were used for diet analysis.

2.4. DNA extraction

All experimental procedures were performed under sterile conditions, as recommended by Hayakawa *et al.* [39]. DNA from each faecal sample was extracted and purified using the QIAmp DNA Fast Stool Mini Kit (Qiagen, Hilden, Germany). DNA from samples stored in the lysis buffer was extracted according to the manufacturer's protocol. Samples stored in RNA*later* were first precipitated and then washed twice with 1 ml of phosphate-buffered saline (pH 7.4; centrifugation speed: 20 000g for 10 min). Each of the processed samples was disrupted using four zirconia beads (3 mm in diameter) and 1 mg zirconia/silica beads (0.1 mm in diameter) in a 2 ml plastic tube at 4200g for 5 min. The DNA samples were then purified using the QIAmp DNA Fast Stool Mini Kit and eluted in 100 µl of Buffer ATE with 30 min of incubation at ambient temperature. The DNA concentrations were estimated with a Qubit dsDNA HS Assay Kit and a Qubit fluorometer (Thermo Fisher Scientific). The purified DNA samples were stored at 4°C.

2.5. Species identification

We used molecular species identification to identify the specific origin of each faecal sample. To accomplish that we designed a 16S rRNA primer pair (16SrRNA_L2513_felid: GCCTGTTTACCAAAAACATCAC; 16SrRNA_H2714_felid: CTCCATAGGGTCTTCTCGTCTT) to amplify a –244 bp (excluding the primers) mitochondrial 16S rRNA gene sequence (table 1). The PCR conditions and programs are provided in the electronic supplementary file. The PCR products were purified by using a High Pure PCR Product Purification Kit (Roche, Basel, Switzerland). Direct sequencing was performed using the Big Dye 3.1 Terminator cycle-sequencing kit (Applied Biosystems, Foster City, CA) according to the manufacturer's instructions. The cycle sequencing products were purified by ethanol precipitation and nucleotide sequences were determined using

an ABI PRISM 3130xl genetic analyzer (Applied Biosystems). Forward and reverse complement sequences were aligned using MEGA11 [47]. The resulting sequences were searched in the GenBank nucleotide (nt) database and species identity was determined based on the matches with the highest similarity scores (95–100%).

2.6. Sex identification

After identification of species as snow leopard, we identified sex of the individual that the sample belonged to. We used one set of four primer targeting introns of zinc-finger in X chromosome; ZFX-PF/PR and DEAD box polypeptide in Y chromosome; DBY7-PF/PR [40] (table 1). The PCR conditions and programs are provided in the electronic supplementary file. PCR products were electrophoresed and visualized on 2.0% agarose gels. The same procedure was repeated at least twice, and the sex was determined only when the results were consistent.

2.7. Library preparation and amplicon sequencing

Library preparation and amplicon sequencing were performed with the MiSeq system (Illumina, Inc., San Diego, CA) according to the manufacturer's protocol with modifications optimized for our sample as follows. Five different marker sets were used to analyse species' diets. A universal vertebrate 12SV5 marker [41]; three universal plant markers: Uniplant [42], rbcL mini-barcode [43,44], trnL-g/h [45]; and finally, one Poaceae-specific marker; ITSPoa [46], to increase the taxonomic resolution for grasses (table 1). These primers were fused with 3-6-mer Ns and specific overhang adapters 5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-(forward primer)-3' and 5'-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG-(reverse primer)-3'. PCR was performed using the KAPA HiFi HotStart ReadyMix PCR Kit (Kapa Biosystems, Inc., Wilmington, MA) with 200 nM of each primer and 25 ng DNA as the template in a total volume of 25 μ l. The PCR conditions and programs are in the electronic supplementary file. When 25 ng DNA was unavailable due to low DNA yield, the available maximum volume of the DNA solution was used in the PCR. The resulting amplicons were visualized on agarose gels.

Each PCR product (20 μ l) including negative controls was purified using 36 μ l Agencourt AMPure XP beads (Beckman Coulter, Inc., Carlsbad, CA) with 80% ethanol washes. Each of the purified PCR products was eluted in 10 mM Tris-HCl (pH 8.5). Using the KAPA HiFi HotStart ReadyMix PCR Kit and the Illumina Nextera XT Index Kit v2, specific dual indices and sequencing adapters were attached to each amplicon by PCR conducted in a 50 μ l solution containing 5 μ l of each of the forward and reverse primers and 5 μ l of the first purified PCR solution. The resulting amplicons were visualized on agarose gels. Each product (45 μ l) was purified using Agencourt AMPure XP beads with 80% ethanol washes. Each of the purified products from the second PCR was eluted in 27.5 μ l of 10 mM Tris-HCl (pH 8.5).

The DNA concentration of each product was measured with a Qubit dsDNA HS Assay Kit. Products were mixed in the same amount of DNA concentrations to form the pooled sequencing library. Fragment size distribution of the library was estimated with an Agilent 2200 TapeStation (Agilent Technologies, Inc., La Jolla, CA, USA). The library was diluted to 15 pM and subjected to a sequencing run mixed with other libraries unrelated to this study and 30% PhiX spike-in on an Illumina MiSeq sequencing platform using the MiSeq Reagent Kit v3 (600 cycles). Sequencing was separately operated in four different runs. The read lengths from the MiSeq run were 301 bp (forward sequences), 8 bp (forward indices), 8 bp (reverse indices) and 301 bp (reverse sequences). Although quality scores of nucleotides at the 3'-end of Illumina sequences are generally low, the amplicon sizes of this study were smaller than the number of cycles of the kit (i.e. 600). Therefore, overlapping regions of the forward and reverse reads were used to restore these low-quality sequences in the following bioinformatics procedure.

2.8. Bioinformatics

As suggested in [48], we converted the raw MiSeq BCL data into FASTQ data by ourselves using the bcl2fastq v. 2.20.422 program distributed by Illumina to prevent the potential de-multiplexing errors, and we then de-multiplexed the FASTQ data using the program Claident v. 0.9.2022.04.28. In

the de-multiplexing and primer-trimming process with Claident, all the sequencing reads containing low-quality index (quality scores <30) sequences were eliminated and no mismatch between input and output index sequences was tolerated. Adapter sequences were trimmed using Skewer; <https://sourceforge.net/projects/skewer> [49], and the forward and reverse sequences were corrected with DADA2; <https://github.com/benjjneb/dada2> [50] package on R programming interface [51]. Reads containing ambiguous bases were removed and trimming lengths were adjusted based on sequence quality profiles, so that Q-scores remained above 30. Error model calculation (for R1F/R2R read pairs and then R2F/ R1R read pairs), read correction, read merging and chimera removal was performed at default settings implemented in DADA2. All the resulting amplicon sequencing variant (ASV) tables were curated with LULU [52] package on R to remove spurious ASVs. As the aim of the present study was to detect and identify species, and not intraspecific variation, we decided to create clusters of sequences, instead of denoising and creating ASV [53,54]. According to the developers of LULU algorithm, incorporation of DADA2 and LULU is a safe pathway for producing reliable and accurate metabarcoding data [52]. The LULU curation requires an external algorithm to produce the match list. Thus, we used VSEARCH v. 2.21.1 as recommended by the developers [52].

The remaining operational taxonomic units (OTUs) were then subjected to molecular taxonomic identification based on the automatic database search algorithm of the query-centric auto-*k*-nearest-neighbor (QCauto) method [55] and subsequent taxonomic assignment with the lowest common ancestor (LCA) algorithm [56] using Claident. Among the filtered databases bundled with Claident, we used the 'animals_mt_genus' and 'animals_mt_species' sub-databases for 12SV5 region; 'plants_rbcL_genus' and 'plants_rbcL_species' sub-databases for rbcL mini-barcode; 'plants_cp_genus' and 'plants_cp_species' sub-databases for trnL g-h; 'overall_genus'; 'overall_species' databases for Uniplant and ITS1Poa. The QCauto search information was then subjected to taxonomic assignment with the LCA algorithm (LCA/genus results). As the default setting of the LCA algorithm sometimes returns conservative results, additional taxonomic assignment was conducted with a relaxed setting tolerating 5% mismatches of taxonomic information among database sequences in the LCA process; relaxed-LCA/genus [55]. The overall identification results were obtained by merging the LCA/species, LCA/genus and relaxed-LCA/genus results in this priority order using the 'clmergeassign' command of Claident. Since the QCauto method is conservative [55], we conducted additional megablast search for 12SV5 marker and complemented the taxonomic assignment. If an OTU was assigned to several species and we knew which candidate species inhabited the study area [57], we assigned the inhabiting species to the OTU. When several local species were assigned with same probability or no species was assigned with >95% match, we kept the QCauto result.

Index hopping rate of MiSeq is estimated to be 0.001 [58]. To ensure that index hopping did not result in false positives, the reads of the OTU in the samples were removed whenever the number of reads of an OTU detected in each sample were <0.001 of the number of reads of the OTU detected in all samples [59]. A recent study showed that a combination of a sample-based threshold with removal of maximum taxon contamination is an optimal method to remove artefacts [60]. Following the suggested filtering process [60], read counts within a sample that are less than a proportion of the total sample read count for that sample were removed. We decided the threshold proportion to 0.01 and 0.05 for the universal markers [43,61,62] and ITS1Poa [63], respectively. Threshold proportion of 12SV5 varied from 0.001 [64] to 0.05 [65], thus we chose 0.01 as other universal markers. In addition, we removed any read count within each OTU that lower than the highest read count within a negative control or blank cells for that OTU [60]. Based on the molecular taxonomic identification results, non-target OTUs (non-vertebrate and human in 12SV5, non-plant for the three universal plant markers, and those not in Poaceae family for ITS1Poa) were excluded. The OTUs from host carnivore species were also excluded from the 12SV5 dataset. The sequencing read set of each sample was rarefied to the minimum coverage rate among the analysed samples [66] using vegan [67] package of R. The coverage rate of each marker was 1.00, 1.00, 0.83, 0.83 and 1.00, for 12SV5, Uniplant, rbcL, trnL and ITS1Poa, respectively.

In order to overcome problems of primer specificity and bias, we integrated information from the four molecular markers used for plant identification using the Python 3.0 script [62]. The script provides a single list of taxa detected per sample controlling for duplications by collapsing less resolved taxa detected by one marker with higher resolved taxa detected using a different marker [62]. The ITS1Poa marker was Poaceae-specific marker to improve the resolution of grasses. Considering that Poaceae are common and faecal samples are often on the ground with grasses, Poaceae-specific amplification may increase the risk of amplification of rare sequences contaminated from the environment. Therefore, data from the ITS1Poa marker were only merged to the samples in which Poaceae sequence was detected by other three universal makers.

2.9. Statistical analysis

Visualization and basic statistical analyses were performed using the phyloseq v.1.26.1 [68] package in R. Dietary data were summarized across samples using two occurrence-based metrics commonly used in molecular dietary data analysis: (i) frequency of occurrence (FOO) and (ii) weighted per cent of occurrence (wPOO) [69]. The number of samples that contain a given food item is expressed as FOO, whereas wPOO weighs each occurrence according to the number of food items in the sample (i.e. lower weights to individual food taxa in a mixed meal), which is considered to be more biologically realistic [69]. Since we merged data from multiple markers, we did not use a sequence abundance-based metric. The samples without any prey or plant OTUs were excluded from subsequent statistical analyses.

A machine learning-based classification approach was applied to clarify the difference in dietary plant composition of snow leopards and other mammals. We used machine learning models using the randomForest package [70] in R to determine which plant genera best discriminated whether a sample came from snow leopard or other sympatric mammals based on the sample–plant matrix [71,72]. RandomForest evaluates an ensemble of decision trees to perform classification [70], in this instance, it classifies snow leopards and other mammals based on the plant composition in their faecal samples. Random forest models are considered to be robust against overfitting and known to have high predictive accuracy [73]. We tuned the random forest models to determine the number of variables (mtry) to try at each node of the tree that resulted in the lowest out-of-bag (OOB) error rates using randomForest function. OOB error is an internal validation method, estimating the prediction error of random forest models by using bootstrap samples not included in the construction of each tree. Since the number of samples were biased toward snow leopard, classwt option with inverse of the ratio of the sample size was used to enforce penalties for errors in minority category. Random forests estimate the variable importance. Thus, we were able to identify which plant genera represents the snow leopard faeces. Random forests provide two indicators for variable importance: mean decrease accuracy (MDA) and mean decrease gini (MDG). MDG is considered to be more stable than MDA [74], therefore we used MDG as the indicator.

A post hoc probabilistic co-occurrence analysis was conducted to show which taxa are simultaneously present in the same faecal samples of predators using package co-occur [75] in R. A prey-specific co-occurrence would indicate secondary predation of prey gut content [76]. In addition, we summed up all prey OTUs as single ‘prey’ OTU and evaluated co-occurrence with each plant OTU that indicated accidental intake from the environment such as grasses on the ground. Conversely, when a plant OTU negatively co-occurred with prey OTUs, it is more likely to be consumed by snow leopards.

The difference in the dietary composition between each sex of snow leopard was assessed by permutational analysis of variance (PERMANOVA) using the adonis2 function with 9999 permutations and visualized by non-metric multi-dimensional scaling (NMDS) using Bray–Curtis dissimilarity, as implemented in vegan [67].

Constrained analysis of principal (CAP) coordinates [77] was performed to evaluate dietary composition differences between male and female snow leopards (model 1: dietary animal composition, model 2: dietary plant composition) while accounting for the effects of sampling season and spatial autocorrelation. Spatial autocorrelation variables were added to consider the effect of spatial proximity on the sample. We first generated a set of Moran’s eigenvectors from the coordinates of each sampling point using distance-based Moran’s eigenvector maps; MEMs [78]. We then identified positive MEMs that significantly ($p < 0.05$) described spatial patterns using the function ‘moranNP.randtest’ using the R package adespatial [79]. The two CAP models were constructed by setting the Bray–Curtis dissimilarity as a response variable. The models included ‘Sex’, ‘Sampling season’, ‘Altitude’ and MEM vectors as explanatory variables. April and May were defined as spring while September and October were defined as autumn. The variables’ variance inflation factors (VIFs) were computed to check the collinearity. If the variable’s VIF was above 20, the variable was excluded from the model [80], resulting in different degrees of freedom in MEMs. The significance of models and explanatory variables were tested using permutational analysis of variance with 9999 permutations.

Table 2. The number of identified taxa for each marker. The total number of plant OTUs after merging the four plant markers is labelled as 'merged'.

marker	order	family	genus	species	OTUs
12SV5	5	7	11	11	13
uniplant	11	17	25	5	69
rbcl	9	13	13	1	34
trnL	20	28	19	0	62
ITS1Poa	1	1	10	1	48
merged	21	29	44	7	141

3. Results

3.1. Summary of sequence data

We collected 150 mammal faecal samples in total out of which, we could genetically identify the host species of 126 samples. These samples (90 snow leopards; 7 wolves; 9 red foxes; 3 brown bears; 9 ibexes; 7 argali; 1 marmot) were used in the dietary analysis. We obtained 16 623 180 raw sequence reads after de-multiplexing (27 567 reads per sample on average; standard deviation 25 858 reads). The 11 453 323 (20 862 reads per sample on average; standard deviation 21 185 reads) that passed the filtering processes were used as curated OTUs in the following analysis. In total, 13 prey animal OTUs were identified, with 11 at the species level, 1 at the family level and 1 at the order level. Additionally, of the 141 plant OTUs, 7 were identified at the species level, 81 at the genus level, 49 at the family level and 1 at the order level.

The results from FOO and wPOO were qualitatively similar, thus we show wPOO-based results for subsequent dietary analysis in the main text. The results from FOO were shown in the electronic supplementary file, figures S1–S4.

For snow leopard samples, on average, 21 367 reads per sample passed the filtering processes for animal, and 15 319 reads per sample passed for plants. We detected 8 animal prey OTUs and 77 plant OTUs. There were 51 samples from male and 27 samples from female snow leopards. We could not determine the sex in the remaining 12 samples.

3.2. Prey

In total, 13 OTUs were found from predator samples (table 2). Wild ungulates and marmots were the frequently detected prey species in Sarychat-Ertash for the larger predators. Ibex was the main prey of snow leopards, whereas wolves and red foxes preyed more on marmots (figure 1, electronic supplementary material, figure S5). We detected smaller mammals mainly in red fox faecal samples.

Remarkably, we found no OTU of argali in samples from female snow leopards (figure 2). However, the result of PERMANOVA ($p = 0.128$) as well as CAP model ($p = 0.576$) did not show a significant difference between male and female for prey composition (table 3). The CAP model controlled the effect of spatial proximity and sampling season and explained 85% of total variance.

3.3. Plant

The Uniplant, rbcl, trnL and ITS1Poa markers detected 69, 34, 62 and 48 OTUs, respectively (table 2). The composition of detected plant taxa was similar among Uniplant and rbcl, but trnL showed different compositions (electronic supplementary material, figures S6–S9). The merged OTU table contained 141 OTUs. The order Caryophyllales was the most common in snow leopard samples from Sarychat-Ertash Reserve. Three plant families, Asteraceae, Tamaricaceae (*Myricaria* sp.) and Poaceae showed high wPOO (0.16, 0.29 and 0.17, respectively) in snow leopard samples (figure 3). While Asteraceae and Poaceae were also detected from




Figure 1. Weighted per cent of occurrence of vertebrate taxa for predators. The number in the parentheses shows the number of faecal samples.

other mammals, Tamaricaceae was rarely detected in other species. Wolf and fox samples often contained Poaceae (wPOO: 0.49 and 0.29, respectively), whereas argali and ibex typically consumed Asteraceae (wPOO: 0.11 and 0.24, respectively), Poaceae (wPOO: 0.22 and 0.31, respectively) and Chenopodiaceae; *Lepidium* spp., *Chenopodium* spp. and *Krascheninnikovia* spp. (wPOO: 0.27 and 0.22, respectively) in autumn.

We achieved a final OOB error rate of 12.61%. The model correctly identified snow leopards in 78% of the samples it labelled as snow leopards (precision), and it correctly found 83% of the actual snow leopard samples in the dataset (recall). *Myricaria* sp. was notably important plant genera to distinguish snow leopard samples from other sympatric mammals with MDG value of 13.3, compared with 3.9 for the next most important genus, and wPOO of this genus was higher in the snow leopard samples (electronic supplementary material, figure S10).

The post hoc co-occurrence analysis showed that *Myricaria* sp., which was a representative plant genus in snow leopard faeces, was negatively co-occurring with prey DNA; $p < 0.001$ (electronic supplementary material, figure S11) while *Festuca*, Rosaceae and *Ephedra* spp. OTUs in snow leopard faeces co-occurred with ibex OTU; $p = 0.002$, 0.006 and 0.006, respectively (figure 4).

Figure 5b shows the NMDS plot of plant composition (stress: 0.067) where we removed outliers to make it easy to interpret. The CAP model explained 49% of the total variances and showed that plant composition was different among sex ($p = 0.008$), as well as the result from PERMANOVA ($p = 0.049$). The effect of the sampling season (spring or autumn) was suggested, though it was not statistically significant (table 3). *Myricaria* sp. was not detected from female samples in autumn, when the presence of *Ephedra* spp., Asteraceae, Poaceae and Crassulaceae increased. In case of males, *Myricaria* sp. was detected during both seasons.




Figure 2. Weighted per cent of occurrence of vertebrate taxa in snow leopard faeces. The caption 'M' represents male samples and 'F' represents female samples. The number in the parentheses shows the number of faecal samples.

Table 3. CAP coordinates for factors structuring the prey and plant composition in snow leopard samples. Models included sex, altitude, sampling season and the MEM vectors (i.e. horizontal spatial structure) as explanatory variables. Model 1 included wPOO-based OTU matrix of prey and model 2 included that of plant. Significant variables are highlighted in bold.

model	OTU matrix	explanatory variable	d.f.	F	p
model 1	prey	sex	1	0.7	0.576
		altitude	1	1.4	0.272
		season	1	1.7	0.197
		MEMs	18	2.1	0.019
model 2	plant	sex	1	2.7	0.008
		altitude	1	0.3	0.984
		season	1	1.6	0.099
		MEMs	21	1.4	0.003

4. Discussion

4.1. Prey

Carnivores in Sarychat-Ertash relied on wild ungulates and marmots as reported in previous micro-histological research [31]. Conducted between June and October in 2009 at the same study site as ours, Jumabay-Uulu [31] reported higher occurrence of argali than ibex in the diets of snow leopard and wolves (18:3 for snow leopards and 12:8 for wolves). In contrast, our study finds occurrence of argali to be lower than that of ibex in the carnivore diets [31]. This variation may be due to differences in methodology (e.g. microhistological versus molecular, sampling season, sampling location) or ecological factors (e.g. changes in relative abundance). Many argali died due to unexpected heavy




Figure 3. Weighted per cent of occurrence of (a) the five most frequent plant genera and (b) the three most frequent plant families in faeces from each mammal. Less-frequent taxa were summarized as ‘other’. The number in the parentheses shows the number of faecal samples.




Figure 4. Co-occurrence matrix of plant OTUs and (a) each prey OTUs, (b) summarized prey OTU. Names of OTUs are positioned to indicate the columns and rows that represent their pairwise relationships with other OTUs. The colour of each cell represents positive, negative and random co-occurrence. Cells that show positive and negative co-occurrence of prey and plant were highlighted.

snowfall in 2022 (Zhumabai-uulu, pers. comm.), which might have influenced the proportion of argali and ibex in carnivores’ diets. One snow leopard sample contained a small number of red fox DNA reads ([red fox]:[snow leopard] = 180:8468), and two red fox samples contained a small number of snow leopard DNA reads ([red fox]:[snow leopard] = 23 892:1194 and 26 974:703). Red fox sometimes scavenges from snow leopard kill, thus snow leopard DNA in red fox samples was probably a by-product of scavenging. Red fox sometimes defecate close to the scrapings of snow leopard [81] and snow leopard is known to kill smaller predators such as red fox to prevent scavenging [82].




Figure 5. NMDS plot of wP00-based Bray–Curtis dissimilarity of (a) vertebrates from snow leopard samples (stress: 0.138) and (b) plants from snow leopard samples (stress: 0.067). Colour and shape correspond to different sex.




Figure 6. Weighted per cent of occurrence of plants from snow leopard samples. The caption ‘M’ represents males and ‘F’ represents females. The numbers in the parentheses are the number of samples collected in spring and autumn, respectively. The colour corresponds to different sampling seasons.

We did not find a significant difference in dietary prey items between sexes in snow leopard samples. One female sample contained *T. himalayensis* and another contained unidentified Artiodactyla (the order to which ungulates belong). Unidentified Artiodactyla OTU was believed to be a by-product of DNA degradation since the sample contained *C. sibirica* OTU as well (figure 5a). The limited diversity of potential prey mammals in the study area might have obscured any sex differences. Notably, only male samples contained traces of argali (figure 2). Although we did not identify individuals, argali was detected from male samples collected in 2018, 2022 and 2023. Females (~43 kg), being slightly smaller than males (~52 kg) might prey on argali; 60–185 kg, less often due to its larger size compared to ibex; 30–100 kg [30,83]. It will require further detailed investigation to determine whether this difference between male and female snow leopard’s consumption of argali was an artefact

of the size difference between male and female snow leopards [30,83], or the possibility that with their larger home ranges [84], male snow leopards were more likely to venture into sub-optimal habitat such as rolling terrain that are used by argali [85].

4.2. Plant

We found various plant taxa from snow leopard samples. The frequent detection of the genus *Myricaria* agreed with previous observation-based reports in the same study sites [31]. We evaluated the potential for secondary consumption by integrating co-occurrence analysis with the examination of dietary plant items present in the faecal samples of prey. The *Festuca*, Rosaceae and *Ephedra* OTUs in snow leopard faeces were non-randomly detected with ibex OTU indicating the possibility of secondary consumption from ibex gut content. Although *Ephedra* spp. was not detected from ibex samples collected in this study, a previous study reported that livestock ate *Ephedra*'s young shoot in early spring in China [86]. In this study, ibex samples were collected in autumn, thus seasonal food plant fluctuation might have prevented the detection of *Ephedra* OTUs. It is also reported that Siberian ibex in eastern Tianshan, China preferred eating forbes (Asteraceae, Gentianaceae, Rosaceae, Fabaceae) that have higher nutritional value than graminoids during the warm season and graminoids occupied a high proportion of their diet during the cold season [87]. Since our co-occurrence analysis is only exploratory, it does not necessarily confirm ecological interactions [76]. However, given that plants that positively co-occurred were reported to be frequently consumed by ungulates, their presence in snow leopard faeces is more likely attributable to secondary consumption rather than voluntary intake by the snow leopards. Despite the lack of significant co-occurrence with prey OTUs, the frequent detection of Asteraceae in prey faecal samples indicates a potential for secondary consumption.

On the other hand, *Myricaria* spp., the representative plant in snow leopards, tended to be detected from samples which did not contain any prey OTUs. This suggests that snow leopards intentionally consumed this bush more frequently, particularly when their digestive tracts were empty. The factors that cause snow leopards to intake *Myricaria* spp. may have some relationship to whether the individual obtained prey or not. In domestic cats, it has been hypothesized that constant availability of food (ad libitum feeding) may reduce the inclination to ingest alternative items such as plastic [88]. When the digestive tracts of felids are empty, they may exhibit a tendency to bite hard objects as a means to compensate for their appetite. One species in this genus, *Myricaria bracteata*, has been used in traditional Tibetan medicine and contains anti-inflammatory compounds [89], although its medicinal effects have not been specifically tested on snow leopards. Therefore, intake of *Myricaria* spp. and the failure to acquire prey may be related to the individual health condition of the snow leopards.

Female samples tended to contain *Ephedra* spp. and Asteraceae as often as *Myricaria* spp. Besides, no *Myricaria* spp. were detected from female samples collected in autumn (figure 6). As previously mentioned, *Ephedra* spp. and Asteraceae were suspected to be instances of secondary consumption. Since snow leopards give birth in mainly early summer [90], this difference could be resulting from the seasonal behavioural differences among males and females. The genus *Myricaria* is not a dominant plant in the study area and is sporadically distributed along rivers. Although there is little information about snow leopard nursing behaviour in the wild, during nursing period, mother may have preferred to stay closer to the cubs than proactively look for the *Myricaria* spp. patch. A female captive snow leopard exhibited a lower frequency of plant-eating behaviour in the year she shared an enclosure with her cub, as compared to the following year when the cub became independent [19]. While our results are indicative, it is important to consider the caveat of small sample size especially in case of female faeces. The CAP model explained only half of the total variance, thus there is a possibility of other factors, not included in this study, affecting the presence of plants in snow leopard diet.

An inter-regional sampling with a specific study design is required to better understand the relationship between snow leopards and plants. *Myricaria* spp. was often detected in snow leopard diet from other countries such as Nepal and India, but frequent containment of feather grass was reported in Mongolia [30]. Comparison of plant repertoire in different regions will provide answers to why snow leopards selectively intake on *Myricaria* spp. in this study area and identifying commonalities will lead to understanding the adaptive significance of plant-eating. In addition, a comprehensive vegetation survey is necessary to evaluate the preference in light of availability. Furthermore, adopting multi-faceted approaches, such as investigating the characteristics of the behaviour and the effects on immunity and digestion, is crucial to unravel the reasons behind plant-eating behaviour.

4.3. Limitation

Due to challenging terrain that limited human access, we could not establish a clear transect for sampling, and we did not identify individual animals for each faecal sample, possibly leading to sampling bias. Seasonal constraints further limited our study; for example, high water levels and deep snow prevented sampling in summer and winter. Additionally, the number of faecal samples from species other than snow leopards was limited, restricting our ability to perform statistical comparisons between species. While metabarcoding is powerful for diet analysis, its high sensitivity can also pick up environmental contamination or accidental intake [76]. We refrained from using host blocking primers to prevent unexpected amplification bias towards prey animals, though this approach might have lowered the sensitivity in detecting prey species. Sometimes, there was amplification bias as we found in plant markers in this study (electronic supplementary material, figures S6–S8). Although we took steps to minimize these biases, they couldn't be entirely eliminated. The resolution of the markers was not enough to identify plant OTUs at the species level, and incompleteness of plant references due to lack of a comprehensive vegetation survey in the area could have led to inflated diversity estimates for plant OTUs.

5. Conclusion

In this study, we applied a molecular-based approach to comprehensively investigate animal and plant in faeces of mammals in the alpine habitat of Kyrgyzstan. Detected prey items from large carnivores agreed with previous studies in the same study site. Red fox, a mesocarnivore, consumed smaller mammals as well. Although statistical significance was not detected, consumption of argali was biased toward male snow leopards indicating the possibility of prey selection according to the predator's body size.

We focused on dietary plants and highlighted the feature of plant repertoire in snow leopard faeces. As mentioned in an observation-based report, the genus *Myricaria* characterized the snow leopard samples. We found the plant was negatively co-occurred with animal prey DNA, indicating the consumption of this bush when the digestive tracts were empty. This suggests the importance of simultaneous investigation of prey and plant in carnivore diet.

Unveiling the relationship between snow leopards and plants, obligate carnivores and plants in general, improves our understanding of not only their behaviour and ecology but also evolution of diet repertoire and animal–plant interaction in the ecosystems. Practically, it aids in the conservation planning of felids, as conserving species interactions is considered important in conjunction with the conservation of individual species [91].

Ethics. This work did not require ethical approval from a human subject or animal welfare committee.

Data accessibility. The raw sequencing data and related metadata have been deposited in the DNA Data Bank of Japan database (BioProject PRJDB16690) [92]. Processed data and R code have been deposited in the Dryad Digital Repository at [93].

Electronic supplementary material is available online at [94].

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors' contributions. H.Y.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, visualization, writing—original draft. T.H.: conceptualization, investigation, methodology, project administration, supervision, writing—review and editing. D.M.K.: conceptualization, investigation, resources, supervision, writing—review and editing. K.Z.: investigation, resources, supervision, writing—review and editing. H.Q.: investigation, methodology, writing—review and editing. T.S.: investigation, methodology, writing—review and editing. K.S.: project administration, resources, supervision, writing—review and editing. K.K.: conceptualization, investigation, project administration, resources, supervision, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. This work was supported by Japan Society for the Promotion of Science (JSPS) Overseas Challenge Program for Young Researcher 202280252 to H.Y., JSPS KAKENHI Grant-in-Aid for Scientific Research 21J23216 to H.Y., 15K18471 to T.S., 20H03008 to K.K., JSPS Bilateral Research Program JPJSBP120209915 to K.K., and Leading Graduate Program in Primatology and Wildlife Science to H.Y.

Acknowledgements. We are deeply grateful to all staff in Snow Leopard Foundation in Kyrgyzstan and Snow Leopard Trust for their generous support. We are grateful to the Ministry of Natural Resources, Ecology and Technical Supervision of the Kyrgyz Republic and the management of Sarychat Ertash Reserve for their support

and necessary permissions for conducting this research. We are also grateful to Park rangers Emil Djaparov, Omurbek Kurmanaliev, Askat Abdykasiev, Urmat Solokov, Ulan Abilgaziev, and Joldosh Bektemirov for their help in collecting data from the field. We also thank all the people involved in the sample collection. We are deeply appreciative to Prof Miho Inoue-Murayama in Wildlife Research Center, Kyoto University for the perspective and input on the topic. We also acknowledge Mr Haruka Kitayama and Mr Ryo Akashi in the Faculty of Environmental Earth Science, Hokkaido University, and Ms Hiromi Kobayashi and Dr Annegret M. Naito-Liederbach in Wildlife Research Center, Kyoto University, and Dr Minori Arahori in Anicom Speciality Medical Institute Inc. for contributing the molecular work. We thank Prof Satoshi Hirata in Wildlife Research Center, Kyoto University for contributing the conceptualization and writing. Computational time was provided by the Supercomputer System, Human Genome Center, Institute of Medical Science, The University of Tokyo.

References

- Tomme P, Warren RAJ, Gilkes NR. 1995 Cellulose hydrolysis by bacteria and fungi. *Adv. Microb. Physiol.* **37**, 1–81. (doi:10.1016/s0065-2911(08)60143-5)
- Watanabe H, Tokuda G. 2001 Animal cellulases. *Cell. Mol. Life Sci.* **58**, 1167–1178. (doi:10.1007/PL00000931)
- Dearing MD, Foley WJ, McLean S. 2005 The influence of plant secondary metabolites on the nutritional ecology of herbivorous terrestrial vertebrates. *Annu. Rev. Ecol. Evol. Syst.* **36**, 169–189. (doi:10.1146/annurev.ecolsys.36.102003.152617)
- Hofmann RR. 1989 Evolutionary steps of ecophysiological adaptation and diversification of ruminants: a comparative view of their digestive system. *Oecologia* **78**, 443–457. (doi:10.1007/BF00378733)
- Vallentine JF. 2001 Grazing Herbivore nutrition. In *Grazing management*, pp. 29–66. Amsterdam: Elsevier. See <https://linkinghub.elsevier.com/retrieve/pii/B9780127100012502422>.
- Légrand-Defretin V. 1994 Differences between cats and dogs: a nutritional view. *Proc. Nutr. Soc.* **53**, 15–24. (doi:10.1079/pns19940004)
- Sunquist M, Sunquist F. 2002 *Wild cats of the world*. Chicago: University of Chicago Press.
- Van Valkenburgh B. 1999 Major patterns in the history of carnivorous mammals. *Annu. Rev. Earth Planet. Sci.* **27**, 463–493. (doi:10.1146/annurev.earth.27.1.463)
- Stevens CE, Hume DI. 2004 *Comparative physiology of the vertebrate digestive system*. Cambridge: Cambridge University Press (CUP).
- Hayami H. 1967 Nutritional differences between animal and plant proteins. *Nutr. Food* **20**, 7–14. (doi:10.4327/jsnfs1949.20.259)
- Bosch G, Hagen-Plantinga EA, Hendriks WH. 2015 Dietary nutrient profiles of wild wolves: insights for optimal dog nutrition? *Br. J. Nutr.* **113**, S40–S4. (doi:10.1017/S0007114514002311)
- Glendinning JI. 1994 Is the bitter rejection response always adaptive? *Physiol. Behav.* **56**, 1217–1227. (doi:10.1016/0031-9384(94)90369-7)
- Jiang P, Josue J, Li X, Glaser D, Li W, Brand JG, Margolskee RF, Reed DR, Beauchamp GK. 2012 Major taste loss in carnivorous mammals. *Proc. Natl. Acad. Sci. USA* **109**, 4956–4961. (doi:10.1073/pnas.1118360109)
- Kim S *et al.* 2016 Comparison of carnivore, omnivore, and herbivore mammalian genomes with a new leopard assembly. *Genome Biol.* **17**, 211. (doi:10.1186/s13059-016-1071-4)
- Li D, Zhang J. 2014 Diet shapes the evolution of the vertebrate bitter taste receptor gene repertoire. *Mol. Biol. Evol.* **31**, 303–309. (doi:10.1093/molbev/mst219)
- McGrane SJ, Gibbs M, Hernangomez de Alvaro C, Dunlop N, Winnig M, Klebansky B, Waller D. 2023 Umami taste perception and preferences of the domestic cat (*Felis catus*), an obligate carnivore. *Chem. Senses* **48**, 1–17. (doi:10.1093/chemse/bjad026)
- Montalvo V, Sáenz-Bolaños C, Cruz JC, Hagnauer I, Carrillo E. 2020 Consumption of wild rice (*Oryza latifolia*) by free-ranging jaguars, pumas, and ocelots (Carnivora-Felidae) in northwestern Costa Rica. *Food Webs* **22**, e00138. (doi:10.1016/j.fooweb.2019.e00138)
- Yoshimura H, Hirata S, Kinoshita K. 2021 Plant-eating carnivores: multispecies analysis on factors influencing the frequency of plant occurrence in obligate carnivores. *Ecol. Evol.* **11**, 10968–10983. (doi:10.1002/ece3.7885)
- Yoshimura H *et al.* 2020 The relationship between plant-eating and hair evacuation in snow leopards (*Panthera uncia*). *PLoS One* **15**, e0236635. (doi:10.1371/journal.pone.0236635)
- de Villa Meza A, Martínez Meyer E, López González CA. 2002 Ocelot (*Leopardus pardalis*) Food Habits in a Tropical Deciduous Forest of Jalisco, Mexico. *Am. Midl. Nat.* **148**, 146–154. (doi:10.1674/0003-0031(2002)148[0146:OLPFIH]2.0.CO;2)
- Nakanishi N, Izawa M. 2016 Importance of frogs in the diet of the Iriomote cat based on stomach content analysis. *Mamm. Res.* **61**, 35–44. (doi:10.1007/s13364-015-0246-9)
- McKinney T, Smith TW. 2007 Diets of sympatric bobcats and coyotes during years of varying rainfall in Central Arizona. *West. N. Am. Nat.* **67**, 8–15. (doi:10.3398/1527-0904(2007)67[8:DOSBAC]2.0.CO;2)
- Melville H, Bothma J du P, Mills MGL. 2004 Prey selection by caracal in the Kgalagadi Transfrontier Park. *South Afr. J. Wildl. Res.* **34**, 67–75.
- Witzuk J, Pagacz S, Gliwicz J, Mills LS. 2015 Niche overlap between sympatric coyotes and bobcats in highland zones of Olympic Mountains, Washington. *J. Zool* **297**, 176–183. (doi:10.1111/jzo.12270)
- Xiong M, Shao X, Long Y, Bu H, Zhang D, Wang D, Li S, Wang R, Yao M. 2016 Molecular analysis of vertebrates and plants in scats of leopard cats (*Prionailurus bengalensis*) in southwest China. *J. Mammal.* **97**, 1054–1064. (doi:10.1093/jmammal/gyw061)

26. Ramesh T Downs CT. Diet of serval (*Leptailurus serval*) on farmlands in the Drakensberg Midlands, South Africa. *Mammalia* **79**, 399–407. (doi:10.1515/mammalia-2014-0053)
27. Herbst M, Mills M. 2010 The feeding habits of the Southern African wildcat, a facultative trophic specialist, in the southern Kalahari (Kgalagadi Transfrontier Park, South Africa/Botswana). *J. Zool.* **280**, 403–413. (doi:10.1111/j.1469-7998.2009.00679.x)
28. Shultz D. 2019 Mystery solved? Why cats eat grass. *Science* (doi:10.1126/science.aaz0485)
29. Lovari S, Minder I, Ferretti F, Mucci N, Randi E, Pellizzi B. 2013 Common and snow leopards share prey, but not habitats: competition avoidance by large predators? *J. Zool.* **291**, 127–135. (doi:10.1111/jzo.12053)
30. Fox JL, Chundawat RS. 2016 What is a Snow Leopard? Behavior and ecology. In *Snow Leopards (biodiversity of the world conservation from genes to landscapes)* (eds T McCarthy, D Mallon), pp. 13–21, London: Academic Press.
31. Jumabay-Uulu K, Wegge P, Mishra C, Sharma K. 2014 Large carnivores and low diversity of optimal prey: a comparison of the diets of snow leopards *Panthera uncia* and wolves *Canis lupus* in Sarychat-Ertash Reserve in Kyrgyzstan. *Oryx* **48**, 529–535. (doi:10.1017/S0030605313000306)
32. Liu G, Zhang S, Zhao X, Li C, Gong M, Analysis AD. 2021 Advances and limitations of next generation sequencing in animal diet analysis. *Genes (Basel)* **12**, 1854. (doi:10.3390/genes12121854)
33. Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E. 2012 Towards next-generation biodiversity assessment using DNA metabarcoding. *Mol. Ecol.* **21**, 2045–2050. (doi:10.1111/j.1365-294X.2012.05470.x)
34. Pompanon F, Deagle BE, Symondson WOC, Brown DS, Jarman SN, Taberlet P. 2012 Who is eating what: diet assessment using next generation sequencing. *Mol. Ecol.* **21**, 1931–1950. (doi:10.1111/j.1365-294X.2011.05403.x)
35. Bernard KMT, Perry TW, Mgqatsa N. 2023 Puma (*Puma concolor*) sex influences diet in Southwest New Mexico. *West. N. Am. Nat.* **83**, 153–164. (doi:10.3398/064.083.0201)
36. Oftedal OT, Gittleman JL. 1989 Patterns of energy output during reproduction in Carnivores. In *Carnivore behavior, ecology, and evolution* pp. 355–378. Boston: Springer. (doi:10.1007/978-1-4757-4716-4)
37. SER. 2007 The Sarychat-Ertash state reserve management plan 2007–2015. Bishkek, Kyrgyzstan.
38. Longmire JL, Maltbie M, Baker RJ, Texas Tech University. 1997 Use of “lysis buffer” in DNA isolation and its implication for museum collections (*Occasional papers*). Lubbock, TX: Museum of Texas Tech University. (doi:10.5962/bhl.title.143318)
39. Hayakawa T et al. 2018 Improving the standards for gut microbiome analysis of fecal samples: insights from the field biology of Japanese macaques on Yakushima Island. *Primates* **59**, 423–436. (doi:10.1007/s10329-018-0671-x)
40. Sugimoto T, Nagata J, Aramilev VV, Belozor A, Higashi S, McCullough DR. 2006 Species and sex identification from faecal samples of sympatric carnivores, Amur leopard and Siberian tiger, in the Russian Far East. *Conserv. Genet.* **7**, 799–802. (doi:10.1007/s10592-005-9071-z)
41. Riaz T, Shehzad W, Viari A, Pompanon FF, Taberlet P, Coissac E. 2011 EcoPrimers: inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic Acids Res.* **39**, e145. (doi:10.1093/nar/gkr732)
42. Moorhouse-Gann RJ, Dunn JC, de Vere N, Goder M, Cole N, Hipperson H, Symondson WOC. 2018 New universal ITS2 primers for high-resolution herbivory analyses using DNA metabarcoding in both tropical and temperate zones. *Sci. Rep.* **8**, 8542. (doi:10.1038/s41598-018-26648-2)
43. Erickson DL, Reed E, Ramachandran P, Bourq NA, McShea WJ, Ottesen A. 2017 Reconstructing a herbivore's diet using a novel *rbcl* DNA mini-barcode for plants. *AoB Plants* **9**, plx015. (doi:10.1093/aobpla/plx015)
44. Kress WJ, Erickson DL. 2007 A two-locus global DNA barcode for land plants: the coding *rbcl* gene complements the non-coding *trnH-psbA* spacer region. *PLoS One* **2**, e508. (doi:10.1371/journal.pone.0000508)
45. Taberlet P et al. 2007 Power and limitations of the chloroplast *trnL* (UAA) intron for plant DNA barcoding. *Nucleic Acids Res.* **35**, e14. (doi:10.1093/nar/gkl938)
46. Ait Baamrane MA, Shehzad W, Ouhammou A, Abbad A, Naimi M, Coissac E, Taberlet P, Znari M. 2012 Assessment of the food habits of the Moroccan dorcas gazelle in M'Sabih Talaa, west central Morocco, using the *trnL* approach. *PLoS One* **7**, e35643. (doi:10.1371/journal.pone.0035643)
47. Tamura K, Stecher G, Kumar S. 2021 MEGA11: molecular evolutionary genetics analysis version 11. *Mol. Biol. Evol.* **38**, 3022–3027. (doi:10.1093/molbev/msab120)
48. Toju H, Tanabe AS, Ishii HS. 2016 Ericaceous plant-fungus network in a harsh alpine-subalpine environment. *Mol. Ecol.* **25**, 3242–3257. (doi:10.1111/mec.13680)
49. Jiang H, Lei R, Ding SW, Zhu S. 2014 Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. *BMC Bioinformatics* **15**, 182. (doi:10.1186/1471-2105-15-182)
50. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016 DADA2: high-resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583. (doi:10.1038/nmeth.3869)
51. R Foundation for Statistical Computing. 2019 R development core team 3.0.1. R: a language and environment for statistical computing.
52. Froslev TG, Kjølner R, Bruun HH, Ejrnæs R, Brunbjerg AK, Pietroni C, Hansen AJ. 2017 Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nat. Commun.* **8**, 1188. (doi:10.1038/s41467-017-01312-x)
53. Antich A, Palacin C, Wangenstein OS, Turon X. 2021 To denoise or to cluster, that is not the question: optimizing pipelines for COI metabarcoding and metaphylogeography. *BMC Bioinformatics* **22**, 177. (doi:10.1186/s12859-021-04115-6)
54. Lynggaard C, Bertelsen MF, Jensen CV, Johnson MS, Froslev TG, Olsen MT, Bohmann K. 2022 Airborne environmental DNA for terrestrial vertebrate community monitoring. *Curr. Biol.* **32**, 701–707.e5. (doi:10.1016/j.cub.2021.12.014)

55. Tanabe AS, Toju H. 2013 Two new computational methods for universal DNA barcoding: a benchmark using barcode sequences of bacteria, archaea, animals, fungi, and land plants. *PLoS One* **8**, e76910. (doi:10.1371/journal.pone.0076910)
56. Huson DH, Auch AF, Qi J, Schuster SC. 2007 MEGAN analysis of metagenomic data. *Genome Res.* **17**, 377–386. (doi:10.1101/gr.5969107)
57. Davletbakov AT, Milko DA, Ostashchenko AN (eds). 2015 In *Cadastre of the genetic fund of Kyrgyzstan* Bishkek, Kyrgyzstan.
58. MacConaill LE *et al.* 2018 Unique, dual-indexed sequencing adapters with UMIs effectively eliminate index cross-talk and significantly improve sensitivity of massively parallel sequencing. *BMC Genomics* **19**, 30. (doi:10.1186/s12864-017-4428-5)
59. Tsukamoto Y, Yonezawa S, Katayama N, Isagi Y. 2021 Detection of endangered aquatic plants in rapid streams using environmental DNA. *Front. Ecol. Evol.* **8**, 530. (doi:10.3389/fevo.2020.622291)
60. Drake LE, Cuff JP, Young RE, Marchbank A, Chadwick EA, Symondson WOC. 2022 An assessment of minimum sequence copy thresholds for identifying and reducing the prevalence of artefacts in dietary metabarcoding data. *Methods Ecol. Evol.* **13**, 694–710. (doi:10.1111/2041-210X.13780)
61. Ando H, Fujii C, Kawanabe M, Ao Y, Inoue T, Takenaka A. 2018 Evaluation of plant contamination in metabarcoding diet analysis of a herbivore. *Sci. Rep.* **8**, 15563. (doi:10.1038/s41598-018-32845-w)
62. da Silva LP, Mata VA, Lopes PB, Pereira P, Jarman SN, Lopes RJ, Beja P. 2019 Advancing the integration of multi-marker metabarcoding data in dietary analysis of trophic generalists. *Mol. Ecol. Resour.* **19**, 1420–1432. (doi:10.1111/1755-0998.13060)
63. de Barba M, Miquel C, Boyer F, Mercier C, Rioux D, Coissac E, Taberlet P. 2014 DNA metabarcoding multiplexing and validation of data accuracy for diet assessment: application to omnivorous diet. *Mol. Ecol. Resour.* **14**, 306–323. (doi:10.1111/1755-0998.12188)
64. Xiong M, Wang D, Bu H, Shao X, Zhang D, Li S, Wang R, Yao M. 2017 Molecular dietary analysis of two sympatric felids in the mountains of Southwest China biodiversity hotspot and conservation implications. *Sci. Rep.* **7**, 41909. (doi:10.1038/srep41909)
65. Shao X, Lu Q, Xiong M, Bu H, Shi X, Wang D, Zhao J, Li S, Yao M. 2021 Prey partitioning and livestock consumption in the world's richest large carnivore assemblage. *Curr. Biol.* **31**, 4887–4897. (doi:10.1016/j.cub.2021.08.067)
66. Chao A, Jost L. 2012 Coverage-based rarefaction and extrapolation: standardizing samples by completeness rather than size. *Ecology* **93**, 2533–2547. (doi:10.1890/11-1952.1)
67. Oksanen J *et al.* 2022 Vegan: community ecology package version 2.6-4
68. McMurdie PJ, Holmes S. 2013 phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* (ed. M Watson), **8**, e61217. (doi:10.1371/journal.pone.0061217)
69. Deagle BE, Thomas AC, McInnes JC, Clarke LJ, Vesterinen EJ, Clare EL, Kartzinel TR, Eveson JP. 2019 Counting with DNA in metabarcoding studies: how should we convert sequence reads to dietary data? *Mol. Ecol.* **28**, 391–406. (doi:10.1111/mec.14734)
70. Liaw A, Wiener M. 2002 Classification and regression by randomForest. *R News* **2**, 18–22.
71. Goldberg AR, Conway CJ, Tank DC, Andrews KR, Gour DS, Waits LP. 2020 Diet of a rare herbivore based on DNA metabarcoding of feces: selection, seasonality, and survival. *Ecol. Evol.* **10**, 7627–7643. (doi:10.1002/ece3.6488)
72. Urban P, Praebel K, Bhat S, Dierking J, Wangenstein OS. 2022 DNA metabarcoding reveals the importance of gelatinous zooplankton in the diet of *Pandalus borealis*, a keystone species in the Arctic. *Mol. Ecol.* **31**, 1562–1576. (doi:10.1111/mec.16332)
73. Breiman L. 2001 Random forests. *Mach. Learn.* **45**, 5–32. (doi:10.1023/A:1010933404324)
74. Calle ML, Urrea V. 2011 Letter to the editor: stability of random forest importance measures. *Brief. Bioinform.* **12**, 86–89. (doi:10.1093/bib/bbq011)
75. Griffith DM, Veech JA, Marsh CJ. 2016 Cooccur: probabilistic species co-occurrence analysis in R. *J. Stat. Softw.* **69**, 1–17. (doi:10.18637/jss.v069.c02)
76. Terce M, Symondson WOC, Cuff JP. 2021 The problem of omnivory: a synthesis on omnivory and DNA metabarcoding. *Mol. Ecol.* **30**, 2199–2206. (doi:10.1111/mec.15903)
77. Anderson MJ, Willis TJ. 2003 Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. *Ecology* **84**, 511–525. (doi:10.1890/0012-9658(2003)084[0511:CAOPCA]2.0.CO;2)
78. Borcard D, Legendre P. 2012 Is the mantel correlogram powerful enough to be useful in ecological analysis? A simulation study. *Ecology* **93**, 1473–1481. (doi:10.1890/11-1737.1)
79. Dray S, Bauman D, Blanchet G, Borcard D, Clappe S, Guenard G, *et al.* 2021 Package “adespatial”. See <https://github.com/sdray/adespatial/issues>.
80. Borcard D, Gillet F, Legendre P. 2011 *Numerical Ecology with R*. (eds R Gentleman, GG Parmigiani, K Hornik). New York, NY: Springer.
81. Janečka JE, Munkhtsog B, Jackson RM, Naranbaatar G, Mallon DP, Murphy WJ. 2011 Comparison of noninvasive genetic and camera-trapping techniques for surveying snow leopards. *J. Mammal.* **92**, 771–783. (doi:10.1644/10-MAMM-A-036.1)
82. Samelius G, Xiao L, Lkhagvajav P, Johansson Ö. Risky business: red foxes killed when scavenging from snow leopard kills. *SLR* **1**, 7–10. (doi:10.56510/slr.v1.8092)
83. University of Michigan Museum of Zoology. 2020 *Animal diversity Web*. See <https://animaldiversity.org/>
84. Johansson Ö, Koehler G, Rauset GR, Samelius G, Andrén H, Mishra C, Lhagvasuren P, McCarthy T, Low M. 2018 Sex-specific seasonal variation in puma and snow leopard home range utilization. *Ecosphere* **9**, e02371. (doi:10.1002/ecs2.2371)
85. Mallon D, Harris RB, Wegge P. 2016 Snow leopard prey and diet. In *Snow leopards (Biodiversity of the world conservation from genes to landscapes)* (eds T McCarthy, D Mallon), pp. 43–55. London: Academic Press. (doi:10.11339/JTM.25.108)
86. Mikage M, Hong H, Cai X. 2008 Studies of ephedra plants in Asia. Part 5. The herbivory damage to ephedra plants by livestock. *J. Tradit. Med.* **25**, 108–111.

87. Han L, Blank D, Wang M, da Silva AA, Yang W, Ruckstuhl K, Alves J. 2020 Diet differences between males and females in sexually dimorphic ungulates: a case study on Siberian ibex. *Eur. J. Wildl. Res.* **66**, 55. (doi:10.1007/s10344-020-01387-w)
88. Demontigny-Bédard I, Beauchamp G, Bélanger MC, Frank D. 2016 Characterization of pica and chewing behaviors in privately owned cats: a case-control study. *J. Feline Med. Surg.* **18**, 652–657. (doi:10.1177/1098612X15591589)
89. Liu JB *et al.* 2015 Anti-inflammatory hydrolyzable tannins from *Myricaria bracteata*. *J. Nat. Prod.* **78**, 1015–1025. (doi:10.1021/np500953e)
90. Johansson Ö, Ausilio G, Low M, Lkhagvajav P, Weckworth B, Sharma K. 2021 The timing of breeding and independence for snow leopard females and their cubs. *Mamm. Biol.* **101**, 173–180. (doi:10.1007/s42991-020-00073-3)
91. Tylianakis JM, Laliberté E, Nielsen A, Bascompte J. 2010 Conservation of species interaction networks. *Biol. Conserv.* **143**, 2270–2279. (doi:10.1016/j.biocon.2009.12.004)
92. Yoshimura H, Hayakawa T, Kikuchi DM, Zhumabai-uulu K, Qi H, Sugimoto T, Kinoshita K. 2023 Metabarcoding of vertebrate and plant in scat of alpine mammals in Kyrgyzstan. DNA Data Bank of Japan, BioProject, PRJDB16690.
93. Yoshimura H, Hayakawa T, Kikuchi DM, Zhumabai Uulu K, Qi H, Sugimoto T, Kinoshita K. 2023 Data from: Metabarcoding analysis provides insight into the link between prey and plant intake in a large alpine cat carnivore, the snow leopard. Dryad Digital Repository (doi:10.5061/dryad.j9kd51cjp)
94. Yoshimura H, Hayakawa T, Kikuchi DM, Zhumabai Uulu K, Qi H, Sugimoto T. 2024 Supplementary material from: Metabarcoding analysis provides insight into the link between prey and plant intake in a large alpine cat carnivore, the snow leopard. Figshare (doi:10.6084/m9.figshare.c.7162451)