

The knot quandles of oriented 2-knots

Yuta Taniguchi

Department of mathematics, Graduate school of science, Osaka University

1 Introduction

The purpose of this note is to give a summery of the results given in the papers [18] and [20]. In this note, an *oriented 1-knot* is an oriented circle smoothly embedded in the 3-sphere S^3 , an *oriented 2-knot* is an oriented 2-sphere smoothly embedded in the 4-sphere S^4 and they are correctively called *oriented knots*.

A quandle [9, 11] is an algebra whose axioms correspond to the Reidemeister moves. It is a useful tool to define the oriented knot invariants. A typical example of knot invariants using quandles is the *knot quandle*. The knot quandle of an oriented knot \mathcal{K} is defined as the homotopy classes of pathes in the knot exterior with a certain operation. Since then, some properties of the knot quandle of an oriented 1-knot have been studied.

In this note, we focus on the knot quandles of oriented 2-knots. First, we discussed the differences between knot quandles and knot groups as invariants for oriented 2-knots. By [9, 11], if two oriented knots have the same knot quandle, then their knot groups are isomorphic as groups. For oriented 1-knots, it is known that there is a pair of oriented 1-knots with the same knot group but different knot quandles. In Section 3, we showed that there are infinitely many triples of oriented 2-knots with the same knot group but different knot quandles. This result implies that the knot quandle is really stronger invariant than the knot group for oriented 2-knots. Second, we study the *quandle homology groups* of knot quandles of oriented 2-knots. In [1], Carter, Jelsovsky, Kamada, Langford and Saito introduced the quandle homology group and defined invariants of oriented knots using them, which are called the *quandle cocycle invariants*. In general, it is difficult to determine the quandle homology group of a quandle. We show that the second quandle homology group of the knot quandle of an oriented 2-knot is trivial in Section 4. As a consequence of this result, we see that the knot quandle of a non-trivial oriented 1-knot can not be realized by the knot quandle of 2-knots.

2 Definitions

A *quandle* X [9, 11] is a non-empty set equipped with a binary operation $*$ satisfying the following conditions:

- For any $x \in X$, we have $x * x = x$.
- For any $y \in X$, the map $S_y : X \rightarrow X; x \mapsto x * y$ is a bijection.

- For any $x, y, z \in X$, we have $(x * y) * z = (x * z) * (y * z)$.

Here are examples of quandles:

Example 2.1. Let G be a group and $f : G \rightarrow G$ a group automorphism. We define the operation $*$ on G by $x * y := f(xy^{-1})y$. Then, $\text{GAlex}(G, f) = (G, *)$ is a quandle, which is called the *generalized Alexander quandle*.

Example 2.2. Let k be an oriented 1-knot. Let $N(k)$ be a tubular neighborhood of k and $E(k) = S^3 \setminus \text{int}N(k)$ an exterior of k . We fix a point $p \in E(k)$. Let $Q(k, p)$ be the set of homotopy classes of all paths in $E(k)$ from a point in $\partial E(k)$ to p . The set $Q(k, p)$ is a quandle with an operation defined by $[\alpha] * [\beta] := [\alpha \cdot \beta^{-1} \cdot m_{\beta(0)} \cdot \beta]$, where $m_{\beta(0)}$ is a meridian loop starting from $\beta(0)$ and going along in the positive direction. We call $Q(k, p)$ the *knot quandle* of k . The isomorphism class of the knot quandle does not depend on the base point p . Thus, we denote the knot quandle simply by $Q(k)$. For an oriented 2-knot F , the *knot quandle* $Q(F)$ of F is defined in the same way as for oriented 1-knots.

The *associated group* of X , denoted by $\text{As}(X)$, is the group defined as

$$\langle x \ (x \in X) \mid x * y = y^{-1}xy \ (x, y \in X) \rangle.$$

The associated group $\text{As}(X)$ acts on X from the right by $x \cdot y := x * y$ for any $x, y \in X$. A quandle X is *connected* if the action of $\text{As}(X)$ on X is transitive.

A map $f : X \rightarrow Y$ between quandles is a *quandle isomorphism* if $f(x * y) = f(x) * f(y)$ for any $x, y \in X$ and f is a bijection. When there is a quandle isomorphism $f : X \rightarrow Y$, we say that X and Y are *quandle isomorphic*.

3 Knot quandles vs Knot groups

In this section, we compare the knot group and the knot quandle. In Subsection 3.1, we review a relation between the knot group and the knot quandle, and introduce our result. In Subsection 3.2, we explain the outline of the proof. This section is a joint work with Kokoro Tanaka.

3.1 Back ground and Main result

Let \mathcal{K} and \mathcal{K}' be oriented knots. We consider the following conditions:

- (i) The knot groups $G(\mathcal{K}')$ and $G(\mathcal{K})$ are group isomorphic.
- (ii) The knot quandles $Q(\mathcal{K}')$ and $Q(\mathcal{K})$ are quandle isomorphic.

Since the associated group $\text{As}(Q(\mathcal{K}))$ is group isomorphic to the knot group $G(\mathcal{K})$ for an oriented knot \mathcal{K} [9, 11], we have (ii) \Rightarrow (i). For oriented 1-knots, the converse does not hold, that is, there are oriented 1-knots with the same knot group but different knot quandles. For example, the square knot $3_1 \# 3_1^*$ and the granny knot $3_1 \# 3_1$ satisfy the condition (i) but does not satisfy the condition (ii). In this section, we consider the case of oriented 2-knots. More precisely, we consider the following question:

Question 3.1. *Are there oriented 2-knots such that these oriented 2-knots satisfy the condition (i) but do not satisfy the condition (ii)?*

In this note, we give an affirmative answer to this question.

Theorem 3.2. *There exist infinitely many triples $\{F_1, F_2, F_3\}$ of oriented 2-knots such that*

- (1) *the knot groups $G(F_1), G(F_2)$ and $G(F_3)$ are mutually group isomorphic, and*
- (2) *no two of knot quandles $Q(F_1), Q(F_2)$ and $Q(F_3)$ are quandle isomorphic.*

The 2-knots F_1, F_2 and F_3 are obtained from the *twist spinning construction*, which is introduced by Zeeman [21]. We review the some properties of a twist spun knot. Let k be an oriented 1-knot in S^3 and n a non-negative integer. We denote by $\tau^n(k)$ the n -twist spun knot of k . Zeeman introduced the n -twist spun knot and showed that if $n \neq 0$, the n -twist spun knot $\tau^n(k)$ is a fibered 2-knot whose fiber is the once punctured M_k^n , where M_k^n is the n -fold cyclic branched covering space of S^3 branched along k . In particular, the 1-twist spun knot $\tau^1(k)$ is a trivial 2-knot for any oriented 1-knot k .

Remark 3.3. (1) It is known that for any oriented 1-knot k , the knot group of $\tau^n(k)$ is a quotient group of the knot group of k . Thus, the knot groups of $\tau^n(3_1 \# 3_1)$ and $\tau^n(3_1 \# 3_1^*)$ are group isomorphic for any non-negative integer n . However, since $\tau^0(3_1 \# 3_1)$ and $\tau^0(3_1 \# 3_1^*)$ are equivalent [6, 16], we see that $\tau^n(3_1 \# 3_1)$ and $\tau^n(3_1 \# 3_1^*)$ satisfy the condition (ii) for any non-negative integer n (cf. [17]).

(2) Let $S(p, q)$ be the 2-bridge knot of type (p, q) . It is known that $\tau^2(S(p, q))$ and $\tau^2(S(p, q'))$ satisfy the condition (i) for any q, q' (see [15]). In [7], Inoue showed that for any q , the knot quandle $Q(\tau^2(S(p, q)))$ is quandle isomorphic to the quandle $\text{GAlex}(\mathbb{Z}/p\mathbb{Z}, \text{Inv})$, where $\text{Inv} : \mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$ is the group automorphism defined by $\text{Inv}(x) = -x$. This implies that $\tau^2(S(p, q))$ and $\tau^2(S(p, q'))$ satisfy the condition (ii) for any q, q' . Hence, 2-twist spun 2-bridge knots do not give an example of such 2-knots.

3.2 Outline of proof of Theorem

Let p, q, r be coprime positive integers. We denote the torus knot of type (m, n) by $t_{m,n}$.

Theorem 3.4. [5] *The knot group $G(\tau^p(t_{q,r}))$ is group isomorphic to $\pi_1(M_{t_{q,r}}^p) \times \mathbb{Z}$.*

Since $M_{t_{q,r}}^p, M_{t_{r,p}}^q$ and $M_{t_{p,q}}^r$ are homeomorphic ([12]), the oriented 2-knots $F_1 := \tau^p(t_{q,r}), F_2 := \tau^q(t_{r,p})$ and $F_3 := \tau^r(t_{p,q})$ satisfy the condition (1).

To show the oriented 2-knots F_1, F_2 and F_3 satisfy the condition (2), we focus on the notion of *type* of a quandle. The *type* of a quandle X , denoted by $\text{type}(X)$, is a minimum positive integer n such that S_x^n is the identity map id_X for any $x \in X$. If there is no such n , we define $\text{type}(X) = \infty$. In general, it is difficult to determine $\text{type}(X)$ for a given quandle X . However, it is easy to compute the type of a generalized Alexander quandle.

Proposition 3.5. *Let G be a group and φ a group automorphism of G . The type of $\text{GAlex}(G, \varphi)$ is equal to the order of φ .*

As we mentioned above, the n -twist spun knot $\tau^n(k)$ is a fibered 2-knot for any positive integer n . Inoue studied the structure of the knot quandle of a fibered 2-knot and showed the following theorem.

Theorem 3.6. [7] *Let F be an oriented fibered 2-knot, M the fiber of $S^4 \setminus F$ and φ the monodromy of $S^4 \setminus F$. Then, the knot quandle $Q(F)$ is quandle isomorphic to $\text{GAlex}(\pi_1(M), \varphi_*)$, where $\varphi_* : \pi_1(M) \rightarrow \pi_1(M)$ is the group automorphism induced by φ .*

By [21], the monodromy φ of $S^4 \setminus \tau^n(k)$ is the canonical covering homeomorphism of M_k^n . In particular, it holds that the order of the induced group automorphism $\varphi_* : \pi_1(M_k^n) \rightarrow \pi_1(M_k^n)$ is n . Thus, we obtain the following theorem.

Theorem 3.7. *Let k be an oriented 1-knot and n a positive integer. Then, we have $\text{type}(Q(\tau^n(k))) = n$.*

Since $F_1 = \tau^p(t_{q,r})$, $F_2 = \tau^q(t_{r,p})$ and $F_3 = \tau^r(t_{p,q})$, we have $\text{type}(Q(F_1)) = p$, $\text{type}(Q(F_2)) = q$ and $\text{type}(Q(F_3)) = r$. This implies that F_1, F_2 and F_3 satisfy the condition (2).

4 Quandle homology groups of knot quandles

In this section, we discuss the quandle homology group of the knot quandle $Q(\mathcal{K})$. We review the quandle homology group [1] in Subsection 4.1 and the f -twisted Alexander matrix [8] in Subsection 4.2. We give the outline of the proof in Subsection 4.3.

4.1 Definition and Main result.

Let X be a quandle. For each positive integer n , we denote by $C_n^R(X)$ the free abelian group whose basis is X^n . We set $C_0^R(X) = 0$. For each $(x_1, \dots, x_n) \in X^n$, let us define an element $\partial(x_1, \dots, x_n) \in C_{n-1}^R(X)$ by

$$\begin{aligned} \partial(x_1, \dots, x_n) &:= \sum_{i=2}^n (-1)^i (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \\ &\quad - \sum_{i=2}^n (-1)^i (x_1 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n). \end{aligned}$$

Using this, we have a group homomorphism $\partial_n : C_n^R(X) \rightarrow C_{n-1}^R(X)$ for $n \geq 2$. We define $\partial_1 : C_1^R(X) \rightarrow C_0^R(X)$ by the zero map. Then, we see that $\partial_{n-1} \circ \partial_n$ is the zero map. Hence, $(C_n^R(X), \partial_n)$ is a chain complex.

Let $C_n^D(X)$ be the subgroup of $C_n^R(X)$ generated by n -tuples (x_1, \dots, x_n) with $x_i = x_{i+1}$ for some i . We can see that $\partial_n(C_n^D(X)) \subset C_{n-1}^D(X)$ for any n . Thus, setting $C_n^Q(X) = C_n^R(X)/C_n^D(X)$, we have a chain complex $(C_n^Q(X), \partial_n)$. The n -th quandle homology group $H_n^Q(X)$ [1] is the n -th homology group of the chain complex $(C_n^Q(X), \partial_n)$.

Problem 4.1. *Determine the quandle homology group $H_n^Q(X)$ for a quandle X .*

We consider this problem for knot quandles. Since a knot quandle is connected, we see that $H_1^Q(Q(\mathcal{K})) = \mathbb{Z}$ for an oriented knot \mathcal{K} (cf. [2]). For a 1-knot k , the second and third quandle homology groups of the knot quandle $Q(k)$ have been calculated.

Theorem 4.2. [3] *Let k be a non-trivial oriented 1-knot. Then we have $H_2^Q(Q(k)) = \mathbb{Z}$.*

Theorem 4.3. [14] *Let k be a non-trivial oriented 1-knot. Then we have $H_3^Q(Q(k)) = \mathbb{Z}$.*

In contrast to oriented 1-knots, there are few knot quandles of oriented 2-knots whose second quandle homology groups are computed.

- It is known that for any oriented 1-knot k , the knot quandle of $\tau^0(k)$ is quandle isomorphic to the knot quandle of the *long knot* corresponding to k . Eisermann showed that $H_2^Q(Q(\hat{k}))$ is trivial for any long knot \hat{k} . Thus, we have $H_2^Q(Q(\tau^0(k))) = 0$.
- Let $S(p, q)$ be the 2-bridge knot of type (p, q) . As we mentioned above, $Q(\tau^2(S(p, q)))$ is quandle isomorphic to $\text{GAlex}(\mathbb{Z}/p\mathbb{Z}, \text{Inv})$. In [13], Mochizuki showed that the second quandle homology group $H_2^Q(\text{GAlex}(\mathbb{Z}/p\mathbb{Z}, \text{Inv}))$ is trivial for any odd prime integer p . Thus, we have $H_2^Q(Q(\tau^2(S(p, q)))) = 0$.

Main result of this section is the following theorem.

Theorem 4.4. *Let F be an oriented 2-knot. Then, we have $H_2^Q(Q(F)) = 0$.*

The proof of Theorem 4.4 is based on the *f-twisted Alexander matrix* introduced by Ishii and Oshiro. The outline of the proof is explained later.

At the end of this section, we discuss a difference between knot quandles of oriented 1-knots and knot quandles of oriented 2-knots. Since the knot group of an oriented 1-knot k is group isomorphic to the knot group of $\tau^0(k)$, we see that

$$\{G(k) \mid k : \text{a 1-knot}\} \subset \{G(F) \mid F : \text{a 2-knot}\}.$$

On the other hand, by Theorem 4.2 and Theorem 4.4, the knot quandle of a non-trivial oriented 1-knot can not be realized by the knot quandle of oriented 2-knots, that is, it holds that

$$\{Q(k) \mid k : \text{a non-trivial 1-knot}\} \cap \{Q(F) \mid F : \text{a 2-knot}\} = \emptyset.$$

4.2 *f*-twisted Alexander matrix

Let X be a quandle, and R a ring with the unity 1. A pair (f_1, f_2) of maps $f_1, f_2 : X \times X \rightarrow R$ is called an *Alexander pair* if f_1 and f_2 satisfy the following conditions:

- For any $x \in X$, we have $f_1(x, x) + f_2(x, x) = 1$.
- For any $x, y \in X$, $f_1(x, y)$ is invertible.
- For any $x, y, z \in X$, we have

$$\begin{aligned} f_1(x * y, z) f_1(x, y) &= f_1(x * z, y * z) f_1(x, z), \\ f_1(x * y, z) f_2(x, y) &= f_2(x * z, y * z) f_1(y, z), \text{ and} \\ f_2(x * y, z) &= f_1(x * z, y * z) f_2(x, z) + f_2(x * z, y * z) f_2(y, z). \end{aligned}$$

Example 4.5. Let X be a quandle and $\mathbb{Z}[t^{\pm 1}]$ the ring of Laurent polynomials with integer coefficients. The maps $f_1, f_2 : X \times X \rightarrow \mathbb{Z}[t^{\pm 1}]$ defined by $f_1(x, y) := t$ and $f_2(x, y) := 1 - t$ give an Alexander pair.

Example 4.6. Let X be a quandle and A an abelian group. A map $\theta : X \times X \rightarrow A$ is a *quandle 2-cocycle* [1] if it satisfies the following conditions:

- For any $x \in X$, we have $\theta(x, x) = 0_A$, where 0_A is the identity element.
- For any $x, y, z \in X$, we have $\theta(x * y, z) + \theta(x, y) = \theta(x * z, y * z) + \theta(x, z)$.

Let $\theta : X \times X \rightarrow A$ be a quandle 2-cocycle and $\mathbb{Z}[A]$ the group ring. We set maps $f_\theta, 0 : X \times X \rightarrow \mathbb{Z}[A]$ by $f_\theta(x, y) := 1 \cdot \theta(x, y)$ and $0(x, y) = 0$. Then, the pair $(f_\theta, 0)$ is an Alexander pair. We call this Alexander pair $(f_\theta, 0)$ the *Alexander pair associated with a quandle 2-cocycle* θ [19].

Next, we review the definition of the f -twisted Alexander matrix. Refer to [8] for more details. Let $FQ(S)$ the free quandle on a finite set $S = \{x_1, \dots, x_n\}$, Q a quandle with a finite presentation of $\langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$ and $\text{pr} : FQ(S) \rightarrow Q$ the canonical projection. In this note, we omit pr to present $\text{pr}(a)$ as a . Let R be a ring with the unity 1 and $f = (f_1, f_2)$ an Alexander pair of maps $f_1, f_2 : Q \times Q \rightarrow R$. For $x_j \in S$, the f -derivative with respect to x_j [8] is a map $\frac{\partial f}{\partial x_j} : FQ(S) \rightarrow R$ satisfies the following conditions:

- For any $x_i \in S$, we have $\frac{\partial f}{\partial x_j}(x_i) = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$
- For any $x, y \in FQ(S)$, we have

$$\frac{\partial f}{\partial x_j}(x * y) = f_1(x, y) \frac{\partial f}{\partial x_j}(x) + f_2(x, y) \frac{\partial f}{\partial x_j}(y).$$

For a relator $r = (r_1, r_2)$, we set $\frac{\partial f}{\partial x_j}(r) := \frac{\partial f}{\partial x_j}(r_1) - \frac{\partial f}{\partial x_j}(r_2)$.

Let A be an $m \times n$ matrix over a commutative ring R . The d -th elementary ideal of A , denoted by $E_d(A)$, is the ideal generated by all $(n - d)$ -minors of A if $n - m \leq d < n$, and $E_d(A) = \begin{cases} 0 & \text{if } d < n - m, \\ R & \text{if } n \leq d. \end{cases}$

Let Q be a quandle with a finite presentation of $\langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$, X a quandle, $\rho : Q \rightarrow X$ a quandle homomorphism, R a ring with the unity 1 and $f = (f_1, f_2)$ an Alexander pair of maps $f_1, f_2 : X \times X \rightarrow R$. We set $f \circ (\rho \times \rho) = (f_1 \circ (\rho \times \rho), f_2 \circ (\rho \times \rho))$. Then the pair $f \circ (\rho \times \rho)$ is also an Alexander pair. The f -twisted Alexander matrix of (Q, ρ) [8], which is denoted by $A(Q, \rho; f_1, f_2)$, is the $m \times n$ matrix defined by

$$A(Q, \rho; f_1, f_2) := \begin{pmatrix} \frac{\partial f \circ (\rho \times \rho)}{\partial x_1}(r_1) & \cdots & \frac{\partial f \circ (\rho \times \rho)}{\partial x_n}(r_1) \\ \vdots & \ddots & \vdots \\ \frac{\partial f \circ (\rho \times \rho)}{\partial x_1}(r_m) & \cdots & \frac{\partial f \circ (\rho \times \rho)}{\partial x_n}(r_m) \end{pmatrix}.$$

Suppose that R is a commutative ring. Let Q' be a finitely presented quandle and $\rho' : Q' \rightarrow X$ a quandle homomorphism. Ishii and Oshiro [8] showed that if there is a quandle isomorphism $\varphi : Q \rightarrow Q'$ such that $\rho = \rho' \circ \varphi$, then we have $E_d(A(Q, \rho; f_1, f_2)) = E_d(A(Q', \rho'; f_1, f_2))$ for all d .

4.3 Outline of proof of Theorem

Let $\theta : X \times X \rightarrow A$ be a quandle 2-cocycle. By the definition, the linear extension $\theta : \mathbb{Z}[X \times X] \rightarrow A$ is a 2-cocycle of $C_2^2(X; A)$. Thus, we can regard θ as a group homomorphism from $H_2^Q(X)$ to A . In [20], we proved the following theorems:

Theorem 4.7. *Let Q be a connected quandle with a finite presentation, X a quandle, A an abelian group and $\theta : X \times X \rightarrow A$ a quandle 2-cocycle. For any quandle homomorphism $\rho : Q \rightarrow X$, we have*

$$E_0(A(Q, \rho; f_\theta, 0)) = (\{1 \cdot a - 1 \cdot 0_A \mid a \in \text{Im}(\theta \circ \rho_*)\}) \subset \mathbb{Z}[A],$$

where $\rho_* : H_2^Q(Q) \rightarrow H_2^Q(X)$ is the group homomorphism induced by ρ .

Theorem 4.8. *Let F be an oriented 2-knot, X a quandle, A an abelian group and $\theta : X \times X \rightarrow A$ a quandle 2-cocycle. For any quandle homomorphism $\rho : Q \rightarrow X$, we have*

$$E_0(A(Q(F), \rho; f_\theta, 0)) = (0).$$

Let F be an oriented 2-knot. Suppose that $H_2^Q(Q(F))$ is non-trivial. We set $A := H_2^Q(Q(F))$, $X := Q(F)$ and $\rho := \text{id} : Q(F) \rightarrow Q(F) = X$. By the universal coefficient theorem, there is a quandle 2-cocycle $\theta : X \times X \rightarrow A$ such that the group homomorphism $\theta \circ \rho_* : H_2^Q(Q(F)) \rightarrow A = H_2^Q(Q(F))$ coincides with the identity map on $H_2^Q(Q(F))$. By the assumption and Theorem 4.7, we have $E_0(A(Q, \rho; f_\theta, 0)) \neq (0)$. This contradicts to Theorem 4.8.

Acknowledgement

The author would like to express his gratitude to Kokoro Tanaka for the collaboration [18] and many discussions. The author was supported by JSPS KAKENHI Grant Number 21J21482.

References

- [1] J S. Carter, D. Jelsovsky, S. Kamada, L. Langford, and M. Saito, *Quandle cohomology and state-sum invariants of knotted curves and surfaces*, Trans. Amer. Math. Soc. 355 (2003), no. 10, 3947–3989.
- [2] J S. Carter, D. Jelsovsky, S. Kamada, and M. Saito, *Quandle homology groups, their Betti numbers, and virtual knots*, J. Pure Appl. Algebra 157 (2001), 135–155.
- [3] M. Eisermann, *Homological characterization of the unknot*, J. Pure Appl. Algebra 177 (2003), no. 2, 131–157.

- [4] R. Fenn and C. Rourke, *Racks and links in codimension two*, J. Knot Theory Ramifications 1 (1992), no. 04, 343–406
- [5] C. McA. Gordon, *Twist-spun torus knots*, Proc. Amer. Math. Soc. 32 (1972), 319–322.
- [6] C. McA. Gordon, *A note on spun knots*, Proc. Amer. Math. Soc. 58 (1976), 361–362.
- [7] A. Inoue, *On the knot quandle of a fibered knot, finiteness and equivalence of knot quandles*, Topology Appl. 265 (2019), 106811, 8.
- [8] A. Ishii and K. Oshiro, *Derivatives with Alexander pairs for quandles*, Fund. Math. 259 (2022), no. 1, 1–31.
- [9] D. Joyce, *A classifying invariant of knots, the knot quandle*, J. Pure Appl. Algebra 23 (1982), no. 1, 37–65.
- [10] S. Kamada, *Surface-knots in 4-space. An introduction*, Springer Monographs in Mathematics, Springer, Singapore, 2017
- [11] S. V. Matveev, *Distributive groupoids in knot theory*, Mat. Sb. 161 (1982), no. 1, 78–88.
- [12] J. Milnor, *On the 3-dimensional Brieskorn manifolds $M(p, q, r)$* , Knots, groups, and 3-manifolds (Papers dedicated to the memory of R. H. Fox), 1975, pp. 175–225.
- [13] T. Mochizuki, *Some calculations of cohomology groups of finite Alexander quandles*, J. Pure Appl. Algebra 179, 287–330 (2003)
- [14] T. Nosaka, *On the fundamental 3-classes of knot group representations*, Geom. Dedicata, 204, 1–24.
- [15] S. P. Plotnick, *The homotopy type of four-dimensional knot complements*, Math. Z. 183 (1983), no. 4, 447–471.
- [16] D. Roseman, *The spun square knot is the spun granny knot*, Bol. Soc. Mat. Mexicana (2) 20 (1975), no. 2, 49–55.
- [17] K. Tanaka, *Inequivalent surface-knots with the same knot quandle*, Topol. Appl. 154 (15) (2007) 2757–2763.
- [18] K. Tanaka and Y. Taniguchi, *2-knots with the same knot group but different knot quandles*, in preparation.
- [19] Y. Taniguchi, *Alexander matrices of link quandles associated to quandle homomorphisms and quandle cocycle invariants*, to appear in Tokyo J. Math..
- [20] Y. Taniguchi, *Twisted Alexander matrices of quandles associated with a certain Alexander pair*, preprint, available at arXiv:2302.11175.
- [21] E. C. Zeeman, *Twisting spun knots*, Trans. Amer. Math. Soc. 115 (1965), 471–495.

Department of mathematics, Graduate school of science,
Osaka University
Osaka 560-0043
JAPAN
E-mail address: u660451k@ecs.osaka-u.ac.jp

大阪大学大学院理学研究科数学専攻 谷口雄大