



# *Geometry & Topology*

Volume 29 (2025)

**Virtual classes via vanishing cycles**

TASUKI KINJO



# Virtual classes via vanishing cycles

TASUKI KINJO

We develop a new method to construct the virtual fundamental classes for quasismooth derived schemes (and more generally, derived 1-Artin stacks) using the perverse sheaves of vanishing cycles on their  $-1$ -shifted cotangent spaces. It is based on the author's previous work that can be regarded as a version of the Thom isomorphism for  $-1$ -shifted cotangent spaces. We use the Fourier–Sato transform to prove that our classes coincide with the virtual fundamental classes introduced by the work of Behrend–Fantechi and Li–Tian, under the schematic and quasiprojectivity assumption. We also discuss an approach to construct DT4 virtual classes for  $-2$ -shifted symplectic derived schemes using the perverse sheaves of vanishing cycles.

14A30; 14N35

1. Introduction	349
2. Preliminaries on sheaf operations	355
3. Specialization functor and specialization map	367
4. Fourier–Sato transform of the specialization	372
5. Virtual fundamental classes via vanishing cycles	378
6. DT4 virtual classes and categorifications	383
7. Postponed proofs	388
References	396

## 1 Introduction

### 1.1 History

In many cases, moduli spaces such as those of stable maps and coherent sheaves have bigger dimensions than the expected dimensions computed by the Riemann–Roch theorem. To define a correct enumerative invariant counting points in such a moduli space, we need to integrate cohomology classes over the *virtual fundamental class* which is a cycle class of the virtual dimension instead of the usual fundamental class which may be ill-defined. In the work of Behrend and Fantechi [1997] and Li and Tian [1998], they introduced the notion of perfect obstruction theory for a scheme (or more generally a Deligne–Mumford stack)  $X$  and constructed the virtual fundamental class

$$[X]^{\text{vir}} \in A_{\text{vdim } X}(X)$$

from that data. Here  $\mathrm{vdim} X$  denotes the virtual dimension. In modern terms, the notion of perfect obstruction theory is understood as a classical shadow of the quasismooth derived structure. Their work can be used to construct Gromov–Witten invariants counting stable maps to smooth projective varieties [Behrend 1997; Li and Tian 1998] and Donaldson–Thomas invariants counting stable sheaves on Calabi–Yau threefolds [Thomas 2000].

Later, Behrend [2009] proved that for a projective variety  $X$  defined over  $\mathbb{C}$  and a perfect obstruction theory on it admitting a certain symmetry (called symmetric perfect obstruction theory), there exists a natural constructible function  $\nu_X$  such that the equality

$$(1-1) \quad \int_{[X]^{\mathrm{vir}}} 1 = \sum_i i \cdot \chi(\nu_X^{-1}(i))$$

holds. The natural perfect obstruction theory on a projective component of the moduli space of stable sheaves on a Calabi–Yau threefold satisfies this assumption. This suggests that Donaldson–Thomas invariants admit sheaf-theoretic categorifications. This was achieved by the work of Joyce and his collaborators [Ben-Bassat et al. 2015; Brav et al. 2015; 2019] which we briefly explain now. Let  $X$  be a derived scheme equipped with a  $-1$ -shifted symplectic structure [Pantev et al. 2013]. It induces a natural symmetric perfect obstruction theory on  $X = t_0(X)$ . It is proved in [Brav et al. 2019] that  $X$  is Zariski locally equivalent to a derived critical locus of a regular function on a smooth scheme. If  $X$  admits an orientation, one can glue certain twists of the locally defined vanishing cycle complexes to define a globally defined perverse sheaf

$$\varphi_X \in \mathrm{Perv}(X).$$

See [Brav et al. 2015] for details. The Euler characteristic of the perverse sheaf  $\varphi_X$  at a point  $x \in X$  is equal to the value of the Behrend function  $\nu_X(x)$ . Therefore the equality (1-1) implies the equality

$$\int_{[X]^{\mathrm{vir}}} 1 = \sum_i (-1)^i H^i(X, \varphi_X).$$

If we take  $X$  to be the derived moduli space of stable sheaves on a Calabi–Yau threefold equipped with the canonical  $-1$ -shifted symplectic structure [Pantev et al. 2013, Theorem 0.1] and the canonical orientation constructed in [Joyce and Upmeyer 2021], the perverse sheaf  $\varphi_X$  can be regarded as a categorification of the Donaldson–Thomas invariant.

This paper aims to connect two mathematical objects we have seen: the virtual fundamental class for an *arbitrary* quasismooth derived scheme defined over  $\mathbb{C}$  (not necessary  $-1$ -shifted symplectic!) and the perverse sheaf associated with an oriented  $-1$ -shifted symplectic derived scheme.

## 1.2 Results

Let  $Y$  be a quasismooth derived scheme over  $\mathrm{Spec} \mathbb{C}$  and

$$\mathbf{T}^*[-1]Y := \mathbf{Spec}_Y(\mathrm{Sym}(\mathbb{L}_Y^\vee[1]))$$

be its  $-1$ -shifted cotangent space. Write  $Y = t_0(Y)$  and  $\tilde{Y} = t_0(\mathbf{T}^*[-1]Y)$  and we let  $\pi: \tilde{Y} \rightarrow Y$  denote the natural projection. It is shown in [Pantev et al. 2013, Proposition 1.21] that

$$\mathbf{T}^*[-1]Y := \mathbf{Spec}_Y(\mathrm{Sym}(\mathbb{L}_Y^\vee[1]))$$

carries a canonical  $-1$ -shifted symplectic structure and one can easily show that it also carries a natural orientation. Therefore we can define a natural perverse sheaf  $\varphi_{\mathbf{T}^*[-1]Y} \in \mathrm{Perv}(\tilde{Y})$ . The following isomorphisms are proved in [Kinjo 2022, Theorem 3.1]:

$$(1-2) \quad \gamma: \pi_! \varphi_{\mathbf{T}^*[-1]Y} \cong \mathbb{Q}_Y[\mathrm{vdim} Y], \quad \bar{\gamma}: \pi_* \varphi_{\mathbf{T}^*[-1]Y} \cong \omega_Y[-\mathrm{vdim} Y].$$

Roughly speaking, the map  $\gamma$  can be regarded as a version of the Thom isomorphism for  $\mathbf{T}^*[-1]Y$  and the map  $\bar{\gamma}$  is an analog of the homotopy invariance for the singular cohomology. Therefore the composition

$$\mathbb{Q}_Y[\mathrm{vdim} Y] \xrightarrow[\cong]{\gamma^{-1}} \pi_! \varphi_{\mathbf{T}^*[-1]Y} \rightarrow \pi_* \varphi_{\mathbf{T}^*[-1]Y} \xrightarrow[\cong]{\bar{\gamma}} \omega_Y[-\mathrm{vdim} Y]$$

can be regarded as the Euler class for  $\mathbf{T}^*[-1]Y$ . We let

$$e_\varphi(\mathbf{T}^*[-1]Y) \in H_{2\mathrm{vdim} Y}^{\mathrm{BM}}(Y)$$

be the class corresponding to the above composition. As the virtual fundamental class is a generalization of the Euler class, it is natural to compare the class  $e_\varphi(\mathbf{T}^*[-1]Y)$  with the virtual fundamental class  $[Y]^{\mathrm{vir}}$ . The aim of this paper is to prove the following result:

**Theorem 1.1** (Theorem 5.3) *Assume  $\mathbb{L}_Y|_Y$  has a global resolution by a two-term complex of locally free sheaves. Then the following equality in  $H_{2\mathrm{vdim} Y}^{\mathrm{BM}}(Y)$  holds:*

$$e_\varphi(\mathbf{T}^*[-1]Y) = (-1)^{\mathrm{vdim} Y \cdot (\mathrm{vdim} Y - 1)/2} \mathrm{cl}_Y([Y]^{\mathrm{vir}}),$$

where  $\mathrm{cl}_Y: A_{\mathrm{vdim} Y}(Y) \rightarrow H_{2\mathrm{vdim} Y}^{\mathrm{BM}}(Y)$  is the cycle class map.

In other words, the above theorem gives a new construction of the virtual fundamental class in the Borel–Moore homology under a mild assumption that is satisfied when  $Y$  is quasiprojective.

### 1.3 Strategy of the proof

Before explaining the strategy of the proof of Theorem 1.1, let us recall the construction of the virtual fundamental class due to [Behrend and Fantechi 1997; Li and Tian 1998]. First, the natural map  $\mathbb{L}_Y|_Y \rightarrow \mathbb{L}_Y$  defines a purely zero-dimensional cone substack  $\mathfrak{C}_Y \subset t_0(\mathbf{T}[1]Y)$  of the classical truncation of the 1-shifted tangent stack called the intrinsic normal cone of  $Y$ . Then the virtual fundamental class is defined to be the image of the class  $[\mathfrak{C}_Y] \in A_0(t_0(\mathbf{T}[1]Y))$  under the homotopy-invariance isomorphism for vector bundle stacks proved by Kresch [1999, Proposition 5.3.2],

$$A_0(t_0(\mathbf{T}[1]Y)) \cong A_{\mathrm{vdim} Y}(Y).$$

The main difficulty of Theorem 5.3 is that proving the statement locally does not imply the global statement since the Borel–Moore homology does not satisfy the sheaf condition (in the 1-categorical

sense). Therefore we need to compare the intrinsic normal cone  $\mathfrak{C}_Y$  and the vanishing cycle complex  $\varphi_{\mathbf{T}^*[-1]Y}$  directly though they live in different spaces.

We overcome this difficulty using the Fourier–Sato transform<sup>1</sup> to relate constructible sheaves on  $\mathbf{T}^*[-1]Y$  and those on  $\mathbf{T}[1]Y$ . More precisely, take a global resolution  $\mathbb{L}_Y|_Y \cong [E^{-1} \rightarrow E^0]$  by locally free sheaves and consider the Fourier–Sato transform [Kashiwara and Schapira 1990, Section 3.7] between the  $\mathbb{R}^+$ -equivariant derived category of sheaves,

$$\mathrm{FS}_{E^{-1}} : D_{\mathbb{R}^+}^b(E^{-1}) \rightarrow D_{\mathbb{R}^+}^b((E^{-1})^\vee).$$

Using the natural inclusion  $t_0(\mathbf{T}^*[-1]Y) \hookrightarrow E^{-1}$ , one can regard  $\varphi_{\mathbf{T}^*[-1]Y}$  as a perverse sheaf on  $E^{-1}$ . Then one can show that the complex  $\mathrm{FS}_{E^{-1}}(\varphi_{\mathbf{T}^*[-1]Y})$  has the support contained in the pullback of the intrinsic normal cone  $p^*\mathfrak{C}_Y$  along the natural map  $p : (E^{-1})^\vee \rightarrow [(E^{-1})^\vee / (E^0)^\vee] \cong t_0(\mathbf{T}[1]Y)$  (see Lemma 5.4). This observation and some sheaf-theoretic analysis imply that Theorem 1.1 holds.

**Remark 1.2** Our method of the proof works for any quasismooth derived algebraic spaces with the resolution property. More generally, we believe that our proof works for any quasismooth derived Deligne–Mumford stacks with the resolution property if we can generalize the Fourier–Sato transform [Kashiwara and Schapira 1990, Section 3.7] for sheaves on Deligne–Mumford stacks.

## 1.4 Microlocal interpretation

The proof of Theorem 5.3 outlined in the previous section is motivated by microlocal geometry. We now briefly explain it.

Let  $X$  be a manifold and  $M \subset X$  be a closed submanifold. Kashiwara and Schapira [1990, Section 4.3] introduced a functor

$$\mu_{M/X} : D^b(X) \rightarrow D_{\mathbb{R}^+}^b(T_M^*X)$$

called the microlocalization functor. It is defined as the Fourier–Sato dual of the specialization functor

$$\mathrm{Sp}_{M/X} : D^b(X) \rightarrow D_{\mathbb{R}^+}^b(T_M X).$$

For a quasismooth derived scheme  $Y$ , the  $-1$ -shifted cotangent scheme  $\mathbf{T}^*[-1]Y$  can be regarded as the “conormal bundle” along the constant map  $Y \rightarrow *$ . Then the following slogan is the main idea behind the proof of Theorem 5.3:

The vanishing cycle complex  $\varphi_{\mathbf{T}^*[-1]Y}$  can be thought of as the microlocalization of the constant sheaf on a point along the constant map  $Y \rightarrow *$ .

In other words, the vanishing cycle complex  $\varphi_{\mathbf{T}^*[-1]Y}$  should be thought of as the “intrinsic microlocalization”. Here the usage of the term “intrinsic” is completely analogous to the one in the intrinsic normal cone. We can relate them using the Fourier–Sato transform.

<sup>1</sup>The author came up with this idea after a private discussion with Adeel Khan where he learned Khan’s ongoing project on derived microlocal geometry. We believe that the technical assumption on the global resolution in Theorem 1.1 can be removed once basic properties of the Fourier–Sato transform is proved in this context.

Unfortunately, the above slogan is not mathematically rigorous (at least now) because we do not have microlocalization functors with respect to arbitrary morphisms between derived schemes. Nevertheless, using a local model for quasismooth derived schemes, we can formulate a local version of the above slogan which will be stated in Theorem 4.1. This local version is enough for our purposes.

Recently, Kendric Schefers [2022] gave a completely different proof of Theorem 4.1. Further, he introduced a microlocalization functor for quasismooth closed immersion and realized the above slogan when  $Y$  can be embedded into a smooth ambient space. In another work, Adeel Khan recently announced a construction of the microlocalization functor for arbitrary morphisms between derived Artin stacks with some technical assumptions. We believe that the above slogan can be made precise using his microlocalization functor.

## 1.5 Joyce's conjecture

The present work is motivated by Joyce's conjecture [Amorim and Ben-Bassat 2017, Conjecture 5.18] which we briefly discuss now. Let  $\tau: L \rightarrow X$  be an oriented Lagrangian of an oriented  $-1$ -shifted symplectic derived Artin stack. Write  $\tau = t_0(\tau)$ ,  $L = t_0(L)$  and  $X = t_0(X)$ . Let  $\varphi_X \in \text{Perv}(X)$  be the vanishing cycle complex associated with the oriented  $-1$ -shifted symplectic structure on  $X$ . Then Joyce's conjecture predicts that there exists a natural map

$$\mu_L: \mathbb{Q}_L[\text{vdim } L] \rightarrow \tau^! \varphi_X.$$

See [Amorim and Ben-Bassat 2017, Section 5.3] for the full statement of this conjecture and [Joyce 2018, Section 7] for important consequences including the construction of cohomological Hall algebras for CY3 categories. When  $X$  is a point equipped with the trivial  $-1$ -shifted symplectic structure, the vanishing cycle complex  $\varphi_X$  is the constant sheaf and the oriented Lagrangian structure for  $\tau$  is identified with an oriented  $-2$ -shifted symplectic structure on  $L$ . In this case, it is expected that the map  $\mu_L$  is given by the DT4 virtual class introduced in the work of Cao and Leung [2014], Borisov and Joyce [2017] and Oh and Thomas [2023]. In other words, the map  $\mu_L$  in the general case should be understood as a relative version of the DT4 virtual class.

Joyce's conjecture suggests that (perverse) sheaf theory may be used to give a new construction of DT4 virtual classes. Since the virtual fundamental class for a quasismooth derived scheme is equal to the DT4 virtual class of its  $-2$ -shifted cotangent space (see [Oh and Thomas 2023, Section 8]), Theorem 1.1 can be regarded as the first step toward this project. In Section 6.1, we propose a conjectural approach to construct DT4 virtual classes assuming some variations of the isomorphism (1-2).

## 1.6 Structure of the paper

The paper is organized as follows.

In Section 2 we recall some sheaf operations and prove some of their basic properties.

In Section 3 we recall the construction of Verdier's specialization functor and use it to construct the specialization morphism of the Borel–Moore homology.

In Section 4 we prove that the Fourier–Sato transform relates the specialization functor and the vanishing cycle functor. This gives a new proof of Davison's dimensional reduction theorem [2017, Theorem A.1] and a proof of Theorem 1.1 for local models of quasismooth derived schemes.

In Section 5 we prove Theorem 1.1.

In Section 6 we discuss some conjectural generalizations of Theorem 1.1.

**Acknowledgements** I am very grateful to Adeel Khan for sharing many ideas on his ongoing project on derived microlocal geometry. Without him, I could have never come up with the idea to use the Fourier–Sato transform to prove Theorem 1.1. I also thank my supervisor Yukinobu Toda for fruitful discussions and careful reading, Hyeonjun Park for teaching me DT4 theory and Alexandre Minets for pointing out that a conjecture in the previous version of this paper was not correct.

This work was supported by the WINGS-FMSP program at the Graduate School of Mathematical Science, University of Tokyo, and JSPS KAKENHI grant JP21J21118.

## Notation and conventions

- All schemes considered in this paper are separated and of finite type over the complex number field.
- For a topological space  $X$ , we let  $D^+(X)$  (resp.  $D^b(X)$ ,  $D_c^b(X)$ ) denote the derived category of lower bounded complexes (resp. bounded complexes, bounded complexes with constructible cohomology) in  $\mathbb{Q}$ -vector spaces.
- For a topological space  $X$  with an  $\mathbb{R}^+$ -action, we let  $D_{\mathbb{R}^+}^+(X) \subset D^+(X)$  denote the full subcategory consisting of  $\mathbb{R}^+$ -equivariant objects following [Kashiwara and Schapira 1990, Definition 3.7.1].
- For a closed subset  $Z \subset X$  and a complex  $\mathcal{F} \in D^+(Z)$ , we let  $\mathcal{F}|_Z$  denote the complex  $(Z \hookrightarrow X)^* \mathcal{F}$ ,  $\mathcal{F}_Z$  denote the complex  $(Z \hookrightarrow X)_* (Z \hookrightarrow X)^* \mathcal{F}$ ,  $\mathcal{F}|_Z^!$  denote the complex  $(Z \hookrightarrow X)^! \mathcal{F}$  and  $\Gamma_Z(\mathcal{F})$  denote the complex  $(Z \hookrightarrow X)_! (Z \hookrightarrow X)^! \mathcal{F}$ .
- For a closed subset  $Z \subset X$  and a complex  $\mathcal{F} \in D^+(Z)$ , we sometimes regard  $\mathcal{F}$  as a complex on  $X$ . Conversely, for a complex  $\mathcal{G} \in D^+(X)$  whose support is contained in  $Z$ , we sometimes regard  $\mathcal{G}$  as a complex on  $Z$ .
- For a continuous map  $f: X \rightarrow Y$  between locally compact Hausdorff spaces such that the functor  $f^!$  is defined, we let  $\omega_{X/Y}$  denote the complex  $f^! \mathbb{Q}_Y$  and  $H^*(X \rightarrow Y)$  denote its cohomology  $H^*(X, \omega_{X/Y})$ . We also write  $\omega_X = \omega_{X/\text{pt}}$  and  $H_*^{\text{BM}}(X) = H^{-*}(X \rightarrow \text{pt})$ . The Verdier-duality functor  $R\mathcal{H}om(-, \omega_X)$  is denoted by  $\mathbb{D}_X$ .



- For a complex separated scheme  $X$  of finite type and its irreducible subscheme  $Z$ , the classes  $[Z] \in A_{\dim Z}(X)$  and  $[Z]_{\text{BM}} \in H_{2\dim Z}^{\text{BM}}(X)$  are defined by the pushforward of the fundamental class of  $Z$  to  $X$  in the Chow group and the Borel–Moore homology group respectively.
- If there is no confusion, we use expressions such as  $f_*$ ,  $f_!$ , and  $\mathcal{H}om$  for the derived functors  $Rf_*$ ,  $Rf_!$ , and  $R\mathcal{H}om$ .

## 2 Preliminaries on sheaf operations

In this section, we recall the definition of the nearby cycle and vanishing cycle functors and the Fourier–Sato transform, and prove some of their basic properties. Readers who are familiar with sheaf theory can safely skip this section. We assume that all topological spaces are locally compact Hausdorff, and for all continuous maps  $f$ , the functor  $f_!$  has finite cohomological dimension.

### 2.1 Base change maps

Consider a commutative diagram of topological spaces

$$(2-1) \quad \begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ g' \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

We have the natural transformations of functors from  $D^+(X)$  to  $D^+(Y')$

$$(2-2) \quad g^* f_* \rightarrow f'_* g'^*,$$

and functors from  $D^+(X)$  to  $D^+(Y')$

$$(2-3) \quad f'_! g'^! \rightarrow g^! f_!.$$

The former map is defined by the composition

$$g^* f_* \rightarrow f'_* f'^* g^* f_* \cong f'_* g'^* f^* f_* \rightarrow f'_* g'^*$$

and the latter map is defined similarly.

If the diagram (2-1) is Cartesian, we have natural transformations of functors from  $D^+(X')$  to  $D^+(Y)$ ,

$$(2-4) \quad f_! g'_* \rightarrow g_* f'_!,$$

functors from  $D^+(X)$  to  $D^+(Y')$ ,

$$(2-5) \quad g^* f_! \xrightarrow{\sim} f'_! g'^*,$$

functors from  $D^+(X)$  to  $D^+(Y')$ ,

$$(2-6) \quad f'_* g'^! \xrightarrow{\sim} g^! f_*,$$

and functors from  $D^+(Y)$  to  $D^+(X')$ ,

$$(2-7) \quad g'^* f^! \rightarrow f'^! g^*.$$

The map (2-4) is defined in [Kashiwara and Schapira 1990, (2.5.7)], (2-5) is defined in [Kashiwara and Schapira 1990, Proposition 2.6.7], (2-6) is defined in [Kashiwara and Schapira 1990, Proposition 3.1.9(ii)] and (2-7) is defined in [Kashiwara and Schapira 1990, Proposition 3.1.9(iii)]. We call the maps (2-2)–(2-7) base change maps.

## 2.2 Natural transformations between four functors

Let  $f: X \rightarrow Y$  be a continuous map between topological spaces. Then we can construct the natural transformations

$$(2-8) \quad \text{id} \rightarrow f_* f^*, \quad f^* f_* \rightarrow \text{id},$$

$$(2-9) \quad \text{id} \rightarrow f^! f_!, \quad f_! f^! \rightarrow \text{id},$$

$$(2-10) \quad f_! \rightarrow f_*.$$

If  $f$  is a closed immersion, we have a natural transformation

$$(2-11) \quad f^! \rightarrow f^*$$

obtained by the composition

$$f^! \cong f^* f_* f^! \cong f^* f_! f^! \rightarrow f^*.$$

If we have an isomorphism  $f^! \mathbb{Q}_Y \cong \mathbb{Q}_X[r]$  (eg a smooth morphism between complex analytic spaces), we have a natural transformation

$$(2-12) \quad f^*[r] \rightarrow f^!.$$

This is a consequence of [Kashiwara and Schapira 1990, Proposition 3.1.11], which constructs a natural transformation

$$(2-13) \quad f^*(-) \otimes f^! \mathbb{Q}_Y \rightarrow f^!(-)$$

for general  $f$ .

Now consider a Cartesian diagram of topological spaces

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ \downarrow g' & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

For a choice of the natural transformations  $\eta$  in (2-8)–(2-11) and a choice of the sheaf operation  $a \in \{(-)^*, (-)_*, (-)^!, (-)_!\}$ , we can verify the commutativity of  $\eta$  and the base change maps for  $a$ .

For example, if we take  $\eta$  to be the unit map  $\text{id} \rightarrow f_* f^*$  and  $a = (-)^!$ , the following diagram commutes:

$$\begin{array}{ccc} g^! & \longrightarrow & g^! f_* f^* \\ \downarrow & & \wr \uparrow \\ f'_* f'^* g^! & \longrightarrow & f'_* g'^! f^* \end{array}$$

Now assume that there exists an isomorphism  $f^! \mathbb{Q}_Y \cong \mathbb{Q}_X[r]$  for some integer  $r$  and the composition

$$(2-14) \quad \mathbb{Q}_{X'}[r] \cong g'^* \mathbb{Q}_X[r] \cong g'^* f^! \mathbb{Q}_Y \rightarrow f'^! g^* \mathbb{Q}_Y \cong f'^! \mathbb{Q}_{Y'}$$

is an isomorphism. Then we can see that for each choice of  $a \in \{(-)^*, (-)_*, (-)^!, (-)_!\}$  the natural transformation (2-12) commutes with the base change maps for  $a$ . For example if we take  $a = (-)^!$ , the following diagram commutes:

$$\begin{array}{ccc} f'^*[r]g^! & \longrightarrow & f'^!g^! \\ \downarrow & & \wr \downarrow \\ g'^!f^*[r] & \longrightarrow & g'^!f^! \end{array}$$

Now remove the assumption that (2-14) is invertible and take  $a \in \{(-)_*, (-)_!\}$  and  $b \in \{(-)^*, (-)^!\}$ . Let  $G_a: D^+(Y') \rightarrow D^+(Y)$  (resp.  $G_b: D^+(Y) \rightarrow D^+(Y')$ ) be the functor corresponding to  $a$  (resp.  $b$ ). Then we can see that the base change maps for the composition transformation  $G_a G_b$  commute with the map (2-12). For example if we take  $a = (-)_*$  and  $b = (-)^*$ , or  $a = (-)_!$  and  $b = (-)^!$ , we have the following commutative diagrams:

$$(2-15) \quad \begin{array}{ccc} f^*[r]g_*g^* & \longrightarrow & f^!g_*g^* \\ \downarrow & & \wr \uparrow \\ g'_*f'^*[r]g^* & & g'_*f'^!g^* \\ \wr \downarrow & & \uparrow \\ g'_*g'^*f^*[r] & \longrightarrow & g'_*g'^*f^! \end{array} \quad \begin{array}{ccc} f^*[r]g_!g^! & \longrightarrow & f^!g_!g^! \\ \wr \downarrow & & \uparrow \\ g'_!f'^*[r]g^! & & g'_!f'^!g^! \\ \downarrow & & \wr \uparrow \\ g'_!g'^!f^*[r] & \longrightarrow & g'_!g'^!f^! \end{array}$$

### 2.3 Vanishing cycle and nearby cycle functors

Let  $X$  be a complex analytic space and  $u: X \rightarrow \mathbb{C}$  be a complex analytic function. We write

$$X_0 := u^{-1}(0), \quad X_{>0} := \{z \in X \mid \text{Re}(u(z)) > 0\}, \quad X_{\leq 0} := X \setminus X_{>0}.$$

We define the vanishing cycle functor and the nearby cycle functor

$$\psi_u, \varphi_u: D^+(X) \rightarrow D^+(X_0)$$

by the formulas

$$\begin{aligned} \psi_u &:= (X_0 \hookrightarrow X)^*(X_{>0} \hookrightarrow X)_*(X_{>0} \hookrightarrow X)^*, \\ \varphi_u &:= (X_0 \hookrightarrow X_{\leq 0})^*(X_{\leq 0} \hookrightarrow X)^!. \end{aligned}$$

If we are given a morphism  $q: T \rightarrow X$  from another complex analytic space, we have natural transformations

$$(2-16) \quad \psi_u q_* \rightarrow q_0_* \psi_{u \circ q}, \quad \varphi_u q_* \rightarrow q_0_* \varphi_{u \circ q},$$

$$(2-17) \quad q_0! \psi_{u \circ q} \rightarrow \psi_u q!, \quad q_0! \varphi_{u \circ q} \rightarrow \varphi_u q!,$$

$$(2-18) \quad q_0^* \psi_u \rightarrow \psi_{u \circ q} q^*, \quad q_0^* \varphi_u \rightarrow \varphi_{u \circ q} q^*,$$

$$(2-19) \quad \psi_{u \circ q} q^! \rightarrow q_0^! \psi_u, \quad \varphi_{u \circ q} q^! \rightarrow q_0^! \varphi_u,$$

where the map  $q_0: T_0 = (u \circ q)^{-1}(0) \rightarrow X_0$  is induced from  $q$ . All of these maps are defined by combining the base change maps (2-2)–(2-7). We call these maps the base change maps for the nearby and vanishing cycle functor. The four maps in (2-16) and (2-17) are invertible when  $q$  is proper, and the four maps in (2-18) and (2-19) are invertible when  $q$  is smooth.

There is another way to define the nearby cycle and vanishing cycle functors as in [Kashiwara and Schapira 1990, Section 8.6]. Let  $\tilde{\mathbb{C}}^*$  be the universal cover of  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ , and  $p: \tilde{\mathbb{C}}^* \rightarrow \mathbb{C}^*$  the natural map. Define objects  $K'_\psi, K'_\varphi \in D^b(\mathbb{C})$  by

$$K'_\psi := p! \mathbb{Q}_{\tilde{\mathbb{C}}^*}, \quad K'_\varphi := \text{Cone}(p! \mathbb{Q}_{\tilde{\mathbb{C}}^*} \xrightarrow{\text{tr}} \mathbb{Q}_{\mathbb{C}}),$$

where the map  $\text{tr}: p! \mathbb{Q}_{\tilde{\mathbb{C}}^*} \rightarrow \mathbb{Q}_{\mathbb{C}}$  is defined by the composition

$$p! \mathbb{Q}_{\tilde{\mathbb{C}}^*} \cong p! p^! \mathbb{Q}_{\mathbb{C}} \rightarrow \mathbb{Q}_{\mathbb{C}}.$$

For a given complex analytic function  $u: X \rightarrow \mathbb{C}$ , we define functors

$$\psi'_u, \varphi'_u: D^+(X) \rightarrow D^+(X_0)$$

by

$$\psi'_u := (X_0 \hookrightarrow X)^* R\mathcal{H}om(u^* K'_\psi, -),$$

$$\varphi'_u := (X_0 \hookrightarrow X)^* R\mathcal{H}om(u^* K'_\varphi, -).$$

Now we want to compare  $\psi_u$  and  $\psi'_u$ , and  $\varphi_u$  and  $\varphi'_u$ . To do this, we first define objects  $K_\psi, K_\varphi \in D^b(\mathbb{C})$  by

$$K_\psi := (\mathbb{C}_{>0} \hookrightarrow \mathbb{C})! \mathbb{Q}_{\mathbb{C}_{>0}},$$

$$K_\varphi := \text{Cone}((\mathbb{C}_{>0} \hookrightarrow \mathbb{C})! \mathbb{Q}_{\mathbb{C}_{>0}} \rightarrow \mathbb{Q}_{\mathbb{C}}) \cong (\mathbb{C}_{\leq 0} \hookrightarrow \mathbb{C})! \mathbb{Q}_{\mathbb{C}_{\leq 0}}.$$

Then we have isomorphisms

$$\begin{aligned} (X_0 \hookrightarrow X)^* R\mathcal{H}om(u^* K_\psi, -) &\cong (X_0 \hookrightarrow X)^* R\mathcal{H}om((X_{>0} \hookrightarrow X)! \mathbb{Q}_{X_{>0}}, -) \\ &\cong (X_0 \hookrightarrow X)^* (X_{>0} \hookrightarrow X)_* R\mathcal{H}om(\mathbb{Q}_{X_{>0}}, (X_{>0} \hookrightarrow X)^*(-)) \\ &\cong (X_0 \hookrightarrow X)^* (X_{>0} \hookrightarrow X)_* (X_{>0} \hookrightarrow X)^* = \psi_u \end{aligned}$$

and

$$\begin{aligned} (X_0 \hookrightarrow X)^* R\mathcal{H}om(u^* K_\varphi, -) &\cong (X_0 \hookrightarrow X)^* R\mathcal{H}om((X_{\leq 0} \hookrightarrow X)! \mathbb{Q}_{X_{\leq 0}}, -) \\ &\cong (X_0 \hookrightarrow X)^* (X_{\leq 0} \hookrightarrow X)_* R\mathcal{H}om(\mathbb{Q}_{X_{\leq 0}}, (X_{\leq 0} \hookrightarrow X)^!(-)) \\ &\cong (X_0 \hookrightarrow X_{\leq 0})^* (X_{\leq 0} \hookrightarrow X)^! = \varphi_u. \end{aligned}$$

If we pick an open immersion  $j : \mathbb{C}_{>0} \hookrightarrow \tilde{\mathbb{C}}^*$  over  $\mathbb{C}$ , the natural map  $j_! \mathbb{Q}_{\mathbb{C}_{>0}} \rightarrow \mathbb{Q}_{\tilde{\mathbb{C}}^*}$  induces maps

$$K_\psi \rightarrow K'_\psi, \quad K_\varphi \rightarrow K'_\varphi,$$

and hence maps

$$\psi'_u \rightarrow \psi_u, \quad \varphi'_u \rightarrow \varphi_u.$$

For a constructible complex  $\mathcal{F} \in D_c^b(X)$ , one can show that the maps

$$(2-20) \quad \psi'_u(\mathcal{F}) \rightarrow \psi_u(\mathcal{F}), \quad \varphi'_u(\mathcal{F}) \rightarrow \varphi_u(\mathcal{F})$$

are invertible (see [Schürmann 2003, Lemma 1.1.1, Example 1.1.3] for details).

Now consider the following natural short exact sequence of complexes on  $\mathbb{C}$ :

$$(2-21) \quad \Delta_1 : 0 \rightarrow \mathbb{Q}_{\mathbb{C}} \rightarrow K'_\varphi \rightarrow K'_\psi[1] \rightarrow 0.$$

Then the distinguished triangle  $(X_0 \hookrightarrow X)^* R\mathcal{H}om(u^* \Delta_1, -)$  is identified with

$$(2-22) \quad \psi'_u[-1] \rightarrow \varphi'_u \rightarrow (X_0 \hookrightarrow X)^* \rightarrow \psi'_u.$$

Next, consider the following short exact sequence of complexes on  $\mathbb{C}$ :

$$(2-23) \quad \Delta_2 : 0 \rightarrow K'_\psi[1] \rightarrow K'_\varphi \rightarrow \text{Cone}((\mathbb{C}^* \hookrightarrow \mathbb{C})_! \mathbb{Q}_{\mathbb{C}^*} \rightarrow \mathbb{Q}_{\mathbb{C}}) \rightarrow 0.$$

Here the first map is defined by the diagram

$$\begin{array}{ccc} p_! \mathbb{Q}_{\tilde{\mathbb{C}}^*} & \xrightarrow{1-T} & p_! \mathbb{Q}_{\tilde{\mathbb{C}}^*} \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathbb{Q}_{\mathbb{C}} \end{array}$$

where  $T$  is the monodromy operator; in other words, if we let  $\alpha : \tilde{\mathbb{C}}^* \xrightarrow{\sim} \tilde{\mathbb{C}}^*$  denote the covering transformation over  $\mathbb{C}^*$  corresponding to a counterclockwise loop, the map  $T$  is given by the map  $\alpha_! \mathbb{Q}_{\tilde{\mathbb{C}}^*} \cong \mathbb{Q}_{\tilde{\mathbb{C}}^*}$ . The second map in  $\Delta_2$  is defined by the following diagram:

$$\begin{array}{ccc} p_! \mathbb{Q}_{\tilde{\mathbb{C}}^*} & \xrightarrow{\text{tr}} & (\mathbb{C}^* \hookrightarrow \mathbb{C})_! \mathbb{Q}_{\mathbb{C}^*} \\ \downarrow & & \downarrow \\ \mathbb{Q}_{\mathbb{C}} & \xrightarrow{\text{id}} & \mathbb{Q}_{\mathbb{C}} \end{array}$$

Since we have an isomorphism  $\text{Cone}((\mathbb{C}^* \hookrightarrow \mathbb{C})_! \mathbb{Q}_{\mathbb{C}^*} \rightarrow \mathbb{Q}_{\mathbb{C}}) \cong \mathbb{Q}_0$ , the distinguished triangle

$$(X_0 \hookrightarrow X)^* R\mathcal{H}om(u^* \Delta_2, -)$$

is identified with

$$(2-24) \quad (X_0 \hookrightarrow X)^! \rightarrow \varphi'_u \rightarrow \psi'_u[-1] \rightarrow (X_0 \hookrightarrow X)^![1].$$

For a constructible complex  $\mathcal{F} \in D_c^b(X)$ , the isomorphisms in (2-20) and the distinguished triangles (2-22) and (2-24) induces the distinguished triangles

$$(2-25) \quad \psi_u(\mathcal{F})[-1] \rightarrow \varphi_u(\mathcal{F}) \rightarrow (X_0 \hookrightarrow X)^* \mathcal{F} \rightarrow \psi_u(\mathcal{F}),$$

$$(2-26) \quad (X_0 \hookrightarrow X)^! \mathcal{F} \rightarrow \varphi_u(\mathcal{F}) \rightarrow \psi_u(\mathcal{F})[-1] \rightarrow (X_0 \hookrightarrow X)^! \mathcal{F}[1].$$

It is clear that the map

$$(2-27) \quad \varphi_u(\mathcal{F}) \rightarrow (X_0 \hookrightarrow X)^* \mathcal{F}$$

in the distinguished triangle (2-25) is equal to the composition

$$\varphi_u(\mathcal{F}) = (X_0 \hookrightarrow X_{\leq 0})^* (X_{\leq 0} \hookrightarrow X)^! \mathcal{F} \rightarrow (X_0 \hookrightarrow X_{\leq 0})^* (X_{\leq 0} \hookrightarrow X)^* \mathcal{F} \cong (X_0 \hookrightarrow X)^* \mathcal{F}$$

where the first map is (2-11), and the map

$$(2-28) \quad (X_0 \hookrightarrow X)^! \mathcal{F} \rightarrow \varphi_u(\mathcal{F})$$

in the distinguished triangle (2-26) is equal to the following composition

$$(X_0 \hookrightarrow X)^! \mathcal{F} \cong (X_0 \hookrightarrow X_{\leq 0})^! (X_{\leq 0} \hookrightarrow X)^! \rightarrow (X_0 \hookrightarrow X_{\leq 0})^* (X_{\leq 0} \hookrightarrow X)^! = \varphi_u(\mathcal{F})$$

where the second map is (2-11).

Let  $q: T \rightarrow X$  be a morphism of complex analytic spaces and  $u$  be a holomorphic function on  $X$ . Then for each choice of  $a \in \{(-)^*, (-)_*, (-)^!, (-)_!\}$ , one can show that the distinguished triangles (2-25) and (2-26) commute with the base change maps for  $a$ . For example, if we take  $a = (-)_*$ , for a complex  $\mathcal{F} \in D_c^b(T)$  the following diagrams commute:

$$(2-29) \quad \begin{array}{ccccccc} \psi_u(q_* \mathcal{F})[-1] & \longrightarrow & \varphi_u(q_* \mathcal{F}) & \longrightarrow & (X_0 \hookrightarrow X)^* q_* \mathcal{F} & \longrightarrow & \psi_u(q_* \mathcal{F}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ q_{0*} \psi_{u \circ q}(\mathcal{F})[-1] & \rightarrow & q_{0*} \varphi_{u \circ q}(\mathcal{F}) & \rightarrow & q_{0*} (T_0 \hookrightarrow T)^* \mathcal{F} & \rightarrow & q_{0*} \psi_{u \circ q}(\mathcal{F}) \end{array}$$

$$(2-30) \quad \begin{array}{ccccccc} (X_0 \hookrightarrow X)^! q_* \mathcal{F} & \longrightarrow & \varphi_u(q_* \mathcal{F}) & \longrightarrow & \psi_u(q_* \mathcal{F})[-1] & \longrightarrow & (X_0 \hookrightarrow X)^! q_* \mathcal{F}[1] \\ \downarrow \wr & & \downarrow & & \downarrow & & \downarrow \wr \\ q_{0*} (T_0 \hookrightarrow T)^! \mathcal{F} & \rightarrow & q_{0*} \varphi_{u \circ q}(\mathcal{F}) & \rightarrow & q_{0*} \psi_{u \circ q}(\mathcal{F})[-1] & \rightarrow & q_{0*} (T_0 \hookrightarrow T)^! \mathcal{F}[1] \end{array}$$

Now let  $Z$  be a complex analytic space and take a constructible object  $\mathcal{F} \in D_c^b(Z)$ . Denote by  $\pi: Z \times \mathbb{C} \rightarrow Z$  and  $u: Z \times \mathbb{C} \rightarrow \mathbb{C}$  the projections. Then it is clear that  $\varphi_u(\pi^* \mathcal{F}) = 0$ . We have the following statement whose proof will be given in Section 7.1:

**Proposition 2.1** Denote by  $i: Z \times \{0\} \hookrightarrow Z \times \mathbb{C}$  the natural inclusion. Then the composition

$$\mathcal{F} \cong i^* \pi^* \mathcal{F} \xrightarrow{\sim} \psi_u(\pi^* \mathcal{F}) \xrightarrow{\sim} i^! \pi^* \mathcal{F}[2] \cong i^! \pi^! \mathcal{F} \cong \mathcal{F}$$

is the identity map. Here the second and third morphisms are constructed in (2-25) and (2-26) respectively, and the third isomorphism is constructed in (2-12).

Now we discuss the commutativity of the vanishing cycle functor and the Verdier duality functor following [Massey 2016]. First, for  $\alpha, \beta \in \mathbb{R}$  such that  $0 \leq \beta - \alpha < 2\pi$ , we define a closed subset  $D^{[\alpha, \beta]} \subset \mathbb{C}$  by

$$D^{[\alpha, \beta]} := \{z = R \cdot e^{i\theta} \in \mathbb{C} \mid R \in \mathbb{R}_{\geq 0}, \theta \in [\alpha, \beta]\}.$$

For a complex analytic space  $X$  and a regular function  $u$  on it, we define a functor

$$\varphi_u^{[\alpha, \beta]}: D^+(X) \rightarrow D^+(X_0)$$

by

$$\varphi_u^{[\alpha, \beta]} := (X_0 \hookrightarrow u^{-1}(D^{[\alpha, \beta]}))^*(u^{-1}(D^{[\alpha, \beta]}) \hookrightarrow X)^!.$$

Clearly, we have  $\varphi_u = \varphi_u^{[\pi/2, 3\pi/2]}$ . For real numbers  $\alpha, \alpha', \beta', \beta \in \mathbb{R}$  such that  $\alpha \leq \alpha' \leq \beta' \leq \beta$  and  $\beta - \alpha < 2\pi$  and a constructible complex  $\mathcal{F} \in D_c^b(X)$ , the natural transform (2-11) induces a map

$$(2-31) \quad \varphi_u^{[\alpha', \beta']}(\mathcal{F}) \rightarrow \varphi_u^{[\alpha, \beta]}(\mathcal{F})$$

which is an isomorphism (see [Schürmann 2003, Lemma 1.1.1, Example 1.1.3]). We define an isomorphism  $T_\pi: \varphi_u \cong \varphi_u^{[3\pi/2, 5\pi/2]}$  by the composition

$$\varphi_u = \varphi_u^{[\pi/2, 3\pi/2]} \xleftarrow{\sim} \varphi_u^{[3\pi/2, 3\pi/2]} \xrightarrow{\sim} \varphi_u^{[3\pi/2, 5\pi/2]}.$$

This map is a half of the monodromy operator. Define closed subsets  $X_{\geq 0}$  and  $X_{\text{Re}=0}$  of  $X$  by

$$X_{\geq 0} := \{z \in X \mid \text{Re}(u(z)) \geq 0\}, \quad X_{\text{Re}=0} := \{z \in X \mid \text{Re}(u(z)) = 0\}.$$

We define an isomorphism

$$(2-32) \quad \mathbb{D}_{X_0} \circ \varphi_u \cong \varphi_u \circ \mathbb{D}_X$$

by the composition

$$\begin{aligned} \mathbb{D}_{X_0} \circ \varphi_u &= \mathbb{D}_{X_0}(X_0 \hookrightarrow X_{\leq 0})^*(X_{\leq 0} \hookrightarrow X)^! \\ &\cong \mathbb{D}_{X_0}(X_0 \hookrightarrow X_{\text{Re}=0})^*(X_{\text{Re}=0} \hookrightarrow X_{\leq 0})^*(X_{\leq 0} \hookrightarrow X)^! \\ &\cong (X_0 \hookrightarrow X_{\text{Re}=0})^!(X_{\text{Re}=0} \hookrightarrow X_{\leq 0})^!(X_{\leq 0} \hookrightarrow X)^! \mathbb{D}_X \\ &\cong (X_0 \hookrightarrow X_{\text{Re}=0})^!(X_{\text{Re}=0} \hookrightarrow X_{\geq 0})^*(X_{\geq 0} \hookrightarrow X)^! \mathbb{D}_X \\ &\cong (X_0 \hookrightarrow X_{\text{Re}=0})^*(X_{\text{Re}=0} \hookrightarrow X_{\geq 0})^*(X_{\geq 0} \hookrightarrow X)^! \mathbb{D}_X \\ &\cong (X_0 \hookrightarrow X_{\geq 0})^*(X_{\geq 0} \hookrightarrow X)^! \mathbb{D}_X = \varphi_u^{[3\pi/2, 5\pi/2]} \circ \mathbb{D}_X \cong \varphi_u \circ \mathbb{D}_X, \end{aligned}$$

where the third isomorphism is the inverse of the base change map (2-7) which is invertible thanks to [Massey 2016, Lemma 2.2], the fourth isomorphism is the natural map (2-11) which is invertible thanks to [Massey 2016, Lemma 2.1], and the final isomorphism is  $T_\pi^{-1} \mathbb{D}_X$ . For a constructible complex  $\mathcal{F} \in D_c^b(X)$  we can show that the diagram

$$(2-33) \quad \begin{array}{ccc} \mathbb{D}_{X_0}(X_0 \hookrightarrow X)^* \mathcal{F} & \longrightarrow & \mathbb{D}_{X_0} \varphi_u(\mathcal{F}) \\ \downarrow \wr & & (2-32) \downarrow \wr \\ (X_0 \hookrightarrow X)^! \mathbb{D}_X \mathcal{F} & \longrightarrow & \varphi_u(\mathbb{D}_X \mathcal{F}) \end{array}$$

commutes, where the top (resp. bottom) horizontal arrow is the map (2-27) (resp. (2-28)). This is essentially a consequence of the Verdier self-duality of the natural transform (2-11).

## 2.4 Fourier–Sato transforms

Let  $X$  be a topological space with an  $\mathbb{R}^+$ -action. A sheaf  $\mathcal{F}$  on  $X$  is called conic if for each  $x \in X$ , the restriction  $\mathcal{F}|_{\mathbb{R}^+ \cdot x}$  is locally constant. We let  $D_{\mathbb{R}^+}^+(X) \subset D^+(X)$  denote the full subcategory consisting of objects whose cohomology sheaves are conic. Now let  $Z$  be a topological space and consider a vector bundle  $E$  on  $Z$ . We equip  $E$  with the scaling  $\mathbb{R}^+$ -action. Denote by  $\pi_E: E \rightarrow Z$  the projection and  $0_E: Z \hookrightarrow E$  the zero section. Then for an object  $\mathcal{F} \in D_{\mathbb{R}^+}^+(E)$ , it is shown in [Kashiwara and Schapira 1990, Proposition 3.7.5] that the maps

$$(2-34) \quad \pi_{E*} \mathcal{F} \rightarrow \pi_{E*} 0_{E*} 0_E^* \mathcal{F} \cong 0_E^* \mathcal{F},$$

$$(2-35) \quad 0_E^! \mathcal{F} \cong \pi_{E!} 0_{E!} 0_E^! \mathcal{F} \rightarrow \pi_{E!} \mathcal{F}$$

are invertible.

Now we define the Fourier–Sato transform following [Kashiwara and Schapira 1990, Section 3.7]. For applications in this paper, we always work with complex vector bundles. Let  $Z$  be a topological space and  $E$  be a complex vector bundle over  $Z$ . Denote by  $E^\vee$  the (complex) dual vector bundle. Define closed subsets  $P, P' \subset E \oplus E^\vee$  by

$$P := \{(v, w) \in E \oplus E^\vee \mid \operatorname{Re} w(v) \geq 0\}, \quad P' := \{(v, w) \in E \oplus E^\vee \mid \operatorname{Re} w(v) \leq 0\}.$$

Consider the diagram

$$\begin{array}{ccccc} & & P' & & \\ & & \downarrow \iota_{P'} & & \\ E & \xleftarrow{p} & E \oplus E^\vee & \xrightarrow{p'} & E^\vee \\ & & \uparrow \iota_P & & \\ & & P & & \end{array}$$

where  $p$  and  $p'$  are natural projections and  $\iota_P$  and  $\iota_{P'}$  are natural inclusions. We define four functors  $\operatorname{FS}_E, \operatorname{FS}'_E, \check{\operatorname{FS}}_E, \check{\operatorname{FS}}'_E: D_{\mathbb{R}^+}^+(E) \rightarrow D_{\mathbb{R}^+}^+(E^\vee)$  by

$$(2-36) \quad \begin{aligned} \operatorname{FS}_E &:= p'_! \iota_{P'}^* \iota_P^* p^*, & \operatorname{FS}'_E &:= p'_* \iota_{P'}^! \iota_P^! p^*, \\ \check{\operatorname{FS}}_E &:= p'_! \iota_{P'}^* \iota_P^! p^!, & \check{\operatorname{FS}}'_E &:= p'_! \iota_P^* \iota_{P'}^! p^!. \end{aligned}$$

The functor  $\operatorname{FS}_E$  is called the Fourier–Sato transform. It is shown in [Kashiwara and Schapira 1990, Theorem 3.7.9] that these functors are equivalences. Now we discuss the relations between these functors. Let  $i_P: P \cap P' \hookrightarrow P$  and  $i_{P'}: P \cap P' \hookrightarrow P'$  be the natural inclusions and consider the composition

$$\begin{aligned} \operatorname{FS}_E &= p'_! \iota_{P'}^* \iota_P^* p^* \xleftarrow{\sim} p'_! \iota_{P'}^* i_{P'}^! i_P^! \iota_P^* p^* \\ &\xleftarrow{\sim} p'_! \iota_P^! i_{P*} i_P^* \iota_P^! p^* \xrightarrow{\sim} p'_! \iota_P^! i_{P*} i_P^* \iota_P^! p^* \xleftarrow{\sim} p'_! \iota_P^! p^* = \operatorname{FS}'_E, \end{aligned}$$



where the first isomorphism follows from (2-35) applied to the projection  $p': E \oplus E^\vee \rightarrow E^\vee$ , the second isomorphism follows from [Kashiwara and Schapira 1990, Exercice II.2] or [Massey 2016, Lemma 2.2], the third isomorphism follows from [Kashiwara and Schapira 1990, Lemma 3.7.6] and the final isomorphism follows from (2-34). Similarly, we have a natural isomorphism

$$\check{\mathrm{FS}}_E \cong \check{\mathrm{FS}}'_E.$$

Let  $a: E \xrightarrow{\sim} E$  be the map multiplying  $-1$ , and for  $\mathcal{F} \in D_{\mathbb{R}+}^+(E)$  we write  $\mathcal{F}^a := a^* \mathcal{F}$ . It is clear that  $\mathrm{FS}_E(\mathcal{F}^a) \cong \mathrm{FS}_E(\mathcal{F})^a$ . We write  $\mathrm{FS}_E^a := \mathrm{FS}_E \circ (-)^a$ , and similarly for  $\mathrm{FS}'_E^a$ ,  $\check{\mathrm{FS}}_E^a$  and  $\check{\mathrm{FS}}'_E^a$ . We have

$$\check{\mathrm{FS}}_E'^a \cong p'_! \iota_{P'}^* \iota_{P'}^! p'^! \cong p'_! \iota_{P'}^* \iota_{P'}^! p'^*[2 \operatorname{rank} E] = \mathrm{FS}_E[2 \operatorname{rank} E],$$

where the second isomorphism is defined using the orientation on  $E$  induced by the complex structure. Therefore we have isomorphisms

$$(2-37) \quad \mathrm{FS}'_E \cong \mathrm{FS}_E \cong \check{\mathrm{FS}}_E'^a[-2 \operatorname{rank} E] \cong \check{\mathrm{FS}}_E^a[-2 \operatorname{rank} E].$$

By definition the functor  $\mathrm{FS}_E$  is naturally left adjoint (hence quasi-inverse) to  $\check{\mathrm{FS}}_{E^\vee}$  and the functor  $\mathrm{FS}'_E$  is naturally right adjoint (hence quasi-inverse) to  $\check{\mathrm{FS}}_{E^\vee}'$  after choosing isomorphisms  $p^! \cong p^*[2 \operatorname{rank} E]$  and  $p'^! \cong p'^*[2 \operatorname{rank} E]$  induced by the orientations of  $E$  and  $E^\vee$  coming from the complex structures. Now consider the following diagram:

$$\begin{array}{ccc} \check{\mathrm{FS}}_{E^\vee} \circ \mathrm{FS}_E & \xleftarrow{\sim} & \mathrm{id} \\ \wr \downarrow & & \parallel \\ \check{\mathrm{FS}}_{E^\vee}' \circ \mathrm{FS}'_E & \xrightarrow{\sim} & \mathrm{id} \end{array}$$

Since we always equip complex vector bundles with the orientations induced from the complex structures, we see that the above diagram commutes up to the sign  $(-1)^{\operatorname{rank} E}$  using [Kashiwara and Schapira 1990, Remark 3.7.11]. This implies that the following diagram also commutes up to the sign  $(-1)^{\operatorname{rank} E}$ :

$$(2-38) \quad \begin{array}{ccc} \check{\mathrm{FS}}_{E^\vee} \circ \mathrm{FS}_E & \xleftarrow{\sim} & \mathrm{id} \\ \wr \downarrow & & \parallel \\ (\mathrm{FS}_{E^\vee}^a[2 \operatorname{rank} E]) \circ (\check{\mathrm{FS}}_E^a[-2 \operatorname{rank} E]) & \xrightarrow{\sim} & \mathrm{id} \end{array}$$

For  $\mathcal{F} \in D_{\mathbb{R}+}^+(E)$ , we define an isomorphism

$$(2-39) \quad \eta(\mathcal{F}): \mathrm{FS}_{E^\vee} \mathrm{FS}_E(\mathcal{F}) \cong \mathcal{F}^a[-2 \operatorname{rank} E]$$

by the composition

$$\mathrm{FS}_{E^\vee} \mathrm{FS}_E(\mathcal{F}) \cong \mathrm{FS}_{E^\vee} \check{\mathrm{FS}}_E^a(\mathcal{F})[-2 \operatorname{rank} E] \cong \mathcal{F}^a[-2 \operatorname{rank} E].$$

Note that this map differs from the composition

$$\mathrm{FS}_{E^\vee} \mathrm{FS}_E(\mathcal{F}) \cong \check{\mathrm{FS}}_{E^\vee}^a \mathrm{FS}_E(\mathcal{F})[-2 \operatorname{rank} E] \cong \mathcal{F}^a[-2 \operatorname{rank} E]$$

by the sign  $(-1)^{\text{rank } E}$ . This implies that the following diagram commutes up to the sign  $(-1)^{\text{rank } E}$ :

$$(2-40) \quad \begin{array}{ccc} \text{FS}_E \text{FS}_{E^\vee} \text{FS}_E(\mathcal{F}) & \xrightarrow{\text{FS}_E(\eta(\mathcal{F}))} & \text{FS}_E(\mathcal{F}^a)[-2 \text{rank } E] \\ \eta(\text{FS}_E(\mathcal{F})) \downarrow & \nearrow & \\ \text{FS}_E(\mathcal{F}^a)[-2 \text{rank } E] & & \end{array}$$

Now we discuss base change maps for Fourier–Sato transforms. Let  $f: Z' \rightarrow Z$  be a continuous map between topological spaces and  $E$  be a complex vector bundle on  $Z$ . We let  $E_{Z'}$  denote the base change of  $E$  to  $Z'$ , and  $f_E: E_{Z'} \rightarrow E$  and  $f_{E^\vee}: E_{Z'}^\vee \rightarrow E^\vee$  denote the base changes of  $f$ . Using base change maps (2-2)–(2-7), we can construct natural maps

$$(2-41) \quad \text{FS}_E \circ f_{E*} \rightarrow f_{E^\vee*} \circ \text{FS}_{E_{Z'}},$$

$$(2-42) \quad f_{E^\vee}^* \circ \text{FS}_E \rightarrow \text{FS}_{E_{Z'}} \circ f_E^*,$$

$$(2-43) \quad f_{E^\vee!} \circ \text{FS}_{E_{Z'}} \rightarrow \text{FS}_E \circ f_{E!},$$

$$(2-44) \quad \text{FS}_{E_{Z'}} \circ f_E^! \rightarrow f_{E^\vee}^! \circ \text{FS}_E.$$

It is shown in [Kashiwara and Schapira 1990, Proposition 3.7.13] that all these maps are invertible. We call these maps the base change maps for the Fourier–Sato transform. We have similar isomorphisms for  $\text{FS}'$ ,  $\tilde{\text{FS}}$ , and  $\tilde{\text{FS}}'$ , and these isomorphisms commute with the maps in (2-37).

Now let  $Z$  be a topological space,  $E_1$  and  $E_2$  be a complex vector bundle over  $Z$ , and  $g: E_1 \rightarrow E_2$  be a morphism of vector bundles. Denote by  ${}^t g: E_2^\vee \rightarrow E_1^\vee$  the dual map of  $g$ . We define

$$\dim g := \text{rank } E_1 - \text{rank } E_2.$$

We have an isomorphism

$$(2-45) \quad g^! \mathbb{Q}_{E_2} \cong \mathbb{Q}_{E_1}[2 \dim g]$$

defined by the composition

$$g^! \mathbb{Q}_{E_2} \cong g^! \pi_{E_2}^! \mathbb{Q}_Z[-2 \text{rank } E_2] \cong \pi_{E_1}^! \mathbb{Q}_Z[-2 \text{rank } E_2] \cong \mathbb{Q}_{E_1}[2 \dim g],$$

where the first and third isomorphisms are defined by the orientations on  $E_1$  and  $E_2$  induced from the complex structures. For  $\mathcal{F} \in D_{\mathbb{R}^+}^+(E_1)$  and  $\mathcal{G} \in D_{\mathbb{R}^+}^+(E_2)$  we will recall the construction of following four isomorphisms defined in [Kashiwara and Schapira 1990, Proposition 3.7.14]:

$$(2-46) \quad \text{FS}_{E_2}(g_! \mathcal{F}) \cong {}^t g^* \text{FS}_{E_1}(\mathcal{F}),$$

$$(2-47) \quad \text{FS}_{E_2}(g_* \mathcal{F}) \cong {}^t g^! \text{FS}_{E_1}(\mathcal{F})[2 \dim g],$$

$$(2-48) \quad \text{FS}_{E_1}(g^! \mathcal{G}) \cong {}^t g_* \text{FS}_{E_2}(\mathcal{G}),$$

$$(2-49) \quad \text{FS}_{E_1}(g^* \mathcal{G}) \cong {}^t g! \text{FS}_{E_2}(\mathcal{G})[-2 \dim g].$$

Consider the following commutative diagram:

$$(2-50) \quad \begin{array}{ccccc} & & E_1^\vee & \xleftarrow{\iota g} & E_2^\vee \\ & p'_1 \uparrow & & & \uparrow \tilde{p}'_1 \\ E_1 \oplus E_1^\vee & \xleftarrow{\iota \tilde{g}} & E_1 \oplus E_2^\vee & \xrightarrow{\tilde{g}} & E_2 \oplus E_2^\vee \\ & \searrow p_1 & \downarrow \tilde{p}_2 & & \downarrow p_2 \\ & & E_1 & \xrightarrow{g} & E_2 \end{array}$$

Here the upper left and lower right squares are Cartesian. Now define closed subsets  $P_1, P'_1 \subset E_1 \oplus E_1^\vee$  and  $P_2, P'_2 \subset E_2 \oplus E_2^\vee$  as before, and closed subsets  $\tilde{P}, \tilde{P}' \subset E_1 \oplus E_2^\vee$  by

$$\tilde{P} := \{(v, w) \in E_1 \oplus E_2^\vee \mid \operatorname{Re} w(g(v)) \geq 0\}, \quad \tilde{P}' := \{(v, w) \in E_1 \oplus E_2^\vee \mid \operatorname{Re} w(g(v)) \leq 0\}.$$

We let  $\iota_{P_1} : P_1 \hookrightarrow E_1 \oplus E_1^\vee$  denote the inclusion map and define the maps  $\iota_{P'_1}, \iota_{P_2}, \iota_{P'_2}, \iota_{\tilde{P}},$  and  $\iota_{\tilde{P}'}$  in the same way. The isomorphism (2-46) is defined by the composition

$$\begin{aligned} \operatorname{FS}_{E_2}(g! \mathcal{F}) &= p'_2! \iota_{P'_2} \iota_{P_2}^* p_2^* g! \mathcal{F} \cong p'_2! \iota_{P'_2} \iota_{P_2}^* \tilde{g}! \tilde{p}_2^* \mathcal{F} \\ &\cong p'_2! \tilde{g}! \iota_{\tilde{P}'} \iota_{\tilde{P}}^* \tilde{p}_2^* \mathcal{F} \\ &\cong \tilde{p}'_1! \iota_{\tilde{P}'} \iota_{\tilde{P}}^* \tilde{g}^* p_1^* \mathcal{F} \\ &\cong \tilde{p}'_1! \tilde{g}^* \iota_{P'_1} \iota_{P_1}^* p_1^* \mathcal{F} \cong {}^t g^* p'_1! \iota_{P'_1} \iota_{P_1}^* p_1^* \mathcal{F} = {}^t g^* \operatorname{FS}_{E_1}(\mathcal{F}). \end{aligned}$$

Here all isomorphisms are defined by using the base change maps. The isomorphism (2-47) is defined by the composition

$$\begin{aligned} \operatorname{FS}_{E_2}(g_* \mathcal{F}) &\cong \operatorname{FS}'_{E_2}(g_* \mathcal{F}) = p'_2! \iota_{P'_2} \iota_{P_2}^! p_2^* g_* \mathcal{F} \\ &\cong p'_2! \iota_{P'_2} \iota_{P_2}^! \tilde{g}_* \tilde{p}_2^* \mathcal{F} \\ &\cong p'_2! \tilde{g}_* \iota_{\tilde{P}'} \iota_{\tilde{P}}^! \tilde{p}_2^* \mathcal{F} \\ &\cong \tilde{p}'_1! \iota_{\tilde{P}'} \iota_{\tilde{P}}^! \tilde{g}^* p_1^* \mathcal{F} \\ &\cong \tilde{p}'_1! \tilde{g}^! \iota_{P'_1} \iota_{P_1}^! p_1^* \mathcal{F}[2 \dim g] \\ &\cong {}^t g^! p'_1! \iota_{P'_1} \iota_{P_1}^! p_1^* \mathcal{F}[2 \dim g] \\ &\cong {}^t g^! p'_1! \iota_{P'_1} \iota_{P_1}^! p_1^* \mathcal{F}[2 \dim g] \\ &= {}^t g^! \operatorname{FS}'_{E_1}(\mathcal{F})[2 \dim g] \cong {}^t g^! \operatorname{FS}_{E_1}(\mathcal{F})[2 \dim g]. \end{aligned}$$

Here the first and final isomorphisms are defined in (2-37), the fifth isomorphism is defined by using (2-45) and (2-12) and other isomorphisms are defined using the base change maps. The isomorphism (2-48) is defined by the composition

$$\begin{aligned} \operatorname{FS}_{E_1}(g^! \mathcal{G}) &\cong \operatorname{FS}_{E_1} g^! \operatorname{FS}_{E_2^\vee} \operatorname{FS}_{E_2}(\mathcal{G}^a)[2 \operatorname{rank} E_2] \\ &\cong \operatorname{FS}_{E_1} \operatorname{FS}_{E_1^\vee} {}^t g_* \operatorname{FS}_{E_2}(\mathcal{G}^a)[2 \operatorname{rank} E_1] \cong {}^t g_* \operatorname{FS}_{E_2}(\mathcal{G}), \end{aligned}$$

where the first and third isomorphisms are defined using  $\eta(-)$  in (2-39) and the second isomorphism is defined using (2-47). Similarly, the isomorphism (2-49) is defined by the composition

$$\begin{aligned} \mathrm{FS}_{E_1}(g^*\mathcal{G}) &\cong \mathrm{FS}_{E_1} g^* \mathrm{FS}_{E_2^\vee} \mathrm{FS}_{E_2}(\mathcal{G}^a)[2\mathrm{rank} E_2] \\ &\cong \mathrm{FS}_{E_1} \mathrm{FS}_{E_1^\vee} {}^t g_! \mathrm{FS}_{E_2}(\mathcal{G}^a)[2\mathrm{rank} E_2] \cong {}^t g_! \mathrm{FS}_{E_2}(\mathcal{G})[-2\dim g]. \end{aligned}$$

Assume further that we are given a continuous map  $f: Z' \rightarrow Z$ . Denote by  $g_{Z'}: (E_1)_{Z'} \rightarrow (E_2)_{Z'}$  the base change of  $g$ . Then for each choice of  $a \in \{(-)^*, (-)_*, (-)!, (-)_!\}$ , one can see that the isomorphisms (2-46)–(2-49) commute with base change maps for  $a$ . For example if we choose  $a = (-)^*$  and (2-48), we see that the following diagram commutes:

$$(2-51) \quad \begin{array}{ccc} f_{E_1^\vee}^* \mathrm{FS}_{E_1}(g^! \mathcal{G}) & \xrightarrow{\sim} & f_{E_1^\vee}^* {}^t g_* \mathrm{FS}_{E_2}(\mathcal{G}) \\ \wr \downarrow & & \downarrow \\ \mathrm{FS}_{(E_1)_Z}(f_{E_1}^* g^! \mathcal{G}) & & {}^t g_{Z'} * f_{E_2^\vee}^* \mathrm{FS}_{E_2}(\mathcal{G}) \\ \downarrow & & \wr \downarrow \\ \mathrm{FS}_{(E_1)_Z}(g_{Z'}^! f_{E_2}^* \mathcal{G}) & \xrightarrow{\sim} & {}^t g_{Z'} * \mathrm{FS}_{(E_2)_Z}(f_{E_2}^* \mathcal{G}) \end{array}$$

We can show that the maps (2-46) and (2-47) are associative in the following sense. Let  $E_3$  be another vector bundle over  $Z$  and  $h: E_2 \rightarrow E_3$  be morphism of vector bundles. For  $\mathcal{F} \in D_{\mathbb{R}^+}^+(E_1)$ , the following two diagrams commute:

$$(2-52) \quad \begin{array}{ccc} \mathrm{FS}_{E_3}((h \circ g)_! \mathcal{F}) & \xrightarrow{\sim} & \mathrm{FS}_{E_3}(h_! \circ g_! \mathcal{F}) \\ \wr \downarrow & & \downarrow \wr \\ & & ({}^t h)^* \mathrm{FS}_{E_2}(g_! \mathcal{F}) \\ & & \downarrow \wr \\ ({}^t g \circ {}^t h)^* \mathrm{FS}_{E_1}(\mathcal{F}) & \xrightarrow{\sim} & ({}^t h)^* ({}^t g)^* \mathrm{FS}_{E_1}(\mathcal{F}) \end{array}$$

$$(2-53) \quad \begin{array}{ccc} \mathrm{FS}_{E_3}((h \circ g)_* \mathcal{F}) & \xrightarrow{\sim} & \mathrm{FS}_{E_3}(h_* \circ g_* \mathcal{F}) \\ \wr \downarrow & & \downarrow \wr \\ & & ({}^t h)^! \mathrm{FS}_{E_2}(g_* \mathcal{F})[2\dim h] \\ & & \downarrow \wr \\ ({}^t g \circ {}^t h)^! \mathrm{FS}_{E_1}(\mathcal{F})[2\dim g + 2\dim h] & \xrightarrow{\sim} & ({}^t h)^! ({}^t g)^! \mathrm{FS}_{E_1}(\mathcal{F})[2\dim g + 2\dim h]. \end{array}$$

We omit the proof as it is a routine argument in sheaf theory. We have similar statements for maps (2-48) and (2-49).

Now we discuss the compatibility between Fourier–Sato transforms and natural transformations discussed in Section 2.2.

**Proposition 2.2** *The following diagram commutes:*

$$\begin{array}{ccc} \mathrm{FS}_{E_2}(g_! \mathcal{F}) & \longrightarrow & \mathrm{FS}_{E_2}(g_* \mathcal{F}) \\ \downarrow \wr & & \downarrow \wr \\ {}^t g^* \mathrm{FS}_{E_1}(\mathcal{F}) & \longrightarrow & {}^t g^! \mathrm{FS}_{E_1}(\mathcal{F})[-2 \dim {}^t g] \end{array}$$

Here the upper horizontal arrow is the map (2-10) and the lower horizontal arrow is defined by using (2-45) and (2-12).

The proof of this proposition will be given in Section 7.2.

**Proposition 2.3** *Consider the following diagrams:*

$$\begin{array}{ccc} \mathrm{FS}_{E_2}(\mathcal{G}) & \longrightarrow & \mathrm{FS}_{E_2}(g_* g^* \mathcal{G}) \\ & \searrow & \downarrow \wr \\ & & {}^t g^! \mathrm{FS}_{E_1}(g^* \mathcal{G})[2 \dim g] \\ & & \downarrow \wr \\ & & {}^t g^! {}^t g^! \mathrm{FS}_{E_2}(\mathcal{G}) \end{array} \quad \begin{array}{ccc} \mathrm{FS}_{E_2}(g_! g^! \mathcal{G}) & \longrightarrow & \mathrm{FS}_{E_2}(\mathcal{G}) \\ \downarrow \wr & & \downarrow \wr \\ {}^t g^* \mathrm{FS}_{E_1}(g^! \mathcal{G}) & & \\ \downarrow \wr & & \nearrow \\ {}^t g^* {}^t g^* \mathrm{FS}_{E_2}(\mathcal{G}) & & \end{array}$$

Then the left diagram commutes up to the sign  $(-1)^{\mathrm{rank} E_1}$  and the right diagram commutes up to the sign  $(-1)^{\mathrm{rank} E_2}$ .

The proof of this proposition will be given in Section 7.3.

### 3 Specialization functor and specialization map

Let  $X$  be a separated scheme of finite type over  $\mathrm{Spec} \mathbb{C}$  and  $Z$  be a closed subscheme of  $X$ . We let  $C_{Z/X}$  denote the normal cone of  $Z$  in  $X$ . In this section, we introduce the specialization functor

$$\mathrm{Sp}_{Z/X}: D^+(X) \rightarrow D^+(C_{Z/X})$$

following Verdier [1983], and define the specialization map for the Borel–Moore homology groups

$$\mathrm{Sp}_{Z/X}^{\mathrm{BM}}: H_*^{\mathrm{BM}}(X) \rightarrow H_*^{\mathrm{BM}}(C_{Z/X})$$

using the specialization functor. Then we prove that the specialization map  $\mathrm{Sp}_{Z/X}^{\mathrm{BM}}$  is compatible with the specialization map of Chow groups defined by Fulton [1984, Section 5].

### 3.1 Specialization functors

Let us recall the definition and basic properties of the specialization functor introduced by Verdier [1983]. Let  $X$  be a separated scheme of finite type over  $\text{Spec } \mathbb{C}$ , and  $Z$  be a closed subscheme. Define schemes  $M_{Z/X}$  and  $M_{Z/X}^0$  by

$$M_{Z/X} := \text{Bl}_{Z \times \{0\}}(X \times \mathbb{A}^1), \quad M_{Z/X}^0 := M_{Z/X} \setminus \text{Bl}_Z(X).$$

Then we have the following commutative diagram:

$$(3-1) \quad \begin{array}{ccccccc} Z & \hookrightarrow & Z \times \mathbb{A}^1 & \hookrightarrow & Z \times (\mathbb{A}^1 \setminus \{0\}) & & \\ \downarrow & & \downarrow i_{Z \times \mathbb{A}^1} & & \downarrow & & \\ C_{Z/X} & \xrightarrow{i_{C_{Z/X}}} & M_{Z/X}^0 & \xrightarrow{j} & X \times (\mathbb{A}^1 \setminus \{0\}) & \xrightarrow{\text{pr}_1} & X \\ \downarrow & & \downarrow p & & \downarrow & & \\ \{0\} & \hookrightarrow & \mathbb{A}^1 & \hookrightarrow & (\mathbb{A}^1 \setminus \{0\}) & & \end{array}$$

**Definition 3.1** [Verdier 1983, Section 8] The specialization functor

$$\text{Sp}_{Z/X}: D^+(X) \rightarrow D_{\mathbb{R}^+}^+(C_{Z/X})$$

is defined by the formula

$$\text{Sp}_{Z/X}(\mathcal{F}) := \psi_p j_! \text{pr}_1^*(\mathcal{F}).$$

Here  $\mathbb{R}^+$ -action on  $C_{Z/X}$  is defined by the scaling of fibers.

The following are basic properties of the specialization functor we use in this paper.

**Proposition 3.2** [Verdier 1983, Section 9] Let  $X$  be a separated scheme of finite type over  $\text{Spec } \mathbb{C}$  and  $Z \subset X$  be a closed subscheme. For a constructible complex  $\mathcal{F} \in D_c^b(X)$ , we have isomorphisms

$$(3-2) \quad \text{Sp}_{Z/X}(\mathcal{F})|_Z \cong \mathcal{F}|_Z,$$

$$(3-3) \quad \text{Sp}_{Z/X}(\mathcal{F})|_Z^! \cong \mathcal{F}|_Z^!.$$

**Proof** First note that we have natural morphisms

$$\text{Sp}_{Z/X}(\mathcal{F})|_Z \rightarrow \text{Sp}_{Z/X}(\mathcal{F}_Z)|_Z, \quad \text{Sp}_{Z/X}(\Gamma_Z(\mathcal{F}))|_Z^! \rightarrow \text{Sp}_{Z/X}(\mathcal{F})|_Z^!.$$

We claim that these maps are isomorphisms. To do this, it is enough to prove that for complexes  $\mathcal{F} \in D_c^b(X)$  with  $\mathcal{F}|_Z = 0$  we have  $\text{Sp}_{Z/X}(\mathcal{F})|_Z = 0$ , and for  $\mathcal{F} \in D_c^b(X)$  with  $\mathcal{F}|_Z^! = 0$  we have  $\text{Sp}_{Z/X}(\mathcal{F})|_Z^! = 0$ . By shrinking  $X$  and embedding it into an affine space so that  $Z$  is cut out by linear equations, we may find the Cartesian diagram

$$\begin{array}{ccc} Z & \hookrightarrow & X \\ \downarrow & & \downarrow \\ \mathbb{A}^n & \hookrightarrow & \mathbb{A}^{n+r} \end{array}$$

for some  $n, r \in \mathbb{Z}_{>0}$ . In particular, we may assume that  $Z$  and  $X$  are smooth. Then these claims follow from [Kashiwara and Schapira 1990, Theorem 4.2.3].

Now consider the composition

$$(3-4) \quad \mathrm{Sp}_{Z/X}(\mathcal{F}_Z)|_Z \rightarrow \psi_{\mathrm{poi}_{Z \times \mathbb{A}^1}}((Z \times \mathbb{A}^1 \rightarrow Z)^*(\mathcal{F}|_Z)) \rightarrow \mathcal{F}|_Z$$

where the first map is (2-16) and the second map is (2-27). It is clear that these maps are isomorphisms; hence we obtain the isomorphism (3-2). Similarly, we have isomorphisms

$$(3-5) \quad \mathrm{Sp}_{Z/X}(\Gamma_Z(\mathcal{F}))|_Z^! \cong \psi_{\mathrm{poi}_{Z \times \mathbb{A}^1}}((Z \times \mathbb{A}^1 \rightarrow Z)^!(\mathcal{F}|_Z^![-2])) \cong \mathcal{F}|_Z^!$$

which imply the isomorphism (3-3).  $\square$

### 3.2 Specialization maps

Consider the natural map defined in (2-13),

$$\mathrm{Sp}_{Z/X}(\mathbb{Q}_X)|_Z \otimes \omega_{Z/C_{Z/X}} \rightarrow \mathrm{Sp}_{Z/X}(\mathbb{Q}_X)|_Z^!.$$

Using the proposition above, this map is identified with a map

$$(3-6) \quad \omega_{Z/C_{Z/X}} \rightarrow \omega_{Z/X}.$$

The isomorphism (2-35) implies  $\omega_{Z/C_{Z/X}} \cong (C_{Z/X} \rightarrow Z)_! \mathbb{Q}_{C_{Z/X}}$ ; hence by adjunction we obtain an element

$$e_{Z/X} \in H^0(C_{Z/X} \rightarrow X).$$

Define a map

$$\mathrm{Sp}_{Z/X}^{\mathrm{BM}}: H_*^{\mathrm{BM}}(X) \rightarrow H_*^{\mathrm{BM}}(C_{Z/X})$$

by composing  $e_{Z/X}$  with elements in  $H_*^{\mathrm{BM}}(X) = H^{-*}(X \rightarrow \mathrm{Spec} \mathbb{C})$ . Now we compare this specialization map for Borel–Moore homology groups and the specialization map for Chow groups

$$\mathrm{Sp}_{Z/X}^{\mathrm{Chow}}: A_*(X) \rightarrow A_*(C_{Z/X})$$

defined by Fulton [1984, Section 5].

**Theorem 3.3** *The diagram*

$$\begin{array}{ccc} A_*(X) & \xrightarrow{\mathrm{Sp}_{Z/X}^{\mathrm{Chow}}} & A_*(C_{Z/X}) \\ \downarrow \mathrm{cl}_X & & \downarrow \mathrm{cl}_{C_{Z/X}} \\ H_*^{\mathrm{BM}}(X) & \xrightarrow{\mathrm{Sp}_{Z/X}^{\mathrm{BM}}} & H_*^{\mathrm{BM}}(C_{Z/X}) \end{array}$$

*commutes, where  $\mathrm{cl}_X$  and  $\mathrm{cl}_{C_{Z/X}}$  are cycle maps.*

This theorem will be proved in two steps: first we construct the Gysin pullback for Borel–Moore homology to a principal divisor using the nearby cycle functor and compare it with the Gysin pullback for Chow groups (Proposition 3.5), and then reduce the theorem to this statement.

Let  $X$  be a separated scheme of finite type over  $\text{Spec } \mathbb{C}$  and  $u: X \rightarrow \mathbb{A}^1$  be a regular function. Write  $D := u^{-1}(0)$  and we let  $i: D \hookrightarrow X$  denote the inclusion map. The composition of morphisms in (2-25) and (2-26) defines the map

$$(D \hookrightarrow X)^* \mathbb{Q}_X \rightarrow \psi_u(\mathbb{Q}_X) \rightarrow (D \hookrightarrow X)^! \mathbb{Q}_X[2],$$

hence an element  $e_u \in H^2(D \rightarrow X)$ . The element  $e_u$  defines a map

$$\psi i^!: H_*^{\text{BM}}(X) \rightarrow H_{*-2}^{\text{BM}}(D).$$

Now let  $f: X' \rightarrow X$  be a proper map and write  $D' := (u \circ f)^{-1}(0)$ . We let  $i': D' \hookrightarrow X'$  denote the inclusion map.

**Lemma 3.4** *The following diagram commutes:*

$$(3-7) \quad \begin{array}{ccc} H_*^{\text{BM}}(X') & \xrightarrow{\psi i'^!} & H_{*-2}^{\text{BM}}(D') \\ \downarrow f_* & & \downarrow (f|_{D'})_* \\ H_*^{\text{BM}}(X) & \xrightarrow{\psi i^!} & H_{*-2}^{\text{BM}}(D) \end{array}$$

**Proof** Consider the following diagram in  $D^b(D)$ :

$$\begin{array}{ccccccc} \mathbb{Q}_D & \xrightarrow{\sim} & i^* \mathbb{Q}_X & \xrightarrow{\quad} & \psi_u(\mathbb{Q}_X) & \xrightarrow{\quad} & i^! \mathbb{Q}_X[2] \\ \downarrow & & \downarrow & \text{(B)} & \downarrow & \text{(C)} & \downarrow \\ & & i^* f_* \mathbb{Q}_{X'} & \xrightarrow{\quad} & \psi_u(f_* \mathbb{Q}_{X'}) & \xrightarrow{\quad} & i^! f_* \mathbb{Q}_{X'}[2] \\ & & \downarrow \wr & \text{(D)} & \downarrow \wr & \text{(E)} & \downarrow \wr \\ (f|_{D'})_* \mathbb{Q}_{D'} & \xrightarrow{\sim} & (f|_{D'})_* i'^* \mathbb{Q}_{X'} & \longrightarrow & (f|_{D'})_* \psi_{u \circ f}(\mathbb{Q}_{X'}) & \longrightarrow & (f|_{D'})_* i'^! \mathbb{Q}_{X'}[2] \end{array}$$

The commutativity of diagrams (A), (B), and (C) is obvious. The commutativity of the diagrams (D) and (E) follows from the fact that the maps in (2-25) and (2-26) commute with the base change maps. By taking the Verdier-dual of the outer square of the above diagram, we obtain the desired claim.  $\square$

**Proposition 3.5** *Assume that  $D \subset X$  is a Cartier divisor (ie  $f^{\text{red}}$  is not identically zero over any irreducible component of  $X$ ). Then the diagram*

$$\begin{array}{ccc} A_*(X) & \xrightarrow{i^!} & A_*(D) \\ \downarrow \text{cl}_X & & \downarrow \text{cl}_D \\ H_*^{\text{BM}}(X) & \xrightarrow{\psi i^!} & H_*^{\text{BM}}(D) \end{array}$$

*commutes, where  $i^!$  is the usual Gysin pullback for Chow groups.*



**Proof** We need to prove

$$\mathrm{cl}_D \circ i^!([V]) = \psi i^!([V]_{\mathrm{BM}})$$

for each irreducible subvariety  $V \subset X$ . Firstly consider the case when  $V$  is included in  $D$ . Then  $i^!([V]) = 0$  since  $D$  is linearly equivalent to 0, and the commutativity of (3-7) implies  $\psi i^!([V]_{\mathrm{BM}}) = 0$ .

Now we may assume  $V$  and  $D$  intersect properly. Replacing  $X$  by  $V$  using the commutativity of (3-7), we may assume  $X = V$ . Further replacing  $X$  by its normalization and using the commutativity of (3-7) again and shrinking  $X$  if necessary, we may assume  $X$  is smooth and  $D$  is irreducible. In this case  $i^!([V]) = d[D^{\mathrm{red}}]$  where  $d$  is the multiplicity of  $D$  in  $X$ . Now we want to compute

$$\psi i^!([X]_{\mathrm{BM}}) \in H_{2\dim X - 2}^{\mathrm{BM}}(D) = \mathbb{Z} \cdot [D^{\mathrm{red}}]_{\mathrm{BM}}.$$

Since this class can be computed analytically locally, we may replace  $X$  by  $\mathbb{A}^n$  and  $u$  by  $z_1^d$  where  $z_1$  is the first projection. Using the commutativity of (3-7) for the map  $f = (z_1^d, \mathrm{id}_{\mathbb{A}^{n-1}})$ , we may assume  $d = 1$ . In this case, the claim follows from Proposition 2.1.  $\square$

**Proof of Theorem 3.3** We use notation from the diagram (3-1). We need to prove

$$\mathrm{cl}_{C_{Z/X}}(\mathrm{Sp}_{Z/X}^{\mathrm{Chow}}([V])) = \mathrm{Sp}_{Z/X}^{\mathrm{BM}}([V]_{\mathrm{BM}})$$

for each irreducible subvariety  $V \subset X$  of dimension  $d$ . We let  $\tilde{V} \subset M_{Z/X}^o$  denote the strict transform of  $V \times \mathbb{A}^1$ . Proposition 3.5 claims the equality

$$\mathrm{cl}_{C_{Z/X}}(\mathrm{Sp}_{Z/X}^{\mathrm{Chow}}([V])) = (\psi i_{C_{Z/X}})^!([\tilde{V}]_{\mathrm{BM}});$$

hence what we need to prove is the equality

$$\mathrm{Sp}_{Z/X}^{\mathrm{BM}}([V]_{\mathrm{BM}}) = (\psi i_{C_{Z/X}})^!([\tilde{V}]_{\mathrm{BM}}).$$

Consider the following diagram:

$$(3-8) \quad \begin{array}{ccccccc} \mathbb{Q}_{C_{Z/X}} & \longrightarrow & \psi_p(\mathbb{Q}_{M_{Z/X}^o}) & \xrightarrow{\psi_p(-\cap[\tilde{V}]_{\mathrm{BM}})} & \psi_p(\omega_{M_{Z/X}^o})[-2d-2] & \longrightarrow & \omega_{C_{Z/X}}[-2d] \\ \parallel & & \uparrow \wr & & \uparrow \wr & & \parallel \\ \mathbb{Q}_{C_{Z/X}} & \longrightarrow & \mathrm{Sp}_{Z/X}(\mathbb{Q}_X) & \xrightarrow{\mathrm{Sp}_{Z/X}(-\cap[V]_{\mathrm{BM}})} & \mathrm{Sp}_{Z/X}(\omega_X)[-2d] & \longrightarrow & \omega_{C_{Z/X}}[-2d] \end{array}$$

Here the middle left and middle right vertical arrows are defined by isomorphisms  $\mathrm{pr}_1^* \mathbb{Q}_X \cong \mathbb{Q}_{X \times (\mathbb{A}^1 \setminus \{0\})}$  and  $\mathrm{pr}_1^* \omega_X \cong \omega_{X \times (\mathbb{A}^1 \setminus \{0\})}[-2]$  respectively; the upper left and upper right horizontal arrows are defined in (2-25) and (2-26) respectively; and the lower left and lower right horizontal arrows are defined so that the left and right squares commute. The commutativity of the middle diagram follows from the equality

$$\mathrm{pr}_1^*[V] = [\tilde{V} \cap (X \times (\mathbb{A}^1 \setminus \{0\}))].$$

The composition of upper horizontal arrows corresponds to the element

$$(\psi i_{C_{Z/X}})^!([\tilde{V}]_{\mathrm{BM}}) \in H_{2d}^{\mathrm{BM}}(C_{Z/X})$$

by definition. Therefore we need to prove that the composition of lower horizontal arrows corresponds to the element  $\mathrm{Sp}_{Z/X}^{\mathrm{BM}}([V]_{\mathrm{BM}})$ . By the definition of  $\mathrm{Sp}_{Z/X}^{\mathrm{BM}}$ , this statement is equivalent to the commutativity of the outer rectangle of the diagram

$$\begin{array}{ccccccc}
 \omega_{Z/C_{Z/X}} & \longrightarrow & \mathrm{Sp}_{Z/X}(\mathbb{Q}_X)|_Z^! & \longrightarrow & \mathrm{Sp}_{Z/X}(\omega_X)[-2d]|_Z^! & \xrightarrow{\sim} & \omega_Z[-2d] \\
 \parallel & & \nwarrow (3-3) & & & & \parallel \\
 \omega_{Z/C_{Z/X}} & \xrightarrow{(3-6)} & \omega_{Z/X} & \xrightarrow{(Z \hookrightarrow X)^!(-\cap[V])} & \omega_Z[-2d] & & 
 \end{array}$$

where the upper horizontal arrows are obtained by applying the functor  $(Z \hookrightarrow C_{Z/X})^!$  to the maps in the lower horizontal arrows of the diagram (3-8). The commutativity of the right quadrilateral is obvious. Note that the composition of (3-3) and (3-6) is obtained by applying the functor  $(Z \hookrightarrow C_{Z/X})^!$  to the composition

$$\mathbb{Q}_{C_{Z/X}} \otimes (Z \hookrightarrow C_{Z/X})_* \omega_{Z/C_{Z/X}} \xrightarrow{(3-2)} \mathrm{Sp}_{Z/X}(\mathbb{Q}_X) \otimes (Z \hookrightarrow C_{Z/X})_* \omega_{Z/C_{Z/X}} \rightarrow \mathrm{Sp}_{Z/X}(\mathbb{Q}_X)$$

where the second map is the  $!$ -counit map. Then the commutativity of the left quadrilateral follows from the construction of the map (3-2).  $\square$

## 4 Fourier–Sato transform of the specialization

The aim of this section is to prove the following statement:

**Theorem 4.1** *Let  $X$  be a separated scheme of finite type over  $\mathrm{Spec} \mathbb{C}$ ,  $E$  be a vector bundle on  $X$ , and  $s \in \Gamma(X, E)$  be a section. Write  $Z = s^{-1}(0)$  and  $E_Z = E|_Z$ . Denote by  $\pi_{E^\vee} : E^\vee \rightarrow X$  the projection and  $\bar{s} : E^\vee \rightarrow \mathbb{A}^1$  the regular function corresponding to  $s$ , ie the composite  $E^\vee \xrightarrow{(s \circ \pi_{E^\vee}, \mathrm{id})} E \times_X E^\vee \xrightarrow{\text{pairing}} \mathbb{A}^1$ . We let  $\iota_{C_{Z/X}} : C_{Z/X} \hookrightarrow E_Z$  denote the natural inclusion from the normal cone. Then for a constructible complex  $\mathcal{F} \in D_c^b(X)$ , we have an isomorphism*

$$(4-1) \quad \mathrm{FS}_{E_Z^\vee}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) \cong \iota_{C_{Z/X},*} \mathrm{Sp}_{Z/X}(\mathcal{F}).$$

As a consequence, we will give a new proof of Davison’s dimensional reduction theorem [2017, Theorem A.1] and prove our main result for local models for quasismooth derived schemes.

### 4.1 Proof of Theorem 4.1

Firstly, we rewrite the complex  $\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}$  so that its Fourier–Sato transform can be easily computed. Let  $\ell_{\leq 0} \subset \mathbb{C}$  denote the closed half line consisting of the nonpositive real numbers. Consider the commutative diagram

$$\begin{array}{ccccc}
 E_Z^\vee & \xrightarrow{i_{E_Z^\vee}} & E^\vee & \xrightarrow{\hat{s}} & X \times \mathbb{A}^1 & \xleftarrow{i_{X \times \ell_{\leq 0}}} & X \times \ell_{\leq 0} \\
 \downarrow \pi_{E_Z^\vee} & & \downarrow \pi_{E^\vee} & \nearrow p_{\mathbb{A}^1} & & \nearrow p_{\ell_{\leq 0}} & \\
 Z & \xrightarrow{i_Z} & X & & & & 
 \end{array}$$

where we define  $\hat{s} := (\pi_{E^\vee}, \bar{s})$ ,  $i_{E_Z^\vee}$ ,  $i_Z$  and  $i_{X \times \ell_{\leq 0}}$  are natural inclusions, and  $\pi_{E_Z^\vee}$ ,  $\pi_{E^\vee}$ ,  $p_{\mathbb{A}^1}$  and  $p_{\ell_{\leq 0}}$  are natural projections. Consider the complex

$$\varphi_{\bar{s}}^{\ell_{\leq 0}}(\pi_{E^\vee}^! \mathcal{F}) := (\bar{s}^{-1}(0) \hookrightarrow \bar{s}^{-1}(\ell_{\leq 0}))^*(\bar{s}^{-1}(\ell_{\leq 0}) \hookrightarrow E^\vee)^! \pi_{E^\vee}^! \mathcal{F}.$$

As we have seen in (2-31), we have a natural isomorphism

$$\varphi_{\bar{s}}^{\ell_{\leq 0}}(\pi_{E^\vee}^! \mathcal{F}) \xrightarrow{\cong} \varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F}).$$

Therefore we have

$$(4-3) \quad \varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee} \cong (\hat{s}^! i_{X \times \ell_{\leq 0}, * p_{\ell_{\leq 0}}^! \mathcal{F})|_{E_Z^\vee}.$$

Applying the functor  $\text{FS}_{E_Z^\vee}$  to both sides, we obtain

$$(4-4) \quad \begin{aligned} \text{FS}_{E_Z^\vee}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) &\cong \text{FS}_{E_Z^\vee}((\hat{s}^! i_{X \times \ell_{\leq 0}, * p_{\ell_{\leq 0}}^! \mathcal{F})|_{E_Z^\vee}) \\ &\cong \text{FS}_{E^\vee}(\hat{s}^! i_{X \times \ell_{\leq 0}, * p_{\ell_{\leq 0}}^! \mathcal{F})|_{E_Z} \\ &\cong (\tilde{s}^* \text{FS}_{X \times \mathbb{A}^1}(i_{X \times \ell_{\leq 0}, * p_{\ell_{\leq 0}}^! \mathcal{F}))|_{E_Z} \end{aligned}$$

where  $\tilde{s}$  is the dual map of  $\hat{s}$ . The second and third isomorphism are defined in (2-42) and (2-48) respectively.

Consider the diagram

$$\begin{array}{ccccccc} E_Z & \xhookrightarrow{i_{E_Z}} & E & \xleftarrow{\tilde{s}} & X \times \mathbb{A}^1 & \xleftarrow{j_{X \times \mathbb{A}^1_{>0}}} & X \times \mathbb{A}^1_{>0} \\ \downarrow \pi_{E_Z} & & \downarrow \pi_E & \nearrow p_{\mathbb{A}^1} & & \nearrow p_{\mathbb{A}^1_{>0}} & \\ Z & \xhookrightarrow{i_Z} & X & & & & \end{array}$$

where  $\mathbb{A}^1_{>0} \subset \mathbb{A}^1$  is the open subset consisting of points whose real part is positive,  $i_{E_Z}$  and  $j_{X \times \mathbb{A}^1_{>0}}$  are natural inclusions and  $\pi_E$ ,  $\pi_{E_Z}$ ,  $p_{\mathbb{A}^1}$  and  $p_{\mathbb{A}^1_{>0}}$  are natural projections.

**Lemma 4.2** *There exists an isomorphism*

$$(4-5) \quad \text{FS}_{X \times \mathbb{A}^1}(i_{X \times \ell_{\leq 0}, * p_{\ell_{\leq 0}}^! \mathcal{F}) \cong j_{X \times \mathbb{A}^1_{>0}, * p_{\mathbb{A}^1_{>0}}^* \mathcal{F}.$$

**Proof** Let  $\mathcal{G}$  be the complex at the left-hand side of the claim. For a point  $\xi \in \mathbb{A}^1 \setminus \mathbb{A}^1_{>0}$ , we claim the vanishing  $\mathcal{G}|_{X \times \{\xi\}}^! = 0$ . Using the isomorphism (2-43), we may assume that  $X$  is a point which will be denoted by  $\text{pt}$  and  $\mathcal{F}$  is a constant sheaf  $\mathbb{Q}_{\text{pt}}$ . Now recall that we have seen in (2-37) that there exists an isomorphism of functors  $\text{FS}_{\mathbb{A}^1} \cong \text{FS}'_{\mathbb{A}^1}$  where the functor  $\text{FS}'_{\mathbb{A}^1}$  is defined in (2-36). We define closed subsets  $H_\xi, H'_\xi \subset \mathbb{A}^1$  by

$$H_\xi := \{t \in \mathbb{A}^1 \mid \text{Re}(t \cdot \xi) \geq 0\}, \quad H'_\xi := \{t \in \mathbb{A}^1 \mid \text{Re}(t \cdot \xi) \leq 0\}.$$

By the base change theorem, we have

$$\text{FS}'_{\mathbb{A}^1}(i_{\ell_{\leq 0}, * p_{\ell_{\leq 0}}^! \mathbb{Q}_{\text{pt}})|_\xi^! \cong R\Gamma(H_\xi, (i_{\ell_{\leq 0}, * p_{\ell_{\leq 0}}^! \mathbb{Q}_{\text{pt}})|_{H_\xi}^!)$$

where  $i_{\ell_{\leq 0}}: \ell_{\leq 0} \hookrightarrow \mathbb{A}^1$  is the natural inclusion. The assumption on  $\xi$  implies that  $H_\xi$  contains  $\ell_{\leq 0}$ , which implies the isomorphisms

$$R\Gamma(H_\xi, (i_{\ell_{\leq 0}, *p_{\ell_{\leq 0}}^! \mathbb{Q}_{\text{pt}})|_{H_\xi}^!) \cong R\Gamma(\ell_{\leq 0}, p_{\ell_{\leq 0}}^! \mathbb{Q}_{\text{pt}}) \simeq H_c^*(\ell_{\leq 0})^\vee = 0.$$

Therefore the unit map

$$(4-6) \quad \mathcal{G} \rightarrow j_{X \times \mathbb{A}_{>0}^1, *}(\mathcal{G}|_{X \times \mathbb{A}_{>0}^1})$$

is an isomorphism.

For  $\xi \in \mathbb{A}_{>0}^1$ , the half line  $\ell_{\leq 0}$  is contained in  $H'_\xi$ . This implies the isomorphism

$$\mathcal{G}|_{X \times \mathbb{A}_{>0}^1} \cong (\text{pr}_{13}^* \text{pr}_{12}^* i_{X \times \ell_{\leq 0}, *} p_{\ell_{\leq 0}}^! \mathcal{F})|_{X \times \mathbb{A}_{>0}^1}.$$

Here  $\text{pr}_{12}, \text{pr}_{13}: X \times \mathbb{A}^1 \times \mathbb{A}^1 \rightarrow X \times \mathbb{A}^1$  are the projections to the first and second, and the first and third factors respectively. By the base change theorem, we see that the right-hand side is isomorphic to  $p_{\mathbb{A}_{>0}^1}^* \mathcal{F}$ . This isomorphism and the isomorphism (4-6) imply the desired result.  $\square$

Combining (4-4) and Lemma 4.2, the complex at the left-hand side of (4-1) is isomorphic to the complex

$$(4-7) \quad (\tilde{s}_* j_{X \times \mathbb{A}_{>0}^1, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z}.$$

Consider the diagram

$$(4-8) \quad \begin{array}{ccccc} X & \xleftarrow{p_{\mathbb{A}_{>0}^1}} & X \times \mathbb{A}_{>0}^1 & \xrightarrow{(s \circ p_{\mathbb{A}_{>0}^1}) \cdot (\tau \circ \text{pr}_2)} & E \xleftarrow{i_{E_Z}} E_Z \\ & & \downarrow ((s \circ p_{\mathbb{A}_{>0}^1}) \cdot (\tau \circ \text{pr}_2), j_{\mathbb{A}_{>0}^1} \circ \text{pr}_2) & \nearrow 0_{E/E \times \mathbb{A}^1} & \\ & & E \times \mathbb{A}^1 & & \end{array}$$

where  $\tau: \mathbb{A}_{>0}^1 \xrightarrow{\sim} \mathbb{A}_{>0}^1$  is the involution taking the inverse,  $j_{\mathbb{A}_{>0}^1}: \mathbb{A}_{>0}^1 \hookrightarrow \mathbb{A}^1$  is the natural open inclusion,  $0_{E/E \times \mathbb{A}^1}: E \hookrightarrow E \times \mathbb{A}^1$  is the zero section, and  $\text{pr}_2: X \times \mathbb{A}_{>0}^1 \rightarrow \mathbb{A}_{>0}^1$  is the projection to the second factor. To simplify the notation, we write

$$\begin{aligned} \tilde{s}_\tau &:= (s \circ p_{\mathbb{A}_{>0}^1}) \cdot (\tau \circ \text{pr}_2): X \times \mathbb{A}_{>0}^1 \rightarrow E, \\ \tilde{s}_{\tau, p} &:= (\tilde{s}_\tau, j_{\mathbb{A}_{>0}^1} \circ \text{pr}_2): X \times \mathbb{A}_{>0}^1 \rightarrow E \times \mathbb{A}^1. \end{aligned}$$

Note that there exists a natural isomorphism

$$(4-9) \quad (\tilde{s}_* j_{X \times \mathbb{A}_{>0}^1, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z} \cong (\tilde{s}_{\tau, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z}.$$

Let  $\pi_{E \times \mathbb{A}^1/E}: E \times \mathbb{A}^1 \rightarrow E$  be the projection. Then the natural transformation

$$\pi_{E \times \mathbb{A}^1/E, *} \rightarrow \pi_{E \times \mathbb{A}^1/E, *} 0_{E/E \times \mathbb{A}^1, *} 0_{E/E \times \mathbb{A}^1}^* \xrightarrow{\sim} 0_{E/E \times \mathbb{A}^1}^*$$

induces a map of complexes on  $E_Z$

$$(4-10) \quad (\tilde{s}_{\tau, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z} \rightarrow (0_{E/E \times \mathbb{A}^1}^* \tilde{s}_{\tau, p, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z}.$$

**Lemma 4.3** *The map (4-10) is an isomorphism.*

**Proof** We let  $\mathcal{L}$  (resp.  $\mathcal{R}$ ) denote the complex at the left-hand side (resp. right-hand side) of the statement. We need to prove the induced map

$$(4-11) \quad H^i(\mathcal{L}_p) \rightarrow H^i(\mathcal{R}_p)$$

is an isomorphism for each  $i \in \mathbb{Z}$  and  $p \in E_Z$ . We have isomorphisms

$$\begin{aligned} H^i(\mathcal{L}_p) &\cong \varinjlim_{p \in V \subset E} H^i(\tilde{s}_\tau^{-1}(V), p_{\mathbb{A}_{>0}^1}^* \mathcal{F}), \\ H^i(\mathcal{R}_p) &\cong \varinjlim_{\substack{p \in V \subset E \\ 0 \in D \subset \mathbb{A}^1}} H^i(\tilde{s}_\tau^{-1}(V) \cap (X \times D), p_{\mathbb{A}_{>0}^1}^* \mathcal{F}), \end{aligned}$$

where  $V$  ranges over all open subsets of  $E$  containing  $p$  and  $D$  ranges over all open subsets of  $\mathbb{A}^1$  containing the origin.

First assume that  $p$  is not contained in the zero section. We will show that for each choice of  $(V, D)$  as above, there exists an open subset  $V' \subset E$  containing  $p$  such that the inclusion

$$\tilde{s}_\tau^{-1}(V') \subset \tilde{s}_\tau^{-1}(V) \cap (X \times D)$$

holds, which implies that the map (4-11) is an isomorphism. To see this, note that the map  $\tilde{s}_\tau|_{X \times (\mathbb{A}_{>0}^1 \setminus D)}$  naturally extends to a map

$$\tilde{s}_{\tau, \mathbb{P}^1 \setminus D}: X \times (\mathbb{P}^1 \setminus D) \rightarrow E.$$

which is proper. It is clear that  $p$  is not contained in the image of  $\tilde{s}_{\tau, \mathbb{P}^1 \setminus D}$  since we assumed  $p \in E_Z \setminus Z$ . Then we can take  $V'$  as  $V \setminus \text{Im}(\tilde{s}_{\tau, \mathbb{P}^1 \setminus D})$ .

Now suppose that  $p$  is contained in the zero section. Let  $V$  be a convex open neighborhood of  $p$  in  $E$  and  $D$  be a convex open neighborhood of 0 in  $\mathbb{A}^1$ . For each point  $q \in X \times \mathbb{A}^1$ , it is clear that  $(\mathbb{R}^+ \cdot q) \cap \tilde{s}_\tau^{-1}(V)$  is contractible. Therefore [Kashiwara and Schapira 1990, Corollary 3.7.3] implies an isomorphism

$$H^i(\tilde{s}_\tau^{-1}(V), p_{\mathbb{A}_{>0}^1}^* \mathcal{F}) \cong H^i(p_{\mathbb{A}_{>0}^1}(\tilde{s}_\tau^{-1}(V)), \mathcal{F}).$$

Similarly, we have an isomorphism

$$H^i(\tilde{s}_\tau^{-1}(V) \cap (X \times D), p_{\mathbb{A}_{>0}^1}^* \mathcal{F}) \cong H^i(p_{\mathbb{A}_{>0}^1}(\tilde{s}_\tau^{-1}(V) \cap (X \times D)), \mathcal{F}).$$

Therefore we have isomorphisms

$$H^i(\mathcal{L}_p) \cong H^i(\mathcal{R}_p) \cong H^i(\mathcal{F}_p),$$

which implies the claim. □

**Lemma 4.4** *There exists an isomorphism*

$$(4-12) \quad (0_{E/E \times \mathbb{A}^1}^* \tilde{s}_{\tau, p, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z} \cong \iota_{C_Z/X, *} \text{Sp}_{Z/X}(\mathcal{F}).$$

**Proof** Consider the commutative diagram

$$(4-13) \quad \begin{array}{ccccc} & C_{Z/X} & \hookrightarrow & M_{Z/X}^o & \longleftarrow X \times \mathbb{A}_{>0}^1 \\ & \downarrow \iota_{C_{Z/X}} & \lrcorner & \downarrow & \swarrow \tilde{s}_{\tau,p} \\ E_Z \times \{0\} & \hookrightarrow & E \times \{0\} & \xrightarrow{0_{E/E \times \mathbb{A}^1}} & E \times \mathbb{A}^1 \end{array}$$

where the middle right vertical map  $M_{Z/X}^o \rightarrow E \times \mathbb{A}^1$  is the natural inclusion induced by the section  $s$  (see [Fulton 1984, Remark 5.1.1]). By the definition of the specialization functor and the nearby cycle functor, we have an isomorphism

$$\mathrm{Sp}_{Z/X}(\mathcal{F}) \cong (C_{Z/X} \hookrightarrow M_{Z/X}^o)^*(X \times \mathbb{A}_{>0}^1 \hookrightarrow M_{Z/X}^o)_* p_{\mathbb{A}_{>0}^1}^* \mathcal{F}.$$

Therefore the assertion follows from the base change theorem.  $\square$

**Proof of Theorem 4.1** We have already seen that the left-hand side of the statement is isomorphic to the complex (4-7), which is isomorphic to the right-hand side of the statement by combining the isomorphism (4-9), and Lemmas 4.3 and 4.4.  $\square$

By the proof, the isomorphism (4-1) in Theorem 4.1 is obtained by the composition

$$(4-14) \quad \begin{aligned} \mathrm{FS}_{E_Z^\vee}(\varphi_{\tilde{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) &\xrightarrow[\sim]{(4-3)} \mathrm{FS}_{E_Z^\vee}((\hat{s}^! i_{X \times \ell \leq 0, *} p_{\ell \leq 0}^! \mathcal{F})|_{E_Z^\vee}) \\ &\xrightarrow[\sim]{(2-42)} \mathrm{FS}_{E^\vee}(\hat{s}^! i_{X \times \ell \leq 0, *} p_{\ell \leq 0}^! \mathcal{F})|_{E_Z} \\ &\xrightarrow[\sim]{(2-48)} (\tilde{s}_* \mathrm{FS}_{X \times \mathbb{A}^1}(i_{X \times \ell \leq 0, *} p_{\ell \leq 0}^! \mathcal{F}))|_{E_Z} \\ &\xrightarrow[\sim]{(4-5)} (\tilde{s}_* j_{X \times \mathbb{A}_{>0}^1, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z} \\ &\xrightarrow[\sim]{(4-9)} (\tilde{s}_{\tau, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z} \\ &\xrightarrow[\sim]{(4-10)} (0_{E/E \times \mathbb{A}^1}^* \tilde{s}_{\tau, p, *} p_{\mathbb{A}_{>0}^1}^* \mathcal{F})|_{E_Z} \xrightarrow[\sim]{(4-12)} \iota_{C_{Z/X}, *} \mathrm{Sp}_{Z/X}(\mathcal{F}). \end{aligned}$$

## 4.2 Dimensional reduction via Fourier–Sato transform

We use the notation as in the previous subsection. Applying the functor  $\pi_{E_Z^\vee, !}$  to the natural map  $\varphi_{\tilde{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee} \rightarrow (\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}$  defined in (2-27), we obtain a map

$$(4-15) \quad \pi_{E_Z^\vee, !}(\varphi_{\tilde{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) \rightarrow \mathcal{F}|_Z$$

which is proven to be an isomorphism by Davison [2017, Theorem A.1]. Similarly, applying the functor  $\pi_{E_Z^\vee, *}$  to the natural map  $(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee} \rightarrow \varphi_{\tilde{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}$  defined in (2-28), we obtain a map

$$(4-16) \quad \mathcal{F}|_Z \rightarrow \pi_{E_Z^\vee, *}(\varphi_{\tilde{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee})[-2 \operatorname{rank} E]$$

which is also an isomorphism. We claim that these isomorphisms are compatible with the isomorphism (4-1). More precisely, the following statement holds:

**Proposition 4.5** (i) The following diagram in  $D_c^b(Z)$  commutes:

$$(4-17) \quad \begin{array}{ccc} \pi_{E_Z^\vee, !}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow[\sim]{(4-15)} & \mathcal{F}|_Z \\ \wr \downarrow (2-46) & & (3-2) \uparrow \wr \\ 0_{E_Z}^* \text{FS}_{E_Z^\vee}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow[\sim]{(-1) \cdot (4-1)} & 0_{E_Z}^* \iota_{C_{Z/X}, *} \text{Sp}_{Z/X}(\mathcal{F}) \end{array}$$

where  $0_{E_Z}: Z \hookrightarrow E_Z$  is the zero section.

(ii) The following diagram in  $D_c^b(Z)$  commutes:

$$(4-18) \quad \begin{array}{ccc} \pi_{E_Z^\vee, *}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee})[-2 \text{rank } E] & \xleftarrow[\sim]{(4-16)} & \mathcal{F}|_Z^! \\ \wr \downarrow (2-47) & & (3-3) \downarrow \wr \\ 0_{E_Z}^! \text{FS}_{E_Z^\vee}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow[\sim]{(-1)^{\text{rank } E+1} \cdot (4-1)} & 0_{E_Z}^! \iota_{C_{Z/X}, *} \text{Sp}_{Z/X}(\mathcal{F}). \end{array}$$

The proof is technical, so we defer it to Section 7.4.

**Remark 4.6** The proof of the above proposition does not depend on Davison's dimensional reduction theorem [2017, Theorem A.1]. Therefore we obtain a new proof of this theorem.

Now assume  $X$  is smooth and consider the composition

$$\mathbb{Q}_Z \cong \pi_{E_Z^\vee, !}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathbb{Q}_X)|_{E_Z^\vee}) \rightarrow \pi_{E_Z^\vee, *}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathbb{Q}_X)|_{E_Z^\vee})^! \cong \omega_Z[2 \text{rank } E - 2 \dim X]$$

where the first isomorphism is the inverse of the map (4-15), the second map exists since the support of  $\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathbb{Q}_X)$  is contained in  $E_Z^\vee$ , and the last isomorphism is the inverse of the map (4-16). We let

$$e_\varphi(E, s) \in H_{2 \dim X - 2 \text{rank } E}^{\text{BM}}(Z)$$

denote the element corresponding to the above composition.

Recall that the localized Euler class  $e_{\text{loc}}(E, s) \in H_{2 \dim X - 2 \text{rank } E}^{\text{BM}}(Z)$  is defined by the image of the fundamental class of the normal cone  $[C_{Z/X}] \in H_{2 \dim X}^{\text{BM}}(C_{Z/X})$  under the following map

$$H_{2 \dim X}^{\text{BM}}(C_{Z/X}) \rightarrow H_{2 \dim X}^{\text{BM}}(E|Z) \simeq H_{2 \dim X - 2 \text{rank } E}^{\text{BM}}(Z).$$

**Corollary 4.7** We have an equality

$$e_\varphi(E, s) = (-1)^{\text{rank } E} e_{\text{loc}}(E, s)$$

where  $e_{\text{loc}}(E, s)$  is the localized Euler class.

**Proof** Consider the composition

$$\mathbb{Q}_Z \cong 0_{E_Z}^* \iota_{C_{Z/X}, *} \text{Sp}_{Z/X}(\mathbb{Q}_X) \rightarrow 0_{E_Z}^! \iota_{C_{Z/X}, *} \text{Sp}_{Z/X}(\mathbb{Q}_X)[2 \text{rank } E] \cong \omega_Z[2 \text{rank } E - 2 \dim X]$$

where the first isomorphism is the inverse of (3-2), the second morphism is (2-12), and the last isomorphism is the inverse of (3-3). Propositions 4.5 and 2.2 imply that this composition corresponds to the element

$(-1)^{\text{rank } E} e_\varphi(E, s)$ . On the other hand, Theorem 3.3 implies that the above composition corresponds to the element  $e_{\text{loc}}(E, s)$ .  $\square$

We will prove the global version of the above corollary in the next section.

## 5 Virtual fundamental classes via vanishing cycles

### 5.1 Virtual fundamental classes for quasismooth derived schemes

We recall the construction of the virtual fundamental class for quasismooth derived schemes following [Behrend and Fantechi 1997; Li and Tian 1998]. Let  $Y$  be a separated quasismooth derived scheme over  $\text{Spec } \mathbb{C}$  and we let  $Y = t_0(Y)$  denote the classical truncation. It is shown in [Brav et al. 2019, Theorem 4.1] that for each  $p \in Y$  we can find a smooth scheme  $U$ , a vector bundle  $E$  on  $U$  and a section  $s \in \Gamma(U, E)$  such that there exists an open immersion

$$Z(s) \hookrightarrow Y$$

where  $Z(s)$  is the derived zero locus of  $s$ . Behrend and Fantechi [1997, Definition 3.10] introduced a conical closed substack of pure dimension zero  $\mathfrak{C}_Y \subset t_0(\mathbf{T}[1]Y)$  of the classical truncation of the shifted tangent stack  $\mathbf{T}[1]Y = \mathbf{Spec}_Y(\text{Sym}(\mathbb{L}_Y[-1]))$  called the intrinsic normal cone characterized by the following property: for each open immersion  $Z(s) \hookrightarrow Y$  as above, the restriction  $\mathfrak{C}_Y|_{t_0(Z(s))}$  is isomorphic to the quotient stack

$$[C_{Z(s)/U}/T_U|_{Z(s)}] \subset [E|_{Z(s)}/T_U|_{Z(s)}] \cong t_0(\mathbf{T}[1]Z(s))$$

where  $Z(s)$  is the classical truncation of  $Z(s)$  and the action of  $T_U|_{Z(s)}$  on the normal cone  $C_{Z(s)/U}$  and the vector bundle  $E|_{Z(s)}$  is defined by the differential of the map  $s: U \rightarrow E$ .

Now assume that  $\mathbb{L}_Y|_Y$  is represented by a two-term complex of locally free sheaves

$$E^\bullet = [E^{-1} \rightarrow E^0]$$

concentrated in degree  $[-1, 0]$ . This assumption is satisfied when  $Y$  is quasiprojective. Write  $E_i := (E^{-i})^\vee$  for  $i = 0, 1$ . For such a resolution  $E^\bullet$ , we define a conical closed subscheme of pure dimension  $\text{rank } E_0$

$$C_{E^\bullet} \subset E_1$$

by the pullback of  $\mathfrak{C}_Y$  along the projection

$$E_1 \rightarrow [E_1/E_0] \cong t_0(\mathbf{T}[1]Y).$$

We define a class  $[Y]_{E^\bullet}^{\text{vir}} \in A_{\text{vdim } Y}(Y)$  by the image of the class  $[C_{E^\bullet}] \in A_{\text{rank } E_0}(E_1)$  under the isomorphism  $A_{\text{rank } E_0}(E_1) \cong A_{\text{vdim } Y}(Y)$ . It is shown in [Behrend and Fantechi 1997, Proposition 5.3] that the class  $[Y]_{E^\bullet}^{\text{vir}}$  is independent of the chosen resolution  $E^\bullet$ , and the class  $[Y]^{\text{vir}} := [Y]_{E^\bullet}^{\text{vir}}$  is called the virtual fundamental class of  $Y$ .



**Example 5.1** Let  $E$  be a vector bundle over a smooth scheme and  $s \in \Gamma(U, E)$  be a section. Then we have the equality

$$\mathrm{cl}_{Z(s)}([Z(s)]_{\mathrm{vir}}) = e_{\mathrm{loc}}(E, s)$$

in  $H_{2 \cdot \mathrm{vdim} Z(s)}^{\mathrm{BM}}(Z(s))$  which easily follows from the definition.

## 5.2 Vanishing cycle complex associated with a $-1$ -shifted cotangent scheme

In [Brav et al. 2015, Theorem 6.9], the authors define natural perverse sheaves associated with oriented d-critical schemes. We do not recall the notion of the d-critical scheme and the construction of this perverse sheaf here. Instead, we explain some properties of the perverse sheaf

$$\varphi_{\mathbf{T}^*[-1]Y} \in \mathrm{Perv}(t_0(\mathbf{T}^*[-1]Y))$$

associated with the natural oriented d-critical structure on the classical truncation  $t_0(\mathbf{T}^*[-1]Y)$  of the  $-1$ -shifted cotangent scheme of a quasismooth derived scheme  $Y$ . As explained in the previous subsection,  $Y$  is locally isomorphic to a derived zero locus  $Z(s)$  where  $s$  is a section of a vector bundle  $E$  on a smooth scheme  $U$ . Note that we have a natural isomorphism

$$t_0(\mathbf{T}^*[-1]Z(s)) \cong \mathrm{Crit}(\bar{s})$$

where  $\bar{s}: E^\vee \rightarrow \mathbb{A}^1$  is the regular function corresponding to the section  $s$ . The first property of  $\varphi_{\mathbf{T}^*[-1]Y}$  that we use in this paper is that for each open inclusion  $\iota: Z(s) \hookrightarrow Y$  as above such that  $E$  and  $\Omega_U$  are trivial vector bundles, we have a natural isomorphism

$$(5-1) \quad \eta_\iota: \varphi_{\mathbf{T}^*[-1]Y}|_{t_0(\mathbf{T}^*[-1]Z(s))} \cong \varphi_{\bar{s}}(\mathbb{Q}_{E^\vee}[\dim U + \mathrm{rank} E]).$$

See [Kinjo 2022, Lemma 2.19] for the proof.

Now write  $\tilde{Y} = t_0(\mathbf{T}^*[-1]Y)$  and  $Y = t_0(Y)$ . We let  $\pi: \tilde{Y} \rightarrow Y$  denote the natural projection. The second property of  $\varphi_{\mathbf{T}^*[-1]Y}$  that we use is the following.

**Theorem 5.2** [Kinjo 2022, Theorem 3.1] *There exist natural isomorphisms which we call the dimensional reduction isomorphisms*

$$\gamma: \pi_* \varphi_{\mathbf{T}^*[-1]Y} \cong \mathbb{Q}_Y[\mathrm{vdim} Y], \quad \bar{\gamma}: \pi_* \varphi_{\mathbf{T}^*[-1]Y} \cong \omega_Y[-\mathrm{vdim} Y]$$

such that for each open inclusion  $\iota: Z(s) \hookrightarrow Y$ , the following diagrams commute:

$$(5-2) \quad \begin{array}{ccc} (\pi_* \varphi_{\mathbf{T}^*[-1]Y})|_{t_0(Z(s))} & \xrightarrow{\gamma|_{t_0(Z(s))}} & \mathbb{Q}_{t_0(Z(s))}[\mathrm{vdim} Y] \\ \downarrow (\pi|_{t_0(\mathbf{T}^*[-1]Z(s))})^*(\eta_\iota) & & \parallel \\ (\pi|_{t_0(\mathbf{T}^*[-1]Z(s))})^* \varphi_{\bar{s}}(\mathbb{Q}_{E^\vee}[\dim U + \mathrm{rank} E]) & \xrightarrow{(4-15)} & \mathbb{Q}_{t_0(Z(s))}[\mathrm{vdim} Y] \end{array}$$

$$(5-3) \quad \begin{array}{ccc} (\pi_* \varphi_{\mathbf{T}^*[-1]Y})|_{t_0(Z(s))} & \xrightarrow{(-1)^{\dim E \cdot (\dim E - 1)/2} \bar{\gamma}|_{t_0(Z(s))}} & \omega_{t_0(Z(s))}[-\mathrm{vdim} Y] \\ \downarrow (\pi|_{t_0(\mathbf{T}^*[-1]Z(s))})^*(\eta_\iota) & & \parallel \\ (\pi|_{t_0(\mathbf{T}^*[-1]Z(s))})^* \varphi_{\bar{s}}(\mathbb{Q}_{E^\vee}[\dim U + \mathrm{rank} E]) & \xleftarrow{(4-16)} & \omega_{t_0(Z(s))}[-\mathrm{vdim} Y] \end{array}$$

The isomorphism  $\gamma$  is constructed in [Kinjo 2022, Theorem 3.1] and the commutativity of the diagram (5-2) follows from its proof. The isomorphism  $\bar{\gamma}$  can be constructed using the Verdier self-duality of  $\varphi_{\mathbf{T}^*[-1]Y}$ , and the commutativity of the diagram (5-3) follows from the Verdier dual of the diagram (5-2), the commutativity of the diagram (2-33), and the discussion after [Kinjo 2022, Theorem 2.17].

Now consider the composition

$$(5-4) \quad \mathbb{Q}_Y[\mathrm{vdim} Y] \xleftarrow[\cong]{\gamma} \pi^! \varphi_{\mathbf{T}^*[-1]Y} \rightarrow \pi_* \varphi_{\mathbf{T}^*[-1]Y} \xrightarrow[\cong]{\bar{\gamma}} \omega_Y[-\mathrm{vdim} Y].$$

We let  $e_\varphi(\mathbf{T}^*[-1]Y) \in H_{2\mathrm{vdim} Y}^{\mathrm{BM}}(Y)$  denote the element corresponding to this composition.

**Theorem 5.3** *Assume that  $\mathbb{L}_Y|_Y$  is represented by a two-term complex of locally free sheaves. Then*

$$(5-5) \quad e_\varphi(\mathbf{T}^*[-1]Y) = (-1)^{\mathrm{vdim} Y \cdot (\mathrm{vdim} Y - 1)/2} \mathrm{cl}_Y([Y]^{\mathrm{vir}}).$$

In other words, the above theorem gives a new construction of the virtual fundamental class in the Borel–Moore homology under a mild assumption that is satisfied when  $Y$  is quasiprojective.

### 5.3 Proof of Theorem 5.3

Take a global resolution  $\mathbb{L}_Y|_Y \cong E^\bullet = [E^{-1} \rightarrow E^0]$  and let  $\iota_{E^\bullet}: t_0(\mathbf{T}^*[-1]Y) \hookrightarrow E^{-1}$  be the natural inclusion. The following lemma is the key ingredient in the proof of Theorem 5.3.

**Lemma 5.4** *The support of the complex  $\mathrm{FS}_{E^{-1}}(\iota_{E^\bullet,*} \varphi_{\mathbf{T}^*[-1]Y})$  is contained in the cone  $C_{E^\bullet}$ .*

**Proof** Take another global resolution  $\mathbb{L}_Y|_Y \cong F^\bullet$  by a two-term complex of locally free sheaves. We use the same notation as above for  $F^\bullet$ . We claim that the statements for  $E^\bullet$  and  $F^\bullet$  are equivalent. Arguing as the proof of [Behrend and Fantechi 1997, Proposition 5.3], we may assume that there exists a strict monomorphism of complexes  $F^\bullet \hookrightarrow E^\bullet$ . Let  $q: E_1 \rightarrow F_1$  be the dual morphism to the map  $F^{-1} \hookrightarrow E^{-1}$  which is smooth surjective by assumption. Then we have an equality  $C_{E^\bullet} = q^{-1}(C_{F^\bullet})$ . Thus the desired equivalence follows from the isomorphism (2-46).  $\square$

Now we return back to the proof of the lemma. Since the statement can be checked locally, we may assume  $Y = Z(s)$  for some section  $s \in \Gamma(U, E)$  of a vector bundle  $E$  on a smooth scheme  $U$ . Further, we may take the canonical resolution  $[E^\vee|_{t_0(Z(s))} \rightarrow \Omega_U|_{t_0(Z(s))}] \simeq \mathbb{L}_{Z(s)}|_{t_0(Z(s))}$  as the global resolution  $E^\bullet$ . Then the claim follows from the isomorphism (5-1) and Theorem 4.1.  $\square$

Let  $0_{C_{E^\bullet}}: Y \hookrightarrow C_{E^\bullet}$  and  $0_{E_1}: Y \hookrightarrow E_1$  be the zero sections and  $i_{C_{E^\bullet}}: C_{E^\bullet} \hookrightarrow E_1$  be the natural inclusion. Consider the diagram in  $D_c^b(Y)$

$$(5-6) \quad \begin{array}{ccc} 0_{C_{E^\bullet}}^*(\mathrm{FS}_{E^{-1}}(\iota_{E^\bullet,*} \varphi_{\mathbf{T}^*[-1]Y})) \otimes 0_{C_{E^\bullet}}^! \mathbb{Q}_{C_{E^\bullet}} & \xrightarrow{(2-13)} & 0_{C_{E^\bullet}}^!(\mathrm{FS}_{E^{-1}}(\iota_{E^\bullet,*} \varphi_{\mathbf{T}^*[-1]Y})) \\ \uparrow (2-46) & & \uparrow (2-47) \\ \pi^! \varphi_{\mathbf{T}^*[-1]Y} \otimes 0_{C_{E^\bullet}}^! \mathbb{Q}_{C_{E^\bullet}} & & \pi_* \varphi_{\mathbf{T}^*[-1]Y}[-2 \mathrm{rank} E_1] \\ \downarrow \gamma \otimes \mathrm{id} & & \downarrow \bar{\gamma} \\ 0_{C_{E^\bullet}}^! \mathbb{Q}_{C_{E^\bullet}}[\mathrm{vdim} Y] & \xrightarrow{\alpha} & \omega_Y[-\mathrm{vdim} Y - 2 \mathrm{rank} E_1] \end{array}$$

where the bottom horizontal arrow  $\alpha$  is defined to make the diagram commutative. Let  $\pi_{C_E^\bullet} : C_E^\bullet \rightarrow Y$  be the natural projection. We have a natural isomorphism  $\pi_{C_E^\bullet}^! \omega_{C_E^\bullet} \cong 0_{C_E^\bullet}^! \omega_{C_E^\bullet}$  constructed in (2-35). Therefore the map  $\alpha$  induces a map

$$\mathbb{Q}_{C_E^\bullet}[\mathrm{vdim} Y] \rightarrow \omega_{C_E^\bullet}[-\mathrm{vdim} Y - 2 \mathrm{rank} E_1]$$

or equivalently an element in  $H_{2 \mathrm{rank} E_0}^{\mathrm{BM}}(C_E^\bullet) \cong H_0^{\mathrm{BM}}(\mathcal{C}_Y)$  which is denoted by  $[\mathcal{C}_Y]_\varphi^{E^\bullet}$ .

**Lemma 5.5** *The class  $[\mathcal{C}_Y]_\varphi^{E^\bullet}$  is independent of the choice of the resolution  $E^\bullet$ .*

**Proof** Arguing as the proof of Lemma 5.4, we need to prove the equality  $[\mathcal{C}_Y]_\varphi^{E^\bullet} = [\mathcal{C}_Y]_\varphi^{F^\bullet}$  for resolutions  $E^\bullet$  and  $F^\bullet$  of the complex  $\mathbb{L}_{Z(s)}|_{Z(s)}$  such that there exists a strict monomorphism  $F^\bullet \hookrightarrow E^\bullet$ . We keep the notation from the proof of Lemma 5.4 and let  $q_C : C_E^\bullet \rightarrow C_F^\bullet$  be the restriction of  $q$ . Consider the diagram in  $D_C^b(Y)$

$$\begin{array}{ccc} 0_{C_E^\bullet}^*(\mathrm{FS}_{E-1}(\iota_{E^\bullet, *}\varphi_{T^*[-1]Y})) \otimes 0_{C_E^\bullet}^! \mathbb{Q}_{C_E^\bullet} & \xrightarrow{\quad} & 0_{C_E^\bullet}^!(\mathrm{FS}_{E-1}(\iota_{E^\bullet, *}\varphi_{T^*[-1]Y})) \\ \downarrow \scriptstyle (A) & & \downarrow \scriptstyle (B) \\ 0_{C_F^\bullet}^*(\mathrm{FS}_{F-1}(\iota_{F^\bullet, *}\varphi_{T^*[-1]Y})) \otimes 0_{C_F^\bullet}^! \mathbb{Q}_{C_F^\bullet}[-2 \dim q] & \rightarrow & 0_{C_F^\bullet}^!(\mathrm{FS}_{F-1}(\iota_{F^\bullet, *}\varphi_{T^*[-1]Y}))[-2 \dim q] \\ \downarrow \scriptstyle \wr & & \downarrow \scriptstyle \wr \\ 0_{C_F^\bullet}^! \mathbb{Q}_{C_F^\bullet}[\mathrm{vdim} Y - 2 \dim q] & \dashrightarrow & \omega_Y[-\mathrm{vdim} Y - 2 \mathrm{rank} E_1] \\ \downarrow \scriptstyle (C) & & \downarrow \\ 0_{C_E^\bullet}^! \mathbb{Q}_{C_E^\bullet}[\mathrm{vdim} Y] & \dashrightarrow & \omega_Y[-\mathrm{vdim} Y - 2 \mathrm{rank} E_1] \end{array}$$

where the inner and outer rectangles are the diagram (5-6) for  $F^\bullet$  and  $E^\bullet$  respectively, the map (A) is defined using the isomorphism (2-46) and the natural isomorphism  $q_C^! \mathbb{Q}_{C_F^\bullet}[-2 \dim q] \cong \mathbb{Q}_{C_E^\bullet}$ , the map (B) is defined using the isomorphism (2-47), and the map (C) is defined by the natural isomorphism  $q_C^! \mathbb{Q}_{C_F^\bullet}[-2 \dim q] \cong \mathbb{Q}_{C_E^\bullet}$  again. The commutativity of the left (resp. right) trapezoid follows from the commutative diagram (2-52) (resp. the diagram (2-53)). The commutativity of the upper trapezoid follows from Proposition 2.2. Therefore we obtain the commutativity of the lower trapezoid, which implies the lemma.  $\square$

**Proposition 5.6** *The class  $[\mathcal{C}_Y]_\varphi^{E^\bullet}$  is equal to  $(-1)^{\mathrm{vdim} Y \cdot (\mathrm{vdim} Y - 1)/2} [\mathcal{C}_Y]$  where  $[\mathcal{C}_Y]$  is the fundamental class of  $\mathcal{C}_Y$ .*

**Proof** Let  $\mathcal{C}_{Y,1}, \dots, \mathcal{C}_{Y,l}$  denote the set of irreducible components of  $\mathcal{C}_Y$ . Since  $\mathcal{C}_Y$  is purely of dimension zero, we have

$$H_0^{\mathrm{BM}}(\mathcal{C}_Y) \cong \bigoplus_{i=1}^l \mathbb{Q} \cdot [\mathcal{C}_{Y,i}].$$

Therefore a class in  $H_0^{\mathrm{BM}}(\mathcal{C}_Y)$  is determined by the multiplicity over each irreducible component. In particular, it is enough to prove the statement locally on  $Y$ .

We may assume  $Y = Z(s)$  where  $s$  is a section of a vector bundle  $E$  on a smooth scheme  $U$ . We write  $Z(s) := t_0(Z(s))$ . Using the previous lemma, we may take the canonical resolution

$$[E^\vee|_{Z(s)} \rightarrow \Omega_U|_{Z(s)}] \simeq \mathbb{L}_{Z(s)|Z(s)}$$

as the global resolution  $E^\bullet$ . In this case, the cone  $C_{E^\bullet}$  is identified with the normal cone  $C_{Z(s)/U} \subset E$  and the embedding  $\iota_{E^\bullet} : t_0(\mathbf{T}^*[-1]Z(s)) \hookrightarrow E^\vee$  is identified with the natural inclusion  $\text{Crit}(\bar{s}) \hookrightarrow E^\vee$ . Therefore the isomorphism (5-1) and Theorem 4.1 implies an isomorphism

$$\text{FS}_{E^\vee}(\iota_{E^\bullet}, * \varphi_{\mathbf{T}^*[-1]Y}) \cong \text{Sp}_{Z(s)/U}(\mathbb{Q}_U[\text{vdim } Z(s)]).$$

Further, Proposition 4.5 and the commutativity of the diagrams (5-2) and (5-3) imply that the outer square of the diagram (5-6) is identified with the following diagram up to the shift by  $\text{vdim } Z(s)$ :

$$(5-7) \quad \begin{array}{ccc} 0_{C_{Z(s)/U}}^*(\text{Sp}_{Z(s)/U}(\mathbb{Q}_U)) \otimes 0_{C_{Z(s)/U}}^! \mathbb{Q}_{C_{Z(s)/U}} & \xrightarrow{(-1)^{\dim E \cdot (\dim E - 1)/2(2-13)}} & 0_{C_{Z(s)/U}}^!(\text{Sp}_{Z(s)/U}(\mathbb{Q}_U)) \\ \wr \downarrow (-1) \cdot (3-2) & & \wr \downarrow (-1)^{\text{rank } E + 1} \cdot (3-3) \\ 0_{C_{Z(s)/U}}^! \mathbb{Q}_{C_{Z(s)/U}} & \xrightarrow{\alpha[-\text{vdim } Z(s)]} & \omega_{Z(s)}[-2 \text{rank } E] \end{array}$$

where  $0_{C_{Z(s)/U}} : Z(s) \hookrightarrow C_{Z(s)/U}$  is the zero section. Therefore Theorem 3.3 and the construction of the map  $\text{Sp}_{Z(s)/U}^{\text{BM}}$  in Section 3.2 implies an equality

$$(C_{Z(s)/U} \rightarrow \mathfrak{C}_Y)^! [\mathfrak{C}_Y]_{\varphi}^{E^\bullet} = (-1)^{\text{vdim } Y \cdot (\text{vdim } Y - 1)/2} \text{Sp}_{Z(s)/U}^{\text{Chow}}([U]) = (-1)^{\text{vdim } Y \cdot (\text{vdim } Y - 1)/2} [C_{Z(s)/U}]$$

which implies the proposition.  $\square$

**Proof of Theorem 5.3** Consider the diagram

$$\begin{array}{ccc} 0_{C_E}^*(\text{FS}_{E^{-1}}(\iota_{E^\bullet}, * \varphi_{\mathbf{T}^*[-1]Y})) \otimes 0_{C_E}^! \mathbb{Q}_{C_E} & \xrightarrow{(2-13)} & 0_{C_E}^! \text{FS}_{E^{-1}}(\iota_{E^\bullet}, * \varphi_{\mathbf{T}^*[-1]Y}) \\ \uparrow & & \wr \uparrow \\ 0_{E_1}^*(\text{FS}_{E^{-1}}(\iota_{E^\bullet}, * \varphi_{\mathbf{T}^*[-1]Y})) \otimes 0_{E_1}^! \mathbb{Q}_{E_1} & \xrightarrow{(2-13)} & 0_{E_1}^! \text{FS}_{E^{-1}}(\iota_{E^\bullet}, * \varphi_{\mathbf{T}^*[-1]Y}) \\ \wr \uparrow (2-46) & & \wr \uparrow (2-47) \\ \pi_! \varphi_{\mathbf{T}^*[-1]Y} \otimes 0_{E_1}^! \mathbb{Q}_{E_1} & & \pi_* \varphi_{\mathbf{T}^*[-1]Y}[-2 \text{rank } E_1] \\ \wr \downarrow \gamma \otimes \text{id} & & \wr \downarrow \bar{\gamma} \\ 0_{E_1}^! \mathbb{Q}_{E_1}[\text{vdim } Y] & \xrightarrow{e_{\varphi}(\mathbf{T}^*[-1]X)[-2 \text{rank } E_1]} & \omega_Y[-\text{vdim } Y - 2 \text{rank } E_1] \\ \downarrow & & \parallel \\ 0_{C_E}^! \mathbb{Q}_{C_E}[\text{vdim } Y] & \xrightarrow{\alpha} & \omega_Y[-\text{vdim } Y - 2 \text{rank } E_1] \end{array}$$

where the outer square is equal to the outer square of (5-6). The commutativity of the upper rectangle and the left and right subdiagrams is obvious, and the commutativity of the middle rectangle follows from Proposition 2.2. This implies the commutativity of the bottom rectangle; hence we conclude that the theorem holds using Proposition 5.6.  $\square$

## 6 DT4 virtual classes and categorifications

In this section, we discuss some possible generalizations of the sheaf-theoretic construction of the virtual fundamental class we have seen in Theorem 5.3. In Section 6.1 we formulate a conjecture closely related to the dimensional reduction theorem [Kinjo 2022, Theorem 3.1] and propose a sheaf-theoretic construction of the DT4 invariant assuming this conjecture. We will see that this construction recovers the square root Euler class up to sign in a special case, which gives evidence of this conjecture. In Section 6.2, we will discuss a categorification of Theorem 5.3 and its relation with the categorical Donaldson–Thomas invariant introduced by Toda [2024].

### 6.1 DT4 invariants

Let  $Y$  be a  $-2$ -shifted symplectic derived scheme over  $\mathrm{Spec} \mathbb{C}$  (eg the fine moduli space of coherent sheaves on a complex Calabi–Yau fourfold). We refer the reader to [Pantev et al. 2013] for the definition of the shifted symplectic structure. The  $-2$ -shifted symplectic structure defines a symmetry of the cotangent complex  $\mathbb{L}_Y^\vee \cong \mathbb{L}_Y[-2]$ . Further assume that  $Y$  is equipped with a choice of orientation. Based on a previous work by Cao and Leung [2014] and Borisov and Joyce [2017], Oh and Thomas [2023] defined a deformation-invariant class

$$[Y]_{\mathrm{vir}}^{\mathrm{DT4}} \in A_{\mathrm{vdim} Y/2}(Y)\left[\frac{1}{2}\right]$$

called the DT4 virtual class under the assumption that  $Y = t_0(Y)$  is quasiprojective. This virtual class can be used to define Donaldson–Thomas type invariants for Calabi–Yau fourfolds.

We now propose a sheaf-theoretic approach to the DT4 virtual class. We define a quadratic function  $q$  on the  $-1$ -shifted cotangent stack  $\tilde{Y} = t_0(\mathbf{T}^*[-1]Y)$  by the composition

$$q: \tilde{Y} \rightarrow t_0(\mathrm{Tot}_Y(\mathbb{L}_Y[-1] \oplus \mathbb{L}_Y[-1])) \rightarrow \mathbb{A}^1$$

where the first map is the diagonal embedding over  $Y$  and the latter map is defined using the  $-2$ -shifted symplectic form on  $Y$ . Let  $\varphi_{\mathbf{T}^*[-1]Y}$  be the perverse sheaf introduced in [Ben-Bassat et al. 2015, Theorem 4.8] associated with the canonical oriented  $-1$ -shifted symplectic structure on  $\mathbf{T}^*[-1]Y$ . We denote by  $\pi: \tilde{Y} \rightarrow Y$  the natural projection and  $0_Y: Y \rightarrow \tilde{Y}$  the zero section.

**Conjecture 6.1** (i) *There exist natural isomorphisms*

$$\gamma: 0_Y^! \varphi_{\mathbf{T}^*[-1]Y} \cong \mathbb{Q}_Y[\mathrm{vdim} Y], \quad \bar{\gamma}: 0_Y^* \varphi_{\mathbf{T}^*[-1]Y} \cong \omega_Y[-\mathrm{vdim} Y].$$

(ii) *There exists a natural isomorphism*

$$\delta: 0_Y^! \varphi_q(\varphi_{\mathbf{T}^*[-1]Y}) \cong \omega_Y$$

*which depends on the chosen orientation.*

**Remark 6.2** Note that the first statement does not follow from Theorem 5.2 since  $Y$  is not quasismooth now. We believe that this statement can be generalized to an arbitrary derived Artin 1-stack whose

cotangent complex is perfect of amplitude  $[-2, 1]$  (or an arbitrary locally finitely presented derived Artin stack once the work [Ben-Bassat et al. 2015] has generalized to higher Artin stacks).

**Remark 6.3** We do not have a natural map  $0_Y^! \varphi_{\mathbf{T}^*[-1]Y} \rightarrow 0_Y^* \varphi_{\mathbf{T}^*[-1]Y}$  in general since  $0_Y$  is not an embedding when  $Y$  is not quasismooth.

Roughly speaking, the DT4 virtual class can be regarded as a “square root” of the virtual fundamental class though we do not know whether we can define virtual fundamental classes for nonquasismooth derived schemes. Therefore we expect that there exists a some Verdier-self-dual complex that sits between  $\mathbb{Q}_Y[\mathrm{vdim} Y]$  and  $\omega_Y[-\mathrm{vdim} Y]$  which recovers DT4 virtual class. Our proposal is that the complex is  $\varphi_q(\varphi_{\mathbf{T}^*[-1]Y})$ , though it lives in  $\tilde{Y}$  rather than  $Y$ .

Now assume that Conjecture 6.1 is true. Consider the composition

$$\mathbb{Q}_Y[\mathrm{vdim} Y] \xrightarrow{\gamma^{-1}} 0_Y^! \varphi_{\mathbf{T}^*[-1]Y} \rightarrow 0_Y^! \varphi_q(\varphi_{\mathbf{T}^*[-1]Y}) \xrightarrow{\delta} \omega_Y$$

where the second map is constructed using the natural transform  $(q^{-1}(0) \hookrightarrow \tilde{Y})^! \rightarrow \varphi_q$ . Let

$$\sqrt{e_\varphi(\mathbf{T}^*[-1]Y)} \in H_{\mathrm{vdim} Y}^{\mathrm{BM}}(Y)$$

be the element corresponding to the above composition.

**Conjecture 6.4** Assume that  $Y$  is quasiprojective. Then there exists a universal sign  $\epsilon_{\mathrm{vdim} Y} \in \{-1, 1\}$  that only depends on  $\mathrm{vdim} Y$  such that we have an equality

$$\epsilon_{\mathrm{vdim} Y} \cdot \sqrt{e_\varphi(\mathbf{T}^*[-1]Y)} = \mathrm{cl}_Y([Y]_{\mathrm{vir}}^{\mathrm{DT4}})$$

where  $\mathrm{cl}_Y : A_{\mathrm{vdim} Y/2}(Y)[\frac{1}{2}] \rightarrow H_{\mathrm{vdim} Y}^{\mathrm{BM}}(Y)$  is the cycle class map (recall that the Borel–Moore homology is taken in the rational coefficient in this paper).

We prove this conjecture in a special case (up to sign). Let  $U$  be an oriented  $-2$ -shifted symplectic derived scheme such that the classical truncation  $U = t_0(U)$  is smooth and that there exists a resolution of the cotangent complex of the form  $\mathbb{L}_U|_U \cong [T_U \xrightarrow{0} E \xrightarrow{0} \Omega_U]$ , where  $E$  is a vector bundle on  $U$ . The oriented  $-2$ -shifted symplectic structure on  $U$  induces a nondegenerate quadratic form  $\tilde{q}$  on  $E$  and a choice of orientation  $o : \mathbb{O}_U \cong \det(E)$ . In this case, the DT4 virtual class  $[Y]_{\mathrm{vir}}^{\mathrm{DT4}}$  is equal to  $\sqrt{e(E)} \cap [U]$  if  $\mathrm{rank} E$  is even where  $\sqrt{e(E)}$  is the Edidin–Graham square root Euler class (see [Edidin and Graham 1995] or [Oh and Thomas 2023, Section 3]) and zero if  $\mathrm{rank} E$  is odd.

**Proposition 6.5** Let  $U$  be a  $-2$ -shifted symplectic derived scheme as above.

- (i) Conjecture 6.1 holds for  $U$ .
- (ii) Assume that  $U$  is connected. Then there exists a choice of sign  $\epsilon \in \{-1, 1\}$  such that

$$\epsilon \cdot \sqrt{e_\varphi(\mathbf{T}^*[-1]U)} = \mathrm{cl}_U([U]_{\mathrm{vir}}^{\mathrm{DT4}}).$$

**Sketch of the proof** Firstly note that we have a natural isomorphism  $t_0(\mathbf{T}^*[-1]U) \cong [E/T_U]$  where  $T_U$  acts on  $E$  trivially. The smoothness of  $t_0(\mathbf{T}^*[-1]U)$  and [Joyce 2015, Example 2.15] imply that the d-critical structure on  $t_0(\mathbf{T}^*[-1]U)$  is trivial. Therefore there exists a local system  $\mathcal{L}$  on  $t_0(\mathbf{T}^*[-1]U)$  such that we have an isomorphism

$$\varphi_{\mathbf{T}^*[-1]U} \cong \mathcal{L}[\text{rank } E].$$

Detailed analysis on the canonical orientation for  $\mathbf{T}^*[-1]U$  as in [Kinjo 2022] shows that the local system  $\mathcal{L}$  is trivial. Therefore the first statement follows immediately.

Now we prove the second statement. We only treat the case when  $\text{rank } E$  is even as the odd case can be proved in a similar way. Let  $0_E : U \hookrightarrow E$  be the zero section. Then we can easily see that the diagram

$$\begin{array}{ccc} 0_E^! \mathbb{Q}_E[2 \text{rank } E] & \longrightarrow & 0_E^! \varphi_{\tilde{q}}(\mathbb{Q}_E)[2 \text{rank } E] \\ \wr \downarrow & & \wr \downarrow \\ \mathbb{Q}_U & \xrightarrow{\cap \sqrt{e_\varphi(\mathbf{T}^*[-1]U)}} & \omega_U[\text{rank } E - 2 \dim U] \end{array}$$

commutes, where the top vertical arrow is induced by the natural transformation  $(\tilde{q}^{-1}(0) \hookrightarrow E)^! \rightarrow \varphi_{\tilde{q}}$ . As proved in [Edidin and Graham 1995, Proposition 5], there exists a smooth proper map  $f : F \rightarrow U$  with the following properties:

- The quadratic bundle  $f^*E$  admits a positive maximal isotropic subbundle  $\Lambda \subset f^*E$ .
- An equality  $f^![U]_{\text{vir}}^{\text{DT4}} = e(\Lambda) \cap [F]$  holds.
- The map  $f^! : H_*^{\text{BM}}(U) \rightarrow H_{*+2 \dim f}^{\text{BM}}(F)$  is injective.

Therefore the proposition follows from the following lemma.

**Lemma 6.6** *Let  $U$  be a connected and separated scheme of finite type over  $\text{Spec } \mathbb{C}$ ,  $(E, q, o)$  be an oriented nondegenerate quadratic vector bundle on  $U$  which admits a positive maximal isotropic subbundle  $\Lambda \subset E$ . Then there exists a choice of sign  $\epsilon \in \{1, -1\}$  such that the following diagram commutes:*

$$\begin{array}{ccc} 0_E^! \mathbb{Q}_E[2 \text{rank } E] & \longrightarrow & 0_E^! \varphi_q(\mathbb{Q}_E)[2 \text{rank } E] \\ \wr \downarrow & & \wr \downarrow \\ \mathbb{Q}_U & \xrightarrow{\epsilon e(\Lambda)} & \mathbb{Q}_U[\text{rank } E] \end{array}$$

**Proof** Write  $r = \text{rank } E$ . Consider the following diagram in  $D^b(E, \mathbb{Z})$ :

$$\begin{array}{ccccccc} & & \mathbb{Z}_U[-2r] & & & & \\ & \swarrow & \downarrow \wr & \searrow & & & \\ & & \varphi_q(\mathbb{Z}_E|_U^!) & & & & \\ & \swarrow & & \searrow & & & \\ \mathbb{Z}_\Lambda[-r] & \xleftarrow{(-1)^{r/2}(U \hookrightarrow \Lambda)_*} & \mathbb{Z}_E|_\Lambda^! & \xrightarrow{\sim} & \varphi_q(\mathbb{Z}_E|_\Lambda^!) & \xrightarrow{\sim} & \varphi_q(\mathbb{Z}_E) & \xrightarrow{\sim} & \mathbb{Z}_U[-r] \\ & \searrow & & \swarrow & & & & & \\ & & (U \hookrightarrow \Lambda)^* & & & & & & \end{array}$$

where the right dashed arrow is defined so that the upper right subdiagram commutes. It is clear that the left subdiagram and the middle triangle commute. Therefore we need to prove the commutativity of the bottom subdiagram up to sign. To do this, we need to show that the natural map

$$(6-1) \quad \varphi_q(\mathbb{Z}_E|_{\Lambda}^!)|_U \rightarrow \varphi_q(\mathbb{Z}_E)|_U$$

is an isomorphism. Take a point  $u \in U$  and let  $j: E \setminus \Lambda \hookrightarrow E$  be the natural open inclusion. Then the claim that the map (6-1) is invertible at  $u$  is equivalent to the vanishing of the complex  $\varphi_q(j_*\mathbb{Z}_{E \setminus \Lambda})_u$  which is further equivalent to the fact that the natural map

$$(6-2) \quad (j_*\mathbb{Z}_{E \setminus \Lambda})_u \rightarrow \psi_q(j_*\mathbb{Z}_{E \setminus \Lambda})_u \cong \psi_q(\mathbb{Z}_E)_u$$

is an isomorphism. Let  $M_{q,u} \subset E$  be a Milnor fiber of  $q$  at  $E$ . Then it is clear that the inclusions

$$E_u \setminus \Lambda_u \hookleftarrow M_{q,u} \cap (E_u \setminus \Lambda_u) \hookrightarrow M_{q,u}$$

are homotopy equivalences. This implies that the map (6-2) is invertible as the stalk of the nearby cycle complex is isomorphic to the cohomology of the Milnor fiber.  $\square$

This completes the proof of Proposition 6.5.  $\square$

## 6.2 Categorification

It is known that the structure sheaf of a quasismooth derived scheme is a categorification of the virtual fundamental class (see [Ciocan-Fontanine and Kapranov 2009] and [Fantechi and Göttsche 2010]). Further, it is expected that there is a categorification of the perverse sheaf associated with  $-1$ -shifted symplectic derived Artin stacks as a dg-category (see [Toda 2024]). Therefore it is natural to expect that Theorem 5.3 has a categorification, which we briefly discuss now. To simplify the discussion, we only consider the schematic case.

Let  $X$  be a  $-1$ -shifted symplectic derived scheme equipped with a fixed orientation. It is conjectured in [Toda 2024] that there is a natural dg-category

$$\mathcal{DT}(X),$$

called the DT-category, locally isomorphic to (a certain twist of) the matrix factorization category. For a quasismooth derived scheme  $Y$  and a conical closed subset  $\mathcal{X} \subset \mathbf{T}^*[-1]Y$ , Toda [2024, Section 3] proposes a definition of the  $\mathbb{C}^*$ -equivariant version of the DT-category as a dg-quotient

$$(6-3) \quad \mathcal{DT}^{\mathbb{C}^*}(\mathbf{T}^*[-1]Y \setminus \mathcal{X}) := D_{\text{coh}}^b(Y)/\mathcal{C}_{\mathcal{X}}$$

where  $\mathcal{C}_{\mathcal{X}}$  is the full dg-subcategory of  $D_{\text{coh}}^b(Y)$  spanned by objects whose singular support is contained in  $\mathcal{X}$ . We refer the reader to [Arinkin and Gaitsgory 2015] for the definition of the singular support.

This definition is motivated by the Koszul duality equivalence stated as follows (see [Toda 2024, Theorem 2.3.3] for this version): Let  $U$  be a smooth scheme,  $E$  be a vector bundle on  $U$ , and  $s \in \Gamma(U, E)$



be a section. We let  $Z(s)$  denote the derived zero locus of  $s$  and  $\bar{s}: E^\vee \rightarrow \mathbb{A}^1$  the regular function corresponding to  $s$ . Then we have an equivalence

$$D_{\text{coh}}^b(Z(s)) \simeq \text{MF}_{\text{coh}}^{\mathbb{C}^*}(U, \bar{s}).$$

The right-hand side is the  $\mathbb{C}^*$ -equivariant matrix factorization category where  $\mathbb{C}^*$  acts on the fiber of  $E \rightarrow U$  by weight two. This equivalence can be regarded as a categorification of Davison's dimensional reduction theorem [2017, Theorem A.1]. Indeed, we have isomorphisms

$$\begin{aligned} \text{HP}_*(D_{\text{coh}}^b(Z(s))) &\cong H_*^{\text{BM}}(t_0(Z(s))) \otimes_{\mathbb{Q}} \mathbb{C}((u)), \\ \text{HP}_*(\text{MF}_{\text{coh}}^{\mathbb{C}^*}(U, \bar{s})) &\cong H^{*+\text{vdim } Z(s)}(\bar{s}^{-1}(0), \varphi_{\bar{s}}) \otimes_{\mathbb{Q}} \mathbb{C}((u)), \end{aligned}$$

where HP denotes the periodic cyclic homology and  $\mathbb{C}((u))$  is the Laurent function field with the formal variable  $u$  with degree 2. The first isomorphism follows from a result of Preygel [2015, Theorem 6.3.2] and the second isomorphism follows from a result of Efimov [2018, Theorem 1.1] (see also [Toda 2024, Lemma 3.3.2]). Similarly, the definition (6-3) can be regarded as a categorification of the global dimensional reduction isomorphism [Kinjo 2022, Theorem 3.1].

Now we can discuss a categorification of Theorem 5.3. As we do not know how to categorify the complex  $\pi_! \varphi_{\mathbf{T}^*[-1]Y}$  directly, we give a slightly different (though equivalent) version of Theorem 5.3. Let  $0_Y: Y = t_0(Y) \hookrightarrow \tilde{Y} = t_0(\mathbf{T}^*[-1]Y)$  be the zero section, and consider the composition

$$(6-4) \quad R\Gamma(Y, \mathbb{Q}_Y[\text{vdim } Y]) \cong R\Gamma(Y, 0_Y^! \varphi_{\mathbf{T}^*[-1]Y}) \rightarrow R\Gamma(\tilde{Y}, \varphi_{\mathbf{T}^*[-1]Y}) \cong R\Gamma(Y, \omega_Y[-\text{vdim } Y])$$

where the first and third isomorphism follows from [Kinjo 2022, Theorem 3.1] and the second isomorphism is the  $!$ -counit map. It is clear that the above composition is equal to the map given by (5-4). The categorification of the complex

$$R\Gamma(Y, 0_Y^! \varphi_{\mathbf{T}^*[-1]Y}) \simeq \text{Cone}(R\Gamma(\tilde{Y}, \varphi_{\mathbf{T}^*[-1]Y}) \rightarrow R\Gamma(\tilde{Y} \setminus Y, \varphi_{\mathbf{T}^*[-1]Y}|_{\tilde{Y} \setminus Y}))[1]$$

is  $\mathcal{C}_{0_Y}(Y)$  and the categorification of the complex  $R\Gamma(\tilde{Y}, \varphi_{\mathbf{T}^*[-1]Y})$  is  $\mathcal{DT}^{\mathbb{C}^*}(\mathbf{T}^*[-1]Y)$ . The categorification of the first isomorphism in (6-4) is

$$\text{Perf}(Y) \simeq \mathcal{C}_{0_Y}(Y)$$

which follows from [Arinkin and Gaitsgory 2015, Theorem 4.2.6], and the categorification of the latter isomorphism in (6-4) is

$$\mathcal{DT}^{\mathbb{C}^*}(\mathbf{T}^*[-1]Y) \simeq D_{\text{coh}}^b(Y)$$

which is nothing but the definition. Therefore the categorification of Theorem 5.3 is a tautological statement that the image of the structure sheaf  $\mathcal{O}_Y$  under the inclusion

$$\text{Perf}(Y) \hookrightarrow D_{\text{coh}}^b(Y)$$

is also  $\mathcal{O}_Y$ .

**Question 6.7** *Can we give another proof of Theorem 5.3 based on the above discussion combined with the virtual Riemann–Roch theorem [Fantechi and Göttsche 2010; Ciocan-Fontanine and Kapranov 2009]?*

**Question 6.8** Can we formulate a categorification of Conjecture 6.4 and define a version of “DT4 virtual structure sheaf”? The first difficulty is defining a categorification of the perverse sheaf  $\varphi_q(\varphi_{\mathbf{T}^*[-1]}Y)$  associated with a  $-2$ -shifted symplectic derived scheme  $Y$ .

## 7 Postponed proofs

In this section, we will give proofs of Propositions 2.1, 2.2, 2.3 and 4.5.

### 7.1 Proof of Proposition 2.1

First consider the composition

$$(7-1) \quad \mathbb{Q}_0[-2] \cong \text{Cone}((\mathbb{C}^* \hookrightarrow \mathbb{C})_! \mathbb{Q}_{\mathbb{C}^*} \rightarrow \mathbb{Q}_{\mathbb{C}})[-2] \rightarrow K'_{\psi} \rightarrow \mathbb{Q}_{\mathbb{C}}.$$

Here, the second (resp. third) map is defined using the short exact sequence (2-23) (resp. (2-21)). We will prove that this is equal to the counit map

$$\mathbb{Q}_0[-2] \cong (\{0\} \hookrightarrow \mathbb{C})_! (\{0\} \hookrightarrow \mathbb{C})^! \mathbb{Q}_{\mathbb{C}} \rightarrow \mathbb{Q}_{\mathbb{C}}$$

later, and now prove the proposition assuming this statement. Consider the diagram

$$\begin{array}{ccc} i^* \mathcal{H}om(\mathbb{Q}_{Z \times \mathbb{C}}, \pi^* \mathcal{F}) & \longrightarrow & i^* \mathcal{H}om(i_! \mathbb{Q}_{Z \times \{0\}}[-2], \pi^* \mathcal{F}) \\ \downarrow \wr & & \downarrow \wr \\ i^* \pi^* \mathcal{F} & \longrightarrow & i^! \pi^* \mathcal{F}[2] \end{array}$$

where the upper horizontal arrow is induced from the counit map  $i_! i^! \rightarrow \text{id}$  and the bottom horizontal arrow is the natural map (2-12). It is clear that this diagram commutes. Therefore what we need to prove is that the composition

$$\mathcal{F} \cong i^* \pi^* \mathcal{F} \rightarrow i^! \pi^* \mathcal{F}[2] \rightarrow i^! \pi^! \mathcal{F} \cong \mathcal{F}$$

where the second and third maps are the natural maps (2-12) is the identity map, but this is obvious since the composition

$$\mathbb{Q}_Z \cong i^! \pi^! \mathbb{Q}_Z \cong i^* \pi^! \mathbb{Q}_Z[-2] \cong i^* \pi^* \mathbb{Q}_Z \cong \mathbb{Q}_Z$$

is the identity map.

Now we prove that (7-1) is given by the counit map. To prove this statement, it is enough to prove the following composition is the identity map:

$$\mathbb{Q} \cong R\Gamma_c(\mathbb{Q}_0) \rightarrow R\Gamma_c(K'_{\psi})[2] \rightarrow R\Gamma_c(\mathbb{Q}_{\mathbb{C}}[2]) \cong \mathbb{Q}.$$

It is clear that the above map is given by

$$\mathbb{Q} \cong R\Gamma_c([p_! \mathbb{Q}_{\tilde{\mathbb{C}}^*} \xrightarrow{1-T} p_! \mathbb{Q}_{\tilde{\mathbb{C}}^*} \xrightarrow{\text{tr}} \mathbb{Q}_{\mathbb{C}}]) \xrightarrow{-\text{pr}} R\Gamma_c(p_! \mathbb{Q}_{\tilde{\mathbb{C}}^*}[2]) \cong \mathbb{Q}$$

where  $\text{pr}$  denotes the canonical projection.

From now we work with  $\mathbb{R}$ -coefficients, which does not affect the conclusion. We compute the above composition using differential forms. Recall that for a differentiable manifold  $M$ , we can take the de Rham resolution of a constant sheaf

$$\mathbb{R}_M \simeq \Omega_M^\bullet = [\Omega_M^0 \xrightarrow{d} \Omega_M^1 \xrightarrow{d} \cdots].$$

Consider the resolution

$$[p_! \mathbb{R}_{\tilde{\mathbb{C}}^*} \xrightarrow{1-T} p_! \mathbb{R}_{\tilde{\mathbb{C}}^*} \xrightarrow{\text{tr}} \mathbb{R}_{\mathbb{C}}] \simeq \text{Tot}([p_! \Omega_{\tilde{\mathbb{C}}^*}^\bullet \xrightarrow{1-(\alpha^{-1})^*} p_! \Omega_{\tilde{\mathbb{C}}^*}^\bullet \xrightarrow{p_*} \Omega_{\mathbb{C}}^\bullet])$$

where  $\text{Tot}$  denotes the totalization and  $\alpha$  is the covering transformation corresponding to a counterclockwise loop. Since all sheaves appearing on the right-hand side are c-soft, the complex

$$C^\bullet := \Gamma_c(\text{Tot}([p_! \Omega_{\tilde{\mathbb{C}}^*}^\bullet \xrightarrow{1-(\alpha^{-1})^*} p_! \Omega_{\tilde{\mathbb{C}}^*}^\bullet \xrightarrow{p_*} \Omega_{\mathbb{C}}^\bullet]))$$

computes the derived functor, hence is isomorphic to  $\mathbb{R}$  concentrated in degree zero. Note that  $C^0$  consists of triples of differential forms  $(f, \omega_1, \omega_2) \in \Omega_{\mathbb{C}}^0 \oplus \Omega_{\tilde{\mathbb{C}}^*}^1 \oplus \Omega_{\tilde{\mathbb{C}}^*}^2$  with compact supports, which is a cocycle if and only if the following identities hold:

$$df = -p_* \omega_1, \quad d\omega_1 = \omega_2 - (\alpha^{-1})^* \omega_2.$$

Now what we need to prove is

$$f(0) = - \int_{\tilde{\mathbb{C}}^*} \omega_2$$

for such a triple.

By assumption, there exists a positive real numbers  $\epsilon, R \in \mathbb{R}_{>0}$  such that the support of  $f$  is contained in a domain  $B_R \setminus B_\epsilon$  and the supports of  $\omega_1$  and  $\omega_2$  is contained in  $p^{-1}(B_R \setminus B_\epsilon)$ . Here  $B_\epsilon$  and  $B_R$  denote the open disks centered at the origin with radii  $\epsilon$  and  $R$  respectively. Let  $l: [0, 1] \rightarrow \mathbb{C}$  be a line segment such that  $l(0) = \epsilon$  and  $l(1) = R$ . We take a lift  $\tilde{l}_0: [0, 1] \rightarrow \tilde{\mathbb{C}}^*$  of the line  $l$ . For an integer  $i$ , we set  $\tilde{l}_i := \alpha^i(\tilde{l}_0)$ . We let  $D_i \subset \tilde{\mathbb{C}}^*$  the segment surrounded by  $\tilde{l}_i, \tilde{l}_{i+1}$  and arcs of  $p^{-1}(\partial B_\epsilon)$  and  $p^{-1}(\partial B_R)$ . By Stokes' theorem, we have the identities

$$\begin{aligned} f(0) &= f(\epsilon) = f(\epsilon) - f(R) = - \int_l df = \int_l p_* \omega_1 = \sum_i \int_{\tilde{l}_i} \omega_1, \\ \int_{D_i} \omega_2 - \int_{D_{i-1}} \omega_2 &= \int_{D_i} (\omega_2 - (\alpha^{-1})^* \omega_2) = \int_{D_i} d\omega_1 = \int_{\tilde{l}_i} \omega_1 - \int_{\tilde{l}_{i+1}} \omega_1. \end{aligned}$$

Set  $A_i := \int_{D_i} \omega_2 - \int_{D_{i-1}} \omega_2 = \int_{\tilde{l}_i} \omega_1 - \int_{\tilde{l}_{i+1}} \omega_1$ . Then we have the identity

$$\sum_{i \in \mathbb{Z}} i \cdot A_i = \sum_{i \in \mathbb{Z}} \int_{D_i} \omega_2 = - \sum_{i \in \mathbb{Z}} \int_{\tilde{l}_i} \omega_1.$$

In particular, we obtain the identity

$$f(0) = \sum_{i \in \mathbb{Z}} \int_{\tilde{l}_i} \omega_1 = - \sum_{i \in \mathbb{Z}} \int_{D_i} \omega_2 = - \int_{\tilde{\mathbb{C}}^*} \omega_2$$

as desired.

## 7.2 Proof of Proposition 2.2

We use the notation as in the diagram (2-50). First, for a complex vector bundle  $E$ , we define two functors

$$\tilde{\mathrm{FS}}_E, \tilde{\mathrm{FS}}'_E: D_{\mathbb{R}+}^+(E) \rightarrow D_{\mathbb{R}+}^+(E^\vee)$$

by

$$\tilde{\mathrm{FS}}_E := p'_! \iota_{P'} * \iota_{P'}^* \iota_{P'}^! \iota_P^! p^*, \quad \tilde{\mathrm{FS}}'_E := p'_! \iota_{P'} * \iota_{P'}^* \iota_{P'}^! \iota_P^! p^*.$$

Then we have the commutative diagram

$$\begin{array}{ccc} \tilde{\mathrm{FS}}_E & \xrightarrow{\sim} & \tilde{\mathrm{FS}}'_E \\ \downarrow \wr & & \uparrow \wr \\ \mathrm{FS}_E & \xrightarrow{\sim} & \mathrm{FS}'_E \end{array}$$

where the upper horizontal arrow is induced from the natural transform  $p'_! \rightarrow p'_*$ , the left vertical arrow is induced from the counit map  $\iota_P^! \iota_P^! \rightarrow \mathrm{id}$  and the right vertical arrow is induced from the unit map  $\mathrm{id} \rightarrow \iota_{P'} * \iota_{P'}^*$ .

By repeating the construction of (2-46) and (2-47), we can construct natural isomorphisms

$$\tilde{\mathrm{FS}}_{E_2}(g^! \mathcal{F}) \cong {}^t g^* \tilde{\mathrm{FS}}_{E_1}(\mathcal{F}), \quad \tilde{\mathrm{FS}}'_{E_2}(g_* \mathcal{F}) \cong {}^t g^! \tilde{\mathrm{FS}}'_{E_1}(\mathcal{F})[2 \dim g]$$

by composing base change maps or the inverse of base change maps. These maps are compatible with the natural transforms  $\mathrm{FS}_E \cong \tilde{\mathrm{FS}}_E$  and  $\mathrm{FS}'_E \cong \tilde{\mathrm{FS}}'_E$  since the  $!$ -counit and  $*$ -unit commute with base change maps as we have seen in Section 2.2. Therefore we need to prove the commutativity of the following diagram:

$$\begin{array}{ccc} \tilde{\mathrm{FS}}_{E_2}(g^! \mathcal{F}) & \xrightarrow{\quad} & \tilde{\mathrm{FS}}'_{E_2}(g_* \mathcal{F}) \\ \downarrow \wr & & \downarrow \wr \\ {}^t g^* \tilde{\mathrm{FS}}_{E_1}(\mathcal{F}) & \xrightarrow{\quad} & {}^t g^! \tilde{\mathrm{FS}}'_{E_1}(\mathcal{F})[-2 \dim {}^t g] \end{array}$$

By definition, the left vertical map is decomposed into isomorphisms

$$\begin{aligned} \tilde{\mathrm{FS}}_{E_2}(g^! \mathcal{F}) &= p'_{2!} \iota_{P'_2} * \iota_{P'_2}^* \iota_{P_2}^! \iota_{P_2}^! p_2^* g^! \mathcal{F} \xleftarrow{\sim} \tilde{p}'_{1!} \iota_{\tilde{P}'} * \iota_{\tilde{P}'}^* \iota_{\tilde{P}}^! \iota_{\tilde{P}}^! \tilde{g}^* p_1^* \mathcal{F} \\ &\xleftarrow{\sim} {}^t g^* p'_1 \iota_{P'_1} * \iota_{P'_1}^* \iota_{P_1}^! \iota_{P_1}^! p_1^* \mathcal{F} = {}^t g^* \tilde{\mathrm{FS}}_{E_1}(\mathcal{F}) \end{aligned}$$

and the right vertical map is decomposed into the isomorphisms

$$\begin{aligned} \tilde{\mathrm{FS}}'_{E_2}(g_* \mathcal{F}) &= p'_{2*} \iota_{P'_2} * \iota_{P'_2}^* \iota_{P_2}^! \iota_{P_2}^! p_2^* g_* \mathcal{F} \\ &\xrightarrow{\sim} \tilde{p}'_{1*} \iota_{\tilde{P}'} * \iota_{\tilde{P}'}^* \iota_{\tilde{P}}^! \iota_{\tilde{P}}^! \tilde{g}^* p_1^* \mathcal{F} \\ &\cong \tilde{p}'_{1*} \iota_{\tilde{P}'} * \iota_{\tilde{P}'}^* \iota_{\tilde{P}}^! \iota_{\tilde{P}}^! \tilde{g}^! p_1^* \mathcal{F}[-2 \dim {}^t g] \\ &\xrightarrow{\sim} {}^t g^! p'_1 \iota_{P'_1} * \iota_{P'_1}^* \iota_{P_1}^! \iota_{P_1}^! p_1^* \mathcal{F}[-2 \dim {}^t g] \\ &= {}^t g^! \tilde{\mathrm{FS}}'_{E_1}(\mathcal{F})[-2 \dim {}^t g]. \end{aligned}$$

Then the claim follows from the fact that the natural transformations (2-10) and (2-12) commute with base change maps and the commutativity of the diagrams (2-15).

### 7.3 Proof of Proposition 2.3

To prove this statement, we need several lemmas.

**Lemma 7.1** *The following diagram commutes up to the sign  $(-1)^{\text{rank } E_1}$ :*

$$(7-2) \quad \begin{array}{ccc} \text{FS}_{E_2^\vee} \text{FS}_{E_2}(g_* \mathcal{F}) & \xrightarrow{\sim} & \text{FS}_{E_2^\vee} {}^t g^! \text{FS}_{E_1}(\mathcal{F})[2 \dim g] \xrightarrow{\sim} g_* \text{FS}_{E_1^\vee} \text{FS}_{E_1}(\mathcal{F})[2 \dim g] \\ \eta(g_* \mathcal{F}) \downarrow \wr & & g_* \eta(\mathcal{F}) \downarrow \wr \\ g_* \mathcal{F}^a[-2 \text{rank } E_2] & \xlongequal{\quad} & g_* \mathcal{F}^a[-2 \text{rank } E_2] \end{array}$$

**Proof** Note that the map  $\text{FS}_{E_2^\vee} {}^t g^! \text{FS}_{E_1}(\mathcal{F}) \rightarrow g_* \text{FS}_{E_1^\vee} \text{FS}_{E_1}(\mathcal{F})$  is defined by the composition

$$\begin{aligned} \text{FS}_{E_2^\vee} {}^t g^! \text{FS}_{E_1}(\mathcal{F}) & \xrightarrow[\sim]{\text{FS}_{E_2^\vee} {}^t g^! (\eta(\text{FS}_{E_1}(\mathcal{F}^a))^{-1})} \text{FS}_{E_2^\vee} {}^t g^! \text{FS}_{E_1} \text{FS}_{E_1^\vee} \text{FS}_{E_1}(\mathcal{F}^a)[2 \text{rank } E_1] \\ & \xrightarrow{\sim} \text{FS}_{E_2^\vee} \text{FS}_{E_2} g_* \text{FS}_{E_1^\vee} \text{FS}_{E_1}(\mathcal{F}^a)[2 \text{rank } E_2] \\ & \xrightarrow[\sim]{\eta(g_* \text{FS}_{E_1^\vee} \text{FS}_{E_1}(\mathcal{F}^a))} g_* \text{FS}_{E_1^\vee} \text{FS}_{E_1}(\mathcal{F}). \end{aligned}$$

We have seen that  $\eta(\text{FS}_{E_1}(\mathcal{F}^a))$  differs from  $\text{FS}_{E_1}(\eta(\mathcal{F}^a))$  by  $(-1)^{\text{rank } E_1}$  in (2-40). Then the claim follows immediately.  $\square$

Take an object  $\mathcal{G} \in D_{\mathbb{R}^+}^+(E_2)$ . A similar argument shows that the following three diagrams commute up to the sign  $(-1)^{\text{rank } E_1}$ :

$$(7-3) \quad \begin{array}{ccc} \text{FS}_{E_2^\vee} \text{FS}_{E_2}(g_! \mathcal{F}) & \xrightarrow{\sim} & \text{FS}_{E_2^\vee} {}^t g^* \text{FS}_{E_1}(\mathcal{F}) \xrightarrow{\sim} g_! \text{FS}_{E_1^\vee} \text{FS}_{E_1}(\mathcal{F})[2 \dim g] \\ \eta(g_! \mathcal{F}) \downarrow \wr & & g_! \eta(\mathcal{F}) \downarrow \wr \\ g_! \mathcal{F}^a[-2 \text{rank } E_2] & \xlongequal{\quad} & g_! \mathcal{F}^a[-2 \text{rank } E_2] \end{array}$$

$$(7-4) \quad \begin{array}{ccc} \text{FS}_{E_1^\vee} \text{FS}_{E_1}(g^* \mathcal{G}) & \xrightarrow{\sim} & \text{FS}_{E_1^\vee} {}^t g_! \text{FS}_{E_2}(\mathcal{G})[-2 \dim g] \xrightarrow{\sim} g^* \text{FS}_{E_2^\vee} \text{FS}_{E_2}(\mathcal{G})[-2 \dim g] \\ \eta(g^* \mathcal{G}) \downarrow \wr & & g^* \eta(\mathcal{G}) \downarrow \wr \\ g^* \mathcal{G}^a[-2 \text{rank } E_1] & \xlongequal{\quad} & g^* \mathcal{G}^a[-2 \text{rank } E_1] \end{array}$$

$$(7-5) \quad \begin{array}{ccc} \text{FS}_{E_1^\vee} \text{FS}_{E_1}(g^! \mathcal{G}) & \xrightarrow{\sim} & \text{FS}_{E_1^\vee} {}^t g_* \text{FS}_{E_2}(\mathcal{G}) \xrightarrow{\sim} g^! \text{FS}_{E_2^\vee} \text{FS}_{E_2}(\mathcal{G})[-2 \dim g] \\ \eta(g^! \mathcal{G}) \downarrow \wr & & g^! \eta(\mathcal{G}) \downarrow \wr \\ g^! \mathcal{G}^a[-2 \text{rank } E_1] & \xlongequal{\quad} & g^! \mathcal{G}^a[-2 \text{rank } E_1] \end{array}$$

**Lemma 7.2** *The following diagram commutes up to the sign  $(-1)^{\dim g}$ :*

$$\begin{array}{ccc}
 \mathcal{F}^a[-2 \operatorname{rank} E_1] & \xrightarrow{\quad} & g^! g^! \mathcal{F}^a[-2 \operatorname{rank} E_1] \\
 \downarrow \wr & & \downarrow \wr \\
 \operatorname{FS}_{E_1^\vee} \operatorname{FS}_{E_1}(\mathcal{F}) & & \\
 \downarrow & & \downarrow \\
 \operatorname{FS}_{E_1^\vee} {}^t g_* {}^t g^* \operatorname{FS}_{E_1}(\mathcal{F}) & \xrightarrow{\sim} & g^! \operatorname{FS}_{E_2^\vee} \operatorname{FS}_{E_2}(g^! \mathcal{F})[-2 \dim g]
 \end{array}$$

**Proof** Using the fact that the diagram (2-38) commutes up to the sign  $(-1)^{\operatorname{rank} E}$ , we need to prove the commutativity of the following diagram:

$$\begin{array}{ccc}
 \mathcal{F} & \xrightarrow{\quad} & g^! g^! \mathcal{F} \\
 \downarrow \wr & & \downarrow \wr \\
 \check{\operatorname{FS}}_{E_1^\vee} \operatorname{FS}_{E_1}(\mathcal{F}) & & \\
 \downarrow & & \downarrow \\
 \check{\operatorname{FS}}_{E_1^\vee} {}^t g_* {}^t g^* \operatorname{FS}_{E_1}(\mathcal{F}) & \xrightarrow{\sim} & g^! \check{\operatorname{FS}}_{E_2^\vee} \operatorname{FS}_{E_2}(g^! \mathcal{F})
 \end{array}$$

Here the natural transformation

$$\check{\operatorname{FS}}_{E_1^\vee} \circ {}^t g_* \xrightarrow{\sim} g^! \circ \check{\operatorname{FS}}_{E_2^\vee}$$

is defined by composing base change maps as in the construction of the map (2-47). Note that all arrows in the diagram except for the horizontal bottom arrow are defined by composing unit maps. Therefore the claim follows from the construction of the base change map (2-6) in [Kashiwara and Schapira 1990, Proposition 3.1.9(ii)].  $\square$

The following statement is a consequence of the above lemma and the commutativity of diagrams (2-38), (7-2) and (7-4) up to a certain choice of sign.

**Corollary 7.3** *The following diagram commutes:*

$$\begin{array}{ccc}
 \mathcal{G}^a[-2 \operatorname{rank} E_2] & \xrightarrow{\quad} & g_* g^* \mathcal{G}^a[-2 \operatorname{rank} E_2] \\
 \downarrow \wr & & \downarrow \wr \\
 \operatorname{FS}_{E_2^\vee} \operatorname{FS}_{E_2}(\mathcal{G}) & & \\
 \downarrow & & \downarrow \\
 \operatorname{FS}_{E_2^\vee} {}^t g^! {}^t g^! \operatorname{FS}_{E_2}(\mathcal{G}) & \xrightarrow{\sim} & g_* \operatorname{FS}_{E_1^\vee} \operatorname{FS}_{E_1}(g^* \mathcal{G})[2 \dim g]
 \end{array}$$

The following two statements can be proved in a similar way using (7-3) and (7-5):

**Lemma 7.4** *The following diagram commutes:*

$$\begin{array}{ccc}
 g^* g_* \mathcal{F}^a[-2 \operatorname{rank} E_1] & \longrightarrow & \mathcal{F}^a[-2 \operatorname{rank} E_1] \\
 \uparrow \wr & & \uparrow \wr \\
 g^* \operatorname{FS}_{E_2^\vee} \operatorname{FS}_{E_2}(g_* \mathcal{F})[-2 \dim g] & \xrightarrow{\sim} & \operatorname{FS}_{E_1^\vee} {}^t g! {}^t g^* \operatorname{FS}_{E_1}(\mathcal{F})
 \end{array}$$

**Corollary 7.5** *The following diagram commutes up to the sign  $(-1)^{\dim g}$ :*

$$\begin{array}{ccc}
 g! g^! \mathcal{G}^a[-2 \operatorname{rank} E_2] & \longrightarrow & \mathcal{G}^a[-2 \operatorname{rank} E_2] \\
 \uparrow \wr & & \uparrow \wr \\
 g! \operatorname{FS}_{E_1^\vee} \operatorname{FS}_{E_1}(g^! \mathcal{G}^a)[2 \dim g] & \xrightarrow{\sim} & \operatorname{FS}_{E_2^\vee} {}^t g^* {}^t g_* \operatorname{FS}_{E_2}(\mathcal{G})
 \end{array}$$

**Proof of Proposition 2.3** We first prove that the left diagram commutes up to the sign  $(-1)^{\operatorname{rank} E_1}$ . Using Corollary 7.3, we need to prove that the following diagram commutes up to the sign  $(-1)^{\operatorname{rank} E_1}$ :

$$\begin{array}{ccc}
 \operatorname{FS}_{E_2} \mathcal{G}^a[-2 \operatorname{rank} E_2] & \longrightarrow & {}^t g! {}^t g_* \operatorname{FS}_{E_2}(\mathcal{G}^a)[-2 \operatorname{rank} E_2] \\
 \operatorname{FS}_{E_2} \eta(\mathcal{G})^{-1} \downarrow \wr & & \downarrow \wr \\
 \operatorname{FS}_{E_2} \operatorname{FS}_{E_2^\vee} \operatorname{FS}_{E_2}(\mathcal{G}) & & \operatorname{FS}_{E_2}(g_* g^* \mathcal{G}^a)[-2 \operatorname{rank} E_2] \\
 \downarrow & & \downarrow \wr \\
 \operatorname{FS}_{E_2} \operatorname{FS}_{E_2^\vee} {}^t g! {}^t g_* \operatorname{FS}_{E_2}(\mathcal{G}) & \xrightarrow{\sim} & \operatorname{FS}_{E_2} g_* \operatorname{FS}_{E_1^\vee} \operatorname{FS}_{E_1}(g^* \mathcal{G})[2 \dim g]
 \end{array}$$

This follows using the commutativity of the diagram (2-40) up to the sign  $(-1)^{\operatorname{rank} E}$  twice and Lemma 7.1.

The commutativity of the right diagram in the statement of Proposition 2.3 up to the sign  $(-1)^{\operatorname{rank} E_2}$  can be proved in a similar manner using Corollary 7.5 and (7-3).  $\square$

## 7.4 Proof of Proposition 4.5

We first prove statement (i). To do this, consider the following diagram in  $D_c^b(E_Z)$ :

$$\begin{array}{ccccc}
 & & & & 0_{E_Z, *}(\mathcal{F}|_Z) \\
 & & & \nearrow (2-48) \sim & \uparrow (3-4) \wr \\
 (7-6) & \operatorname{FS}_{E_Z^\vee}((\pi_E^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow{(2-42) \sim} & \operatorname{FS}_{E^\vee}(\pi_E^! \mathcal{F})|_{E_Z} & \iota_{C_Z/X, *} \operatorname{Sp}_{Z/X}(\mathcal{F}_Z) \\
 & \nearrow (2-27) & & & \uparrow \\
 & \operatorname{FS}_{E_Z^\vee}(\varphi_{\bar{s}}(\pi_E^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow{(-1) \cdot (4-1) \sim} & \iota_{C_Z/X, *} \operatorname{Sp}_{Z/X}(\mathcal{F})
 \end{array}$$

We claim that this diagram is commutative. The most complicated map in this diagram is of course (4-1) as it is defined by the composition of seven morphisms (4-14). To deal with this map, consider the following composition “parallel” to the composition (4-14):

$$\begin{aligned}
 (7-7) \quad \mathrm{FS}_{E_Z^\vee}((\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) &\xrightarrow{\sim} \mathrm{FS}_{E_Z^\vee}((\hat{s}^! p_{\mathbb{A}^1}^! \mathcal{F})|_{E_Z^\vee}) \\
 &\xrightarrow[\sim]{(2-42)} \mathrm{FS}_{E^\vee}(\hat{s}^! p_{\mathbb{A}^1}^! \mathcal{F})|_{E_Z} \\
 &\xrightarrow[\sim]{(2-48)} (\tilde{s}_* \mathrm{FS}_{X \times \mathbb{A}^1}(p_{\mathbb{A}^1}^! \mathcal{F}))|_{E_Z} \\
 &\xrightarrow[\sim]{(2-48)} (\tilde{s}_* 0_{X/X \times \mathbb{A}^1, *} \mathcal{F})|_{E_Z} \\
 &\cong 0_{E_Z, *}(\mathcal{F}|_Z) = 0_{E_Z, *}(\mathcal{F}|_Z) = 0_{E_Z, *}(\mathcal{F}|_Z)
 \end{aligned}$$

where  $0_{X/X \times \mathbb{A}^1}: X \hookrightarrow X \times \mathbb{A}^1$  in the fourth line is the zero section. Here what we mean by this composition is “parallel” to the composition (4-14) is that each complex appearing in (4-14) has a natural map to the complex appearing in the above composition at the same stage. It is clear that these morphisms commute with morphisms appearing in (4-14) and (7-7) except for the ones in the fourth rows. To prove the commutativity of (7-6) we need to show that the maps in the fourth rows of (4-14) and (7-7) commute with the natural maps from the complexes at the third and fourth rows of (4-14) to the complexes at the same stages of (7-7) up to the sign  $-1$ , ie we need to show the commutativity of the following diagram:

$$\begin{aligned}
 (7-8) \quad &\mathrm{FS}_{X \times \mathbb{A}^1}(i_{X \times \ell \leq 0, *} (p_{\ell \leq 0}^! \mathcal{F})) \longrightarrow \mathrm{FS}_{X \times \mathbb{A}^1}(p_{\mathbb{A}^1}^! \mathcal{F}) \\
 &\quad \wr \downarrow (4-5) \qquad \qquad \qquad \wr \downarrow (-1) \cdot (2-48) \\
 &j_{X \times \mathbb{A}^1_{>0}, *} p_{\mathbb{A}^1_{>0}}^* \mathcal{F} \longrightarrow 0_{X/X \times \mathbb{A}^1, *} \mathcal{F}
 \end{aligned}$$

To do this, consider the following larger diagram:

$$\begin{aligned}
 (7-9) \quad &\mathrm{FS}_{X \times \mathbb{A}^1}(0_{X/X \times \mathbb{A}^1, *} \mathcal{F}) \longrightarrow \mathrm{FS}_{X \times \mathbb{A}^1}(i_{X \times \ell \leq 0, *} p_{\ell \leq 0}^! \mathcal{F}) \longrightarrow \mathrm{FS}_{X \times \mathbb{A}^1}(p_{\mathbb{A}^1}^! \mathcal{F}) \\
 &\quad \wr \downarrow (2-46) \qquad \qquad \qquad \wr \downarrow (4-5) \qquad \qquad \qquad \wr \downarrow (-1) \cdot (2-48) \\
 &p_{\mathbb{A}^1}^* \mathcal{F} \longrightarrow j_{X \times \mathbb{A}^1_{>0}, *} p_{\mathbb{A}^1_{>0}}^* \mathcal{F} \longrightarrow 0_{X/X \times \mathbb{A}^1, *} \mathcal{F}
 \end{aligned}$$

The commutativity of the left rectangle is clear from the construction of the maps (2-46) and (4-5) and the commutativity of the outer rectangle follows from Proposition 2.3. As the lower horizontal arrows become invertible after restricting to the zero section  $X \times \{0\} \subset X \times \mathbb{A}^1$ , we conclude the commutativity of the right rectangle. Now the proof of the commutativity of (7-6) is over.

Next consider the following diagram in  $D_c^b(Z)$ :

$$\begin{aligned}
 (7-10) \quad &\begin{array}{ccccc}
 \pi_{E_Z^\vee, !}(\varphi_{\tilde{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow{(2-27)} & \pi_{E_Z^\vee, !}((\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow{(2-48)} & \mathcal{F}|_Z \\
 \downarrow \wr (2-46) & & \swarrow (2-46) & & \nearrow (2-48) \\
 0_{E_Z}^* \mathrm{FS}_{E_Z^\vee}(\varphi_{\tilde{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow{(2-27)} & 0_{E_Z}^* \mathrm{FS}_{E_Z^\vee}((\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}) & \xrightarrow[\sim]{(2-42)} & 0_{E_Z}^*(\mathrm{FS}_{E^\vee}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z})
 \end{array}
 \end{aligned}$$



The commutativity of the left quadrilateral is clear and the commutativity of the right quadrilateral follows from the commutativity of the diagram (2-51) and Proposition 2.3.

Finally combining the commutativity of the diagram (7-6) restricted to the zero section  $Z \subset E_Z$  and the commutativity of the diagram (7-10), we obtain the commutativity of the diagram (4-17).

Now we move on to the proof of statement (ii). The proof is similar to the proof of statement (i). Consider the following diagram in  $D_c^b(E_Z)$ :

$$(7-11) \quad \begin{array}{ccc} & & 0_{E_Z,*}(\mathcal{F}|_Z^!) \\ & \nearrow (2-48) \sim & \downarrow (3-5) \wr \\ \text{FS}_{E_Z^\vee}((\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}^!) & \xleftarrow{\sim (2-43)} & \text{FS}_{E^\vee}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z}^! & \rightarrow \iota_{C_Z/X,*} \text{Sp}_{Z/X}(\Gamma_Z(\mathcal{F})) \\ & \nwarrow (2-28) & & \downarrow \\ \text{FS}_{E_Z^\vee}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}^!) & \xrightarrow[\sim]{(-1) \cdot (4-1)} & \iota_{C_Z/X,*} \text{Sp}_{Z/X}(\mathcal{F}) \end{array}$$

We will prove the commutativity of this diagram in the same way as the proof of the commutativity of the diagram (7-6). Consider the composition

$$(7-12) \quad \begin{aligned} \text{FS}_{E_Z^\vee}((\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}^!) &\xrightarrow{\sim} \text{FS}_{E_Z^\vee}((\hat{s}^! 0_{X/X \times \mathbb{A}^1,*} \mathcal{F})|_{E_Z^\vee}^!) \\ &\xrightarrow[\sim]{(2-43)} \text{FS}_{E_Z^\vee}(\hat{s}^! 0_{X/X \times \mathbb{A}^1,*} \mathcal{F})|_{E_Z}^! \\ &\xrightarrow[\sim]{(2-48)} (\tilde{s}_* \text{FS}_{X \times \mathbb{A}^1}(0_{X/X \times \mathbb{A}^1,*} \mathcal{F}))|_{E_Z}^! \\ &\xrightarrow[\sim]{(2-46)} (\tilde{s}_* p_{\mathbb{A}^1}^* \mathcal{F})|_{E_Z}^! \cong 0_{E_Z,*}(\mathcal{F}|_Z^!) = 0_{E_Z,*}(\mathcal{F}|_Z^!) = 0_{E_Z,*}(\mathcal{F}|_Z^!). \end{aligned}$$

We have natural maps from the complexes appearing in (7-12) to the complexes appearing in (4-14) at the same stages such that these maps commute with maps appearing in (7-12) and (4-14). Furthermore, the commutativity of the outer rectangle of the diagram (7-9) shows that the composition (7-12) is equal to the following composition after multiplying by  $-1$ :

$$\text{FS}_{E_Z^\vee}((\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}^!) \xleftarrow[\sim]{(2-43)} \text{FS}_{E^\vee}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z}^! \xrightarrow[\sim]{(2-48)} 0_{E_Z,*}(\mathcal{F}|_Z^!).$$

These claims imply the commutativity of the diagram (7-11).

Next consider the following diagram:

$$(7-13) \quad \begin{array}{ccccc} \pi_{E_Z^\vee,*} \varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}^![-2 \text{ rank } E] & \xleftarrow{(2-28)} & \pi_{E_Z^\vee,*}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}^![-2 \text{ rank } E] & \xleftarrow{\sim} & \mathcal{F}|_Z^! \\ & & \nwarrow (2-47) & & \nearrow (-1)^{\text{rank } E} \cdot (2-48) \sim \\ & \wr (2-47) & 0_{E_Z}^! \text{FS}_{E_Z^\vee}((\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}^!) & \xleftarrow[\sim]{(2-43)} & 0_{E_Z}^! \text{FS}_{E^\vee}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z}^! \\ & \downarrow & \nwarrow (2-28) & & \\ 0_{E_Z}^! \text{FS}_{E_Z^\vee}(\varphi_{\bar{s}}(\pi_{E^\vee}^! \mathcal{F})|_{E_Z^\vee}^!) & & & & \end{array}$$

The commutativity of the left quadrilateral is obvious, and the commutativity of the right quadrilateral follows from Propositions 2.2 and 2.3.

Finally, combining the commutativity of diagrams (7-11) and (7-13), we obtain the commutativity of diagram (4-18).

## References

- [Amorim and Ben-Bassat 2017] **L Amorim, O Ben-Bassat**, *Perversely categorified Lagrangian correspondences*, Adv. Theor. Math. Phys. 21 (2017) 289–381 MR Zbl
- [Arinkin and Gaiitsgory 2015] **D Arinkin, D Gaiitsgory**, *Singular support of coherent sheaves and the geometric Langlands conjecture*, Selecta Math. 21 (2015) 1–199 MR Zbl
- [Behrend 1997] **K Behrend**, *Gromov–Witten invariants in algebraic geometry*, Invent. Math. 127 (1997) 601–617 MR Zbl
- [Behrend 2009] **K Behrend**, *Donaldson–Thomas type invariants via microlocal geometry*, Ann. of Math. 170 (2009) 1307–1338 MR Zbl
- [Behrend and Fantechi 1997] **K Behrend, B Fantechi**, *The intrinsic normal cone*, Invent. Math. 128 (1997) 45–88 MR Zbl
- [Ben-Bassat et al. 2015] **O Ben-Bassat, C Brav, V Bussi, D Joyce**, *A “Darboux theorem” for shifted symplectic structures on derived Artin stacks, with applications*, Geom. Topol. 19 (2015) 1287–1359 MR Zbl
- [Borisov and Joyce 2017] **D Borisov, D Joyce**, *Virtual fundamental classes for moduli spaces of sheaves on Calabi–Yau four-folds*, Geom. Topol. 21 (2017) 3231–3311 MR Zbl
- [Brav et al. 2015] **C Brav, V Bussi, D Dupont, D Joyce, B Szendrői**, *Symmetries and stabilization for sheaves of vanishing cycles*, J. Singul. 11 (2015) 85–151 MR Zbl
- [Brav et al. 2019] **C Brav, V Bussi, D Joyce**, *A Darboux theorem for derived schemes with shifted symplectic structure*, J. Amer. Math. Soc. 32 (2019) 399–443 MR Zbl
- [Cao and Leung 2014] **Y Cao, N C Leung**, *Donaldson–Thomas theory for Calabi–Yau 4-folds*, preprint (2014) arXiv 1407.7659
- [Ciocan-Fontanine and Kapranov 2009] **I Ciocan-Fontanine, M Kapranov**, *Virtual fundamental classes via dg-manifolds*, Geom. Topol. 13 (2009) 1779–1804 MR Zbl
- [Davison 2017] **B Davison**, *The critical CoHA of a quiver with potential*, Q. J. Math. 68 (2017) 635–703 MR Zbl
- [Edidin and Graham 1995] **D Edidin, W Graham**, *Characteristic classes and quadric bundles*, Duke Math. J. 78 (1995) 277–299 MR Zbl
- [Efimov 2018] **A I Efimov**, *Cyclic homology of categories of matrix factorizations*, Int. Math. Res. Not. 2018 (2018) 3834–3869 MR Zbl
- [Fantechi and Göttsche 2010] **B Fantechi, L Göttsche**, *Riemann–Roch theorems and elliptic genus for virtually smooth schemes*, Geom. Topol. 14 (2010) 83–115 MR Zbl
- [Fulton 1984] **W Fulton**, *Intersection theory*, Ergebnisse der Math. 2, Springer (1984) MR Zbl
- [Joyce 2015] **D Joyce**, *A classical model for derived critical loci*, J. Differential Geom. 101 (2015) 289–367 MR Zbl

- [Joyce 2018] **D Joyce**, *Shifted symplectic derived algebraic geometry and generalizations of Donaldson–Thomas theory*, lecture notes (2018) Available at <https://people.maths.ox.ac.uk/joyce/SeoulHandout3.pdf>
- [Joyce and Upmeyer 2021] **D Joyce, M Upmeyer**, *Orientation data for moduli spaces of coherent sheaves over Calabi–Yau 3-folds*, *Adv. Math.* 381 (2021) art.id. 107627 MR Zbl
- [Kashiwara and Schapira 1990] **M Kashiwara, P Schapira**, *Sheaves on manifolds*, *Grundle. Math. Wissen.* 292, Springer (1990) MR Zbl
- [Kinjo 2022] **T Kinjo**, *Dimensional reduction in cohomological Donaldson–Thomas theory*, *Compos. Math.* 158 (2022) 123–167 MR Zbl
- [Kresch 1999] **A Kresch**, *Cycle groups for Artin stacks*, *Invent. Math.* 138 (1999) 495–536 MR Zbl
- [Li and Tian 1998] **J Li, G Tian**, *Virtual moduli cycles and Gromov–Witten invariants of algebraic varieties*, *J. Amer. Math. Soc.* 11 (1998) 119–174 MR Zbl
- [Massey 2016] **D Massey**, *Natural commuting of vanishing cycles and the Verdier dual*, *Pacific J. Math.* 284 (2016) 431–437 MR Zbl
- [Oh and Thomas 2023] **J Oh, R P Thomas**, *Counting sheaves on Calabi–Yau 4-folds, I*, *Duke Math. J.* 172 (2023) 1333–1409 MR Zbl
- [Pantev et al. 2013] **T Pantev, B Toën, M Vaquié, G Vezzosi**, *Shifted symplectic structures*, *Publ. Math. Inst. Hautes Études Sci.* 117 (2013) 271–328 MR Zbl
- [Preygel 2015] **A Preygel**, *Ind-coherent complexes on loop spaces and connections*, from “Stacks and categories in geometry, topology, and algebra” (T Pantev, C Simpson, B Toën, M Vaquié, G Vezzosi, editors), *Contemp. Math.* 643, *Amer. Math. Soc.*, Providence, RI (2015) 289–323 MR Zbl
- [Schefers 2022] **K Schefers**, *An equivalence between vanishing cycles and microlocalization*, preprint (2022) arXiv 2205.12436
- [Schürmann 2003] **J Schürmann**, *Topology of singular spaces and constructible sheaves*, *Math. Inst. Polish Acad. Sci. Math. Monogr.* 63, Birkhäuser, Basel (2003) MR Zbl
- [Thomas 2000] **R P Thomas**, *A holomorphic Casson invariant for Calabi–Yau 3-folds, and bundles on  $K3$  fibrations*, *J. Differential Geom.* 54 (2000) 367–438 MR Zbl
- [Toda 2024] **Y Toda**, *Categorical Donaldson–Thomas theory for local surfaces*, *Lecture Notes in Math.* 2350, Springer (2024)
- [Verdier 1983] **J-L Verdier**, *Spécialisation de faisceaux et monodromie modérée*, from “Analysis and topology on singular spaces, II–III”, *Astérisque* 101–102, *Soc. Math. France*, Paris (1983) 332–364 MR Zbl

Research Institute for Mathematical Sciences, Kyoto University  
Kyoto, Japan

`tkinjo@kurims.kyoto-u.ac.jp`

Proposed: Gang Tian

Seconded: Richard P Thomas, Jim Bryan

Received: 3 October 2022

Revised: 31 December 2023

