# Convex integration method and non-uniqueness of weak solutions for viscous fluids

Yachun Li<sup>a</sup>, Peng Qu<sup>b</sup>, Zirong Zeng<sup>a</sup>, Deng Zhang<sup>a</sup>

<sup>a</sup>School of Mathematical Sciences, Shanghai Jiao Tong University. <sup>b</sup>School of Mathematical Sciences, Fudan University.

#### Abstract

We review our recent results concerning the non-uniqueness of weak solutions to viscous fluid dynamics, based on the technique of convex integration. Concerning the hyperdissipative incompressible Navier-Stokes and MHD equations, we proved the sharp non-uniqueness of weak solutions in supercritical regimes with respect to the Ladyženskaja-Prodi-Serrin (LPS) criteria. For the hypo-viscous compressible Naver-Stokes equations, we proved that there exist infinitely many weak solutions with the same initial data, which provides the first non-uniqueness result of weak solutions to viscous compressible fluid.

### 1 Background

We are concerned with the three-dimensional incompressible Navier-Stokes equations (INS for short)

$$\begin{cases} \partial_t u - \nu \Delta u + (u \cdot \nabla) u + \nabla P = 0, \\ \operatorname{div} u = 0, \end{cases}$$
(1.1)

where  $u = (u_1, u_2, u_3)^{\top}(t, x) \in \mathbb{R}^3$  and  $P = P(t, x) \in \mathbb{R}$  represent the velocity field and pressure of the fluid, respectively,  $\nu > 0$  is the viscosity coefficient. In the groundbreaking paper [62], Leray demonstrated the existence of weak solutions to INS in the space  $L_t^{\infty} L_x^2 \cap L_t^2 \dot{H}_x^1$  which obey the energy inequality

$$\|u(t)\|_{L^2}^2 + 2\nu \int_{t_0}^t \|\nabla u(s)\|_{L^2}^2 \mathrm{d}s \le \|u(t_0)\|_{L^2}^2.$$
(1.2)

This solution is now referred to as Leray-Hopf solutions, also due to Hopf [51] in the case of bounded domains. Since then, the uniqueness of Leray-Hopf solutions has remained an challenging open problem in the theory of INS.

One criterion for the well-posedness/ill-posedness is the scaling exponent. It gives a useful classification of subcritical, critical and supercritical spaces. A general philosophy is that equations are well-posed in the subcritical spaces, while solutions may exhibit ill-posedness phenomena in the supercritical spaces. For the INS, the equation is invariant under the scaling

$$u(t,x) \mapsto \lambda u(\lambda^2 t, \lambda x), \quad P(t,x) \mapsto \lambda^2 P(\lambda^2 t, \lambda x),$$
(1.3)

and the critical scaling in the mixed Lebsgue space  $L_t^{\gamma} L_x^p$  is the so-called Ladyženskaja-Prodi-Serrin (LPS for short) condition

$$\frac{2}{\gamma} + \frac{d}{p} = 1 \tag{1.4}$$

with d being the underlying spatial dimension.

It is well-known that weak solutions in the (sub)critical spaces  $L_t^{\gamma} L_x^p$  with  $2/\gamma + 3/p \leq 1$  are unique and even are Leray-Hopf solutions. See [28, 40, 41] and references therein. See also [58, 59] for comprehensive discussions of scaling exponents for general nonlinear PDEs. In contrast, on the flexible side (super-critical regime), based on the convex integration scheme, Buckmaster-Vicol [16] first proved the non-uniqueness of weak solutions to INS. The convex integration approach was introduced to 3D Euler equations by De Lellis and Székelyhidi in the pioneering papers [35, 36]. There have been significant progresses towards the non-uniqueness problem for various fluid models based on convex integrations. One milestone is the resolution of the flexible part of Onsager's conjecture for incompressible Euler equations, developed in [10, 12, 13, 37, 38] and finally settled by Isett [52] and Buckmaster-De Lellis-Székelyhidi-Vicol [14].

The main ingredient introduced by Buckmaster-Vicol [16] is the intermittent building blocks, which in particular permit to control the strong viscosity in INS. The intermittent convex integration also has been applied successfully to various other models. We refer, e.g., to [71] for hyperdissipative INS (1.1), [70] for 2D hypoviscous INS, [73] for stationary INS, and [32, 39] for the non-uniqueness of Leray solutions to hypodissipative INS. See also the surveys [17, 18] for other interesting applications. In particular, in the recent works [15, 77], the intermittent convex integration enables to achieve the regularity close to 1/2, beyond the Onsager regularity 1/3, for non-conservative solutions of 3D Euler equations, and provides a new proof of the flexible side of Onsager's conjecture. See also [49, 50] for the proof of the strong Onsager's conjecture.

In the recent remarkable paper [28], Cheskidov-Luo proved the sharp non-uniqueness of weak solutions in view of the LPS condition to INS in the spaces  $L_t^{\gamma} L_x^{\infty}$  for any  $1 \leq \gamma < 2$ . Moreover, in the 2D case, they also proved the sharp non-uniqueness in  $L_t^{\infty} L_x^p$ , where 1 , which is another endpoint caseof the LPS condition. Very recently, for the 3D hyperdissipative INS with viscosity beyond the Lionsexponent 5/4, we proved the sharp non-uniqueness at two endpoints of the LPS condition [63].

We also refer to another programme by Jia and Šverák [53, 54] towards the non-uniqueness of Leray-Hopf solutions, under a certain assumption for the linearized Navier-Stokes operator. Recently, Albritton-Brué-Colombo [1] constructed the non-uniqueness of Leray solutions to the forced incompressible INS. See also [2] for the case of bounded domain and [3] for the case of forced hypo-viscous INS in dimension two.

The compressible models is also one of the famed models of fluid dynamics and has been extensively studied in literature. For the Euler equations of inviscid compressible fluid, although the uniqueness and stability results have been well established in the 1D case with small BV initial data and mild assumptions (see, e.g., [8, 69]), it was a long standing open problem in the multi-dimensional case until it was observed by De Lellis-Székelyhidi [36], that non-unique bounded entropy solutions can be constructed for multi-dimensional compressible Euler equations. Afterwards, several achievements have been obtained and we refer to [9, 22, 29–31, 48, 60, 72, 74] and the references therein for the flexibility of compressible Euler equations.

In contrast to the above extensive studies of compressible Euler equations and INS, the non-uniqueness of weak solutions to compressible Navier-Stokes equations remains a challenging problem.

Besides the Navier-Stokes equations, non-uniqueness and related turbulence of MHD equations also attract significant interests in literature. Faraco-Lindberg [42] first constructed non-vanishing smooth strict subsolutions to 3D ideal MHD. Afterwards, Faraco-Lindberg-Székelyhidi [44, 45] constructed infinitly many bounded solutions with prescribed total energy and cross helicity. Based on the convex integration via staircase laminates, they also solve the conjecture that  $L_{t,x}^3$  is the threshold for magnetic helicity conservation in [45]. Based on the intermittent convex integration scheme, Beekie-Buckmaster-Vicol [6] gave the first example of weak solutions with non-conserved magnetic helicity. It was also proved in [6] that Taylor's conjecture fails for the ideal MHD equations.

Concerning the non-uniqueness problem for the viscous and resistive MHD, different from the case of INS, the specific geometry of MHD restricts the choice of oscillation directions, and thus, in particular, limits the spatial intermittency of building blocks in the convex integration scheme. So, the control of viscosity and resistivity in (4.1) becomes significantly hard. In [66], we provided the first examples of sharp non-uniqueness for MHD equations near one endpoint of the critical-scaling LPS condition. This in particular extends the recent sharp non-uniqueness result in [28] in the context of Navier-Stokes equations. Another interesting phenomena observed in [66] is that the scaling-invariant LPS condition

serves as the criterion to detect non-uniqueness: the uniqueness would fail even in the high viscous and resistive regime beyond the Lions exponent, if the (sub)criticality of state space is violated. We also refer to [34] for the case of Hall MHD equations, [65] for MHD equations with the viscosity and resistivity up to the Lions exponent and [76] for the non-uniqueness in  $C_t L_x^2$  space. See also [24] and [25] for other ill-posedness phenomena.

**Organization of paper.** In  $\S2$  we present our recent results on the sharp non-uniqueness of weak solutions in view of the LPS criterion to hyperdissipative Navier-Stokes equations [63]. Then,  $\S3$  is devoted to the results of the non-uniqueness of weak solutions to hypoviscous compressible Navier-Stokes equations [64]. Finally, in  $\S4$  we review the results on the non-uniqueness of weak solutions to viscous and resistive MHD equations [65, 66].

## 2 Incompressible Navier-Stokes equations

In this section, we review our recent result concerning the sharp non-uniqueness of weak solutions to the following hyperdissipative Navier-Stokes equations [63] on the torus  $\mathbb{T}^3 := [-\pi, \pi]^3$ :

$$\begin{cases} \partial_t u + \nu (-\Delta)^{\alpha} u + (u \cdot \nabla) u + \nabla P = 0, \\ \operatorname{div} u = 0, \end{cases}$$
(2.1)

where  $u = (u_1, u_2, u_3)^{\top}(t, x) \in \mathbb{R}^3$  and  $P = P(t, x) \in \mathbb{R}$  represent the velocity field and pressure of the fluid, respectively,  $\nu > 0$  is the viscosity coefficient,  $\alpha \in [1, 2)$  is the viscosity exponent, and the hyperdissipation operator  $(-\Delta)^{\alpha}$  is defined on the flat torus via the Fourier transform

$$\mathcal{F}((-\Delta)^{\alpha}u)(\xi) = |\xi|^{2\alpha}\mathcal{F}(u)(\xi), \quad \xi \in \mathbb{Z}^3.$$

To begin with, let us formulate precisely the definition of weak solutions to equation (2.1).

**Definition 2.1.** (Weak solutions) Given any weakly divergence-free initial datum  $u_0 \in L^2(\mathbb{T}^3)$ , we say that  $u \in L^2([0,T] \times \mathbb{T}^3)$  is a weak solution for the hyperdissipative Navier-Stokes equations (2.1) if u is divergence-free for a.e.  $t \in [0,T]$ , and

$$\int_{\mathbb{T}^3} u_0 \varphi(0, x) \mathrm{d}x = -\int_0^T \int_{\mathbb{T}^3} u(\partial_t \varphi - \nu(-\Delta)^\alpha \varphi + (u \cdot \nabla)\varphi) \mathrm{d}x \mathrm{d}t$$
(2.2)

for any divergence-free test function  $\varphi \in C_0^{\infty}([0,T] \times \mathbb{T}^3)$ .

The uniqueness of weak solutions to (2.1) in the critical mixed Lebesgue space  $L_t^{\gamma} W_x^{s,p}$  is contained in Theorem 2.2 below.

**Theorem 2.2** (Uniqueness of weak solutions in  $L_t^{\gamma} W_x^{s,p}$ , [63]). Let  $u \in L_t^{\gamma} W_x^{s,p}$  and  $(s, \gamma, p)$  satisfy

$$\frac{2\alpha}{\gamma} + \frac{3}{p} = 2\alpha - 1 + s, \qquad (2.3)$$

with  $\alpha \ge 1$ ,  $s \ge 0$ ,  $2 \le \gamma \le \infty$ ,  $1 \le p \le \infty$  and  $0 \le \frac{1}{p} - \frac{s}{3} \le \frac{1}{2}$ . If u is a weak solution to (2.1) in the sense of Definition 2.1, then u is a unique Leray-Hopf solution.

**Main results.** Our mains results include the sharp non-uniqueness at two LPS endpoints, the strong nonuniqueness for the high dissipativity beyond the Lions exponent, the partial regularity of weak solutions, and the vanishing viscosity limit.

(i) Sharp non-uniqueness at two LPS endpoints. We focus on the non-uniqueness of weak solutions in the following two supercritical regimes, whose borderlines contain two endpoints of the generalized LPS condition (2.3).

More precisely, in the case  $\alpha \in [5/4, 2)$  we consider the supercritical regime  $\mathcal{A}_1$  given by

$$\mathcal{A}_1 := \left\{ (s, \gamma, p) \in [0, 3) \times [1, \infty] \times [1, \infty] : 0 \le s < \frac{4\alpha - 5}{\gamma} + \frac{3}{p} + 1 - 2\alpha \right\},\tag{2.4}$$

and in the case  $\alpha \in [1,2)$  we consider supercritical regime  $\mathcal{A}_2$  given by

$$\mathcal{A}_{2} := \left\{ (s, \gamma, p) \in [0, 3) \times [1, \infty] \times [1, \infty] : 0 \le s < \frac{2\alpha}{\gamma} + \frac{2\alpha - 2}{p} + 1 - 2\alpha \right\}.$$
 (2.5)

The supercritical regimes  $\mathcal{A}_1$  and  $\mathcal{A}_2$  in the case s = 0 can be seen in Figure 1 below.



Figure 1: The supercritical regimes  $\mathcal{A}_1$  and  $\mathcal{A}_2$  ( $\alpha \in [\frac{5}{4}, 2), s = 0$ )

Concerning the case of  $\alpha = 1$ , Cheskidov-Luo [28] proved that for any  $\gamma < 2$ , there exist nonunique solutions in  $L_t^{\gamma} L_x^{\infty}$  to INS in all dimensions  $d \geq 2$ , which is sharp at the endpoint case of  $(s, \gamma, p) = (0, 2, \infty)$ . The sharp non-uniqueness for the other endpoint case  $(s, \gamma, p) = (0, \infty, 2)$  was also achieved in [26] for the 2D INS.

When  $\alpha \geq 1$ , the sharp non-unqueness of hyperdissipative Navier-Stokes equations (2.1) is presented in Theorem 2.3 below.

**Theorem 2.3.** (Sharp non-uniqueness, [63]) Let  $\alpha \in [5/4, 2)$ . Then, for any weak solution  $\tilde{u}$  to (2.1), there exists a different weak solution  $u \in L_t^{\gamma} W_x^{s,p}$  to (2.1) with the same initial data, where  $(s, \gamma, p) \in \mathcal{A}_1 \cup \mathcal{A}_2$ .

In particular, when  $\alpha = 5/4$ , Theorem 2.3 provides the non-uniqueness of weak solutions in  $C_t L_x^p$  to (2.1) for any p < 2, which is sharp in view of Theorem 2.2.

Furthermore, from the viewpoint of the LPS criteria (2.3), Theorem 2.3 also provides the sharp nonuniqueness for the hyperdissipative INS (2.1) at two endpoints, i.e.,  $(3/p + 1 - 2\alpha, \infty, p)$  for  $\alpha \in [5/4, 2)$ , and  $(2\alpha/\gamma + 1 - 2\alpha, \gamma, \infty)$  for  $\alpha \in (1, 2)$ .

We would also expect the non-uniqueness in the supercritical regimes where  $\alpha \in [2, 5/2)$  or near the endpoint  $(3/p+1-2\alpha, \infty, p)$  when  $\alpha \in [1, 5/4)$ , which probably require a new convex integration scheme different from the existing  $L^2_{t,x}$ -critical convex integration. The reason is that, in the present  $L^2_{t,x}$ -critical scheme, the temporal intermittency has been explored to upgrade temporal integrability by allowing  $\gamma > 2$ , but this in turn requires to reduce the spatial integrability with p < 2. However, in the endpoint

82

case  $(0, \infty, 3/(2\alpha - 1))$  where  $\alpha < 5/4$ , both the temporal and spatial integrability exponents are larger than two, which is thus beyond the present  $L_{t,x}^2$  scheme.

(ii) Strong non-uniqueness for the high dissipativity beyond the Lions exponent. Besides the sharp non-uniqueness, the strong non-uniqueness of weak solutions to (2.1) also holds in the hyperdissipative case where  $\alpha \in [5/4, 2)$ , which is formulated in Theorem 2.4 below.

**Theorem 2.4.** (Strong non-uniqueness, [63]) Let  $\alpha \in [5/4, 2)$ , for every divergence-free  $L_x^2$  initial data, there exist infinitely many weak solutions in  $L_t^{\gamma} W_x^{s,p}$  to (2.1) which are smooth almost everywhere in time.

It is folklore that one has global solvability in the high dissipative case when  $\alpha \geq 5/4$ . In fact, based on the works [11, 68, 71], the critical threshold for the  $C_t L_x^2$  well-posedness for (2.1) is precisely at  $\alpha = 5/4$ . This indicates that weak solutions in  $C_t L_x^2$  to (2.1) are unique if  $\alpha$  is greater than or equal to 5/4, but they become non-unique if  $\alpha$  is less than 5/4.

Theorem 2.4 shows that the non-unique weak solutions still exhibit in the high-dissipative regime where  $\alpha \geq 5/4$ . More specifically, it reveals that the uniqueness breaks down in the space  $L_t^{\gamma} W_x^{s,p}$ , if the exponents  $(s, \gamma, p)$  lies in the supercritical regime  $\mathcal{A}_1 \cup \mathcal{A}_2$ , defined in (2.4) and (2.5), respectively.

(iii) Partial regularity of weak solutions. In the pioneering paper [62], Leray proves that the Leray-Hopf solutions to INS are smooth outside a closed singular set of times, which has zero Hausdorff  $\mathcal{H}^{1/2}$  measure. This provides an alternative approach to address the global existence problem. Following the works of Scheffer [79, 80], Caffarelli-Kohn-Nirenberg [19] proved a space-time regularity version and showed the existence of global Leray-Hopf solutions which have singular sets in  $\mathbb{R}^+ \times \mathbb{R}^3$  of zero Hausdorff  $\mathcal{H}^1$  measure. The simplified proofs were obtained by Lin [67] and Vasseur [83]. In [47], Giga established partial regularity of solutions in mixed  $L_t^q L_x^p$  spaces. For hyperdissipative INS with  $\alpha \in (1, 5/4]$ , Katz-Pavlović [57] proved that the Hausdorff dimension of singular set at the time of first blow-up is at most  $5 - 4\alpha$ . More recently, Colombo-De Lellis-Massaccesi [33] proved a stronger version of the Katz-Pavlović result and establish the existence of Leray-Hopf solutions with singular space-time sets of zero Hausdorff  $\mathcal{H}^{5-4\alpha}$  measure, thus extending the Caffarelli-Kohn-Nirenberg theorem to hyper-dissipative INS.

Concerning the partial regularity of weak solutions to hyper-dissipative INS, in [63, Theorem 1.2] we prove that, in the highly dissipative regime  $\alpha \in [5/4, 2)$ , for any  $\eta_* > 0$ , there exist weak solutions to (2.1) in any small  $L_t^{\gamma} W_x^{s,p}$ -neighborhood of Leray-Hopf solutions, where  $(s, \gamma, p) \in \mathcal{A}_1 \cup \mathcal{A}_2$ . The weak solutions coincide with Leray-Hopf solutions near t = 0, and have singular temporal sets with zero Hausdorff  $\mathcal{H}^{\eta_*}$ measure.

(iv) Vanishing viscosity limit. Theorem 2.5 below contains the vanishing viscosity limit result, which relates INS and Euler equations.

**Theorem 2.5.** (Strong vanishing viscosity limit) Let  $\alpha \in (1,2)$  and  $u \in H^{\widetilde{\beta}}([-2T,2T] \times \mathbb{T}^3)$  ( $\widetilde{\beta} > 0$ ) be any mean-free weak solution to the Euler equation,

$$\begin{cases} \partial_t u + (u \cdot \nabla)u + \nabla P = 0, \\ \operatorname{div} u = 0. \end{cases}$$
(2.6)

Then, there exist  $\beta' \in (0, \tilde{\beta})$  and a sequence of weak solutions  $u^{(\nu_n)} \in H_{t,x}^{\beta'}$  to (2.1), where  $\nu_n$  is the viscosity coefficient, such that as  $\nu_n \to 0$ ,

$$u^{(\nu_n)} \to u \quad strongly \ in \ H^{\beta'}_{t,x}.$$
 (2.7)

Theorem 2.5 shows that, in the  $H_{t,x}^{\tilde{\beta}}$  topology, the set of accumulation points of weak solutions to hyper-dissipative INS (2.1) contain all the weak solutions in  $H_{t,x}^{\tilde{\beta}}$  to the Euler equations, where  $\tilde{\beta}$  can be any small positive constant. Thus, as in the INS context [16], being a strong limit of weak solutions to the hyper-disspative INS cannot serve as a selection criteria for weak solutions to Euler equations.

## 3 Compressible Navier-Stokes equations

This section presents the results in [64] about the non-uniqueness of weak solutions to hypo-viscous compressible Navier-Stokes equations. We consider the *d*-dimensional isentropic hypo-viscous compressible Navier-Stokes equations (CNS for short) on the torus  $\mathbb{T}^d := [-\pi, \pi]^d$ ,

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \mu(-\Delta)^{\alpha} u - (\mu + \nu) \nabla \operatorname{div} u + \operatorname{div}(\rho u \otimes u) + \nabla P(\rho) = 0. \end{cases}$$
(3.1)

where  $d \geq 2$ ,  $\rho : [0,T] \times \mathbb{T}^d \to (0,\infty)$  and  $u : [0,T] \times \mathbb{T}^d \to \mathbb{R}^d$  are the mass density and velocity of fluid, respectively. The pressure  $P(\rho)$  is  $C^2$  with respect to  $\rho$ . The coefficients  $\mu$  and  $\nu$  are constants,  $\mu$  is the shear viscosity coefficient,  $\nu + \frac{2}{d}\mu$  is the bulk viscosity coefficient satisfying the physical assumptions

$$\mu > 0, \quad \nu + \frac{2}{d}\mu \ge 0.$$
 (3.2)

The hypo-viscosity  $(-\Delta)^{\alpha}$ ,  $\alpha \in (0,1)$ , is defined by the Fourier transform

$$\mathcal{F}((-\Delta)^{\alpha}u)(\xi) = |\xi|^{2\alpha}\mathcal{F}(u)(\xi), \quad \xi \in \mathbb{Z}^d.$$

We note that, (3.1) is the classical compressible Navier-Stokes equations when  $\alpha = 1$ , and when the viscosities vanish ( $\mu = \nu = 0$ ), (3.1) reduces to the compressible Euler equations

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla P(\rho) = 0. \end{cases}$$
(3.3)

It would also be convenient to formulate (3.1) in terms of the density and momentum  $m := \rho u$  as follows

$$\begin{cases} \partial_t \rho + \operatorname{div} m = 0, \\ \partial_t m + \mu(-\Delta)^{\alpha}(\rho^{-1}m) - (\mu + \nu)\nabla \operatorname{div}(\rho^{-1}m) + \operatorname{div}(\rho^{-1}m \otimes m) + \nabla P(\rho) = 0. \end{cases}$$
(3.4)

The weak solutions to (3.4) is understood in the distributional sense.

**Definition 3.1.** (Weak solutions) Let  $0 < T < \infty$ . Given any initial data  $\rho_0 \in L^{\infty}(\mathbb{T}^d)$ ,  $\rho_0 > 0$ , and  $u_0 \in L^2(\mathbb{T}^d)$ , we say that  $(\rho, m) \in L^{\infty}([0, T] \times \mathbb{T}^d) \times L^2((0, T) \times \mathbb{T}^d)$  is a weak solution to (3.4) if

•  $\rho \geq 0$  a.e. and

$$\int_{\mathbb{T}^d} \rho_0(x)\varphi(0,x)\mathrm{d}x = -\int_0^T \int_{\mathbb{T}^d} \rho \partial_t \varphi + (m \cdot \nabla)\varphi \mathrm{d}x \mathrm{d}t$$

for any test function  $\varphi \in C_0^{\infty}([0,T] \times \mathbb{T}^d)$ .

• m = 0 whenever  $\rho = 0$ , and

$$\int_{\mathbb{T}^d} m_0 \varphi(0, x) \mathrm{d}x = -\int_0^T \int_{\mathbb{T}^d} m \cdot \partial_t \varphi - \frac{1}{\rho} m \cdot (\mu(-\Delta)^\alpha \varphi - (\mu + \nu) \nabla \mathrm{div}\varphi) + (\frac{1}{\rho} m \otimes m) : \nabla \varphi + P \mathrm{div}\varphi \mathrm{d}x \mathrm{d}t$$

for any test function  $\varphi \in C_0^{\infty}([0,T) \times \mathbb{T}^d)$ , where  $m_0 := \rho_0 u_0$ .

**Main results** Our main results include the non-uniqueness of weak solutions, the sharp viscosity threshold for  $L_t^2 C_x$  well-posedness, and the vanishing viscosity limit.

(i) Non-uniqueness of weak solutions. The non-uniqueness of weak solutions to the hypo-viscous CNS (3.4) is formulated in Theorem 3.2 below.

**Theorem 3.2** (Non-uniqueness for hypo-viscous CNS). There exist  $\rho_0 \in L^{\infty}(\mathbb{T}^d)$ ,  $\rho_0 > 0$ , and  $m_0 \in L^2(\mathbb{T}^d)$ , such that for any exponents (p, s) satisfying

$$\alpha + s - \frac{2\alpha}{p} < 0, \quad (p, s) \in [1, 2] \times [0, 1),$$
(3.5)

there exist infinitely many weak solutions  $(\rho, m) \in C_t C_x^1 \times L_t^p C_x^s$  to the hypo-viscous CNS (3.4) with the same initial data  $(\rho_0, m_0)$ .

To the best of our knowledge, Theorem 3.2 provides the first result for the non-uniqueness of weak solutions to the compressible fluid with viscosity. It also shows that, the hypo-vsicosity cannot rule out the non-uniqueness mechanism, which, actually, coincides with the Ladyženskaja-Prodi-Serrin criteria (see also result (ii) below).

The proof of Theorem 3.2 in [64] also applies to the case of compressible Euler equations (3.3).

**Theorem 3.3** (Non-uniqueness for compressible Euler equations, [64]). There exist  $\rho_0 \in L^{\infty}(\mathbb{T}^d)$  and  $m_0 \in L^2(\mathbb{T}^d)$ , such that for any exponents (p, s) satisfying (3.5) with any  $\alpha \in (0, 1)$ , there exist infinitely many weak solutions  $(\rho, m) \in C_t C_x^1 \times L_t^p C_x^s$  to the compressible Euler equations (3.3) with the initial data  $(\rho_0, m_0)$ .

Theorem 3.3 provides new examples of anomalous dissipation in the class  $L_t^1 C_x^{1-}$  for the compressible Euler equations, by taking p = 1 and  $\alpha$  close to 1. We note that a similar result for the incompressible Euler equations was provided in [28] with a different convex integration scheme.

(ii) Sharp viscosity threshold for  $L_t^2 C_x$  well-posedness. As in the proof of Theorem 3.2, we also have the non-uniqueness result for hypo-viscous INS.

**Theorem 3.4** (Non-uniqueness for hypo-viscous INS). Consider the incompressible Navier-Stokes equations

$$\begin{cases} \partial_t u + \mu (-\Delta)^{\alpha} u + \operatorname{div}(u \otimes u) + \nabla P = 0, \\ \operatorname{div} u = 0, \end{cases}$$
(3.6)

where  $\mu > 0$ ,  $\alpha \in (0,1)$ . Then, there exists  $u_0 \in L^2(\mathbb{T}^d)$  such that for any exponents (p,s) satisfying (3.5), there exist infinitely many weak solutions  $u \in L^p_t C^s_x$  with the initial datum  $u_0$ .

When (p, s) = (2, 0), Theorem 3.4 provides the non-uniqueness of weak solutions in the space  $L_t^2 C_x$ for the hypo-viscous INS, which is sharp due to the well-posedness in  $L_t^2 L_x^{\infty}$ , in view of the Ladyženskaja-Prodi-Serrin criteria.

Thus, Theorem 3.4 reveals that  $\alpha = 1$  is the sharp threshold viscosity for the  $L_t^2 C_x$  well-posedness for the INS. It would serve as a counterpart of the recent results in [11, 71] regarding the sharp threshold of the Lions exponent  $\alpha = 5/4$  for the  $C_t L_x^2$  well-posedness of the 3D INS. It is worth noting that, both the spaces  $L_t^2 C_x$  and  $C_t L_x^2$  are the endpoint spaces of the Ladyženskaja-Prodi-Serrin criteria.

This also complements the very recent work [3], where the non-uniqueness of Leray-Hopf solutions is proved for the 2-D hypo-viscous INS with a force.

(iii) Vanishing viscosity limit. The following strong vanishing viscosity limit relates the hypoviscous CNS with compressible Euler equations.

**Theorem 3.5** (Strong vanishing viscosity limit for hypo-viscous CNS). Let  $\alpha \in (0,1)$  and  $\rho, m \in C_{t,x}^{\tilde{\beta}}$ ,  $\tilde{\beta} > 0$ , be any weak solution to the compressible Euler equations (3.3), such that  $c_1 \leq \rho \leq c_2$  for some

constants  $c_1, c_2 > 0$ . Then, there exist  $\beta' \in (0, \tilde{\beta})$  and a sequence of weak solutions  $(\rho^{(n)}, m^{(n)}) \in C_{t,x} \times H_t^{\beta'} C_x$  to the hypo-viscous CNS (3.1) with viscous coefficients  $\mu = \kappa_n \mu_*$  and  $\nu = \kappa_n \nu_*$ , such that as  $\kappa_n \to 0$ ,

 $\rho^{(n)} \to \rho \quad \text{strongly in } C_{t,x}, \quad \text{and} \quad m^{(n)} \to m \quad \text{strongly in } H_t^{\beta'} C_x.$ (3.7)

Regarding the compressible case, the 1D case with artificial viscosity was solved by Bianchini-Bressan [7]. But it is still open for the multi-dimensional case with general initial data. By virtue of Theorem 3.5, we see that in the vanishing viscosity limit, the set of accumulation points of weak solutions to the hypo-viscous CNS (3.4) contains all the Hölder continuous weak solutions to the compressible Euler equations (3.3).

In particular, the entropy solutions in the Hölder spaces recently constructed in [48] may be obtained as the strong vanishing viscosity limits of weak solutions to the hypo-viscous CNS (3.1) when d = 3.

#### 4 Incompressible MHD equations

In this section, we review our recent results in [65, 66] concerning the non-uniqueness of weak solutions to the following three-dimensional viscous and resistive magnetohydrodynamic (MHD for short) system on the torus  $\mathbb{T}^3 := [-\pi, \pi]^3$ :

$$\begin{cases} \partial_t u - \nu_1 \Delta u + (u \cdot \nabla) u - (B \cdot \nabla) B + \nabla P = 0, \\ \partial_t B - \nu_2 \Delta B + (u \cdot \nabla) B - (B \cdot \nabla) u = 0, \\ \operatorname{div} u = 0, \quad \operatorname{div} B = 0, \end{cases}$$
(4.1)

where  $u = (u_1, u_2, u_3)^{\top}(t, x) \in \mathbb{R}^3$ ,  $B = (B_1, B_2, B_3)^{\top}(t, x) \in \mathbb{R}^3$  and  $P = P(t, x) \in \mathbb{R}$  correspond to the velocity field, magnetic field and pressure of the fluid, respectively, and  $\nu_1, \nu_2 \ge 0$  are the viscous and resistive coefficients, respectively. In particular, in the case without magnetic fields, system (4.1) reduces to the classical INS.

In order to formulate the main results more generally, we consider the MHD equations with hyper viscosity and resistivity:

$$\begin{cases} \partial_t u + \nu_1 (-\Delta)^{\alpha} u + (u \cdot \nabla) u - (B \cdot \nabla) B + \nabla P = 0, \\ \partial_t B + \nu_2 (-\Delta)^{\alpha} B + (u \cdot \nabla) B - (B \cdot \nabla) u = 0, \\ \operatorname{div} u = 0, \quad \operatorname{div} B = 0, \end{cases}$$
(4.2)

where  $\alpha \in [1, 3/2)$ . Note that, the viscosity and resistivity exponents can be larger than the Lions exponent 5/4.

System (4.2) is invariant under the scaling

$$u(t,x) \mapsto \lambda^{2\alpha-1} u(\lambda^{2\alpha} t, \lambda x), \quad B(t,x) \mapsto \lambda^{2\alpha-1} B(\lambda^{2\alpha} t, \lambda x), \tag{4.3}$$

and  $P(t, x) \mapsto \lambda^{4\alpha-2} P(\lambda^{2\alpha}t, \lambda x)$ . The critical exponent  $(s, \gamma, p)$  of the mixed Lebesgue spaces  $L_t^{\gamma} W_x^{s, p}$  satisfies the generalized Ladyženskaja-Prodi-Serrin condition

$$\frac{2\alpha}{\gamma} + \frac{3}{p} = 2\alpha - 1 + s. \tag{4.4}$$

We show that the non-uniqueness of weak solutions also exhibit in the spaces  $L_t^{\gamma} W_x^{s,p}$ , where the exponents  $(s, \gamma, p)$  lie in the following two supercritical regimes, with respect to the scaling (4.3):

$$S_1 := \left\{ (s, \gamma, p) \in [0, 3) \times [1, \infty] \times [1, \infty] : 0 \le s < \frac{2\alpha}{\gamma} + \frac{2\alpha - 2}{p} + 1 - 2\alpha \right\},\tag{4.5}$$

and

$$S_2 := \left\{ (s, \gamma, p) \in [0, 3) \times [1, \infty] \times [1, \infty] : 0 \le s < \frac{4\alpha - 4}{\gamma} + \frac{2}{p} + 1 - 2\alpha \right\}.$$
(4.6)

The weak solutions to (4.2) are taken similarly in the distributional sense.

**Definition 4.1** (Weak solutions). Given any divergence free initial data  $(u_0, B_0) \in L^2(\mathbb{T}^3)$ , we say that  $(u, B) \in L^2([0, T] \times \mathbb{T}^3)$  is a weak solution to (4.2) if

- For all  $t \in [0,T]$ ,  $(u(t,\cdot), B(t,\cdot))$  are divergence free in the sense of distributions and have zero spatial mean.
- Equations (4.2) hold in the sense of distributions, i.e., for any divergence-free test functions  $\varphi \in C_0^{\infty}([0,T] \times \mathbb{T}^3)$ ,

$$\int_{\mathbb{T}^3} u_0 \varphi(0, x) \mathrm{d}x + \int_0^T \int_{\mathbb{T}^3} \partial_t \varphi \cdot u - \nu_1 (-\Delta)^\alpha \varphi \cdot u + \nabla \varphi : (u \otimes u - B \otimes B) \mathrm{d}x \mathrm{d}t = 0,$$
  
$$\int_{\mathbb{T}^3} B_0 \varphi(0, x) \mathrm{d}x + \int_0^T \int_{\mathbb{T}^3} \partial_t \varphi \cdot B - \nu_2 (-\Delta)^\alpha \varphi \cdot B + \nabla \varphi : (B \otimes u - u \otimes B) \mathrm{d}x \mathrm{d}t = 0.$$

Concerning the weak solutions to (4.1) in the sense of Definition 4.1, we have the following sharp non-uniqueness result.

**Theorem 4.2** (Sharp non-uniqueness for MHD). Consider the viscous and resistive MHD (4.1). Then, for any  $1 \leq \gamma < 2$ , there exist infinitely many weak solutions in  $L_t^{\gamma} L_x^{\infty}$  with the same initial data.

**Remark 4.3.** One significant difference between the INS and MHD is the impact of the strong coupling between velocity and magnetic fields in MHD, which limits the available choices for oscillating directions. As a result, constructing spatial building blocks with strong spatial intermittency that align with the specific geometric structure of MHD becomes quite challenging.

It should be mentioned that, the 1D spatial intermittency and intermittent flows were first constructed in [6]. In [65, 66], the 2D intermittent flows were constructed. However, both intermittent flows are not strong enough to control the viscosity  $-\Delta$ . The novelty in [66] is to explore the additional temporal intermittency.

Another distinction between INS and MHD can be seen in the construction of the amplitudes of velocity and magnetic perturbations. This distinction arises from the anti-symmetry of magnetic nonlinearity, which requires a second geometric lemma in a small neighbourhood of the null matrix. This is different from the geometric lemma used for velocity perturbations, which holds in the neighborhood of the identity matrix. Additionally, when constructing velocity perturbations, a new matrix  $\mathring{G}^B$  needs to be introduced, which does not appear in the context of the Navier-Stokes equations. These differences lead to specific algebraic identities for the nonlinear effects associated with magnetic and velocity perturbations.

Besides the non-uniqueness results, it also holds the following strong vanishing viscosity and resistivity limits for the MHD equations (4.1). Its relationship to the Taylor's conjecture will be discussed in Remark 4.5 below.

**Theorem 4.4** (Strong vanishing viscosity and resistivity limit). Let  $\alpha \in [1, 3/2)$  and  $\tilde{\beta} > 0$ . Let  $(u, B) \in H_{t,x}^{\tilde{\beta}} \times H_{t,x}^{\tilde{\beta}}$  be any mean-free weak solution to the ideal MHD. Then, there exist  $\beta' \in (0, \tilde{\beta})$  and a sequence of weak solutions  $(u^{(\nu_n)}, B^{(\nu_n)}) \in H_{t,x}^{\beta'} \times H_{t,x}^{\beta'}$  to the hyper viscous and resistive MHD (4.2), where  $\nu_n = (\nu_{1,n}, \nu_{2,n})$  and  $\nu_{1,n}, \nu_{2,n}$  are the viscosity and resistivity coefficients, respectively, such that as  $\nu_n \to 0$ ,

$$(u^{(\nu_n)}, B^{(\nu_n)}) \to (u, B) \quad strongly \ in \ H^{\beta'}_{t,x} \times H^{\beta'}_{t,x}. \tag{4.7}$$

**Remark 4.5.** It would be interesting to see the relationship between the Taylor's conjecture and the vanishing viscosity and resistivity limit in Theorem 4.4.

The smooth solution to the ideal MHD has several global invariants:

- The total energy:  $\mathcal{E}(t) = \frac{1}{2} \int_{\mathbb{T}^3} |u(t,x)|^2 + |B(t,x)|^2 dx;$
- The cross helicity:  $\mathcal{H}_{\omega,B}(t) = \int_{\mathbb{T}^3} u(t,x) \cdot B(t,x) dx;$
- The magnetic helicity:  $\mathcal{H}_{B,B}(t) := \int_{\mathbb{T}^3} A(t,x) \cdot B(t,x) dx.$

Here A is a mean-free periodic vector field satisfying  $\operatorname{curl} A = B$ .

It is widely acknowledged in the field of plasma physics that magnetic helicity remains conserved as the conductivity approaches infinity. This phenomenon is known as Taylor's conjecture. It has been proved by Faraco-Lindberg [43] that Taylor's conjecture holds true when dealing with the weak limits of Leray-Hopf solutions MHD in simply connected, magnetically closed domain. See also Faraco-Lindberg-MacTaggart-Valli [46] in the case of multiply connected domains.

On the other hand, Beekie-Buckmaster-Vicol [6] demonstrated that Taylor's conjecture does not holds by constructing distributional solutions in  $C_t L_x^2$  breaking the magnetic helicity conservation. The delicate point here is that the conservation of magnetic helicity demands a level of regularity that is much milder than that of total energy and cross helicity. Specifically, the weak solution in  $B_{3,\infty}^{\alpha}$  with  $\alpha > 1/3$  or  $B_{3,c(\mathbb{N})}^{1/3}$ conserve the energy and cross helicity, while the magnetic helicity is conserved in the less regular space  $B_{3,\infty}^{\alpha}$  with  $\alpha > 0$  or in the endpoint space  $L_{t,x}^3$ . See [4, 20, 42, 55].

Theorem 4.4 shows that, in contrast to weak ideal limits, even for the hyper viscous and resistive MHD beyond the Lions exponent, there exists certain sequence of non-Leray-Hopf weak solutions such that the Taylor's conjecture fails in the limit along this sequence.

Acknowledgment. Y. Li thanks the support by NSFC under grants 11831011, 12161141004, and 12371221. P. Qu thanks the supports by NSFC under grants 12122104 and 11831011, and by Shanghai Science and Technology Programs 21ZR1406000 and 21JC1400600. Z. Zeng thanks the support by NSFC under grants 12271352 and 12371221. D. Zhang thanks the supports by NSFC under grants 12271352, 12322108, and 12161141004, and by Shanghai Rising-Star Program 21QA1404500. Y. Li and D. Zhang are also grateful for the supports by Shanghai Frontiers Science Center of Modern Analysis and the Fundamental Research Funds for the Central Universities.

#### References

- D. Albritton, E. Brué, and M. Colombo. Non-uniqueness of Leray solutions of the forced Navier-Stokes equations. Ann. of Math.(2), 196(1):415-455, 2022.
- [2] D. Albritton, E. Brué, and M. Colombo. Gluing non-unique Navier-Stokes solutions. arXiv:2209.03530, 2022.
- [3] D. Albritton and M. Colombo. Non-uniqueness of Leray solutions to the hypodissipative Navier-Stokes equations in two dimensions. arXiv:2209.08713, 2022.
- [4] H. Aluie. Hydrodynamic and magnetohydrodynamic turbulence: Invariants, cascades, and locality. Ph. D. thesis, Johns Hopkins University, 2009.
- [5] C. Bardos, P. Gwiazda, A. Świerczewska-Gwiazda, E. S. Titi, and E. Wiedemann. Onsager's conjecture in bounded domains for the conservation of entropy and other companion laws. *Proc. A.*, 475(2230):20190289, 18 pp, 2019.

- [6] R. Beekie, T. Buckmaster, and V. Vicol. Weak solutions of ideal MHD which do not conserve magnetic helicity. Ann. PDE, 6(1):Paper No. 1, 40, 2020.
- [7] S. Bianchini and A. Bressan. Vanishing viscosity solutions of nonlinear hyperbolic systems. Ann. of Math. (2), 161(1):223-342, 2005.
- [8] A. Bressan, G. Crasta, and B. Piccoli. Well-posedness of the Cauchy problem for n × n systems of conservation laws. Mem. Amer. Math. Soc., 146(694): viii+134 pp, 2000.
- [9] J. Březina, O. Kreml, and V. Mácha. Non-uniqueness of delta shocks and contact discontinuities in the multi-dimensional model of Chaplygin gas. NoDEA Nonlinear Differential Equations Appl., 28(2): Paper No. 13, 24 pp, 2021.
- [10] T. Buckmaster. Onsager's conjecture almost everywhere in time. Comm. Math. Phys., 333(3):1175– 1198, 2015.
- [11] T. Buckmaster, M. Colombo, and V. Vicol. Wild solutions of the Navier-Stokes equations whose singular sets in time have Hausdorff dimension strictly less than 1. J. Eur. Math. Soc., 24(9):3333-3378, 2021.
- [12] T. Buckmaster, C. De Lellis, P. Isett, and L. Székelyhidi, Jr. Anomalous dissipation for 1/5-Hölder Euler flows. Ann. of Math. (2), 182(1):127–172, 2015.
- [13] T. Buckmaster, C. De Lellis, and L. Székelyhidi, Jr. Dissipative Euler flows with Onsager-critical spatial regularity. Comm. Pure Appl. Math., 69(9):1613–1670, 2016.
- [14] T. Buckmaster, C. De Lellis, L. Székelyhidi, Jr., and V. Vicol. Onsager's conjecture for admissible weak solutions. *Comm. Pure Appl. Math.*, 72(2):229–274, 2019.
- [15] T. Buckmaster, N. Masmoudi, M. Novack, and V. Vicol. Non-conservative  $H^{\frac{1}{2}}$  weak solutions of the incompressible 3D Euler equations. arXiv:2101.09278, 2021.
- [16] T. Buckmaster and V. Vicol. Nonuniqueness of weak solutions to the Navier-Stokes equation. Ann. of Math. (2), 189(1):101–144, 2019.
- [17] T. Buckmaster and V. Vicol. Convex integration and phenomenologies in turbulence. EMS Surv. Math. Sci., 6(1-2):173–263, 2019.
- [18] T. Buckmaster and V. Vicol. Convex integration constructions in hydrodynamics. Bull. Amer. Math. Soc. (N.S.), 58(1):1–44, 2021.
- [19] L. Caffarelli, R. Kohn, and L. Nirenberg. Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Comm. Pure Appl. Math.*, 35(6):771–831, 1982.
- [20] R. E. Caflisch, I. Klapper, and G. Steele. Remarks on singularities, dimension and energy dissipation for ideal hydrodynamics and MHD. Comm. Math. Phys., 184(2): 443–455, 1997.
- [21] M. Cannone. Harmonic analysis tools for solving the incompressible Navier-Stokes equations. Handbook of mathematical fluid dynamics. Vol. III, 161–244, North-Holland, Amsterdam, 2004.
- [22] R. Chen, A. Vasseur, and C. Yu. Global ill-posedness for a dense set of initial data to the isentropic system of gas dynamics. Adv. Math., 393:108057, 2021.
- [23] A. Cheskidov and M. Dai. Norm inflation for generalized Navier-Stokes equations. Indiana Univ. Math. J., 63(3):869–884, 2014.
- [24] A. Cheskidov and M. Dai. Norm inflation for generalized magneto-hydrodynamic system. Nonlinearity, 28(1): 129–142, 2015.

- [25] A. Cheskidov and M. Dai. Discontinuity of weak solutions to the 3D NSE and MHD equations in critical and supercritical spaces. J. Math. Anal. Appl., 481(2): 123493, 16 pp., 2020.
- [26] A. Cheskidov and X. Luo.  $L^2$ -critical nonuniqueness for the 2D Navier-Stokes equations. arXiv:2105.12117, 2021.
- [27] A. Cheskidov and X. Luo. Nonuniqueness of weak solutions for the transport equation at critical space regularity. Ann. PDE, 7(1):Paper No. 2, 45, 2021.
- [28] A. Cheskidov and X. Luo. Sharp nonuniqueness for the Navier-Stokes equations. Invent. Math., 229(3):987–1054, 2022.
- [29] E. Chiodaroli, C. De Lellis, and O. Kreml. Global ill-posedness of the isentropic system of gas dynamics. Comm. Pure Appl. Math., 68(7):1157-1190, 2015.
- [30] E. Chiodaroli and E. Feireisl. On the density of "wild" initial data for the barotropic Euler system. arXiv:2208.04810, 2022.
- [31] E. Chiodaroli, O. Kreml, V. Mácha, and S. Schwarzacher. Non-uniqueness of admissible weak solutions to the compressible Euler equations with smooth initial data. *Trans. Am. Math. Soc.*, 374(4):2269-2295, 2021.
- [32] M. Colombo, C. De Lellis, and L. De Rosa. Ill-posedness of Leray solutions for the hypodissipative Navier-Stokes equations. *Comm. Math. Phys.*, 362(2):659–688, 2018.
- [33] M. Colombo, C. De Lellis, and A. Massaccesi. The generalized Caffarelli-Kohn-Nirenberg theorem for the hyperdissipative Navier-Stokes system. *Comm. Pure Appl. Math.*, 73(3):609–663, 2020.
- [34] M. Dai. Non-uniqueness of Leray-Hopf weak solutions of the 3d Hall-MHD system, SIAM J. Math. Anal., 53(5): 5979–6016, 2021.
- [35] C. De Lellis and L. Székelyhidi, Jr. The Euler equations as a differential inclusion. Ann. of Math. (2), 170(3):1417–1436, 2009.
- [36] C. De Lellis and L. Székelyhidi, Jr. On admissibility criteria for weak solutions of the Euler equations. Arch. Ration. Mech. Anal., 195(1):225–260, 2010.
- [37] C. De Lellis and L. Székelyhidi, Jr. Dissipative continuous Euler flows. Invent. Math., 193(2):377–407, 2013.
- [38] C. De Lellis and L. Székelyhidi, Jr. Dissipative Euler flows and Onsager's conjecture. J. Eur. Math. Soc., 16(7):1467–1505, 2014.
- [39] L. De Rosa. Infinitely many Leray-Hopf solutions for the fractional Navier-Stokes equations. Comm. Partial Differential Equations, 44(4): 335–365, 2019.
- [40] L. Escauriaza, G. A. Seregin, and V. Šverák.  $L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness. Uspekhi Mat. Nauk, 58(2(350)):3–44, 2003.
- [41] E. B. Fabes, B. F. Jones, and N. M. Rivière. The initial value problem for the Navier-Stokes equations with data in L<sup>p</sup>. Arch. Ration. Mech. Anal., 45:222–240, 1972.
- [42] D. Faraco and S. Lindberg. Magnetic helicity and subsolutions in ideal MHD. arXiv:1801.04896, 2018.
- [43] D. Faraco and S. Lindberg. Proof of Taylor's conjecture on magnetic helicity conservation. Comm. Math. Phys., 373(2): 707–738, 2020.

- [44] D. Faraco, S. Lindberg, and L. Székelyhidi, Jr. Bounded solutions of ideal MHD with compact support in space-time. Arch. Ration. Mech. Anal., 239(1):51–93, 2021.
- [45] D. Faraco, S. Lindberg, and L. Székelyhidi, Jr. Magnetic helicity, weak solutions and relaxation of ideal MHD. Comm. Pure Appl. Math., DOI: 10.1002/cpa.22168.
- [46] D. Faraco, S. Lindberg, D. MacTaggart, and A. Valli. On the proof of Taylor's conjecture in multiply connected domains. *Appl. Math. Lett.*, 124 (2022), Paper No. 107654, 7 pp
- [47] Y. Giga. Regularity criteria for weak solutions of the Navier-Stokes system. Nonlinear functional analysis and its applications, Part 1 (Berkeley, Calif., 1983), 449–453, Proc. Sympos. Pure Math., 45, Part 1, Amer. Math. Soc., Providence, RI, 1986.
- [48] V. Giri and H. Kwon. On Non-uniqueness of continuous entropy solutions to the isentropic compressible Euler equations. Arch. Ration. Mech. Anal., 245(2):1213-1283, 2022.
- [49] V. Giri, H. Kwon, and M. Novack. The L<sup>3</sup>-based strong Onsager theorem . arXiv:2305.18509, 2023.
- [50] V. Giri and H. Kwon. A wavelet-inspired L<sup>3</sup>-based convex integration framework for the Euler equations . arXiv:2305.18142, 2023.
- [51] E. Hopf. Uber die Anfangswertaufgabe f
  ür die hydrodynamischen Grundgleichungen. Math. Nachr., 4:213–231, 1951.
- [52] P. Isett. A proof of Onsager's conjecture. Ann. of Math. (2), 188(3):871–963, 2018.
- [53] H. Jia and V. Sverák. Local-in-space estimates near initial time for weak solutions of the Navier-Stokes equations and forward self-similar solutions. *Invent. Math.*, 196(1):233–265, 2014.
- [54] H. Jia and V. Šverák. Are the incompressible 3d Navier-Stokes equations locally ill-posed in the natural energy space? J. Funct. Anal., 268(12):3734–3766, 2015.
- [55] E. Kang and J. Lee. Remarks on the magnetic helicity and energy conservation for ideal magnetohydrodynamics. *Nonlinearity*, 20(11): 2681–2689, 2007.
- [56] T. Kato. Strong L<sup>p</sup>-solutions of the Navier-Stokes equation in ℝ<sup>m</sup>, with applications to weak solutions. Math. Z., 187(4):471–480, 1984.
- [57] N. H. Katz and N. Pavlović. A cheap Caffarelli-Kohn-Nirenberg inequality for the Navier-Stokes equation with hyper-dissipation. *Geom. Funct. Anal.*, 12(2):355–379, 2002.
- [58] S. Klainerman. PDE as a unified subject. *Geom. Funct. Anal.*, Number Special Volume, Part I, pages 279–315, 2000. GAFA 2000 (Tel Aviv, 1999).
- [59] S. Klainerman. On Nash's unique contribution to analysis in just three of his papers. Bull. Amer. Math. Soc. (N.S.), 54(2):283–305, 2017.
- [60] C. Klingenberg, O. Kreml, V. Mácha, and S. Markfelder. Shocks make the Riemann problem for the full Euler system in multiple space dimensions ill-posed. *Nonlinearity*, 33(12):6517-6540, 2020.
- [61] P.G. Lemarié-Rieusset. The Navier-Stokes problem in the 21st century. CRC Press, Boca Raton, FL, xxii+718 pp, 2016.
- [62] J. Leray. Sur le mouvement d'un liquide visqueux emplissant l'espace. Acta Math., 63(1):193–248, 1934.
- [63] Y. Li, P. Qu, Z. Zeng, and D. Zhang. Sharp non-uniqueness for the 3D hyperdissipative Navier-Stokes equations: above the Lions exponent. arXiv:2205.10260, 2022.

- [64] Y. Li, P. Qu, Z. Zeng, and D. Zhang. Non-uniqueness for the hypo-viscous compressible Navier-Stokes equations . arXiv:2212.05844, 2022.
- [65] Y. Li, Z. Zeng, and D. Zhang. Non-uniqueness of weak solutions to 3D magnetohydrodynamic equations. J. Math. Pures Appl., 165(9):232-285, 2022.
- [66] Y. Li, Z. Zeng, and D. Zhang. Sharp non-uniqueness of weak solutions to 3D magnetohydrodynamic equations. arXiv: 2208.00624, 2022.
- [67] F. Lin. A new proof of the Caffarelli-Kohn-Nirenberg theorem. Comm. Pure Appl. Math., 51(3):241– 257, 1998.
- [68] J.-L. Lions. Quelques méthodes de résolution des problèmes aux limites non linéaires. Dunod; Gauthier-Villars, Paris, 1969.
- [69] T.-P. Liu, and T. Yang. Well-posedness theory for hyperbolic conservation laws. Comm. Pure Appl. Math., 52(12):1553-1586, 1999.
- [70] T. Luo and P. Qu. Non-uniqueness of weak solutions to 2D hypoviscous Navier-Stokes equations. J. Differential Equations, 269(4):2896–2919, 2020.
- [71] T. Luo and E.S. Titi. Non-uniqueness of weak solutions to hyperviscous Navier-Stokes equations: on sharpness of J.-L. Lions exponent. *Calc. Var. Partial Differential Equations*, 59(3):Paper No. 92, 15, 2020.
- [72] T. Luo, C. Xie, and Z. Xin. Non-uniqueness of admissible weak solutions to compressible Euler systems with source terms. Adv. Math., 291:542-583, 2016.
- [73] X. Luo. Stationary solutions and nonuniqueness of weak solutions for the Navier-Stokes equations in high dimensions. Arch. Ration. Mech. Anal., 233(2):701–747, 2019.
- [74] S. Markfelder and C. Klingenberg. The Riemann problem for the multidimensional isentropic system of gas dynamics is ill-posed if it contains a shock. Arch. Ration. Mech. Anal., 227(3):967-994, 2018.
- [75] Y. Meyer. Wavelets, paraproducts, and Navier-Stokes equations. Current developments in mathematics, 1996 (Cambridge, MA), 105-212, Int. Press, Boston, MA, 1997.
- [76] C. Miao and W. Ye. On the weak solutions for the MHD systems with controllable total energy and cross helicity. arXiv:2208.08311, 2022.
- [77] M. Novack and V. Vicol. An intermittent Onsager theorem. Invent. Math. 233(1):223–323, 2023.
- [78] H. Schmeisser and H. Triebel. Topics in Fourier analysis and function spaces. A Wiley-Interscience Publication. John Wiley & Sons, Ltd., Chichester, 300 pp, 1987.
- [79] V. Scheffer. Partial regularity of solutions to the Navier-Stokes equations. Pacific J. Math., 66(2):535-552, 1976.
- [80] V. Scheffer. Hausdorff measure and the Navier-Stokes equations. Comm. Math. Phys., 55 (1977), no. 2, 97-112.
- [81] J. B. Taylor. Relaxation of toroidal plasma and generation of reverse magnetic fields. *Physical Review Letters*, 33(19): 1139, 1974.
- [82] J. B. Taylor. Relaxation and magnetic reconnection in plasmas. *Reviews of Modern Physics*, 58(3): 741, 1986.
- [83] A. Vasseur. A new proof of partial regularity of solutions to Navier-Stokes equations. NoDEA Nonlinear Differential Equations Appl., 14(5-6):753-785, 2007.

[84] J. Wu. Generalized MHD equations. J. Differential Equations, 195(2): 284–312, 2003.

[85] J. Wu. Lower bounds for an integral involving fractional Laplacians and the generalized Navier-Stokes equations in Besov spaces. Comm. Math. Phys., 263(3):803-831, 2006.

Yachun Li School of Mathematical Sciences, CMA-Shanghai, MOE-LSC, and SHL-MAC Shanghai Jiao Tong University Shanghai 200240 PR China E-mail address: ycli@sjtu.edu.cn

Peng Qu School of Mathematical Sciences & Shanghai Key Laboratory for Contemporary Applied Mathematics Fudan University Shanghai 200433 PR China E-mail address: pqu@fudan.edu.cn

Zirong Zeng School of Mathematical Sciences, Shanghai Jiao Tong University Shanghai 200240 PR China E-mail address: beckzzr@sjtu.edu.cn

Deng Zhang School of Mathematical Sciences, CMA-Shanghai Shanghai Jiao Tong University Shanghai 200240 PR China E-mail address: dzhang@sjtu.edu.cn