

## Weierstrass semigroups of a pair of points whose first non-gaps are three

Jiryō Komeda (米田二良)  
(Collaboration with Seon Jeong Kim)  
Kanagawa Institute of Technology (神奈川工科大学)

### 1 Introduction

Let  $\mathbb{N}$  be the additive semigroup of non-negative integers. Let  $C$  be a complete nonsingular curve of genus  $g \geq 2$  over an algebraically closed field  $k$  of characteristic 0, which is called a *curve* in this talk, and  $K(C)$  the field of rational functions on  $C$ .

**Definition 1.1.** For a point  $P$  of  $C$ , we set

$$H(P) := \{\alpha \in \mathbb{N} \mid \text{there exists } f \in K(C) \text{ with } (f)_\infty = \alpha P\},$$

which is called the *Weierstrass semigroup of the point  $P$* . An integer  $n$  is called the *first non-gap* of  $P$  if it is the minimum positive integer in  $H(P)$ .

**Definition 1.2.** For distinct points  $P$  and  $Q$  of  $C$ , we set

$$H(P, Q) := \{(\alpha, \beta) \in \mathbb{N} \times \mathbb{N} \mid \text{there exists } f \in K(C) \text{ with } (f)_\infty = \alpha P + \beta Q\},$$

which is called the *Weierstrass semigroup of the pair  $(P, Q)$  of points*.

**Fact 1.3.** (Kim [2]) If  $C$  is a hyperelliptic curve, i.e., a double covering of the projective line, then the semigroup  $H(P, Q)$  is determined explicitly.

**Fact 1.4.** (Kim-Komeda [4]) If  $C$  is of genus 3, then the semigroup  $H(P, Q)$  is determined explicitly.

**Aim 1.5.** Let  $P$  and  $Q$  be distinct points of  $C$  of genus  $g \geq 4$ . If the first non-gaps of  $P$  and  $Q$  are three, we determine the semigroup  $H(P, Q)$  explicitly. Moreover, we give examples of two pointed curves  $(C, P, Q)$  whose semigroups are the given

## 2 Possible Weierstrass semigroups of genus $\geq 5$

First, let us review some Kim's results. Let  $C$  be a curve of genus  $g$  and  $P$  its point.

**Definition 2.1.** We set

$$G(P) := \mathbb{N} \setminus H(P) = \{l_1 < l_2 < \dots < l_g\}.$$

The integer  $l_g$  is called the *last gap* at  $P$ .

Let  $Q$  be another point of  $C$  which is distinct from  $P$ . For each  $l \in G(P)$ , the integer  $\min\{\beta \mid (l, \beta) \in H(P, Q)\}$  must be equal to some element in  $G(Q)$ , say  $\sigma(l)$ , and this correspondence  $\sigma$  gives a bijective map between the sets  $G(P)$  and  $G(Q)$ .

**Definition 2.2.** We set

$$\Gamma(P, Q) := \{(l, \sigma(l)) \mid l \in G(P)\}.$$

**Fact 2.3.** (Kim [2]) The semigroup  $H(P, Q)$  is completely determined by the bijective correspondence  $\sigma$ , i.e.,

$$G(P, Q) = \bigcup_{l \in G(P)} (\{(l, \beta) \mid \beta = 0, 1, \dots, \sigma(l) - 1\} \\ \cup \{(\alpha, \sigma(l)) \mid \alpha = 0, 1, \dots, l - 1\}),$$

where we set

$$G(P, Q) = \mathbb{N} \times \mathbb{N} \setminus H(P, Q).$$

Thus, it suffices to determine the graph  $\Gamma(P, Q)$  of  $\sigma$  for describing the semigroup  $H(P, Q)$ .

We consider the case where the first non-gaps of distinct points  $P$  and  $Q$  are three.

**Theorem 2.4.**  $\sigma(l_g) = 1$  or  $\sigma(l_g) = 2$ , i.e.,  $(l_g, 1) \in \Gamma(P, Q)$  or  $(l_g, 2) \in \Gamma(P, Q)$ .

*Proof.* Assume that  $(l_g, \beta) \in \Gamma(P, Q)$  with  $\beta \geq 4$ . Then

$$(\alpha_1, 1) \in \Gamma(P, Q), 1 \leq \alpha_1 < l_g \text{ and } (\alpha_2, 2) \in \Gamma(P, Q), 1 \leq \alpha_2 < l_g.$$

Now

$$\beta \equiv i \pmod{3} \text{ for some } i = 1, 2 \text{ and } (\alpha_i, \beta) = (\alpha_i, i) + (0, 3k) \text{ for some } k.$$

Since  $(\alpha_i, i) \in \Gamma(P, Q)$  and  $(0, 3k) \in H(P, Q)$ , we have  $(\alpha_i, \beta) \in H(P, Q)$ . But  $(l_g, \beta) \in \Gamma(P, Q)$  and  $\alpha_i < l_g$ . This contradicts Fact 2.3. Q.E.D.

From now on we assume that  $g \geq 5$ . By the theory of curves we can find some integer  $n$  with

$$\frac{g-1}{3} \leq n \leq \frac{g}{2}$$

such that

$$S(H(P)) = \{3, 3n+2, 3(g-n)+1\}$$

or

$$S(H(P)) = \{3, 3n+1, 3(g-n)+2\},$$

where  $S(H(P)) = \{3, s_1, s_2\}$ , which is called the *standard basis* for  $H(P)$ , with

$$s_i = \text{Min}\{s \in H(P) | s \equiv i \pmod{3}\} \text{ for } i = 1, 2.$$

Moreover, since there exists  $f \in K(C)$  such that  $(f) = 3P - 3Q$ , the standard basis  $S(H(Q))$  must be one of the above two.

**Definition 2.5.** The point  $P$  is said to be *of the  $n$ -th kind* if

$$S(H(P)) = \{3, 3n+2, 3(g-n)+1\}$$

or

$$S(H(P)) = \{3, 3n+1, 3(g-n)+2\},$$

**Definition 2.6.** The point  $P$  of the  $n$ -th kind is said to be *of type I* (resp. *II*) if

$$n \neq \frac{g}{2} \text{ and } 3n+2 \in S(H(P)) \text{ (resp. } 3n+1 \in S(H(P))).$$

We note that if  $n = \frac{g}{2}$ , then  $S(H(P)) = \{3, 3n+1, 3n+2\}$ .

Using the types of the points  $P$  and  $Q$  we can determine whether

$$(l_g, 1) \in \Gamma(P, Q) \text{ or } (l_g, 2) \in \Gamma(P, Q).$$

**Theorem 2.7.** Let  $P$  and  $Q$  be two distinct points of the  $n$ -th kind and  $n \neq \frac{g}{2}$ .

i) If  $P$  and  $Q$  are of type II, then

$$(l_g, 2) \in \Gamma(P, Q) \text{ and } (3n-2, 1) \in \Gamma(P, Q).$$

ii) If  $P$  (resp.  $Q$ ) is of type I (resp. II), then

$$(l_g, 1) \in \Gamma(P, Q) \text{ and } (3n-2, 2) \in \Gamma(P, Q).$$

iii) If  $P$  and  $Q$  are of type I, then

$$(l_g, 1) \in \Gamma(P, Q) \text{ and } (3n-1, 2) \in \Gamma(P, Q).$$

Moreover, if  $(\alpha_1, 1)$  and  $(\alpha_2, 2)$  belong to  $\Gamma(P, Q)$ , then

$$\Gamma(P, Q) = \left\{ (\alpha_1 - 3k, 1 + 3k) \mid 0 \leq k < \frac{\alpha_1}{3} \right\} \cup \left\{ (\alpha_2 - 3k, 2 + 3k) \mid 0 \leq k < \frac{\alpha_2}{3} \right\}.$$

In some case with  $n = \frac{g}{3}$  we have no candidate of the semigroup  $H(P, Q)$ .

**Proposition 2.8.** *Let  $P$  and  $Q$  be points of the  $\frac{g}{3}$ -th kind. If they are of type II, then  $P = Q$ . In this case,  $H(P)$  is generated by 3 and  $g + 1$  with  $g \equiv 0 \pmod{3}$ .*

There are two possibilities in the case  $n = \frac{g}{2}$ .

**Proposition 2.9.** *Let  $P$  and  $Q$  be distinct points of the  $\frac{g}{2}$ -th kind, i.e.,*

$$S(H(P)) = S(H(Q)) = \{3, 3n + 1, 3n + 2\}.$$

Then

$$(3n - 1, 1) \in \Gamma(P, Q) \text{ or } (3n - 1, 2) \in \Gamma(P, Q).$$

If  $(3n - 1, 1) \in \Gamma(P, Q)$  (resp.  $(3n - 1, 2) \in \Gamma(P, Q)$ ), then

$$\begin{aligned} \Gamma(P, Q) &= \{(\alpha, 3n - \alpha) \mid \alpha \in G(P)\} \\ &\text{(resp. } \{(3k - 2, (3n - 1) - (3k - 2)) \mid k = 1, \dots, n\} \\ &\quad \cup \{(3k - 1, (3n + 1) - (3k - 1)) \mid k = 1, \dots, n\}). \end{aligned}$$

### 3 The existence of two pointed curves

In the previous section we determined the possible Weierstrass semigroups  $H$  of a pair of points on a curve of genus  $\geq 5$  whose first non-gaps are three. In this section for each such a semigroup  $H$  we give two pointed curves  $(C, P, Q)$  such that  $H(P, Q) = H$ .

Let  $C$  be the curve whose function field  $K(C) = k(x, y)$  is defined by the equation

$$y^3 = (x - c_1) \cdots (x - c_{i_1})(x - c_{i_1+1})^2 \cdots (x - c_{i_1+i_2})^2,$$

where  $c_1, \dots, c_{i_1+i_2}$  are distinct elements of  $k$  and  $i_1 + 2i_2$  is not divisible by 3. We note that the genus  $g$  of the curve  $C$  is  $i_1 + i_2 - 1$  by Riemann-Hurwitz formula.

Let  $\pi : C \rightarrow \mathbf{P}^1$  be the morphism corresponding to the inclusion  $k(x) \subset K(C)$ , i.e.,  $\pi(P) = (1 : x(P))$ , where  $\mathbf{P}^1$  denotes the projective line. We set

$$\{P_\infty\} = \pi^{-1}(0 : 1) \text{ and } \{P_s\} = \pi^{-1}(1 : c_s) \text{ for } s = 1, \dots, i_1 + i_2.$$

Then we have  $S(H(P_\infty)) = \{3, i_1 + 2i_2, 2i_1 + i_2\}$  (For example, see Kim-Komeda [3]). If  $i_1 + 2i_2 \equiv 1 \pmod{3}$ ,

$$S(H(P_s)) = \begin{cases} \{3, i_1 + 2i_2 + 1, 2i_1 + i_2 - 1\} & \text{if } 1 \leq s \leq i_1 \\ S(H(P_\infty)) = \{3, i_1 + 2i_2, 2i_1 + i_2\} & \text{if } i_1 + 1 \leq s \leq i_1 + i_2 \end{cases}$$

If  $i_1 + 2i_2 \equiv 2 \pmod{3}$ ,

$$S(H(P_s)) = \begin{cases} S(H(P_\infty)) = \{3, i_1 + 2i_2, 2i_1 + i_2\} & \text{if } 1 \leq s \leq i_1 \\ \{3, i_1 + 2i_2 - 1, 2i_1 + i_2 + 1\} & \text{if } i_1 + 1 \leq s \leq i_1 + i_2 \end{cases}$$

From now on we assume that  $g \geq 5$ . By the above formula we get the following examples :

- Example 3.1.** i) Let  $\frac{g}{3} < n < \frac{g}{2}$ . If  $i_1 = 2g - 3n + 1 > n + 1$  and  $i_2 = 3n - g > 0$ , then the points  $P_\infty$  and  $P_s$  ( $i_1 + 1 \leq s \leq i_1 + i_2$ ) are of the  $n$ -th kind of type II.  
 ii) Let  $\frac{g}{3} \leq n \leq \frac{g-1}{2}$ . If  $i_1 = 2g - 3n + 1 > n + 3$  and  $i_2 = 3n - g \geq 0$ , then the points  $P_\infty$  is of the  $n$ -th kind of type II and the points  $P_s$  ( $1 \leq s \leq i_1$ ) are of the  $n$ -th kind of type I.  
 iii) Let  $\frac{g-1}{3} \leq n \leq \frac{g-1}{2}$ . If  $i_1 = 2g - 3n \geq n + 2$  and  $i_2 = 3n - g + 1 \geq 0$ , then the points  $P_\infty$  and  $P_s$  ( $1 \leq s \leq i_1$ ) are of the  $n$ -th kind of type I.

In the case of Proposition 2.9 we get the following examples:

**Example 3.2.** Let  $n \geq 3$ . If  $i_1 = n + 1$  and  $i_2 = n$ , then  $g = 2n$  and the points  $P_\infty$  and  $P_s$  are of the  $\frac{g}{2}$ -th kind. Then  $S(H(P_\infty)) = S(H(P_s)) = \{3, 3n + 1, 3n + 2\}$ . Moreover,

$$\Gamma(P_\infty, P_s) \ni (3n - 1, 1) \text{ for } 1 \leq s \leq i_1$$

and

$$\Gamma(P_\infty, P_s) \ni (3n - 1, 2) \text{ for } i_1 + 1 \leq s \leq i_1 + i_2.$$

## 4 Weierstrass semigroups of genus 4

In this section we treat the curves  $C$  of genus 4 with point  $P$  whose first non-gap is 3. Then we have  $S(H(P)) = \{3, 5, 10\}$  or  $\{3, 7, 8\}$ .

**Remark 4.1.** Let  $S(H(P)) = \{3, 5, 10\}$ . If  $Q$  is another point of  $C$  whose first non-gap is three, then  $S(H(Q)) = \{3, 5, 10\}$  and there exists  $f \in \mathbf{K}(C)$  such that  $(f) = 3P - 3Q$ .

**Proposition 4.2.** *Let  $P$  and  $Q$  be two distinct points such that  $S(H(P)) = S(H(Q)) = \{3, 5, 10\}$ . Then  $\Gamma(P, Q) = \{(1, 7), (2, 2), (4, 4), (7, 1)\}$ . For example, such pointed curves are given by*

$$y^3 = (x - c_1) \cdots (x - c_5), P = P_\infty \text{ and } Q = P_s \text{ for } s = 1, \dots, 5$$

where we use the notations in Section 3.

**Proposition 4.3.** *Let  $S(H(P)) = \{3, 7, 8\}$  and  $Q$  another point of  $C$  whose first non-gap is three. Suppose that there exists  $f \in \mathbf{K}(C)$  such that  $(f) = 3P - 3Q$ .*

i)  $(5, 1) \in \Gamma(P, Q)$  or  $(5, 2) \in \Gamma(P, Q)$ .

ii) If  $(5, 1) \in \Gamma(P, Q)$ , then  $\Gamma(P, Q) = \{(5, 1), (4, 2), (2, 4), (1, 5)\}$ . For example, such pointed curves are given by

$$y^3 = (x - c_1)(x - c_2)(x - c_3)(x - c_4)^2(x - c_5)^2, P = P_\infty \text{ and } Q = P_s \text{ for } s = 1, 2, 3.$$

iii) If  $(5, 2) \in \Gamma(P, Q)$ , then  $\Gamma(P, Q) = \{(5, 2), (4, 1), (1, 4), (2, 5)\}$ . Such pointed curves are given by the same equations as above,  $P = P_\infty$  and  $Q = P_s$  for  $s = 4, 5$ .

**Proposition 4.4.** *Let  $S(H(P)) = \{3, 7, 8\}$  and  $Q$  another point of  $C$  whose first non-gap is three. Suppose that there is no  $f \in \mathbf{K}(C)$  such that  $(f) = 3P - 3Q$ . Then  $(5, 1) \in \Gamma(P, Q)$  and  $\Gamma(P, Q) = \{(5, 1), (4, 4), (2, 2), (1, 5)\}$ . Such curves  $C$  are also given by the equations*

$$y^3 = (x - c_1)(x - c_2)(x - c_3)(x - c_4)^2(x - c_5)^2.$$

Using the result of Kato [1] we get our desired points  $P$  and  $Q$ .

## References

- [1] T. Kato, *On Weierstrass points whose first non-gaps are three*. J. Reine Angew. Math. **316** (1980), 99-109.
- [2] S.J. Kim, *On the index of the Weierstrass semigroup of a pair of points on a curve*. Arch. Math. **62** (1994), 73-82.
- [3] S.J. Kim and J. Komeda, *Numerical semigroups which cannot be realized as semigroups of Galois Weierstrass points*. To appear in Arch. Math.
- [4] S.J. Kim and J. Komeda, *The Weierstrass semigroup of a pair and moduli in  $\mathcal{M}_3$* . To appear in Bol. Soc. Bras. Mat.